


# Computability and Complexity

## COSC 4200

### DFA Minimization


When can we simplify a DFA?



$q_1$  and  $q_2$  are equivalent.


The two states are equivalent because starting from  $q_1$  or  $q_2$ , exactly the same strings are accepted.

When can we simplify a DFA?




$q_1$  and  $q_2$  are equivalent.

The two states are equivalent because starting from  $q_1$  or  $q_2$ , exactly the same strings are accepted.





$q_1$  and  $q_2$  are equivalent. We may collapse these states.




We will develop an algorithm for:


- 1 Determining which states are equivalent.
- 2 Building the collapsed automaton.










$q_1$  is equivalent to  $q_2$  and  $q_3$  is equivalent to  $q_4$ . These two pairs may be collapsed:



















Each state is equivalent with the opposite state in the diagram:  
 $q_0$  with  $q_3$ ,  $q_1$  with  $q_4$ ,  $q_2$  with  $q_5$ . Collapsing yields:



Each state is equivalent with the opposite state in the diagram:  
 $q_0$  with  $q_3$ ,  $q_1$  with  $q_4$ ,  $q_2$  with  $q_5$ . Collapsing yields:



# Equivalent States

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA.

Intuitively, two states are equivalent if they “remember” the same information. We can say that two states  $p$  and  $q$  are equivalent if for every  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \Leftrightarrow \delta^*(q, x) \in F.$$

In other words,  $p$  and  $q$  are equivalent if exactly the same strings are accepted when we start in  $p$  or  $q$ . We write  $p \approx q$  to denote that  $p$  is equivalent to  $q$ .

Observe that  $\approx$  is an equivalence relation:

- *reflexive*:  $p \approx p$  for all  $p$
- *symmetric*:  $p \approx q$  implies  $q \approx p$
- *transitive*: if  $p \approx q$  and  $q \approx r$ , then  $p \approx r$



Observe that  $\approx$  is an equivalence relation:

- *reflexive*:  $p \approx p$  for all  $p$
- *symmetric*:  $p \approx q$  implies  $q \approx p$
- *transitive*: if  $p \approx q$  and  $q \approx r$ , then  $p \approx r$

For each  $p \in Q$ , let

$$[p] = \{q \in Q \mid q \approx p\}$$

be the equivalence class of  $p$  under  $\approx$ .

Observe that  $\approx$  is an equivalence relation:

- *reflexive*:  $p \approx p$  for all  $p$
- *symmetric*:  $p \approx q$  implies  $q \approx p$
- *transitive*: if  $p \approx q$  and  $q \approx r$ , then  $p \approx r$

For each  $p \in Q$ , let

$$[p] = \{q \in Q \mid q \approx p\}$$

be the equivalence class of  $p$  under  $\approx$ .

These equivalence classes partition  $Q$  (so each state belongs to exactly one class), and we have

$$p \approx q \Leftrightarrow [p] = [q].$$

# Quotient Automaton

We use  $\approx$  to define the *quotient automaton*

$$M/\approx = (Q', \Sigma, \delta', q'_0, F')$$

where

- $Q' = \{[p] \mid p \in Q\}$

# Quotient Automaton

We use  $\approx$  to define the *quotient automaton*

$$M/\approx = (Q', \Sigma, \delta', q'_0, F')$$

where

- $Q' = \{[p] \mid p \in Q\}$
- $\delta'([p], a) = [\delta(p, a)]$  for all  $[p] \in Q'$ ,  $a \in \Sigma$

# Quotient Automaton

We use  $\approx$  to define the *quotient automaton*

$$M/\approx = (Q', \Sigma, \delta', q'_0, F')$$

where

- $Q' = \{[p] \mid p \in Q\}$
- $\delta'([p], a) = [\delta(p, a)]$  for all  $[p] \in Q', a \in \Sigma$
- $q'_0 = [q_0]$

# Quotient Automaton

We use  $\approx$  to define the *quotient automaton*

$$M/\approx = (Q', \Sigma, \delta', q'_0, F')$$

where

- $Q' = \{[p] \mid p \in Q\}$
- $\delta'([p], a) = [\delta(p, a)]$  for all  $[p] \in Q', a \in \Sigma$
- $q'_0 = [q_0]$
- $F' = \{[p] \mid p \in F\}$

# Quotient Automaton

We use  $\approx$  to define the *quotient automaton*

$$M/\approx = (Q', \Sigma, \delta', q'_0, F')$$

where

- $Q' = \{[p] \mid p \in Q\}$
- $\delta'([p], a) = [\delta(p, a)]$  for all  $[p] \in Q', a \in \Sigma$
- $q'_0 = [q_0]$
- $F' = \{[p] \mid p \in F\}$

We claim that  $M/\approx$  is equivalent to  $M$ , that is  $L(M/\approx) = L(M)$ .  
First, we need to check that the transition function  $\delta'$  is well-defined.

The transition function  $\delta'$  is defined by

$$\delta'([p], a) = [\delta(p, a)]$$

for all  $[p] \in Q'$ ,  $a \in \Sigma$ .

For  $\delta'$  to be well-defined, it shouldn't matter which equivalence class representative of  $[p]$  we use to apply  $\delta$ . The following lemma says this is the case.

### Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$ .*



## Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$  for all  $a \in \Sigma$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$  for all  $a \in \Sigma$ .*

**Proof.** Suppose  $p \approx q$ . Then for all  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \iff \delta^*(q, x) \in F.$$

## Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$  for all  $a \in \Sigma$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$  for all  $a \in \Sigma$ .*

**Proof.** Suppose  $p \approx q$ . Then for all  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \iff \delta^*(q, x) \in F.$$

Let  $a \in \Sigma$ . For all  $y \in \Sigma^*$ ,

$$\delta^*(\delta(p, a), y) \in F \iff \delta^*(p, ay) \in F \quad \text{property of } \delta^*$$

## Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$  for all  $a \in \Sigma$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$  for all  $a \in \Sigma$ .*

**Proof.** Suppose  $p \approx q$ . Then for all  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \iff \delta^*(q, x) \in F.$$

Let  $a \in \Sigma$ . For all  $y \in \Sigma^*$ ,

$$\begin{aligned} \delta^*(\delta(p, a), y) \in F &\iff \delta^*(p, ay) \in F && \text{property of } \delta^* \\ &\iff \delta^*(q, ay) \in F && \text{since } p \approx q \end{aligned}$$

## Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$  for all  $a \in \Sigma$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$  for all  $a \in \Sigma$ .*

**Proof.** Suppose  $p \approx q$ . Then for all  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \iff \delta^*(q, x) \in F.$$

Let  $a \in \Sigma$ . For all  $y \in \Sigma^*$ ,

$$\begin{aligned} \delta^*(\delta(p, a), y) \in F &\iff \delta^*(p, ay) \in F && \text{property of } \delta^* \\ &\iff \delta^*(q, ay) \in F && \text{since } p \approx q \\ &\iff \delta^*(\delta(q, a), y) \in F && \text{property of } \delta^*. \end{aligned}$$

## Lemma

*If  $p \approx q$ , then  $\delta(p, a) \approx \delta(q, a)$  for all  $a \in \Sigma$ .*

*Equivalently, if  $[p] = [q]$ , then  $[\delta(p, a)] = [\delta(q, a)]$  for all  $a \in \Sigma$ .*

**Proof.** Suppose  $p \approx q$ . Then for all  $x \in \Sigma^*$ ,

$$\delta^*(p, x) \in F \iff \delta^*(q, x) \in F.$$

Let  $a \in \Sigma$ . For all  $y \in \Sigma^*$ ,

$$\begin{aligned} \delta^*(\delta(p, a), y) \in F &\iff \delta^*(p, ay) \in F && \text{property of } \delta^* \\ &\iff \delta^*(q, ay) \in F && \text{since } p \approx q \\ &\iff \delta^*(\delta(q, a), y) \in F && \text{property of } \delta^*. \end{aligned}$$

Therefore  $\delta(p, a) \approx \delta(q, a)$ . □

The quotient automaton  $M/\approx$  has the same language as the original automaton  $M$ .

Theorem

$$L(M/\approx) = L(M).$$

The quotient automaton  $M/\approx$  has the same language as the original automaton  $M$ .

### Theorem

$$L(M/\approx) = L(M).$$

The proof follows from these two lemmas:

### Lemma A

$$p \in F \Leftrightarrow [p] \in F'.$$

### Lemma B

$$\text{For all } x \in \Sigma^*, \delta'^*([p], x) = [\delta^*(p, x)].$$

### Lemma A

$$p \in F \iff [p] \in F'.$$

**Proof.** The  $\Rightarrow$  direction is immediate from the definition of  $F'$ .



## Lemma A

$$p \in F \iff [p] \in F'.$$

**Proof.** The  $\Rightarrow$  direction is immediate from the definition of  $F'$ .

For the  $\Leftarrow$  direction, we need to show that if  $p \approx q$  and  $p \in F$ , then  $q \in F$ .

## Lemma A

$$p \in F \iff [p] \in F'.$$

**Proof.** The  $\Rightarrow$  direction is immediate from the definition of  $F'$ .

For the  $\Leftarrow$  direction, we need to show that if  $p \approx q$  and  $p \in F$ , then  $q \in F$ . Using  $x = \epsilon$  in the definition of  $p \approx q$ , we have

$$p \in F \iff \delta^*(p, \epsilon) \in F \quad \text{definition of } \delta^*$$

## Lemma A

$$p \in F \iff [p] \in F'.$$

**Proof.** The  $\Rightarrow$  direction is immediate from the definition of  $F'$ .

For the  $\Leftarrow$  direction, we need to show that if  $p \approx q$  and  $p \in F$ , then  $q \in F$ . Using  $x = \epsilon$  in the definition of  $p \approx q$ , we have

$$\begin{aligned} p \in F &\iff \delta^*(p, \epsilon) \in F && \text{definition of } \delta^* \\ &\iff \delta^*(q, \epsilon) \in F && \text{since } p \approx q \end{aligned}$$

## Lemma A

$$p \in F \iff [p] \in F'.$$

**Proof.** The  $\Rightarrow$  direction is immediate from the definition of  $F'$ .

For the  $\Leftarrow$  direction, we need to show that if  $p \approx q$  and  $p \in F$ , then  $q \in F$ . Using  $x = \epsilon$  in the definition of  $p \approx q$ , we have

$$\begin{aligned} p \in F &\iff \delta^*(p, \epsilon) \in F && \text{definition of } \delta^* \\ &\iff \delta^*(q, \epsilon) \in F && \text{since } p \approx q \\ &\iff q \in F && \text{definition of } \delta^*. \quad \square \end{aligned}$$

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings.

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

Assume as our inductive hypothesis the claim

$$\delta'^*([p], x) = [\delta^*(p, x)]$$

holds for some  $x \in \Sigma^*$ . Let  $a \in \Sigma$ . For the inductive step, we must show that

$$\delta'^*([p], xa) = [\delta^*(p, xa)].$$

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

Assume as our inductive hypothesis the claim

$$\delta'^*([p], x) = [\delta^*(p, x)]$$

holds for some  $x \in \Sigma^*$ . Let  $a \in \Sigma$ . For the inductive step, we must show that

$$\delta'^*([p], xa) = [\delta^*(p, xa)].$$

We have

$$\delta'^*([p], xa) = \delta'(\delta'^*([p], x), a) \quad \text{definition of } \delta'^*$$



## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

Assume as our inductive hypothesis the claim

$$\delta'^*([p], x) = [\delta^*(p, x)]$$

holds for some  $x \in \Sigma^*$ . Let  $a \in \Sigma$ . For the inductive step, we must show that

$$\delta'^*([p], xa) = [\delta^*(p, xa)].$$

We have

$$\begin{aligned}\delta'^*([p], xa) &= \delta'(\delta'^*([p], x), a) && \text{definition of } \delta'^* \\ &= \delta'([\delta^*(p, x)], a) && \text{inductive hypothesis}\end{aligned}$$

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

Assume as our inductive hypothesis the claim

$$\delta'^*([p], x) = [\delta^*(p, x)]$$

holds for some  $x \in \Sigma^*$ . Let  $a \in \Sigma$ . For the inductive step, we must show that

$$\delta'^*([p], xa) = [\delta^*(p, xa)].$$

We have

$$\begin{aligned}\delta'^*([p], xa) &= \delta'(\delta'^*([p], x), a) && \text{definition of } \delta'^* \\ &= \delta'([\delta^*(p, x)], a) && \text{inductive hypothesis} \\ &= [\delta(\delta^*(p, x), a)] && \text{definition of } \delta'\end{aligned}$$

## Lemma B

For all  $x \in \Sigma^*$ ,  $\delta'^*([p], x) = [\delta^*(p, x)]$ .

**Proof.** The proof is by induction on strings. In the base case, we have  $x = \epsilon$ :

$$\begin{aligned}\delta'^*([p], \epsilon) &= [p] && \text{definition of } \delta'^* \\ &= [\delta^*(p, \epsilon)] && \text{definition of } \delta^*.\end{aligned}$$

Assume as our inductive hypothesis the claim

$$\delta'^*([p], x) = [\delta^*(p, x)]$$

holds for some  $x \in \Sigma^*$ . Let  $a \in \Sigma$ . For the inductive step, we must show that

$$\delta'^*([p], xa) = [\delta^*(p, xa)].$$

We have

$$\begin{aligned}\delta'^*([p], xa) &= \delta'(\delta'^*([p], x), a) && \text{definition of } \delta'^* \\ &= \delta'([\delta^*(p, x)], a) && \text{inductive hypothesis} \\ &= [\delta(\delta^*(p, x), a)] && \text{definition of } \delta' \\ &= [\delta^*(p, xa)] && \text{definition of } \delta^*. \quad \square\end{aligned}$$

We now use Lemmas A and B to prove the Theorem.

### Theorem

$$L(M/\approx) = L(M).$$

**Proof.** For any  $x \in \Sigma^*$ ,

$$x \in L(M/\approx) \iff \delta'^*(q'_0, x) \in F' \quad \text{definition of acceptance}$$

We now use Lemmas A and B to prove the Theorem.

### Theorem

$$L(M/\approx) = L(M).$$

**Proof.** For any  $x \in \Sigma^*$ ,

$$\begin{aligned} x \in L(M/\approx) &\iff \delta'^*(q'_0, x) \in F' && \text{definition of acceptance} \\ &\iff \delta'^*([q_0], x) \in F' && \text{definition of } q'_0 \end{aligned}$$

We now use Lemmas A and B to prove the Theorem.

### Theorem

$$L(M/\approx) = L(M).$$

**Proof.** For any  $x \in \Sigma^*$ ,

$$\begin{aligned} x \in L(M/\approx) &\iff \delta'^*(q'_0, x) \in F' && \text{definition of acceptance} \\ &\iff \delta'^*([q_0], x) \in F' && \text{definition of } q'_0 \\ &\iff [\delta^*(q_0, x)] \in F' && \text{Lemma B} \end{aligned}$$

We now use Lemmas A and B to prove the Theorem.

### Theorem

$$L(M/\approx) = L(M).$$

**Proof.** For any  $x \in \Sigma^*$ ,

$x \in L(M/\approx)$	$\iff$	$\delta'^*(q'_0, x) \in F'$	definition of acceptance
	$\iff$	$\delta'^*([q_0], x) \in F'$	definition of $q'_0$
	$\iff$	$[\delta^*(q_0, x)] \in F'$	Lemma B
	$\iff$	$\delta^*(q_0, x) \in F$	Lemma A

We now use Lemmas A and B to prove the Theorem.

### Theorem

$$L(M/\approx) = L(M).$$

**Proof.** For any  $x \in \Sigma^*$ ,

$x \in L(M/\approx)$	$\iff$	$\delta'^*(q'_0, x) \in F'$	definition of acceptance
	$\iff$	$\delta'^*([q_0], x) \in F'$	definition of $q'_0$
	$\iff$	$[\delta^*(q_0, x)] \in F'$	Lemma B
	$\iff$	$\delta^*(q_0, x) \in F$	Lemma A
	$\iff$	$x \in L(M)$	definition of acceptance. $\square$



## Theorem

$$L(M/\approx) = L(M).$$

It turns out that  $M/\approx$  cannot be collapsed any further. If we apply the quotient construction a second time, the DFA will not change.

How can we determine whether  $p \approx q$ ?

## Minimization Algorithm

- 1 Make a table of all pairs  $\{p, q\}$ , initially unmarked.

How can we determine whether  $p \approx q$ ?

## Minimization Algorithm

- 1 Make a table of all pairs  $\{p, q\}$ , initially unmarked.
- 2 Mark  $\{p, q\}$  if  $(p \in F \text{ and } q \notin F)$  or  $(p \notin F \text{ and } q \in F)$ .

How can we determine whether  $p \approx q$ ?

## Minimization Algorithm

- 1 Make a table of all pairs  $\{p, q\}$ , initially unmarked.
- 2 Mark  $\{p, q\}$  if  $(p \in F \text{ and } q \notin F)$  or  $(p \notin F \text{ and } q \in F)$ .
- 3 Repeat the following until there are no more changes:  
for each unmarked  $\{p, q\}$   
if  $\{\delta(p, a), \delta(q, a)\}$  is marked for some  $a \in \Sigma$ , then

How can we determine whether  $p \approx q$ ?

## Minimization Algorithm



- ① Make a table of all pairs  $\{p, q\}$ , initially unmarked.
- ② Mark  $\{p, q\}$  if  $(p \in F \text{ and } q \notin F)$  or  $(p \notin F \text{ and } q \in F)$ .
- ③ Repeat the following until there are no more changes:  
    for each unmarked  $\{p, q\}$   
        if  $\{\delta(p, a), \delta(q, a)\}$  is marked for some  $a \in \Sigma$ , then  
            mark  $\{p, q\}$ .

How can we determine whether  $p \approx q$ ?


### Minimization Algorithm

- ① Make a table of all pairs  $\{p, q\}$ , initially unmarked.
- ② Mark  $\{p, q\}$  if  $(p \in F \text{ and } q \notin F)$  or  $(p \notin F \text{ and } q \in F)$ .
- ③ Repeat the following until there are no more changes:  
for each unmarked  $\{p, q\}$   
if  $\{\delta(p, a), \delta(q, a)\}$  is marked for some  $a \in \Sigma$ , then  
mark  $\{p, q\}$ .
- ④ Now  $p \approx q$  if and only if  $\{p, q\}$  is not marked.

# Example



# Initialization



## Initialization:

Mark  $\{p, q\}$  if


$(p \in F \text{ and } q \notin F)$  or

$(p \notin F \text{ and } q \in F)$ .

0					
✓	1				
✓		2			
	✓	✓	3		
	✓	✓		4	
✓			✓	✓	5



# First Pass




## First pass:

- Mark  $\{1, 5\}$  because  $\{1, 5\} \xrightarrow{a} \{3, 5\}$  is marked
- Mark  $\{2, 5\}$  because  $\{2, 5\} \xrightarrow{a} \{4, 5\}$  is marked

0					
✓	1				
✓		2			
	✓	✓	3		
	✓	✓		4	
✓	✓	✓	✓	✓	5

## Second Pass




### Second pass:

- Mark  $\{0, 3\}$  because  $\{0, 3\} \xrightarrow{a} \{1, 5\}$  is marked
- Mark  $\{0, 4\}$  because  $\{0, 4\} \xrightarrow{b} \{2, 5\}$  is marked

0					
✓	1				
✓		2			
✓	✓	✓	3		
✓	✓	✓		4	
✓	✓	✓	✓	✓	5

# Third Pass



**Third pass:** no changes


**Output of algorithm:**

- $1 \approx 2, 3 \approx 4$
- $[0] = \{0\},$   
 $[1] = \{1, 2\} = [2],$   
 $[3] = \{3, 4\} = [4],$   
 $[5] = \{5\}$

0					
✓	1				
✓		2			
✓	✓	✓	3		
✓	✓	✓		4	
✓	✓	✓	✓	✓	5

# Quotient Automaton


**Original DFA:**



$$[0] = \{0\}, [1] = \{1, 2\} = [2], [3] = \{3, 4\} = [4], [5] = \{5\}$$


# Quotient Automaton

## Original DFA:




$$[0] = \{0\}, [1] = \{1, 2\} = [2], [3] = \{3, 4\} = [4], [5] = \{5\}$$

## Quotient Automaton:




# Quotient Automaton

## Original DFA:



$$[0] = \{0\}, [1] = \{1, 2\} = [2], [3] = \{3, 4\} = [4], [5] = \{5\}$$


## Quotient Automaton:



- $\delta'([0], a) = [\delta(0, a)] = [1]$


# Quotient Automaton

## Original DFA:



$$[0] = \{0\}, [1] = \{1, 2\} = [2], [3] = \{3, 4\} = [4], [5] = \{5\}$$


## Quotient Automaton:



- $\delta'([0], a) = [\delta(0, a)] = [1]$
- $\delta'([0], b) = [\delta(0, b)] = [2] = [1]$


# Quotient Automaton

## Original DFA:



$$[0] = \{0\}, [1] = \{1, 2\} = [2], [3] = \{3, 4\} = [4], [5] = \{5\}$$



## Quotient Automaton:



- $\delta'([0], a) = [\delta(0, a)] = [1]$
- $\delta'([0], b) = [\delta(0, b)] = [2] = [1]$
- $\delta'([1], b) = [\delta(1, b)] = [4] = [3]$




# Example



# Initialization

## Initialization:

Mark  $\{p, q\}$  if  
( $p \in F$  and  $q \notin F$ ) or  
( $p \notin F$  and  $q \in F$ ).




0					
	1				
✓	✓	2			
		✓	3		
		✓		4	
✓	✓		✓	✓	5

# First Pass

## First pass:

- Mark  $\{0, 1\}$  because  $\{0, 1\} \xrightarrow{a} \{1, 2\}$  is marked
- Mark  $\{0, 4\}$  because  $\{0, 4\} \xrightarrow{a} \{1, 5\}$  is marked
- Mark  $\{1, 3\}$  because  $\{1, 3\} \xrightarrow{a} \{2, 4\}$  is marked
- Mark  $\{3, 4\}$  because  $\{3, 4\} \xrightarrow{a} \{4, 5\}$  is marked



0					
✓	1				
✓	✓	2			
	✓	✓	3		
✓		✓	✓	4	
✓	✓		✓	✓	5


## Second Pass

### Second pass:

- No changes.

### Output of algorithm:


- $0 \approx 3, 1 \approx 4, 2 \approx 5$
- $[0] = \{0, 3\} = [3],$   
 $[1] = \{1, 4\} = [4],$   
 $[2] = \{2, 5\} = [5]$



0					
✓	1				
✓	✓	2			
	✓	✓	3		
✓		✓	✓	4	
✓	✓		✓	✓	5

# Quotient Automaton

**Original DFA:**




$$[0] = \{0, 3\} = [3]$$

$$[1] = \{1, 4\} = [4]$$

$$[2] = \{2, 5\} = [5]$$

# Quotient Automaton

## Original DFA:




$$[0] = \{0, 3\} = [3]$$

$$[1] = \{1, 4\} = [4]$$

$$[2] = \{2, 5\} = [5]$$

## Quotient Automaton:



$$\delta'([0], a) = [\delta(0, a)] = [1]$$

$$\delta'([1], a) = [\delta(1, a)] = [2]$$

$$\delta'([2], a) = [\delta(2, a)] = [3] = [0]$$