


Computability and Complexity  
COSC 4200

NP-Complete Problems II

Plan for NP-completeness reductions:



# Hamiltonian Cycle

Recall that a Hamiltonian cycle in a graph is a cycle that visits all the vertices without repetition. Not all graphs have a Hamiltonian cycle, and determining if a graph has such a cycle is NP-complete.

The decision problem is

$$\text{HAM-CYCLE} = \{G \mid G \text{ has a Hamiltonian cycle}\}.$$


## Theorem

HAM-CYCLE *is* NP-complete.

**Proof.** We have already seen that HAM-CYCLE  $\in$  NP. We will do a reduction from 3SAT. Let


$$\phi = C_1 \wedge \dots \wedge C_m$$

be a formula over  $n$  variables  $x_1, \dots, x_n$ .




For each  $x_i$ , we construct a ladder  $H_i$  of  $12m + 7$  vertices.

- We have  $4m + 2$  rungs with 3 vertices in each rung, and a vertex at the bottom and at the top.
- The rungs of the ladders are composed of  $u_{i,j}$ ,  $v_{i,j}$ , and  $w_{i,j}$ , where  $i$  corresponds to  $H_i$ , and  $j$  corresponds to the rung.
- There is an edge from  $u_{i,j}$  to the  $u$  above and below it (or to  $y_i$  or  $y_{i+1}$ ) and an edge to  $v_{i,j}$ . Similarly with  $w_{i,j}$ .



This is actually  $12m + 8$  vertices, but we hook this ladders together at the top and bottom, so we only count, for example, the bottom vertex as belonging to  $H_i$ , while the top one belongs to  $H_{i+1}$ . The top of the top ladder connects to the bottom of the bottom ladder.

There are two ways to traverse a ladder, either starting by going to the left, or going to the right. Intuitively, going left in  $H_i$  corresponds to setting  $x_i$  true, and going right corresponds to setting  $x_i$  false.





For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

This completes the construction of the graph.

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.




•  $z_2$


•  $z_1$

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$







$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$




$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$




$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$




$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



•  $z_2$





•  $z_1$

For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$




For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$






For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$




For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$






For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .


$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$



For each clause  $C_j$ , we add a new vertex  $z_j$  such that

- If  $C_j$  contains the literal  $x_i$ , we connect  $z_j$  to  $u_{i,4j-1}$ , and  $u_{i,4j-2}$ .
- If  $C_j$  contains the literal  $\neg x_i$ , we connect  $z_j$  to  $w_{i,4j-1}$  and  $w_{i,4j-2}$ .

$$\phi = (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_2)$$

Observe that:

- If we go left in a Hamiltonian cycle in ladder  $H_i$ , we will traverse
  - the even edges  $(u_{i,2}, u_{i,3}), (u_{i,4}, u_{i,5}), \dots$  on the left side, and
  - the odd edges  $(w_{i,1}, w_{i,1}), (w_{i,3}, w_{i,4}), \dots$  on the right side.



Observe that:

- If we go left in a Hamiltonian cycle in ladder  $H_i$ , we will traverse
  - the even edges  $(u_{i,2}, u_{i,3}), (u_{i,4}, u_{i,5}), \dots$  on the left side, and
  - the odd edges  $(w_{i,1}, w_{i,1}), (w_{i,3}, w_{i,4}), \dots$  on the right side.

If  $x_i$  appears as a literal in clause  $C_j$ , then  $z_j$  will be connected to the endpoints of one of the even edges on the left:

$$(u_{i,4j-2}, u_{i,4j-1}).$$

The cycle could safely visit  $z_j$  instead of taking this edge. In this way, going left in  $H_i$  corresponds to setting  $x_i = T$ .

- Similarly, if we go right in a Hamiltonian cycle in ladder  $H_i$ , we will traverse
  - the odd edges  $(u_{i,1}, u_{i,2}), (u_{i,3}, u_{i,4}), \dots$  on the left side, and
  - the even edges  $(w_{i,2}, w_{i,3}), (w_{i,4}, w_{i,5}), \dots$  on the right side.

- Similarly, if we go right in a Hamiltonian cycle in ladder  $H_i$ , we will traverse
  - the odd edges  $(u_{i,1}, u_{i,2}), (u_{i,3}, u_{i,4}), \dots$  on the left side, and
  - the even edges  $(w_{i,2}, w_{i,3}), (w_{i,4}, w_{i,5}), \dots$  on the right side.

If  $\neg x_i$  appears as a literal in clause  $C_j$ , then  $z_j$  will be connected to the endpoints of one of the even edges on the right:

$$(w_{i,4j-2}, w_{i,4j-1}).$$

The cycle could safely visit  $z_j$  instead of taking this edge. In this way, going left in  $H_i$  corresponds to setting  $x_i = F$ .

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.

Suppose  $\phi$  is satisfiable and fix a satisfying assignment. We use this assignment to define a cycle that goes left at  $H_i$  if  $x_i = T$  and goes right if  $x_i = F$ .

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.

Suppose  $\phi$  is satisfiable and fix a satisfying assignment. We use this assignment to define a cycle that goes left at  $H_i$  if  $x_i = T$  and goes right if  $x_i = F$ .

- The  $z_j$ 's are connected to the ladders so that if  $C_j$  contains  $\neg x_i$ , then vertex  $z_j$  can be visited by a cycle that goes right in ladder  $H_i$ .

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.

Suppose  $\phi$  is satisfiable and fix a satisfying assignment. We use this assignment to define a cycle that goes left at  $H_i$  if  $x_i = T$  and goes right if  $x_i = F$ .

- The  $z_j$ 's are connected to the ladders so that if  $C_j$  contains  $\neg x_i$ , then vertex  $z_j$  can be visited by a cycle that goes right in ladder  $H_i$ .
- Similarly, if  $C_j$  contains  $x_i$ , then vertex  $z_j$  can be visited by a cycle that goes left in ladder  $H_i$ .

Our claim is that  $\phi$  is satisfiable if and only if the graph has a Hamiltonian cycle.

Suppose  $\phi$  is satisfiable and fix a satisfying assignment. We use this assignment to define a cycle that goes left at  $H_i$  if  $x_i = T$  and goes right if  $x_i = F$ .

- The  $z_j$ 's are connected to the ladders so that if  $C_j$  contains  $\neg x_i$ , then vertex  $z_j$  can be visited by a cycle that goes right in ladder  $H_i$ .
- Similarly, if  $C_j$  contains  $x_i$ , then vertex  $z_j$  can be visited by a cycle that goes left in ladder  $H_i$ .

Thus we can extend the cycle to visit all the  $z_j$ 's. Therefore if  $\phi$  is satisfiable, then there is a Hamiltonian cycle.



For the other direction, suppose there is a Hamiltonian cycle.

For the other direction, suppose there is a Hamiltonian cycle.

- We set  $x_i = T$  if the cycle goes left at  $H_i$  and set  $x_i = F$  if it goes right.

For the other direction, suppose there is a Hamiltonian cycle.

- We set  $x_i = T$  if the cycle goes left at  $H_i$  and set  $x_i = F$  if it goes right.
- The cycle visits all the  $z_j$ 's, so it follows from the construction that this assignment satisfies all clauses of  $\phi$ .

Therefore if there is a Hamiltonian cycle, then  $\phi$  is satisfiable.


For the other direction, suppose there is a Hamiltonian cycle.

- We set  $x_i = T$  if the cycle goes left at  $H_i$  and set  $x_i = F$  if it goes right.
- The cycle visits all the  $z_j$ 's, so it follows from the construction that this assignment satisfies all clauses of  $\phi$ .

Therefore if there is a Hamiltonian cycle, then  $\phi$  is satisfiable.

The graph we constructed can be computed in quadratic time in the size of the formula. Therefore  $3SAT \leq_P HAM-CYCLE$ .  $\square$

Plan for NP-completeness reductions:



# Traveling Salesman Problem

In the traveling salesman problem, a salesman wishes to visit a collection of cities at the minimum possible cost.

- For each pair of cities,  $A$  and  $B$ , there is a cost associated with traveling from  $A$  to  $B$ .
- The goal is to find a tour of the cities of the minimum total cost.

# Traveling Salesman Problem

In the traveling salesman problem, a salesman wishes to visit a collection of cities at the minimum possible cost.

- For each pair of cities,  $A$  and  $B$ , there is a cost associated with traveling from  $A$  to  $B$ .
- The goal is to find a tour of the cities of the minimum total cost.

We model an instance of this problem as follows:

- $n$  = number of cities. The cities are numbered from 1 to  $n$ .
- $c : \{1, \dots, n\} \times \{1, \dots, n\} \rightarrow \mathbb{N}^+$  is a cost function.  
 $c(i, j)$  is the cost of traveling from city  $i$  to city  $j$ .

# Traveling Salesman Problem

A tour of the  $n$  cities is a sequence  $(a_1, \dots, a_n, a_{n+1})$  where

- $a_i \in \{1, \dots, n\}$  for all  $1 \leq i \leq n$ ,
- $a_{n+1} = a_1$ , and
- for each  $j \in \{1, \dots, n\}$ , there is some  $i \in \{1, \dots, n\}$  such that  $a_i = j$ .



# Traveling Salesman Problem


A tour of the  $n$  cities is a sequence  $(a_1, \dots, a_n, a_{n+1})$  where

- $a_i \in \{1, \dots, n\}$  for all  $1 \leq i \leq n$ ,
- $a_{n+1} = a_1$ , and
- for each  $j \in \{1, \dots, n\}$ , there is some  $i \in \{1, \dots, n\}$  such that  $a_i = j$ .

The cost of the tour is

$$\sum_{i=1}^n c(a_i, a_{i+1})$$

Sometimes this is also called the length of the tour.



A TSP tour of the continental U.S. state capitals.

# Traveling Salesman Problem

The traveling salesman decision problem is

$$\text{TSP} = \left\{ \langle n, c, k \rangle \mid \begin{array}{l} c \text{ is a cost function and there} \\ \text{is a tour with cost at most } k \end{array} \right\}$$

It is easy to see that  $\text{TSP} \in \text{NP}$ . The witnesses are the tours. We can easily check if the cost of a tour is at most  $k$ .

## Theorem

TSP *is* NP-complete.

Before we prove this theorem, let's discuss the consequences.

In the TSP decision problem, we only have to decide if a tour with cost at most  $k$  exists. In practice what we really want is to find a minimum cost tour.

## Theorem

*TSP is NP-complete.*

Before we prove this theorem, let's discuss the consequences.

In the TSP decision problem, we only have to decide if a tour with cost at most  $k$  exists. In practice what we really want is to find a minimum cost tour. A corollary of the theorem is the following.

## Corollary

*If  $P \neq NP$ , then there is no polynomial-time algorithm that finds shortest TSP tours.*

## Theorem

*TSP is NP-complete.*

Before we prove this theorem, let's discuss the consequences.

In the TSP decision problem, we only have to decide if a tour with cost at most  $k$  exists. In practice what we really want is to find a minimum cost tour. A corollary of the theorem is the following.

## Corollary

*If  $P \neq NP$ , then there is no polynomial-time algorithm that finds shortest TSP tours.*

**Proof.** Suppose we have a polynomial-time algorithm for finding the shortest tours. Then given an instance  $\langle n, c, k \rangle$  of TSP, we can compute the shortest tour and see if its cost is at most  $k$ . This shows that  $TSP \in P$ , so  $P = NP$  because TSP is NP-complete.  $\square$

## Theorem

TSP *is* NP-complete.

**Proof.** We will reduce HAM-CYCLE to TSP. Let  $G = (V, E)$  be a graph.

## Theorem

TSP is NP-complete.

**Proof.** We will reduce HAM-CYCLE to TSP. Let  $G = (V, E)$  be a graph.

Define  $n = |V|$ ,  $k = n$ , and

$$c(i, j) = \begin{cases} 1 & \text{if } (i, j) \in E \\ n + 1 & \text{if } (i, j) \notin E \end{cases}$$

for all  $i, j \in \{1, \dots, n\}$ .



## Theorem

TSP is NP-complete.

**Proof.** We will reduce HAM-CYCLE to TSP. Let  $G = (V, E)$  be a graph.

Define  $n = |V|$ ,  $k = n$ , and

$$c(i, j) = \begin{cases} 1 & \text{if } (i, j) \in E \\ n + 1 & \text{if } (i, j) \notin E \end{cases}$$

for all  $i, j \in \{1, \dots, n\}$ .

We claim that

$$G \in \text{HAM-CYCLE} \iff \langle n, c, k \rangle \in \text{TSP}.$$

Suppose that  $G$  has a Hamiltonian cycle. Then that cycle is a tour of length  $n = k$  in the TSP instance. Therefore  $G \in \text{HAM-CYCLE}$  implies  $\langle n, c, k \rangle \in \text{TSP}$ .

Suppose that  $G$  has a Hamiltonian cycle. Then that cycle is a tour of length  $n = k$  in the TSP instance. Therefore  $G \in \text{HAM-CYCLE}$  implies  $\langle n, c, k \rangle \in \text{TSP}$ .

For the converse, suppose that there is a tour with cost  $\leq k = n$ . Then that tour never goes from a city  $i$  to a city  $j$  with a cost of  $n + 1$ . Therefore, there is an edge in  $G$  between every consecutive pair of cities on the tour, so the tour is a Hamiltonian cycle in  $G$ . Therefore  $\langle n, c, k \rangle \in \text{TSP}$  implies  $G \in \text{HAM-CYCLE}$ .

Suppose that  $G$  has a Hamiltonian cycle. Then that cycle is a tour of length  $n = k$  in the TSP instance. Therefore  $G \in \text{HAM-CYCLE}$  implies  $\langle n, c, k \rangle \in \text{TSP}$ .


For the converse, suppose that there is a tour with cost  $\leq k = n$ . Then that tour never goes from a city  $i$  to a city  $j$  with a cost of  $n + 1$ . Therefore, there is an edge in  $G$  between every consecutive pair of cities on the tour, so the tour is a Hamiltonian cycle in  $G$ . Therefore  $\langle n, c, k \rangle \in \text{TSP}$  implies  $G \in \text{HAM-CYCLE}$ .

Since we can compute  $\langle n, c, k \rangle$  from  $G$  in polynomial time, we have shown that

$$\text{HAM-CYCLE} \leq_P \text{TSP}.$$



Plan for NP-completeness reductions:



# Subset Sum

The subset sum decision problem is

$$\text{SUBSET-SUM} = \left\{ \langle L, S \rangle \left| \begin{array}{l} L = (a_1, \dots, a_n) \text{ is a list of } n \text{ integers} \\ \text{and} \\ \exists I \subseteq \{1, \dots, n\} \text{ such that } \sum_{i \in I} a_i = S \end{array} \right. \right\}$$

## Theorem

SUBSET-SUM *is* NP-complete.

**Proof.** We've seen that SUBSET-SUM  $\in$  NP. We'll reduce 3SAT to SUBSET-SUM.

Let  $\phi = C_1 \wedge \dots \wedge C_m$  be a 3CNF formula over  $n$  variables  $x_1, \dots, x_n$ .

## Theorem

SUBSET-SUM *is NP-complete*.

**Proof.** We've seen that SUBSET-SUM  $\in$  NP. We'll reduce 3SAT to SUBSET-SUM.

Let  $\phi = C_1 \wedge \dots \wedge C_m$  be a 3CNF formula over  $n$  variables  $x_1, \dots, x_n$ .

We will define a list of  $2n + 2m$  integers that will all have values between 0 and  $10^{n+m}$ , and a target sum.

We will write them as decimal numbers using exactly  $n + m$  digits (using leading 0's when necessary).



For any integer  $r$ , let  $r[k]$  be the  $k^{\text{th}}$  most significant digit in the decimal representation of  $r$ . So if  $r = 0156$ , then  $r[1] = 0$  and  $r[4] = 6$ .

For any integer  $r$ , let  $r[k]$  be the  $k^{\text{th}}$  most significant digit in the decimal representation of  $r$ . So if  $r = 0156$ , then  $r[1] = 0$  and  $r[4] = 6$ .

We define  $S$ , our target sum, by

$$S[k] = \begin{cases} 1 & \text{if } 1 \leq k \leq n \\ 3 & \text{if } n+1 \leq k \leq n+m. \end{cases}$$

In other words,  $S$  is the number

$$S = \underbrace{11 \dots 1}_n \underbrace{33 \dots 3}_m.$$

For each  $1 \leq j \leq n$  we define two numbers  $b_{j,0}$  and  $b_{j,1}$  as follows.

For each  $1 \leq j \leq n$  we define two numbers  $b_{j,0}$  and  $b_{j,1}$  as follows.

- For all  $1 \leq k \leq n$ ,

$$b_{j,0}[k] = b_{j,1}[k] = \begin{cases} 1 & \text{if } k = j \\ 0 & \text{if } k \neq j. \end{cases}$$

For each  $1 \leq j \leq n$  we define two numbers  $b_{j,0}$  and  $b_{j,1}$  as follows.

- For all  $1 \leq k \leq n$ ,

$$b_{j,0}[k] = b_{j,1}[k] = \begin{cases} 1 & \text{if } k = j \\ 0 & \text{if } k \neq j. \end{cases}$$

- For all  $1 \leq k \leq m$ ,

$$b_{j,0}[n+k] = \begin{cases} 1 & \text{if } \neg x_j \text{ occurs in } C_k \\ 0 & \text{otherwise.} \end{cases}$$

For each  $1 \leq j \leq n$  we define two numbers  $b_{j,0}$  and  $b_{j,1}$  as follows.

- For all  $1 \leq k \leq n$ ,

$$b_{j,0}[k] = b_{j,1}[k] = \begin{cases} 1 & \text{if } k = j \\ 0 & \text{if } k \neq j. \end{cases}$$

- For all  $1 \leq k \leq m$ ,

$$b_{j,0}[n+k] = \begin{cases} 1 & \text{if } \neg x_j \text{ occurs in } C_k \\ 0 & \text{otherwise.} \end{cases}$$

- For all  $1 \leq k \leq m$ ,

$$b_{j,1}[n+k] = \begin{cases} 1 & \text{if } x_j \text{ occurs in } C_k \\ 0 & \text{otherwise.} \end{cases}$$

For each  $1 \leq i \leq m$  we define two numbers  $c_{i,0}$  and  $c_{i,1}$  by

$$c_{i,0}[k] = c_{i,1}[k] = \begin{cases} 1 & \text{if } k = n + i \\ 0 & \text{otherwise} \end{cases}$$

for all  $1 \leq k \leq n + m$ .

For each  $1 \leq i \leq m$  we define two numbers  $c_{i,0}$  and  $c_{i,1}$  by

$$c_{i,0}[k] = c_{i,1}[k] = \begin{cases} 1 & \text{if } k = n + i \\ 0 & \text{otherwise} \end{cases}$$

for all  $1 \leq k \leq n + m$ .

Let  $L$  be the list of these numbers:

$$L = (b_{1,0}, b_{2,0}, \dots, b_{n,0}, b_{1,1}, \dots, b_{n,1}, c_{1,0}, \dots, c_{n,0}, c_{1,1}, \dots, c_{n,1})$$

Note that  $(L, S)$  can be computed in polynomial time from  $\phi$ .



For each  $1 \leq i \leq m$  we define two numbers  $c_{i,0}$  and  $c_{i,1}$  by

$$c_{i,0}[k] = c_{i,1}[k] = \begin{cases} 1 & \text{if } k = n + i \\ 0 & \text{otherwise} \end{cases}$$

for all  $1 \leq k \leq n + m$ .

Let  $L$  be the list of these numbers:

$$L = (b_{1,0}, b_{2,0}, \dots, b_{n,0}, b_{1,1}, \dots, b_{n,1}, c_{1,0}, \dots, c_{n,0}, c_{1,1}, \dots, c_{n,1})$$

Note that  $(L, S)$  can be computed in polynomial time from  $\phi$ .

Claim

$$\phi \in 3\text{SAT} \iff (L, S) \in \text{SUBSET-SUM}.$$

As an example, consider

$$\phi = (\neg x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3).$$

Then  $n = m = 3$ ,  $S = 111333$ , and

$b_{1,0}$	$=$	100101	$c_{1,0}$	$=$	000100
$b_{1,1}$	$=$	100010	$c_{1,1}$	$=$	000100
$b_{2,0}$	$=$	010011	$c_{2,0}$	$=$	000010
$b_{2,1}$	$=$	010100	$c_{2,1}$	$=$	000010
$b_{3,0}$	$=$	001101	$c_{3,0}$	$=$	000001
$b_{3,1}$	$=$	001010	$c_{3,1}$	$=$	000001

Observe that for any  $k$ ,  $n < k \leq n + m$ , there are at most 5 numbers in  $L$  with a 1 in the  $k^{\text{th}}$  digit. This is because each clause has at most 3 literals, and there are  $2 + (\text{the number of literals in the } k^{\text{th}} \text{ clause})$  many 1's in the  $k^{\text{th}}$  digits of numbers in  $L$ .

Observe that for any  $k$ ,  $n < k \leq n + m$ , there are at most 5 numbers in  $L$  with a 1 in the  $k^{\text{th}}$  digit. This is because each clause has at most 3 literals, and there are  $2 + (\text{the number of literals in the } k^{\text{th}} \text{ clause})$  many 1's in the  $k^{\text{th}}$  digits of numbers in  $L$ .

Also, for any  $k$ ,  $1 \leq k \leq n$ , there are exactly two numbers with a 1 in the  $k^{\text{th}}$  digit, namely  $b_{k,0}$  and  $b_{k,1}$ .

Observe that for any  $k$ ,  $n < k \leq n + m$ , there are at most 5 numbers in  $L$  with a 1 in the  $k^{\text{th}}$  digit. This is because each clause has at most 3 literals, and there are  $2 + (\text{the number of literals in the } k^{\text{th}} \text{ clause})$  many 1's in the  $k^{\text{th}}$  digits of numbers in  $L$ .

Also, for any  $k$ ,  $1 \leq k \leq n$ , there are exactly two numbers with a 1 in the  $k^{\text{th}}$  digit, namely  $b_{k,0}$  and  $b_{k,1}$ .

Therefore to get a sum of  $S$  we need to choose a subset of the numbers that

- has exactly three numbers with a 1 in the  $k^{\text{th}}$  position for  $n < k \leq n + m$  and
- has either  $b_{k,0}$  or  $b_{k,1}$  for  $1 \leq k \leq n$ .

$$\phi = (\neg x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3).$$

$b_{1,0}$	=	100101	$c_{1,0}$	=	000100
$b_{1,1}$	=	100010	$c_{1,1}$	=	000100
$b_{2,0}$	=	010011	$c_{2,0}$	=	000010
$b_{2,1}$	=	010100	$c_{2,1}$	=	000010
$b_{3,0}$	=	001101	$c_{3,0}$	=	000001
$b_{3,1}$	=	001010	$c_{3,1}$	=	000001
$S$	=	111333			

Satisfying assignment:  $x_1 = F$ ,  $x_2 = F$ , and  $x_3 = T$

$$\phi = (\neg x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3).$$

$b_{1,0}$	$=$	100101	$c_{1,0}$	$=$	000100
$b_{1,1}$	$=$	100010	$c_{1,1}$	$=$	000100
$b_{2,0}$	$=$	010011	$c_{2,0}$	$=$	000010
$b_{2,1}$	$=$	010100	$c_{2,1}$	$=$	000010
$b_{3,0}$	$=$	001101	$c_{3,0}$	$=$	000001
$b_{3,1}$	$=$	001010	$c_{3,1}$	$=$	000001
$S$	$=$	111333			

Satisfying assignment:  $x_1 = F$ ,  $x_2 = F$ , and  $x_3 = T$

Choose:

$b_{1,0}$	$=$	100101	$(x_1 = F)$
$b_{2,0}$	$=$	010011	$(x_2 = F)$
$b_{3,1}$	$=$	<u>001010</u>	$(x_3 = T)$
		111122	

$$\phi = (\neg x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3).$$

$b_{1,0}$	=	100101	$c_{1,0}$	=	000100
$b_{1,1}$	=	100010	$c_{1,1}$	=	000100
$b_{2,0}$	=	010011	$c_{2,0}$	=	000010
$b_{2,1}$	=	010100	$c_{2,1}$	=	000010
$b_{3,0}$	=	001101	$c_{3,0}$	=	000001
$b_{3,1}$	=	001010	$c_{3,1}$	=	000001
$S$	=	111333			

Satisfying assignment:  $x_1 = F$ ,  $x_2 = F$ , and  $x_3 = T$

Choose:

$b_{1,0}$	=	100101	$(x_1 = F)$
$b_{2,0}$	=	010011	$(x_2 = F)$
$b_{3,1}$	=	<u>001010</u>	$(x_3 = T)$
		111122	

The last three digits of this sum, 122, correspond to the number of literals satisfied in each clause:  $\neg x_1$  is satisfied  $C_1$ ,  $\neg x_2$  and  $x_3$  are satisfied  $C_2$ , and  $\neg x_1$  and  $\neg x_2$  are satisfied  $C_3$ .



$$\phi = (\neg x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3).$$

$b_{1,0}$	=	100101	$c_{1,0}$	=	000100
$b_{1,1}$	=	100010	$c_{1,1}$	=	000100
$b_{2,0}$	=	010011	$c_{2,0}$	=	000010
$b_{2,1}$	=	010100	$c_{2,1}$	=	000010
$b_{3,0}$	=	001101	$c_{3,0}$	=	000001
$b_{3,1}$	=	001010	$c_{3,1}$	=	000001
$S$	=	111333			

Satisfying assignment:  $x_1 = F$ ,  $x_2 = F$ , and  $x_3 = T$

Choose:

$b_{1,0}$	=	100101	$(x_1 = F)$
$b_{2,0}$	=	010011	$(x_2 = F)$
$b_{3,1}$	=	<u>001010</u>	$(x_3 = T)$
		111122	

The last three digits of this sum, 122, correspond to the number of literals satisfied in each clause:  $\neg x_1$  is satisfied  $C_1$ ,  $\neg x_2$  and  $x_3$  are satisfied  $C_2$ , and  $\neg x_1$  and  $\neg x_2$  are satisfied  $C_3$ .

To finish, we also choose  $c_{1,0}$ ,  $c_{1,1}$ ,  $c_{2,0}$ , and  $c_{3,0}$  to reach  $S$ .

We must show that  $\phi \in 3\text{SAT} \Rightarrow (L, S) \in \text{SUBSET-SUM}$ . So assume that  $\phi$  is satisfied by an assignment  $\tau$  to the variables  $x_1, \dots, x_n$ . We define a sublist  $L'$  of  $L$  as follows.

We must show that  $\phi \in 3\text{SAT} \Rightarrow (L, S) \in \text{SUBSET-SUM}$ . So assume that  $\phi$  is satisfied by an assignment  $\tau$  to the variables  $x_1, \dots, x_n$ . We define a sublist  $L'$  of  $L$  as follows.

For each  $j$ ,  $1 \leq j \leq n$ ,

- If  $\tau(x_j) = F$ , then put  $b_{j,0}$  in  $L'$ .
- If  $\tau(x_j) = T$ , then put  $b_{j,1}$  in  $L'$ .

We must show that  $\phi \in 3\text{SAT} \Rightarrow (L, S) \in \text{SUBSET-SUM}$ . So assume that  $\phi$  is satisfied by an assignment  $\tau$  to the variables  $x_1, \dots, x_n$ . We define a sublist  $L'$  of  $L$  as follows.

For each  $j$ ,  $1 \leq j \leq n$ ,

- If  $\tau(x_j) = F$ , then put  $b_{j,0}$  in  $L'$ .
- If  $\tau(x_j) = T$ , then put  $b_{j,1}$  in  $L'$ .

For each  $i$ ,  $1 \leq i \leq m$ ,

- If  $\tau$  satisfies 3 literals in  $C_i$ , do nothing.
- If  $\tau$  satisfies 2 literals in  $C_i$ , put  $c_{i,0}$  in  $L'$ .
- If  $\tau$  satisfies 1 literal in  $C_i$ , put both  $c_{i,0}$  and  $c_{i,1}$  in  $L'$ .

We must show that  $\phi \in 3\text{SAT} \Rightarrow (L, S) \in \text{SUBSET-SUM}$ . So assume that  $\phi$  is satisfied by an assignment  $\tau$  to the variables  $x_1, \dots, x_n$ . We define a sublist  $L'$  of  $L$  as follows.

For each  $j$ ,  $1 \leq j \leq n$ ,

- If  $\tau(x_j) = F$ , then put  $b_{j,0}$  in  $L'$ .
- If  $\tau(x_j) = T$ , then put  $b_{j,1}$  in  $L'$ .

For each  $i$ ,  $1 \leq i \leq m$ ,

- If  $\tau$  satisfies 3 literals in  $C_i$ , do nothing.
- If  $\tau$  satisfies 2 literals in  $C_i$ , put  $c_{i,0}$  in  $L'$ .
- If  $\tau$  satisfies 1 literal in  $C_i$ , put both  $c_{i,0}$  and  $c_{i,1}$  in  $L'$ .

Then  $\sum_{a \in L'} a = S$ . Therefore  $(L, S) \in \text{SUBSET-SUM}$ .

Now we must show that  $(L, S) \in \text{SUBSET-SUM} \Rightarrow \phi \in 3\text{SAT}$ .  
Suppose that  $L'$  is a sublist of  $L$  that sums to  $S$ .

Now we must show that  $(L, S) \in \text{SUBSET-SUM} \Rightarrow \phi \in 3\text{SAT}$ .  
Suppose that  $L'$  is a sublist of  $L$  that sums to  $S$ .

We make two observations.

- We know that for each  $j$ ,  $1 \leq j \leq n$ ,  $L'$  has exactly one of  $b_{j,0}$  or  $b_{j,1}$ . There's no other way to get a one in each column. Picking neither gives us a 0, and picking both gives us a 2.

Now we must show that  $(L, S) \in \text{SUBSET-SUM} \Rightarrow \phi \in 3\text{SAT}$ .  
Suppose that  $L'$  is a sublist of  $L$  that sums to  $S$ .

We make two observations.

- We know that for each  $j$ ,  $1 \leq j \leq n$ ,  $L'$  has exactly one of  $b_{j,0}$  or  $b_{j,1}$ . There's no other way to get a one in each column. Picking neither gives us a 0, and picking both gives us a 2.
- Furthermore, we don't have to worry about carry-over from another column because we are working in decimal notation and the largest possible sum in a column is 5.



Now we must show that  $(L, S) \in \text{SUBSET-SUM} \Rightarrow \phi \in 3\text{SAT}$ .  
Suppose that  $L'$  is a sublist of  $L$  that sums to  $S$ .

We make two observations.

- We know that for each  $j$ ,  $1 \leq j \leq n$ ,  $L'$  has exactly one of  $b_{j,0}$  or  $b_{j,1}$ . There's no other way to get a one in each column. Picking neither gives us a 0, and picking both gives us a 2.
- Furthermore, we don't have to worry about carry-over from another column because we are working in decimal notation and the largest possible sum in a column is 5.

We define our assignment  $\tau$  as follows.

$$\tau(x_j) = \begin{cases} T & \text{if } L' \text{ contains } b_{j,1} \\ F & \text{if } L' \text{ contains } b_{j,0} \end{cases}$$

We now verify that  $\tau$  satisfies  $\phi$ .

For each  $i$ ,  $1 \leq i \leq m$ , the  $(n+i)^{\text{th}}$  digit in the sum of  $L'$  is 3, so in  $L'$  there must be some  $b_{j,0}$  or  $b_{j,1}$  that has the  $(n+i)^{\text{th}}$  digit equal to 1. This is because from  $c_{i,0}$  and  $c_{i,1}$  we can get at most a sum of 2 in the  $(n+i)^{\text{th}}$  column, so we must get at least one 1 from the  $b_{j,k}$ 's.

We now verify that  $\tau$  satisfies  $\phi$ .

For each  $i$ ,  $1 \leq i \leq m$ , the  $(n+i)^{\text{th}}$  digit in the sum of  $L'$  is 3, so in  $L'$  there must be some  $b_{j,0}$  or  $b_{j,1}$  that has the  $(n+i)^{\text{th}}$  digit equal to 1. This is because from  $c_{i,0}$  and  $c_{i,1}$  we can get at most a sum of 2 in the  $(n+i)^{\text{th}}$  column, so we must get at least one 1 from the  $b_{j,k}$ 's.

- Suppose it is a  $b_{j,0}$ . Then  $\neg x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = F$ , so our assignment satisfies that clause.

We now verify that  $\tau$  satisfies  $\phi$ .

For each  $i$ ,  $1 \leq i \leq m$ , the  $(n+i)^{\text{th}}$  digit in the sum of  $L'$  is 3, so in  $L'$  there must be some  $b_{j,0}$  or  $b_{j,1}$  that has the  $(n+i)^{\text{th}}$  digit equal to 1. This is because from  $c_{i,0}$  and  $c_{i,1}$  we can get at most a sum of 2 in the  $(n+i)^{\text{th}}$  column, so we must get at least one 1 from the  $b_{j,k}$ 's.

- Suppose it is a  $b_{j,0}$ . Then  $\neg x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = F$ , so our assignment satisfies that clause.
- Similarly, suppose it is a  $b_{j,1}$ . Then  $x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = T$ , so our assignment satisfies that clause.

We now verify that  $\tau$  satisfies  $\phi$ .

For each  $i$ ,  $1 \leq i \leq m$ , the  $(n+i)^{\text{th}}$  digit in the sum of  $L'$  is 3, so in  $L'$  there must be some  $b_{j,0}$  or  $b_{j,1}$  that has the  $(n+i)^{\text{th}}$  digit equal to 1. This is because from  $c_{i,0}$  and  $c_{i,1}$  we can get at most a sum of 2 in the  $(n+i)^{\text{th}}$  column, so we must get at least one 1 from the  $b_{j,k}$ 's.

- Suppose it is a  $b_{j,0}$ . Then  $\neg x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = F$ , so our assignment satisfies that clause.
- Similarly, suppose it is a  $b_{j,1}$ . Then  $x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = T$ , so our assignment satisfies that clause.

Therefore,  $\tau$  satisfies all the clauses of  $\phi$ , so  $\phi$  is satisfiable, and  $\phi \in 3\text{SAT}$ .

We now verify that  $\tau$  satisfies  $\phi$ .

For each  $i$ ,  $1 \leq i \leq m$ , the  $(n+i)^{\text{th}}$  digit in the sum of  $L'$  is 3, so in  $L'$  there must be some  $b_{j,0}$  or  $b_{j,1}$  that has the  $(n+i)^{\text{th}}$  digit equal to 1. This is because from  $c_{i,0}$  and  $c_{i,1}$  we can get at most a sum of 2 in the  $(n+i)^{\text{th}}$  column, so we must get at least one 1 from the  $b_{j,k}$ 's.

- Suppose it is a  $b_{j,0}$ . Then  $\neg x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = F$ , so our assignment satisfies that clause.
- Similarly, suppose it is a  $b_{j,1}$ . Then  $x_j$  is a literal in clause  $C_i$  and  $\tau(x_j) = T$ , so our assignment satisfies that clause.

Therefore,  $\tau$  satisfies all the clauses of  $\phi$ , so  $\phi$  is satisfiable, and  $\phi \in 3\text{SAT}$ .

Since the construction is polynomial-time computable, we have shown that  $3\text{SAT} \leq_P \text{SUBSET-SUM}$ . □

# NP-Completeness Summary

Let us review the method of showing a problem  $B$  is NP-complete.

- 1 Show that  $B \in \text{NP}$ . It is usually easy to show this, but it is a step that shouldn't be overlooked.

# NP-Completeness Summary

Let us review the method of showing a problem  $B$  is NP-complete.

- ① Show that  $B \in \text{NP}$ . It is usually easy to show this, but it is a step that shouldn't be overlooked.
- ② Choose some NP-complete problem  $A$  that is a candidate for reduction to  $B$ . Often we try to find a problem that is similar to  $B$  or use 3SAT.



# NP-Completeness Summary

Let us review the method of showing a problem  $B$  is NP-complete.

- ① Show that  $B \in \text{NP}$ . It is usually easy to show this, but it is a step that shouldn't be overlooked.
- ② Choose some NP-complete problem  $A$  that is a candidate for reduction to  $B$ . Often we try to find a problem that is similar to  $B$  or use 3SAT.
- ③ Reduce  $A$  to  $B$ .

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

Then we need to prove that for every instance  $I$  of  $A$ ,

$$I \in A \iff I' \in B.$$

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

Then we need to prove that for every instance  $I$  of  $A$ ,

$$I \in A \iff I' \in B.$$

This involves showing two implications:  $I \in A \implies I' \in B$  and  $I' \in B \implies I \in A$ .

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

Then we need to prove that for every instance  $I$  of  $A$ ,

$$I \in A \iff I' \in B.$$

This involves showing two implications:  $I \in A \implies I' \in B$  and  $I' \in B \implies I \in A$ .

- To show that  $I \in A \implies I' \in B$ , assume that  $I \in A$ . Then there is some witness  $w$  for  $I$  that shows  $I$  is a member of  $A$ . We use this witness to construct a witness  $w'$  that shows  $I' \in B$ .

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

Then we need to prove that for every instance  $I$  of  $A$ ,

$$I \in A \iff I' \in B.$$

This involves showing two implications:  $I \in A \implies I' \in B$  and  $I' \in B \implies I \in A$ .

- To show that  $I \in A \implies I' \in B$ , assume that  $I \in A$ . Then there is some witness  $w$  for  $I$  that shows  $I$  is a member of  $A$ . We use this witness to construct a witness  $w'$  that shows  $I' \in B$ .
- To show the converse, assume that  $I' \in B$ . Then there is some witness  $w'$  for  $I'$ . We use  $w'$  to construct a witness  $w$  that shows  $I \in A$ .

③ Reduce  $A$  to  $B$ .

For this, we need to define a mapping of instances  $I$  of  $A$  to instances  $I'$  of  $B$ .

$$I \mapsto I'$$

Then we need to prove that for every instance  $I$  of  $A$ ,

$$I \in A \iff I' \in B.$$

This involves showing two implications:  $I \in A \implies I' \in B$  and  $I' \in B \implies I \in A$ .

- To show that  $I \in A \implies I' \in B$ , assume that  $I \in A$ . Then there is some witness  $w$  for  $I$  that shows  $I$  is a member of  $A$ . We use this witness to construct a witness  $w'$  that shows  $I' \in B$ .
- To show the converse, assume that  $I' \in B$ . Then there is some witness  $w'$  for  $I'$ . We use  $w'$  to construct a witness  $w$  that shows  $I \in A$ .

Finally, we have to verify that the mapping is polynomial-time computable. Usually this doesn't require formal analysis of the algorithm.

Summary of NP-completeness reductions:

