

Computability and Complexity

COSC 4200

The Myhill-Nerode Theorem

The Relation \equiv_M

Let $A \subseteq \Sigma^*$ be a regular language, and let $M = (Q, \Sigma, \delta, q_0, F)$ be a DFA for A with no inaccessible states.

M induces an equivalence relation \equiv_M on Σ^* defined by

$$x \equiv_M y \iff \delta^*(q_0, x) = \delta^*(q_0, y)$$

for all $x, y \in \Sigma^*$.

Properties of \equiv_M

Besides the reflexive, symmetric, and transitive properties, \equiv_M satisfies a few other useful properties:

Properties of \equiv_M

Besides the reflexive, symmetric, and transitive properties, \equiv_M satisfies a few other useful properties:

- ① It is a *right congruence*: for all $x, y \in \Sigma^*$ and $a \in \Sigma$,

$$x \equiv_M y \implies xa \equiv_M ya.$$

Properties of \equiv_M

Besides the reflexive, symmetric, and transitive properties, \equiv_M satisfies a few other useful properties:

- 1 It is a *right congruence*: for all $x, y \in \Sigma^*$ and $a \in \Sigma$,

$$x \equiv_M y \implies xa \equiv_M ya.$$

- 2 It *refines* A : for all $x, y \in \Sigma^*$,

$$x \equiv_M y \implies (x \in A \Leftrightarrow y \in A).$$

Properties of \equiv_M

Besides the reflexive, symmetric, and transitive properties, \equiv_M satisfies a few other useful properties:

- 1 It is a *right congruence*: for all $x, y \in \Sigma^*$ and $a \in \Sigma$,

$$x \equiv_M y \implies xa \equiv_M ya.$$

- 2 It *refines* A : for all $x, y \in \Sigma^*$,

$$x \equiv_M y \implies (x \in A \Leftrightarrow y \in A).$$

- 3 It is of *finite index*: it has only finitely many equivalence classes.

Myhill-Nerode Relations

Definition. Let $A \subseteq \Sigma^*$. An equivalence relation \equiv on Σ^* is called a *Myhill-Nerode relation* for A if it has the three properties from the previous slide: that is, it is a right congruence of finite index that refines A .

Myhill-Nerode Relations

Definition. Let $A \subseteq \Sigma^*$. An equivalence relation \equiv on Σ^* is called a *Myhill-Nerode relation* for A if it has the three properties from the previous slide: that is, it is a right congruence of finite index that refines A .

- We saw that every DFA M for A yields a Myhill-Nerode relation \equiv_M for A .

Myhill-Nerode Relations

Definition. Let $A \subseteq \Sigma^*$. An equivalence relation \equiv on Σ^* is called a *Myhill-Nerode relation* for A if it has the three properties from the previous slide: that is, it is a right congruence of finite index that refines A .

- We saw that every DFA M for A yields a Myhill-Nerode relation \equiv_M for A .
- We will now show that every Myhill-Nerode relation \equiv for A yields a DFA M_{\equiv} for A .

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Because \equiv is a right congruence, δ is well defined.

From Myhill-Nerode Relation \equiv to the DFA M_{\equiv}

Let $A \subseteq \Sigma^*$ and let \equiv be an arbitrary Myhill-Nerode relation for A .

The \equiv -class of a string $x \in \Sigma^*$ is

$$[x] = \{y \mid y \equiv x\}.$$

There are finitely many \equiv -classes since \equiv has finite index.

We define the DFA $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Because \equiv is a right congruence, δ is well defined.

Since \equiv refines A , we have $x \in A \Leftrightarrow [x] \in F$ for all $x \in \Sigma^*$.

Lemma

For all $x, y \in \Sigma^$, $\delta^*([x], y) = [xy]$.*

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Inductive hypothesis. Suppose $\delta^*([x], y) = [xy]$ holds for some string $y \in \Sigma^*$.

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Inductive hypothesis. Suppose $\delta^*([x], y) = [xy]$ holds for some string $y \in \Sigma^*$.

Inductive step. We must show $\delta^*([x], ya) = [xya]$ for all $a \in \Sigma$.

$$\delta^*([x], ya)$$

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Inductive hypothesis. Suppose $\delta^*([x], y) = [xy]$ holds for some string $y \in \Sigma^*$.

Inductive step. We must show $\delta^*([x], ya) = [xya]$ for all $a \in \Sigma$.

$$\delta^*([x], ya) = \delta(\delta^*([x], y), a) \quad \text{definition of } \delta^*$$

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Inductive hypothesis. Suppose $\delta^*([x], y) = [xy]$ holds for some string $y \in \Sigma^*$.

Inductive step. We must show $\delta^*([x], ya) = [xya]$ for all $a \in \Sigma$.

$$\begin{aligned}\delta^*([x], ya) &= \delta(\delta^*([x], y), a) && \text{definition of } \delta^* \\ &= \delta([xy], a) && \text{inductive hypothesis}\end{aligned}$$

Lemma

For all $x, y \in \Sigma^*$, $\delta^*([x], y) = [xy]$.

Proof. By induction on y .

Base case. $y = \epsilon$:

$$\delta^*([x], \epsilon) = [x] = [x\epsilon].$$

Inductive hypothesis. Suppose $\delta^*([x], y) = [xy]$ holds for some string $y \in \Sigma^*$.

Inductive step. We must show $\delta^*([x], ya) = [xya]$ for all $a \in \Sigma$.

$$\begin{aligned}\delta^*([x], ya) &= \delta(\delta^*([x], y), a) && \text{definition of } \delta^* \\ &= \delta([xy], a) && \text{inductive hypothesis} \\ &= [xya] && \text{definition of } \delta\end{aligned}$$

□

Recall \equiv is a Myhill-Nerode relation for A and $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Theorem

$$L(M_{\equiv}) = A.$$

Recall \equiv is a Myhill-Nerode relation for A and $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Theorem

$$L(M_{\equiv}) = A.$$

Proof. For any $x \in \Sigma^*$,

$$x \in L(M_{\equiv}) \iff \delta^*([\epsilon], x) \in F \quad \text{definition of acceptance}$$

Recall \equiv is a Myhill-Nerode relation for A and $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Theorem

$$L(M_{\equiv}) = A.$$

Proof. For any $x \in \Sigma^*$,

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \delta^*([\epsilon], x) \in F && \text{definition of acceptance} \\ &\iff [x] \in F && \text{Lemma} \end{aligned}$$

Recall \equiv is a Myhill-Nerode relation for A and $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Theorem

$$L(M_{\equiv}) = A.$$

Proof. For any $x \in \Sigma^*$,

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \delta^*([\epsilon], x) \in F && \text{definition of acceptance} \\ &\iff [x] \in F && \text{Lemma} \\ &\iff x \in A && \equiv \text{refines } A \end{aligned}$$

Recall \equiv is a Myhill-Nerode relation for A and $M_{\equiv} = (Q, \Sigma, \delta, q_0, F)$ where

- $Q = \{[x] \mid x \in \Sigma^*\}$
- $q_0 = [\epsilon]$
- $F = \{[x] \mid x \in A\}$
- $\delta([x], a) = [xa]$ for all $x \in \Sigma^*$ and $a \in \Sigma$

Theorem

$$L(M_{\equiv}) = A.$$

Proof. For any $x \in \Sigma^*$,

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \delta^*([\epsilon], x) \in F && \text{definition of acceptance} \\ &\iff [x] \in F && \text{Lemma} \\ &\iff x \in A && \equiv \text{refines } A \end{aligned}$$

Therefore $L(M_{\equiv}) = A$. \square

The following lemma states the mappings $M \rightarrow \equiv_M$ and $\equiv \rightarrow M_{\equiv}$ are inverses of each other, up to isomorphism of automaton.

Lemma

The following lemma states the mappings $M \rightarrow \equiv_M$ and $\equiv \rightarrow M_\equiv$ are inverses of each other, up to isomorphism of automaton.

Lemma

- 1 If \equiv is a Myhill-Nerode relation for A , and if we apply the construction $\equiv \rightarrow M_\equiv$ and then apply the construction $M_\equiv \rightarrow \equiv_{M_\equiv}$, the relation \equiv_{M_\equiv} is identical to \equiv .

$$\equiv \rightarrow M_\equiv \rightarrow \equiv_{M_\equiv} = \equiv$$

The following lemma states the mappings $M \rightarrow \equiv_M$ and $\equiv \rightarrow M_\equiv$ are inverses of each other, up to isomorphism of automaton.

Lemma

- 1 If \equiv is a Myhill-Nerode relation for A , and if we apply the construction $\equiv \rightarrow M_\equiv$ and then apply the construction $M_\equiv \rightarrow \equiv_{M_\equiv}$, the relation \equiv_{M_\equiv} is identical to \equiv .

$$\equiv \rightarrow M_\equiv \rightarrow \equiv_{M_\equiv} = \equiv$$

- 2 If M is a DFA for A with no inaccessible states, and if we apply the construction $M \rightarrow \equiv_M$ and then apply the construction $\equiv_M \rightarrow M_{\equiv_M}$, the resulting DFA M_{\equiv_M} is isomorphic to M .

$$M \rightarrow \equiv_M \rightarrow M_{\equiv_M} \cong M$$

Refining Relations

A relation \equiv_1 *refines* a relation \equiv_2 if

$$x \equiv_1 y \Rightarrow x \equiv_2 y.$$

Refining Relations

A relation \equiv_1 *refines* a relation \equiv_2 if

$$x \equiv_1 y \Rightarrow x \equiv_2 y.$$

- If \equiv_1 and \equiv_2 are equivalence relations, this says that all of \equiv_1 's equivalence classes are contained in equivalence classes of \equiv_2 .

Refining Relations

A relation \equiv_1 *refines* a relation \equiv_2 if

$$x \equiv_1 y \Rightarrow x \equiv_2 y.$$

- If \equiv_1 and \equiv_2 are equivalence relations, this says that all of \equiv_1 's equivalence classes are contained in equivalence classes of \equiv_2 .
- If \equiv_1 refines \equiv_2 , then we say \equiv_2 is *coarser* than \equiv_1 .

Let $A \subseteq \Sigma^*$. We define an equivalence relation \equiv_A on Σ^* by

$$x \equiv_A y \iff (\forall z \in \Sigma^*) (xz \in A \iff yz \in A)$$

for all $x, y \in \Sigma^*$.

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof. For any $x, y \in \Sigma^*$, we have

$$x \equiv_A y \Rightarrow (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$$

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof. For any $x, y \in \Sigma^*$, we have

$$\begin{aligned} x \equiv_A y &\Rightarrow (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A) \\ &\Rightarrow (\forall a \in \Sigma) (\forall w \in \Sigma^*) (xaw \in A \Leftrightarrow yaw \in A) \end{aligned}$$

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof. For any $x, y \in \Sigma^*$, we have

$$\begin{aligned} x \equiv_A y &\Rightarrow (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A) \\ &\Rightarrow (\forall a \in \Sigma) (\forall w \in \Sigma^*) (xaw \in A \Leftrightarrow yaw \in A) \\ &\Rightarrow (\forall a \in \Sigma) xa \equiv_A ya, \end{aligned}$$

so \equiv_A is a right congruence.

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof. For any $x, y \in \Sigma^*$, we have

$$\begin{aligned} x \equiv_A y &\Rightarrow (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A) \\ &\Rightarrow (\forall a \in \Sigma) (\forall w \in \Sigma^*) (xaw \in A \Leftrightarrow yaw \in A) \\ &\Rightarrow (\forall a \in \Sigma) xa \equiv_A ya, \end{aligned}$$

so \equiv_A is a right congruence.

To see that \equiv_A refines A , use $z = \epsilon$ in the definition of \equiv_A :

$$x \equiv_A y \Rightarrow (x \in A \Leftrightarrow y \in A).$$

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof continued. Let \equiv be any equivalence relation on Σ^* that is a right congruence refining A . We have

$$x \equiv y \Rightarrow (\forall z) \, xz \equiv yz \qquad (\equiv \text{ is a right congruence})$$

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof continued. Let \equiv be any equivalence relation on Σ^* that is a right congruence refining A . We have

$$\begin{aligned} x \equiv y &\Rightarrow (\forall z) \, xz \equiv yz && (\equiv \text{ is a right congruence}) \\ &\Rightarrow (\forall z) \, (xz \in A \Leftrightarrow yz \in A) && (\equiv \text{ refines } A) \end{aligned}$$

Lemma

Let $A \subseteq \Sigma^$. The relation \equiv_A is a right congruence refining A and is the coarsest such relation on Σ^* .*

Proof continued. Let \equiv be any equivalence relation on Σ^* that is a right congruence refining A . We have

$$\begin{aligned} x \equiv y &\Rightarrow (\forall z) xz \equiv yz && (\equiv \text{ is a right congruence}) \\ &\Rightarrow (\forall z) (xz \in A \Leftrightarrow yz \in A) && (\equiv \text{ refines } A) \\ &\Rightarrow x \equiv_A y, && (\text{definition of } \equiv_A) \end{aligned}$$

so \equiv_A is coarser than \equiv . \square

Myhill-Nerode Theorem

Let $A \subseteq \Sigma^*$. The following are equivalent:

- 1 A is regular.
- 2 There exists a Myhill-Nerode relation for A .
- 3 The relation \equiv_A is of finite index.

Myhill-Nerode Theorem

Let $A \subseteq \Sigma^*$. The following are equivalent:

- 1 A is regular.
- 2 There exists a Myhill-Nerode relation for A .
- 3 The relation \equiv_A is of finite index.

Proof. 1 \Rightarrow 2: Given a DFA M for A , the construction $M \rightarrow \equiv_M$ yields a Myhill-Nerode relation for A .

Myhill-Nerode Theorem

Let $A \subseteq \Sigma^*$. The following are equivalent:

- 1 A is regular.
- 2 There exists a Myhill-Nerode relation for A .
- 3 The relation \equiv_A is of finite index.

Proof. 1 \Rightarrow 2: Given a DFA M for A , the construction $M \rightarrow \equiv_M$ yields a Myhill-Nerode relation for A .

2 \Rightarrow 3: Let \equiv be a Myhill-Nerode relation for A . Then \equiv is a right congruence that refines A , so \equiv_A is coarser than \equiv by the Lemma. Since \equiv is of finite index, this implies \equiv_A is also of finite index.

Myhill-Nerode Theorem

Let $A \subseteq \Sigma^*$. The following are equivalent:

- 1 A is regular.
- 2 There exists a Myhill-Nerode relation for A .
- 3 The relation \equiv_A is of finite index.

Proof. 1 \Rightarrow 2: Given a DFA M for A , the construction $M \rightarrow \equiv_M$ yields a Myhill-Nerode relation for A .

2 \Rightarrow 3: Let \equiv be a Myhill-Nerode relation for A . Then \equiv is a right congruence that refines A , so \equiv_A is coarser than \equiv by the Lemma. Since \equiv is of finite index, this implies \equiv_A is also of finite index.

3 \Rightarrow 1: If \equiv_A is of finite index, then it is a Myhill-Nerode relation for A by the Lemma. The construction $\equiv_A \rightarrow M_{\equiv_A}$ produces a DFA for A . \square

Corollary

For any regular language A , the DFA M_{\equiv_A} is the minimal DFA for A .

Proof. If A is regular, then \equiv_A is a Myhill-Nerode relation for A . Also, it is the coarsest such relation: it has the fewest equivalence classes of any Myhill-Nerode relation for A . Therefore M_{\equiv_A} has the fewest states in a DFA for A . \square

Applications

We can use the Myhill-Nerode theorem to prove nonregularity.

Example. $A = \{0^n 1^n \mid n \geq 0\}$ is not regular.

Applications

We can use the Myhill-Nerode theorem to prove nonregularity.

Example. $A = \{0^n 1^n \mid n \geq 0\}$ is not regular.

Proof. If $k \neq m$, then $0^k \not\equiv_A 0^m$ since

$$0^k 1^k \in A \text{ and } 0^m 1^k \notin A.$$

Applications

We can use the Myhill-Nerode theorem to prove nonregularity.

Example. $A = \{0^n 1^n \mid n \geq 0\}$ is not regular.

Proof. If $k \neq m$, then $0^k \not\equiv_A 0^m$ since

$$0^k 1^k \in A \text{ and } 0^m 1^k \notin A.$$

Therefore \equiv_A has infinitely many equivalence classes, at least one for each 0^k , $k \geq 0$. By the Myhill-Nerode Theorem, A is not regular. \square

Example. $B = \{0^{n^2} \mid n \geq 0\}$ is not regular.

Applications

Example. $B = \{0^{n^2} \mid n \geq 0\}$ is not regular.

Proof. For any $0 \leq k < j$, $0^{k^2} \not\equiv_B 0^{j^2}$ since

$$0^{k^2}0^{2k+1} \in B \text{ and } 0^{j^2}0^{2k+1} \notin B.$$

- $0^{k^2}0^{2k+1} \in B$ because $k^2 + 2k + 1 = (k + 1)^2$.

Applications

Example. $B = \{0^{n^2} \mid n \geq 0\}$ is not regular.

Proof. For any $0 \leq k < j$, $0^{k^2} \not\equiv_B 0^{j^2}$ since

$$0^{k^2}0^{2k+1} \in B \text{ and } 0^{j^2}0^{2k+1} \notin B.$$

- $0^{k^2}0^{2k+1} \in B$ because $k^2 + 2k + 1 = (k + 1)^2$.
- $0^{j^2}0^{2k+1} \notin B$ because $j^2 < j^2 + 2k + 1 < j^2 + 2j + 1 = (j + 1)^2$ implies $j^2 + 2k + 1$ is not a square.

Applications

Example. $B = \{0^{n^2} \mid n \geq 0\}$ is not regular.

Proof. For any $0 \leq k < j$, $0^{k^2} \not\equiv_B 0^{j^2}$ since

$$0^{k^2}0^{2k+1} \in B \text{ and } 0^{j^2}0^{2k+1} \notin B.$$

- $0^{k^2}0^{2k+1} \in B$ because $k^2 + 2k + 1 = (k + 1)^2$.
- $0^{j^2}0^{2k+1} \notin B$ because $j^2 < j^2 + 2k + 1 < j^2 + 2j + 1 = (j + 1)^2$ implies $j^2 + 2k + 1$ is not a square.

Therefore \equiv_B has infinitely many equivalence classes, so B is not regular by the Myhill-Nerode Theorem. \square

Applications

Example. $F = \{a^m b^n c^l \mid m, n, l \geq 0 \text{ and if } m = 1 \text{ then } n = l\}$

Recall that we showed F satisfies the conclusion of the Pumping Lemma. We can use the Myhill-Nerode Theorem to prove F is not regular.

Applications

Example. $F = \{a^m b^n c^l \mid m, n, l \geq 0 \text{ and if } m = 1 \text{ then } n = l\}$

Recall that we showed F satisfies the conclusion of the Pumping Lemma. We can use the Myhill-Nerode Theorem to prove F is not regular.

Proof. For all $i < j$, $ab^i \not\equiv_F ab^j$ since

$$ab^i c^i \in F \text{ and } ab^j c^i \notin F.$$

Therefore \equiv_F has infinitely many equivalence classes, so F is not regular by the Myhill-Nerode Theorem. \square

Let A be a regular language and let $M = (Q, \Sigma, \delta, q_0, F)$ be the minimal DFA for A .

Recall that for every $x, y \in \Sigma^*$,

$$x \equiv_A y \text{ if } (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$$

and

$$x \equiv_M y \text{ if } \delta^*(q_0, x) = \delta^*(q_0, y).$$

Let A be a regular language and let $M = (Q, \Sigma, \delta, q_0, F)$ be the minimal DFA for A .

Recall that for every $x, y \in \Sigma^*$,

$$x \equiv_A y \text{ if } (\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$$

and

$$x \equiv_M y \text{ if } \delta^*(q_0, x) = \delta^*(q_0, y).$$

From the proof of the Myhill-Nerode Theorem, we know that \equiv_A and \equiv_M are the same relation. In other words, the states of M “remember” the equivalence of x with respect to \equiv_A :

$$\delta^*(q_0, x) \text{ is essentially } [x]_A.$$

Let $A = \{x00 \mid x \in \{0, 1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

Let $A = \{x00 \mid x \in \{0, 1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1 \quad \epsilon \equiv_A 11 \quad \epsilon \equiv_A 01$

Let $A = \{x00 \mid x \in \{0, 1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

Let $A = \{x00 \mid x \in \{0,1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0,1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0,1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

Let $A = \{x00 \mid x \in \{0,1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0,1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0,1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

② What is \equiv_A -equivalent to 0?

- $0 \equiv_A 10$ $0 \equiv_A 1010$ $0 \equiv_A 0010$

Let $A = \{x00 \mid x \in \{0,1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0,1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0,1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

② What is \equiv_A -equivalent to 0 ?

- $0 \equiv_A 10$ $0 \equiv_A 1010$ $0 \equiv_A 0010$
- $0 \not\equiv_A 100$ ($0 \notin A$ but $100 \in A$)
- $0 \not\equiv_A 1$ ($00 \in A$ but $10 \notin A$)

$$[0]_A = \{x \in \{0,1\}^* \mid x \text{ ends in exactly one } 0\}$$

Let $A = \{x00 \mid x \in \{0,1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0,1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0,1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

② What is \equiv_A -equivalent to 0?

- $0 \equiv_A 10$ $0 \equiv_A 1010$ $0 \equiv_A 0010$
- $0 \not\equiv_A 100$ ($0 \notin A$ but $100 \in A$)
- $0 \not\equiv_A 1$ ($00 \in A$ but $10 \notin A$)

$$[0]_A = \{x \in \{0,1\}^* \mid x \text{ ends in exactly one } 0\}$$

③ What is \equiv_A -equivalent to 00?

- $00 \equiv_A 100$ $00 \equiv_A 10100$ $00 \equiv_A 10000$

Let $A = \{x00 \mid x \in \{0, 1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0, 1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0, 1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

② What is \equiv_A -equivalent to 0?

- $0 \equiv_A 10$ $0 \equiv_A 1010$ $0 \equiv_A 0010$
- $0 \not\equiv_A 100$ ($0 \notin A$ but $100 \in A$)
- $0 \not\equiv_A 1$ ($00 \in A$ but $10 \notin A$)

$$[0]_A = \{x \in \{0, 1\}^* \mid x \text{ ends in exactly one } 0\}$$

③ What is \equiv_A -equivalent to 00?

- $00 \equiv_A 100$ $00 \equiv_A 10100$ $00 \equiv_A 10000$
- $00 \not\equiv_A 10$ ($00 \in A$ but $10 \notin A$)
- $00 \not\equiv_A 001$ ($00 \in A$ but $001 \notin A$)

Let $A = \{x00 \mid x \in \{0,1\}^*\}$.

Recall $x \equiv_A y$ if $(\forall z \in \Sigma^*) (xz \in A \Leftrightarrow yz \in A)$.

① What is \equiv_A -equivalent to ϵ ?

- $\epsilon \equiv_A 1$ $\epsilon \equiv_A 11$ $\epsilon \equiv_A 01$
- $\epsilon \not\equiv_A 0$ ($\epsilon 0 \notin A$ but $00 \in A$)
- $\epsilon \not\equiv_A 00$ ($\epsilon \notin A$ but $00 \in A$)

$$\begin{aligned} [\epsilon]_A &= \{\epsilon\} \cup \{x \in \{0,1\}^* \mid x \text{ ends in } 1\} \\ &= \{x \in \{0,1\}^* \mid x \text{ does not end in } 0\} \end{aligned}$$

② What is \equiv_A -equivalent to 0?

- $0 \equiv_A 10$ $0 \equiv_A 1010$ $0 \equiv_A 0010$
- $0 \not\equiv_A 100$ ($0 \notin A$ but $100 \in A$)
- $0 \not\equiv_A 1$ ($00 \in A$ but $10 \notin A$)

$$[0]_A = \{x \in \{0,1\}^* \mid x \text{ ends in exactly one } 0\}$$


③ What is \equiv_A -equivalent to 00?

- $00 \equiv_A 100$ $00 \equiv_A 10100$ $00 \equiv_A 10000$
- $00 \not\equiv_A 10$ ($00 \in A$ but $10 \notin A$)
- $00 \not\equiv_A 001$ ($00 \in A$ but $001 \notin A$)

$$[00]_A = \{x \in \{0,1\}^* \mid x \text{ ends in two or more } 0\text{'s}\}$$

Example

$$A = \{x00 \mid x \in \{0, 1\}^*\}$$



Equivalence classes for \equiv_M and \equiv_A :

$[\epsilon]_M = \{x \mid \delta^*(q_0, x) = q_0\} = \epsilon \cup (0 \cup 1)^*1$	$= \left\{ x \mid \begin{array}{l} x \text{ does not} \\ \text{end in } 0 \end{array} \right\}$	$= [\epsilon]_A$
$[0]_M = \{x \mid \delta^*(q_0, x) = q_1\} = 0 \cup (0 \cup 1)^*10$	$= \left\{ x \mid \begin{array}{l} x \text{ ends in} \\ \text{exactly one } 0 \end{array} \right\}$	$= [0]_A$
$[00]_M = \{x \mid \delta^*(q_0, x) = q_2\} = (0 \cup 1)^*00$	$= \left\{ x \mid \begin{array}{l} x \text{ ends in at} \\ \text{least two } 0\text{'s} \end{array} \right\}$	$= [00]_A$

$$A = [00]_A \text{ and } A^c = [\epsilon]_A \cup [0]_A$$