

Computability and Complexity

COSC 4200

2SAT is in P

Theorem

2SAT \in P.

Proof. Given a 2CNF formula ϕ , we construct a directed graph G as follows.


- The vertices of G are the variables of ϕ and their negations.
- There is an edge from vertex α to vertex β if there is a clause $(\neg\alpha \vee \beta)$ or $(\beta \vee \neg\alpha)$ in ϕ .

In other words, we put an edge from α to β if the implication $\alpha \rightarrow \beta$ is represented in ϕ .

For an example, for the formula

$$\phi = (x_1 \vee x_2) \wedge (x_1 \vee \neg x_3) \wedge (\neg x_1 \vee x_2) \wedge (x_2 \vee x_3)$$

we construct the following graph:



There is a nice symmetry in the graph, in that if there is a path from α to β , there is also a path from $\neg\beta$ to $\neg\alpha$.

Claim

ϕ is unsatisfiable if and only if there is a variable x such that there are paths from x to $\neg x$ and from $\neg x$ to x in G .

Proof of Claim. Suppose that for some variable x , there is a path from x to $\neg x$ and from $\neg x$ to x . Suppose also that ϕ is satisfiable by some assignment τ .

Proof of Claim. Suppose that for some variable x , there is a path from x to $\neg x$ and from $\neg x$ to x . Suppose also that ϕ is satisfiable by some assignment τ . There are two cases:

Case 1: $\tau(x) = T$. Since there is a path in G from x to $\neg x$, and $\neg x = F$ under τ , there must be some edge (α, β) along the path with $\tau(\alpha) = T$ but $\tau(\beta) = F$. But then $(\neg\alpha \vee \beta)$ is a clause in ϕ and it is not satisfied by τ , a contradiction.

Proof of Claim. Suppose that for some variable x , there is a path from x to $\neg x$ and from $\neg x$ to x . Suppose also that ϕ is satisfiable by some assignment τ . There are two cases:

Case 1: $\tau(x) = T$. Since there is a path in G from x to $\neg x$, and $\neg x = F$ under τ , there must be some edge (α, β) along the path with $\tau(\alpha) = T$ but $\tau(\beta) = F$. But then $(\neg\alpha \vee \beta)$ is a clause in ϕ and it is not satisfied by τ , a contradiction.

Case 2: $\tau(x) = F$. Since there is a path in G from $\neg x$ to x and $x = F$ under τ , there must be some edge (α, β) along the path with $\tau(\alpha) = T$ but $\tau(\beta) = F$. But then $(\neg\alpha \vee \beta)$ is a clause in ϕ and it is not satisfied by τ , a contradiction.

Proof of Claim. Suppose that for some variable x , there is a path from x to $\neg x$ and from $\neg x$ to x . Suppose also that ϕ is satisfiable by some assignment τ . There are two cases:

Case 1: $\tau(x) = T$. Since there is a path in G from x to $\neg x$, and $\neg x = F$ under τ , there must be some edge (α, β) along the path with $\tau(\alpha) = T$ but $\tau(\beta) = F$. But then $(\neg\alpha \vee \beta)$ is a clause in ϕ and it is not satisfied by τ , a contradiction.

Case 2: $\tau(x) = F$. Since there is a path in G from $\neg x$ to x and $x = F$ under τ , there must be some edge (α, β) along the path with $\tau(\alpha) = T$ but $\tau(\beta) = F$. But then $(\neg\alpha \vee \beta)$ is a clause in ϕ and it is not satisfied by τ , a contradiction.

In either case we have a contradiction, so ϕ must be unsatisfiable.

Now assume that for all x , there is no path from x to $\neg x$ and from $\neg x$ to x .

We want to show that ϕ is satisfiable.

Now assume that for all x , there is no path from x to $\neg x$ and from $\neg x$ to x .

We want to show that ϕ is satisfiable. To do so, we repeat the following until all vertices have a T/F assignment:

- Pick a vertex α in G whose truth value has not been defined, such that there is no path from α to $\neg\alpha$. We know such a vertex exists by our assumption. Initially we can take any variable and use either it or its negation.

Now assume that for all x , there is no path from x to $\neg x$ and from $\neg x$ to x .

We want to show that ϕ is satisfiable. To do so, we repeat the following until all vertices have a T/F assignment:

- Pick a vertex α in G whose truth value has not been defined, such that there is no path from α to $\neg\alpha$. We know such a vertex exists by our assumption. Initially we can take any variable and use either it or its negation.
- Assign α , and all vertices reachable from α , the value T .

Now assume that for all x , there is no path from x to $\neg x$ and from $\neg x$ to x .

We want to show that ϕ is satisfiable. To do so, we repeat the following until all vertices have a T/F assignment:

- Pick a vertex α in G whose truth value has not been defined, such that there is no path from α to $\neg\alpha$. We know such a vertex exists by our assumption. Initially we can take any variable and use either it or its negation.
- Assign α , and all vertices reachable from α , the value T .
- Assign the negation of all those vertices the value F . In other words, assign $\neg\alpha$, and all vertices from which $\neg\alpha$ can be reached, the value F .

We observe that we can always assign all vertices reachable from α the value of T . If a vertex β can be reached from α and it were already assigned F , then by our algorithm, we would have already assigned α F because β can be reached from it.

We observe that we can always assign all vertices reachable from α the value of T . If a vertex β can be reached from α and it were already assigned F , then by our algorithm, we would have already assigned α F because β can be reached from it.

Also, the negation of every node reachable from α can be assigned F . If both β and $\neg\beta$ are reachable from α , then by symmetry, there are paths to $\neg\alpha$ from both β and $\neg\beta$. This means there is a path from α to $\neg\alpha$, which contradicts our assumption.

Since for all x , there is no path from x to $\neg x$ or there is no path from $\neg x$ to x , all vertices will eventually be assigned a truth value. For every edge (α, β) in the graph, we have either

- $(\alpha = T) \rightarrow (\beta = T)$,
- $(\alpha = F) \rightarrow (\beta = F)$, or
- $(\alpha = F) \rightarrow (\beta = T)$.

This is because the construction ensures the false implication, $(\alpha = T) \rightarrow (\beta = F)$, never occurs. Therefore all clauses of ϕ are satisfied by this assignment. □ Claim

Now we can describe an algorithm for 2SAT.

- We input ϕ and construct the graph G .
- For each variable x in ϕ , check to see if there is a path from x to $\neg x$ and a path from $\neg x$ to x .
- If there is such a set of paths for any variable, we reject. Otherwise, we accept.

Since reachability can be computed in polynomial time (e.g. using BFS, DFS, etc.), we can solve 2SAT in polynomial time. \square

Theorem

$2SAT \in P$.

In fact, a stronger result is true:

Theorem

$2SAT \in NL$.

Theorem

$2SAT \in P$.

In fact, a stronger result is true:

Theorem

$2SAT \in NL$.

Proof. The following algorithm shows $2SAT^c \in NL$.

- We input ϕ , guess a variable x .
- Guess paths from x to $\neg x$ and from $\neg x$ to x .
- If both paths are found, accept.
- Otherwise, reject.

Theorem

$2SAT \in P$.

In fact, a stronger result is true:

Theorem

$2SAT \in NL$.

Proof. The following algorithm shows $2SAT^c \in NL$.

- We input ϕ , guess a variable x .
- Guess paths from x to $\neg x$ and from $\neg x$ to x .
- If both paths are found, accept.
- Otherwise, reject.

This algorithm is correct by the analysis in the previous proof.

Thus $2SAT \in coNL$. Since $NL = coNL$, we have $2SAT \in NL$. \square

Theorem

2SAT is NL-complete.

Proof. We will show $\text{PATH}^c \leq_L \text{2SAT}$. This will establish that $A \leq_L \text{2SAT}$ for every $A \in \text{coNL}$. Since $\text{NL} = \text{coNL}$, it follows that 2SAT is NL-complete.

Theorem

2SAT is NL-complete.

Proof. We will show $\text{PATH}^c \leq_L 2\text{SAT}$. This will establish that $A \leq_L 2\text{SAT}$ for every $A \in \text{coNL}$. Since $\text{NL} = \text{coNL}$, it follows that 2SAT is NL-complete.

Let $G = (V, E)$, s , and t be an instance of PATH^c . Our formula ϕ will have V as variables and have $|E| + 2$ clauses: (s) , $(\neg t)$, and $(\neg u \vee v)$ for each $(u, v) \in E$.

$$\phi = (s) \wedge (\neg t) \wedge \bigwedge_{(u,v) \in E} (\neg u \vee v)$$

Note that ϕ is computable by a log-space transducer.

Suppose there is a path from s to t :

$$s \rightarrow u_1 \rightarrow u_2 \cdots u_k \rightarrow t$$

Suppose there is a path from s to t :

$$s \rightarrow u_1 \rightarrow u_2 \cdots u_k \rightarrow t$$

Then

$$(\neg s \vee u_1), (\neg u_1 \vee u_2), \dots, (\neg u_{k-1} \vee u_k), (\neg u_k \vee t)$$

are all clauses in ϕ .

Suppose there is a path from s to t :

$$s \rightarrow u_1 \rightarrow u_2 \cdots u_k \rightarrow t$$

Then

$$(\neg s \vee u_1), (\neg u_1 \vee u_2), \dots, (\neg u_{k-1} \vee u_k), (\neg u_k \vee t)$$

are all clauses in ϕ .

If these clauses are all satisfied by some assignment τ , then $(\neg s \vee t)$ is satisfied by τ as well.

Suppose there is a path from s to t :

$$s \rightarrow u_1 \rightarrow u_2 \cdots u_k \rightarrow t$$

Then

$$(\neg s \vee u_1), (\neg u_1 \vee u_2), \dots, (\neg u_{k-1} \vee u_k), (\neg u_k \vee t)$$

are all clauses in ϕ .

If these clauses are all satisfied by some assignment τ , then $(\neg s \vee t)$ is satisfied by τ as well.

Thus τ does not satisfy ϕ since it has clauses (s) and $(\neg t)$.
Therefore ϕ is unsatisfiable.

Suppose there is a path from s to t :

$$s \rightarrow u_1 \rightarrow u_2 \cdots u_k \rightarrow t$$

Then

$$(\neg s \vee u_1), (\neg u_1 \vee u_2), \dots, (\neg u_{k-1} \vee u_k), (\neg u_k \vee t)$$

are all clauses in ϕ .

If these clauses are all satisfied by some assignment τ , then $(\neg s \vee t)$ is satisfied by τ as well.

Thus τ does not satisfy ϕ since it has clauses (s) and $(\neg t)$.
Therefore ϕ is unsatisfiable.

This shows $\langle G, s, t \rangle \in \text{PATH} \Rightarrow \phi \notin \text{2SAT}$.

Now suppose there is no path from s to t . Let

$$S = \{v \in V \mid v \text{ is reachable from } s\}$$

$$T = \{v \in V \mid t \text{ is reachable from } v\}$$

$$M = V - (S \cup T)$$

Then S , T , and M are disjoint. There are no edges from S to $M \cup T$ and there are no edges from $S \cup M$ to T .

Now suppose there is no path from s to t . Let

$$S = \{v \in V \mid v \text{ is reachable from } s\}$$

$$T = \{v \in V \mid t \text{ is reachable from } v\}$$

$$M = V - (S \cup T)$$

Then S , T , and M are disjoint. There are no edges from S to $M \cup T$ and there are no edges from $S \cup M$ to T .

Define an assignment τ by

$$\tau(v) = \begin{cases} T & \text{if } v \in S \\ T & \text{if } v \in M \\ F & \text{if } v \in T \end{cases}$$

This assignment satisfies ϕ because it sets $\tau(s) = T$, $\tau(t) = F$, and for any edge (u, v) the clause $(\neg u \vee v)$ is satisfied because $\tau(u) = \tau(v)$.

Now suppose there is no path from s to t . Let

$$S = \{v \in V \mid v \text{ is reachable from } s\}$$

$$T = \{v \in V \mid t \text{ is reachable from } v\}$$

$$M = V - (S \cup T)$$

Then S , T , and M are disjoint. There are no edges from S to $M \cup T$ and there are no edges from $S \cup M$ to T .

Define an assignment τ by

$$\tau(v) = \begin{cases} T & \text{if } v \in S \\ T & \text{if } v \in M \\ F & \text{if } v \in T \end{cases}$$

This assignment satisfies ϕ because it sets $\tau(s) = T$, $\tau(t) = F$, and for any edge (u, v) the clause $(\neg u \vee v)$ is satisfied because $\tau(u) = \tau(v)$.

This proves $\langle G, s, t \rangle \notin \text{PATH} \Rightarrow \phi \in 2\text{SAT}$.

We have shown

$$\langle G, s, t \rangle \in \text{PATH} \Leftrightarrow \phi \notin 2\text{SAT}.$$

Equivalently,

$$\langle G, s, t \rangle \in \text{PATH}^c \Leftrightarrow \phi \in 2\text{SAT}.$$

Thus $\text{PATH}^c \leq_L 2\text{SAT}$.

□