

Nov 2023

S.E. Sem III (E-2019) "C Scheme" AI & DS

21/11/2023

(Time: 3 Hours)

Max. Marks: 80

- N.B. (1) Question No. 1 is compulsory.  
 (2) Answer any three questions from Q.2 to Q.6.  
 (3) Use of Statistical Tables permitted.  
 (4) Figures to the right indicate full marks

- Q1.  
 (a) Find the Laplace transform of  $\frac{\cos 2t \sin t}{e^t}$  [5]  
 (b) Find k such that  $f(z) = \frac{1}{2} \log(x^2 + y^2) + i \tan^{-1} \frac{kx}{y}$  is analytic [5]  
 (c) Calculate the Spearman's rank correlation coefficient R [5]  
 $X : 10, 12, 18, 18, 15, 40.$   
 $Y : 12, 18, 25, 25, 50, 25.$   
 (d) Find the inverse Laplace transform of  $\log\left(\frac{s^2 + a^2}{s^2 + b^2}\right)$ . [5]

- Q2.  
 (a) A continuous random variable has probability density function  
 $f(x) = k(x - x^2), 0 \leq x \leq 1.$   
 $f(x) = 0$  otherwise  
 Find k, mean and variance.

- (b) Find the Laplace transform of  $e^{-3t} \int_0^t u \sin 3u du$ . [6]  
 (c) Obtain the Fourier series to represent  $f(x) = x^2$  in  $(0, 2\pi)$  [6]  
 Hence show that  $\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} \dots$  [8]

- Q3.  
 (a) If the imaginary part of the analytic function  $w = u + iv = f(z)$  is  
 $V = x^2 - y^2 + \frac{x}{x^2 + y^2}$ , then show that  $u = -2xy + \frac{y}{x^2 + y^2}$ . [6]  
 (b) Find inverse Laplace transform of  $\frac{2s^2 - 6s + 5}{(s^3 - 6s^2 + 11s - 6)}$  [6]  
 (c) Fit a second-degree parabolic curve and estimate y when  $x = 10$   
 $x : 1, 2, 3, 4, 5, 6, 7, 8, 9,$   
 $y : 2, 6, 7, 8, 10, 11, 11, 10, 9.$  [8]

- Q4.  
 (a) Obtain the Fourier series to represent  $f(x) = x^3$  in  $(-\pi, \pi)$ . [6]  
 (b) Find (i) the equation of the lines of Regression (ii) coefficient of correlation for the following data  
 $X : 65, 66, 67, 67, 68, 69, 70, 72.$   
 $Y : 67, 68, 65, 66, 72, 72, 69, 71.$  [6]  
 (c) Prove that  $\int_0^\infty e^{-\sqrt{2}t} \frac{\sin t \sinh t}{t} dt = \frac{\pi}{8}.$  [8]

Q5.

(a) Find the orthogonal trajectories of the family of curves  $x^3y - xy^3 = c$ .

[6]

(b) Find the moment generating function of the distribution

X	:	-2	3	1
P (X=x)	:	$\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{6}$

hence find first four central moments.

[6]

(c) Obtain the half range cosine series of  $f(x) = x$  in  $(0, 2)$

Hence show that  $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

[8]

Q6.(a) Using convolution theorem Find the inverse Laplace transform of  $\left[ \frac{s^2}{(s^2+2^2)^2} \right]$

[6]

(b) The probability density function of a random variable X is

X	:	1	2	3	4	5	6	7
P (X=x)	:	k	2k	3k	$k^2$	$k^2 + k$	$2k^2$	$4k^2$

Find k,  $p(X < 5)$ ,  $P(X > 5)$

[6]

(c) If  $v = 3x^2y + 6xy - y^3$ , show that v is harmonic function

And find the corresponding analytic function.

[8]