How Stable Are Regression Conclusions Across Subpopulations?

A Brief for Preprint Review

Research summary

September 2026

Question and motivation

Applied researchers commonly fit one regression to a pooled sample and interpret the resulting coefficients as if they describe every relevant subgroup. This is efficient, but it can be misleading. A pooled coefficient may conceal different subgroup relationships, and a model that performs acceptably on average may not transfer to every group.

This project asks a focused diagnostic question: *when does one pooled linear regression provide a stable summary of relationships observed across predefined subpopulations?* The aim is not to introduce a new estimator or replace stratified regression. Instead, subgroup-specific fits are used to assess whether the pooled model preserves coefficient direction, coefficient magnitude, and predictive performance across groups.

Approach

The study combines a controlled simulation with one empirical illustration. The simulation varies the number of groups, total sample size, group-size imbalance, covariate distributions, and the degree of coefficient heterogeneity. All four scenarios use the same 140-setting grid and 30 replications per setting. Four settings with fewer than three observations in at least one subgroup are excluded from aggregate comparisons, leaving 136 directly comparable settings per scenario.

The analysis compares pooled, group-adjusted, interaction, and separate subgroup regressions. It evaluates directional agreement, standardized coefficient movement, a covariate-context adjustment to that movement, cross-group prediction loss, and covariate-distribution separation. The guiding distinction is between apparent disagreement created by finite samples or different predictor distributions and disagreement that remains when subgroup regression relationships genuinely differ.

Mathematically, if subgroup slopes differ, the pooled population target is a design-weighted compromise:

$$\beta^{\dagger}_{\text{pool}} = \left(\sum_{g=1}^{G} X_g^{\top} X_g \right)^{-1} \sum_{g=1}^{G} X_g^{\top} X_g \beta_g.$$

Even when all β_g are equal, subgroup estimates vary because their precision depends on $X_g^{\top} X_g$. When slopes differ, transporting a model from group a to group b incurs excess target-group risk proportional to

$$(\beta_a - \beta_b)^{\top} \Sigma_b (\beta_a - \beta_b).$$

Main simulation findings

The unified simulation separates three interpretable patterns. Scenario 1 is the finite-sample baseline. Even with no true heterogeneity, splitting observations into many uneven groups increases subgroup coefficient dispersion. Scenario 2 changes predictor distributions while holding slopes fixed. It produces much larger covariate separation

and substantial reduction after context normalization, but little additional prediction loss. Scenario 3 changes slopes while holding predictor distributions broadly comparable. It produces large residual coefficient movement and much worse cross-group prediction. Scenario 4 combines both mechanisms.

Scenario	Drift	Context drift	Prediction gap	Covariate shift	Shrinkage
No heterogeneity	3.81	3.46	0.005	0.064	5.9%
Covariate shift only	3.98	2.63	0.004	0.498	33.1%
Coefficient heterogeneity only	24.12	21.75	0.063	0.058	5.5%
Combined heterogeneity	25.45	16.12	0.065	0.495	32.7%

Relative to the no-heterogeneity baseline, coefficient heterogeneity increases median drift by 6.34 times and the median prediction gap by 12.69 times. The combined scenario has the largest raw drift and prediction gap, and the weakest average sign stability. Its lower context-normalized drift than the coefficient-only case is itself informative: part of its raw movement is associated with covariate separation, but a large residual discrepancy remains.

Empirical illustration and contribution

The empirical analysis uses substantively defined age groups in a health-survey regression. The pooled model masks a predictor whose coefficient changes direction in later age groups. The pattern remains after a sensitivity analysis excluding the smallest group, and group-specific models improve in-group predictive performance across the age strata. This is an illustration, not a causal claim or a universal biomedical conclusion.

The contribution is therefore diagnostic and empirical. The paper provides a structured way to ask whether subgroup regression results justify a single pooled interpretation, and to distinguish three cases: finite-sample disagreement, disagreement associated with covariate context, and persistent relationship heterogeneity. The intended use is as a practical complement to interaction models and stratified regression, not as a substitute for either.

Current limits and request

The work is limited to low-dimensional linear models, a restricted set of simulated mechanisms, one empirical illustration, and primarily in-sample predictive comparisons. The context adjustment is descriptive rather than a formal causal decomposition. In addition, 30 replications support broad comparative patterns but provide only coarse confidence intervals for individual simulation settings. The next technical extensions are higher-replication runs, out-of-sample validation, and additional empirical settings.

The manuscript is now internally consistent, fully compiled, and supported by a unified four-scenario simulation design. I am seeking feedback on whether the framing and evidence are appropriate for posting as a cautious preprint, and whether the contribution is sufficiently clear to support an endorsement.