

Vesper's

BACTOCARE OA +

बैक्टोकैर ओ ए +

Selective Biotechnological Feed Acidifier



Vesper

SINCE 1978

 **VISION** *The Care Takers*
Catalyzes Research
Creates Products
Cares Customers

Organic acids for health and performance. The poultry industry keeps searching for strategies to reduce antibiotic usage while maintaining health and performance of the birds thus ensuring profitability.

Organic acids are used globally by the feed industry for management of gut microbiology improved performance and health. The use of organic acids ensures better nutrients utilization and use of enzymes like phytase, giving better body weights, improved feed conversion and reduced colonization of gut by the pathogens, there by reducing disease and disturbance in the flock.

Poultry Performance profile of organic acids

Acid	Concentration (%)	Effects
Fumaric acid	0.50-1.00	Improvement in weight gain of broilers. Improved feed efficiency in both broilers and layers.
Buffered propionic acid	0.15-0.20	Increase in dressing percentage in female broilers and reduction in abdominal fat for males.
Malic acid	0.50-2.00	Increase in weight gain
Sorbic acid	1.12	Improves feed efficiency
Tartaric acid	0.33	Increase in weight gain
Lactic acid	2.00	Feed to gain ratio significantly improved. Body weight gain.
Formic acid	0.50-1.00	Reduction of caecal pH and bactericidal effect on <i>Salmonella</i>
Benzoic acid	0.20	Positive influence on growth
Butyric acid	0.15	Maintain the beneficial micro flora. Increase the proliferation and maturation of intestinal cells

There is an increased public awareness about the risk of developing cross-resistance of pathogens to antibiotics. Organic acids, such as propionic acid, have been used for more than 30 years to reduce bacterial growth and mould in feedstuffs and thus preserve hygienic quality. Health and performance promoting effects have been demonstrated for a number of organic acids, including formic, fumaric, citric and lactic acid and their salts. In animal rearing, higher feed conversion rates and daily gain as well as reduced incidence of diarrhea, enhanced economic return by lowering feed costs and shorter time to market may be achieved by organic acid supplementation.

The term "organic acid" refers to a broad class of compounds used in fundamental metabolic processes of the body. Chemically, organic acids share the common features of water solubility, acidity, and ninhydrin-negativity (no primary or secondary amines). The term is generally considered to include all carboxylic acids, with or without keto, hydroxyl, or other non-amino functional groups, but does not include most amino acids.

pH, resident time of feed and microbial population in gastrointestinal tract of chicken

Microflora 3 weeks	Parts	Gastro- intestinal tract	pH	Resident time Minutes	Microflora Adult
Streptococci ¹ Coliform ¹ Lactobacilli ¹	Crop		4.5 - 5.3	45	Streptococci ² Coliform ³ Lactobacilli ³
Streptococci ¹ Coliform ¹ Lactobacilli ¹	Proventriculus Gizzard		2.0 - 4.5	70	Streptococci ² Coliform ³ Lactobacilli ³
Streptococci ¹ Coliform ¹ Lactobacilli ¹	Ileum		5.6 - 7.9	160 - 200	Streptococci ¹ Coliform ¹ Lactobacilli ¹
Streptococci ¹ Coliform ¹ Lactobacilli ²	Caeca		5.8 - 6.8	120	Bacteroidesi ¹ Bifidobacteria ¹ Peptostreptococci ¹ Clostridia ¹ Propionic bacteria ¹ Eubacteria ¹
Streptococci ¹ Coliform ¹ Lactobacilli ²	Colon Rectum		6.3 - 7.7	30-50	Mixture: Ileal and caecal bacteria ³

Amino nitrogen and reactive protein stimulate macrophage activation and improve cell-mediated immunity in the birds.

LPO (Lactoperoxidase) is bacteriostic and bactericidal, so it supports probiotics. **Bactocare OA+** provides a maximized comprehensive probiotic activity to competitively inhibit, antagonize and exclude pathogens for improved production and profit. There by reducing dependence on the AGP (Antibiotic Growth Promoters).

Composition

ORGANIC ACIDS (Formic acid, Sorbic acid, Propionic Acid, Acetic acid, Ascorbic acid and Lactic acid) Ammonium propionate, Ammonium Formate along with fatty acids, yeast extract Amino Nitrogen, LPO, reactive Proteins in a fortified base

INDICATIONS

Reduces the gut pH
Reduces Multiplication of the pathogenic gram-ve Microorganisms and fungus
Increase tissue response and improves nutrient utilization
Reduces disease incidences
the best alternative to antibiotic growth promoters
Reduces the duration and cost of treatment.

USAGE INSTRUCTIONS

Regular Use:- 250 gm - 500 gm/ton of feed.

During Stress :- 2 kg/ton of feed.
Or as recommended by the Veterinarian.

Presentation

1 kg x 25 packets in a bag.



Manufactured by

VESPER PHARMACEUTICALS

Singapura, Vidyaranyapura Post,
Bangalore - 560 097 INDIA

Customer Care No. 080 23254277

Email : info@vespergroupvet.com

Website : www.vespergroupvet.com

An ISO 9001-2015 & GMP Certified Company