

Vesperi's

K-Cox Forte

के-कोक्स फ़ोर्ट



Since 1978



Catalyzes Research
Creates Products
Cares Customers
The Care Takers

Vitamin K helps in blood clotting process, coagulation/clotting time will increase in vitamin K deficient birds, they will profusely bleed, will cause shock and death.


Vitamin K is required for syntheses of prothrombin (factor II) plasma clotting factor VII (Proconvertin) christmas factor (IX) and stuart-power factor (X), are all synthesized in the liver as inactive precursors forms, later conjugated to biologically active proteins by the action of Vitamin K to prevent hemorrhage.

Vitamin K starts to work in approx. 6 hours time after its consumption.

The Mechanism of Blood Clotting

INTRINSIC SYSTEM

EXTRINSIC SYSTEM



II-prothrombin
 V-accelerator
 VII-proconvertin
 VII-antihemophilic globulin
 IX-Christmas factor

X-Stuart-prower factor
 XIII-fibrin stabilizing factor
 a-activated
 PL-phospholipid

Adapted from Scott et al., (1982) and Mc Dowell (1989).

NRC (1994) recommends a requirement of Vitamin K ranging between 0.4 to 1.75 mg/kg of diet for poultry.

Subclinical and Clinical Coccidiosis.

Different Stages of coccidia in chickens appear both within the host as well as outside. The developmental stages in the chicken give rise to a microscopic egg (called an oocyst) that is passed out in the droppings. Under proper conditions of temperature and moisture, the oocyst develops within two days, to form a sporulated oocyst, which is capable of infecting other chickens. At this stage, the oocyst contains eight bodies (called sporozoites), each of which is capable of entering a cell in the chicken's intestine after the oocyst is eaten. When sporozoites enter the cells, they divide many times producing either a few or many offspring (merozoites).

The numbers produced depend on the species of coccidia involved. Each merozoite, in turn, may enter another intestinal cell.

This cycle may be repeated several times, because of this cyclic multiplication, large numbers of intestinal cells are destroyed causing hemorrhages. Oocysts remain viable in litter for months. In this way, they can contaminate a farm from year to year. Oocysts are killed by freezing, extreme dryness, high temperatures and by use of effective drugs.



Vitamin K1 is not utilized by the feed industry, due to cost and lack of a stabilized form. Instead, water-soluble (Vitamin K3) salts are used to provide Vitamin K activity in feeds. Because of poor stability, menadione is not used as the pure vitamin, but is produced as water-soluble salts. Water-soluble derivatives of menadione, including menadione sodium bisulfite (MSB), are the principal forms of Vitamin K included in commercial diets. The menadione concentration is 50% for MSB, 45.4% for MPB, and 33% for MSBC.

Menadione (Vitamin K3) Concentrations from Menadione Salts for Feed Use

Menadione (K3) Salt	Menadione (K3) Concentration (%)	Amount (gm) of Menadione Salt to Provide 1 gm of Menadione
MSBC (menadione sodium bisulfite complex)	33.0	3.0
MSB (menadione sodium bisulfite)	50.0	2.0
MPB (menadione dimethylpyrimidinot bisulfite)	45.4	2.2

Debeaking Causes Hemorrhage/Bleeding

Debeaking, also called beak trimming, is the partial removal of the beak of poultry, especially layer hens and breeders although it may also be performed on quail, turkeys and ducks. Most commonly, the beak is shortened permanently, although regrowth can occur,... Beak trimming is most common in egg-laying strains of chickens.



COMPOSITION

Each 50 gm contains

Vitamin K ₃	500 mg
Carriers	q.s.

INDICATIONS

- During debeaking and outbreak of Coccidiosis.
- To Prevent Vitamin K deficiency syndrome in poultry. e.g. prolonged bleeding, large hematoma & severe hemorrhagic anemia etc.
- To Prevent hemorrhage in case of coccidiosis, necrotic enteritis, ulceration etc.
- After treatment with sulpha drugs.

USAGE INSTRUCTIONS

Chick & Growers : 50 gm for 3000 birds in drinking water for 5-7 days.

During Coccidiosis : 1 gm in 5 liters of drinking water for 7-10 days along with treatment. Or as recommended by the Veterinarian.

PRESENTATION: 50 gm pouch.



Manufactured by:-

Vesper Pharmaceuticals

Singapura, Vidyaranyapura Post, Bangalore, Karnataka, 560097, Ph: 080-23254255 Fax: 080-23254244
Email: vespergroupvet@gmail.com Web: www.vespergroupvet.com