

Vesper's

Toxol[®]

टोक्सोल

Nutritional Liver & Kidney Supplement






Since 1978



Vesperi's

Protects and stimulates Liver Under Stress due to Exogenous and Endogenous Toxins




Metabolically, liver is the most active organ of the chicken's body. A major portion of body's total metabolic reactions, be it assimilation of nutrients, catabolism of drugs or detoxification of toxins, take place in liver. For this purpose, liver cells are packed with a variety of enzymes that catalyze different reactions. While kidneys are the main organ for Excretion of waste and metabolised products.

Unfortunately, most liver disorders result in fat accumulation in liver. Depending on the degree of accumulation, the capacity of liver cells to perform the regular metabolic functions goes down, affecting growth, production and detoxification processes. At later stages, the condition also affects kidneys due to increased toxicity. More importantly, the clinical signs will be seen only after about 75 % of the liver is damaged. Lipo tropic agents may not be of much help due to their insignificant contribution towards the transport of lipids from liver cells to outside. What is needed in this situation is a product that accelerates the rate of removal of lipid molecules from liver cells.

Combats the ill effects of Hepatopathies & Nephropathies

Lipid Metabolism and Unique Action of Toxol



COMMERCIAL BROILERS AT 36 DAYS OF AGE, WITH AND WITHOUT TOXOL

Treatment TOXOL	T.No	No. of Chicks	Avg Wt of Chicks in gms	36th day body wt	Mortality No. %		Feed consumed	FCR
1000 gm	3	600	44	2.165	14	2.33	3.560	1.644
500 gm	2	600	42	2	18	3.0	3.366	1.683
CONTROL	1	600	43	1.925	21	3.5	3.388	1.760

Result: Body weights increased with Toxol dosage. FCR improved compared to control. Mortality reduced in the treatment groups.

Toxol[®]

टोक्सोल

Nutritional Liver & Kidney Supplement

COMPOSITION

Toxol Feed Premix: Choline Chloride, Methionine, Inositol, Vitamin B₁₂, Base fortified with Predigested Protein and Aminonitrogen.

Toxol Liquid : Each 10 ml contains: Tricholine Citrate 100 mg, Methionine 10 mg, Liver extract 50 mg, Protein Hydrolysate 50 mg, Inositol 2 mg, Vitamin B₁₂ 0.33 mcg Base fortified with Predigested Protein and Aminonitrogen.

INDICATIONS

Improves Liver & Kidney Health.

Protect from damaging effect of Mycotoxins.

Improves Nutrient Utilization.

USAGE INSTRUCTIONS

Toxol Feed: Normal Mix - 500 gms to 1 kg per ton of feed.

Under Stress / Mycotoxins - Add 2 to 2.5 kgs per ton of feed.

Or as recommended by the Veterinarian.

Toxol Liquid : 5-15 ml / 100 birds.

Or as recommended by the Veterinarian.



PRESENTATION

Toxol Feed Premix : 1 kg, 5 kg, 10 kg, 20 kg bags.

Toxol Liquid : 500 ml, 1 Liter, 5 Liter and 30 Liter Drum.



Since 1978

Manufactured by:

VESPER PHARMACEUTICALS

Singapura, Vidyananyapura Post, Bengaluru - 560 097, Karnataka, INDIA

Ph: 080- 23254255/23254266/23254277. Fax: 080-23254244

Email: info@vespergroupvet.com . Website: www.vespergroupvet.com

An ISO 9001-2008 Certified Company

VISION  Catalyzes Research
Creates Products
Cares Customers
The Care Takers