



CS215 DISCRETE MATH

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

Solving Linear Recurrence Relations of degree k

- Consider an arbitrary linear homogeneous relation of degree k with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

Theorem If this CE has k distinct roots r_i , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all $n \geq 0$, where the α_i 's are constants.



The Case of Degenerate Roots

- **Theorem** If the CE $r^2 - c_1 r - c_2 = 0$ has **only 1** root r_0 , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all $n \geq 0$ and two constants α_1 and α_2 .



The Case of Degenerate Roots

- **Theorem** If the CE $r^2 - c_1 r - c_2 = 0$ has **only 1** root r_0 , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all $n \geq 0$ and two constants α_1 and α_2 .

- **Theorem** [Theorem 4, p.519] Suppose that there are t roots r_1, \dots, r_t with multiplicities m_1, \dots, m_t . Then

$$a_n = \sum_{i=1}^t \left(\sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all $n \geq 0$ and constants $\alpha_{i,j}$.



Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms $F(n)$ that depend only on n

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$ is called the *associated homogeneous recurrence relation*.

Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms $F(n)$ that depend only on n

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$ is called the *associated homogeneous recurrence relation*.

Fact: Assume that the sequence b_n satisfies the recurrence. Then another sequence a_n satisfies the non-homogeneous recurrence if and only if $h_n = a_n - b_n$ is a sequence that satisfies the associated homogeneous recurrence.



Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms $F(n)$ that depend only on n

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$ is called the *associated homogeneous recurrence relation*.

Fact: Assume that the sequence b_n satisfies the recurrence. Then another sequence a_n satisfies the non-homogeneous recurrence if and only if $h_n = a_n - b_n$ is a sequence that satisfies the associated homogeneous recurrence.

Idea: We already know how to find h_n . For many common $F(n)$, a solution b_n to the non-homogeneous recurrence is similar to $F(n)$. We then need find $a_n = b_n + h_n$ to the non-homogeneous recurrence that satisfies both recurrence and initial conditions.



Linear Nonhomogeneous Recurrence Relations

- **Theorem** If $a_n = p(n)$ is any particular solution to the linear nonhomogeneous relation with constant coefficients,

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n),$$

Then all its solutions are of the form

$$a_n = p(n) + h(n),$$

where $a_n = h(n)$ is any solution to the associated homogeneous recurrence relation

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}.$$

Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- t polynomial as the particular solution $p(n)$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- t polynomial as the particular solution $p(n)$.

Let $p(n) = cn + d$, then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means } (2c + 2)n + (2d - 3c) = 0.$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example** $a_n = 3a_{n-1} + 2n$. Which solution has $a_1 = 3$?

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 3r = 0$. Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- t polynomial as the particular solution $p(n)$.

Let $p(n) = cn + d$, then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means} \\ (2c + 2)n + (2d - 3c) = 0.$$

We get $c = -1$ and $d = -3/2$. Thus,

$$p(n) = -n - 3/2$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Tower of Hanoi.** $H(n) = 2H(n - 1) + 1$ with $H(1) = 1$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Tower of Hanoi.** $H(n) = 2H(n - 1) + 1$ with $H(1) = 1$.
The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $H(n) = \alpha 2^n + p(n)$.

Solving Linear Nonhomogeneous Recurrence Relations

- **Example Tower of Hanoi.** $H(n) = 2H(n - 1) + 1$ with $H(1) = 1$.
The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $H(n) = \alpha 2^n + p(n)$.

We try a constant c as the particular solution $p(n)$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Tower of Hanoi.** $H(n) = 2H(n - 1) + 1$ with $H(1) = 1$.
The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $H(n) = \alpha 2^n + p(n)$.

We try a constant c as the particular solution $p(n)$.

Let $p(n) = c$, then

$$c = 2c + 1, \text{ which means } c = -1.$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Tower of Hanoi.** $H(n) = 2H(n-1) + 1$ with $H(1) = 1$. The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $H(n) = \alpha 2^n + p(n)$.

We try a constant c as the particular solution $p(n)$.

Let $p(n) = c$, then

$$c = 2c + 1, \text{ which means } c = -1.$$

We get $H(n) = \alpha 2^n - 1$. With the initial condition $H(1) = 1$, we have $\alpha = 1$. Thus, $H(n) = 2^n - 1$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Merge Sort.** $T(n) = 2T(n/2) + n - 1$ with $T(1) = 0$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Merge Sort.** $T(n) = 2T(n/2) + n - 1$ with $T(1) = 0$.

Let $T(n) = T(2^k) = M(k)$, then we have

$M(k) = 2M(k-1) + 2^k - 1$ with $M(0) = 0$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Merge Sort.** $T(n) = 2T(n/2) + n - 1$ with $T(1) = 0$.

Let $T(n) = T(2^k) = M(k)$, then we have
 $M(k) = 2M(k-1) + 2^k - 1$ with $M(0) = 0$.

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $M(k) = \alpha 2^k + p(k)$.



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Merge Sort.** $T(n) = 2T(n/2) + n - 1$ with $T(1) = 0$.

Let $T(n) = T(2^k) = M(k)$, then we have
 $M(k) = 2M(k-1) + 2^k - 1$ with $M(0) = 0$.

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $M(k) = \alpha 2^k + p(k)$.

We try $c_1 k \cdot 2^k + c_2$ as the particular solution $p(k)$.

Let $p(k) = c_1 k \cdot 2^k + c_2$, then

$$c_1 k \cdot 2^k + c_2 = 2c_1(k-1)2^{k-1} + 2c_2 + 2^k - 1, \text{ which gives} \\ c_1 = c_2 = 1.$$



Solving Linear Nonhomogeneous Recurrence Relations

- **Example Merge Sort.** $T(n) = 2T(n/2) + n - 1$ with $T(1) = 0$.

Let $T(n) = T(2^k) = M(k)$, then we have
 $M(k) = 2M(k-1) + 2^k - 1$ with $M(0) = 0$.

The *characteristic equation* of the associated linear homogeneous recurrence relation is $r^2 - 2r = 0$. Thus, the solution to the original problem are all of the form $M(k) = \alpha 2^k + p(k)$.

We try $c_1 k \cdot 2^k + c_2$ as the particular solution $p(k)$.

Let $p(k) = c_1 k \cdot 2^k + c_2$, then

$$c_1 k \cdot 2^k + c_2 = 2c_1(k-1)2^{k-1} + 2c_2 + 2^k - 1, \text{ which gives} \\ c_1 = c_2 = 1.$$

We get $M(k) = \alpha 2^k + k \cdot 2^k + 1$. With the initial condition $M(0) = 0$, we have $\alpha = -1$. Thus, $M(k) = k \cdot 2^k - 2^k + 1$ and
 $T(n) = n \log n - n + 1$.



Generating Functions

- We may use *generating functions* to **characterize** sequences.



Generating Functions

- We may use *generating functions* to **characterize** sequences.
 - ◇ The sequence $\{a_k\}$ with $a_k = 3$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

- ◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

- ◇ The sequence $\{a_k\}$ with $a_k = 2^k$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence $\{a_k\}$ with $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$



Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence $\{a_k\}$ with $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence $\{a_k\}$ with $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$

Definition The *generating function* for the sequence $a_0, a_1, \dots, a_k, \dots$ of real numbers is the infinite series

$$G(x) = a_0 + a_1x + \dots + a_kx^k + \dots = \sum_{k=0}^{\infty} a_kx^k$$



Generating Functions

- “*Generating functions* are a bridge between *discrete mathematics* , on one hand and *continuous analysis* (particularly complex variable theory) on the other. It is possible to study them solely as tools for solving discrete problems.”
– Herbert S. Wilf



Q -Ary Non-Overlapping Codes: A Generating Function Approach

Geyang Wang^{ID} and Qi Wang^{ID}, *Member, IEEE*

Abstract—Non-overlapping codes are a set of codewords in $\bigcup_{n \geq 2} \mathbb{Z}_q^n$, where $\mathbb{Z}_q = \{0, 1, \dots, q-1\}$, such that the prefix of each codeword is not a suffix of any codeword in the set, including itself; and for variable-length codes, a codeword does not contain any other codeword as a subword. In this paper, we investigate a generic method to generalize binary codes to q -ary ones for $q > 2$, and analyze this generalization on the two constructions given by Levenshtein (also by Gilbert; Chee, Kiah, Purkayastha, and Wang) and Bilotta, respectively. The generalization on the former construction gives large non-expandable fixed-length non-overlapping codes whose size can be explicitly determined; the generalization on the latter construction is the first attempt to generate q -ary variable-length non-overlapping codes. More importantly, this generic method allows us to utilize the generating function approach to analyze the cardinality of the underlying q -ary non-overlapping codes. The generating function approach not only enables us to derive new results, e.g., recurrence relations on their cardinalities, new combinatorial interpretations for the constructions, and the superior limit of their cardinalities for some special cases, but also greatly simplifies the arguments for these results. Furthermore, we give an exact formula for the number of fixed-length words that do not contain the codewords in a variable-length non-overlapping code as subwords. This thereby solves an open problem by Bilotta and induces a recursive upper bound on the maximum size of variable-length non-overlapping codes.

Index Terms—Non-overlapping code, variable-length code, generating function.

- (1) No non-empty prefix of each codeword is a suffix of any one, including itself;
- (2) For all distinct $u, v \in S$, u does not contain v as a subword.

We say that S is a fixed-length non-overlapping code if $S \subseteq \mathbb{Z}_q^n$, otherwise it is called a variable-length non-overlapping code. In this paper, we consider both fixed-length and variable-length cases. Fixed-length non-overlapping codes have been intensively studied in the literature. Let $C(n, q)$ be the maximum size of a q -ary non-overlapping codes of length n . The main research problems are to construct non-overlapping codes as large as possible in size and to bound $C(n, q)$. The first construction was proposed by Levenshtein in 1964 [2], [3] (Construction 1, see also [4]–[6]). Following the work by de Lind van Wijngaarden and Willink [7] in 2000, Bajic and Stojanovic [8] independently rediscovered binary fixed-length non-overlapping codes (under the name *cross-bifix-free codes*) in 2004. In 2012, Bilotta *et al.* [9] provided a binary construction based on Dyck paths, by which the code size is smaller than Levenshtein's. However, it reveals an interesting connection between non-overlapping codes and other combinatorial objects. In 2013, Chee *et al.* [6] rediscovered Levenshtein's construction (Construction 1), and verified that it is optimal for $q =$

Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence a_0, a_1, \dots, a_m , with $a_k = C(m, k)$?



Generating Functions for Finite Sequences

- A finite sequence a_0, a_1, \dots, a_n can be easily extended by setting $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function* $G(x)$ of this infinite sequence $\{a_n\}$ is a polynomial of degree n , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence a_0, a_1, \dots, a_m , with $a_k = C(m, k)$?

$$G(x) = C(m, 0) + \dots + C(m, m)x^m = (1 + x)^m$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$1, 1, 1, 1, 1, \dots$

Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇ $G(x) = 1/(1 - ax)$ for $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$

$$1, 2, 3, 4, 5, \dots$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$
 $1, 1, 1, 1, 1, \dots$

Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$
 $1, 1, 1, 1, 1, \dots$

$$G(x) = 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1$$



Examples

$$\diamond G(x) = 1/(1-x) \text{ for } |x| < 1$$
$$1, 1, 1, 1, 1, \dots$$

$$G(x) = 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1$$
$$-xG(x) = -x - x^2 - x^3 - \dots - x^n - \dots$$

$$G(x) - xG(x) = 1$$



Examples

$$\diamond G(x) = 1/(1-x) \text{ for } |x| < 1$$
$$1, 1, 1, 1, 1, \dots$$

$$G(x) = 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1$$
$$-xG(x) = -x - x^2 - x^3 - \dots - x^n - \dots$$

$$G(x) - xG(x) = 1$$

$$\diamond G(x) = 1/(1-x)^2 \text{ for } |x| < 1$$
$$1, 2, 3, 4, 5, \dots$$



Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$
 $1, 1, 1, 1, 1, \dots$

$$\begin{aligned} G(x) &= 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1 \\ -xG(x) &= -x - x^2 - x^3 - \dots - x^n - \dots \end{aligned}$$

$$G(x) - xG(x) = 1$$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$
 $1, 2, 3, 4, 5, \dots$

$$G(x) = 1 + 2x + 3x^2 + 4x^3 + \dots + (n + 1)x^n + \dots \text{ for } |x| < 1$$

Examples

◇ $G(x) = 1/(1 - x)$ for $|x| < 1$
 $1, 1, 1, 1, 1, \dots$

$$\begin{aligned} G(x) &= 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1 \\ -xG(x) &= -x - x^2 - x^3 - \dots - x^n - \dots \end{aligned}$$

$$G(x) - xG(x) = 1$$

◇ $G(x) = 1/(1 - x)^2$ for $|x| < 1$
 $1, 2, 3, 4, 5, \dots$

$$\begin{aligned} G(x) &= 1 + 2x + 3x^2 + 4x^3 + \dots + (n+1)x^n + \dots \text{ for } |x| < 1 \\ -xG(x) &= -x - 2x^2 - 3x^3 + \dots - nx^n + \dots \end{aligned}$$

Examples

$$\diamond G(x) = 1/(1-x) \text{ for } |x| < 1$$
$$1, 1, 1, 1, 1, \dots$$

$$G(x) = 1 + x + x^2 + x^3 + \dots + x^n + \dots \text{ for } |x| < 1$$
$$-xG(x) = -x - x^2 - x^3 - \dots - x^n - \dots$$

$$G(x) - xG(x) = 1$$

$$\diamond G(x) = 1/(1-x)^2 \text{ for } |x| < 1$$
$$1, 2, 3, 4, 5, \dots$$

$$G(x) = 1 + 2x + 3x^2 + 4x^3 + \dots + (n+1)x^n + \dots \text{ for } |x| < 1$$
$$-xG(x) = -x - 2x^2 - 3x^3 + \dots - nx^n + \dots$$

$$G(x) - xG(x) = 1 + x + x^2 + x^3 + \dots = 1/(1-x)$$

Operations of Generating Functions

- **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1-x)^2$ for $|x| < 1$



Operations of Generating Functions

- **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1-x)^2$ for $|x| < 1$

$$f(x) = 1/(1-x), \quad g(x) = 1/(1-x)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k 1 \right) x^k = \sum_{k=0}^{\infty} (k+1) x^k$$



Operations of Generating Functions

■ **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1 - ax)^2$ for $|ax| < 1$



Operations of Generating Functions

- **Theorem** Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$, and $g(x) = \sum_{k=0}^{\infty} b_k x^k$.
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇ $G(x) = 1/(1 - ax)^2$ for $|ax| < 1$

$$f(x) = 1/(1 - ax), \quad g(x) = 1/(1 - ax)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a^j a^{k-j} \right) x^k = \sum_{k=0}^{\infty} (k+1) a^k x^k$$



Counting with Generating Functions

- **Convolution Rule.** Let $A(x)$ be the generating function for selecting items from a set A , and let $B(x)$ be the generating function for selecting items from a set B disjoint from A . Then the generating function for selecting items from the union $A \cup B$ is the product $A(x) \cdot B(x)$.



Counting with Generating Functions

- **Convolution Rule.** Let $A(x)$ be the generating function for selecting items from a set A , and let $B(x)$ be the generating function for selecting items from a set B **disjoint** from A . Then **the generating function for selecting items from the union $A \cup B$ is the product $A(x) \cdot B(x)$.**

Example. There are **two** types of balls, **blue** and **red**. Let d_k be the $\#$ of ways to select k **blue** or **red** balls, and $D(x)$ be its generating function.



Counting with Generating Functions

- **Convolution Rule.** Let $A(x)$ be the generating function for selecting items from a set A , and let $B(x)$ be the generating function for selecting items from a set B **disjoint** from A . Then **the generating function for selecting items from the union $A \cup B$ is the product $A(x) \cdot B(x)$.**

Example. There are **two** types of balls, **blue** and **red**. Let d_k be the $\#$ of ways to select k **blue** or **red** balls, and $D(x)$ be its generating function.

$$d_k = k + 1, D(x) = 1/(1 - x)^2.$$



Counting with Generating Functions

- **Convolution Rule.** Let $A(x)$ be the generating function for selecting items from a set A , and let $B(x)$ be the generating function for selecting items from a set B disjoint from A . Then the generating function for selecting items from the union $A \cup B$ is the product $A(x) \cdot B(x)$.

Example. There are **two** types of balls, **blue** and **red**. Let d_k be the # of ways to select k **blue** or **red** balls, and $D(x)$ be its generating function.

$$d_k = k + 1, D(x) = 1/(1 - x)^2.$$

This can be generalized to counting the # of ways to select k balls with **n** colors.

$$D(x) = 1/(1 - x)^n, \text{ what is } d_k?$$



Counting with Generating Functions

- This can be generalized to counting the # of ways to select k balls with n colors.

$$D(x) = 1/(1 - x)^n, \text{ what is } d_k?$$



Counting with Generating Functions

- This can be generalized to counting the # of ways to select k balls with n colors.

$D(x) = 1/(1 - x)^n$, what is d_k ?

$$d_k = [x^k](1/(1 - x)^n) = \binom{n+k-1}{k}$$



Counting with Generating Functions

- This can be generalized to counting the # of ways to select k balls with n colors.

$D(x) = 1/(1-x)^n$, what is d_k ?

$$d_k = [x^k](1/(1-x)^n) = \binom{n+k-1}{k}$$

Theorem (Maclaurin's Theorem).

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

$$\frac{d^k}{d^k x} (1/(1-x)^n) = n(n+1) \cdots (n+k-1)(1-x)^{-(n+k)}$$

$$[x^k](1/(1-x)^n) = \frac{n(n+1) \cdots (n+k-1)(1-0)^{-(n+k)}}{k!} = \binom{n+k-1}{k}$$



Counting with Generating Functions

- This can be generalized to counting the # of ways to select k balls with n colors.

$D(x) = 1/(1-x)^n$, what is d_k ?

$$d_k = [x^k](1/(1-x)^n) = \binom{n+k-1}{k}$$

Theorem (Maclaurin's Theorem).

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

$$\frac{d^k}{dx^k}(1/(1-x)^n) = n(n+1)\dots(n+k-1)(1-x)^{-(n+k)}$$

$$[x^k](1/(1-x)^n) = \frac{n(n+1)\dots(n+k-1)(1-0)^{-(n+k)}}{k!} = \binom{n+k-1}{k}$$

Alternatively, apply the **extended binomial theorem**:

$$\binom{-n}{k} = \frac{(-n)(-n-1)\dots(-n-k+1)}{k!} = (-1)^k \frac{(n+k-1)\dots(n+1)n}{k!} = (-1)^k \binom{n+k-1}{k}$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$



Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$

$$\frac{1}{(1-x)^2} = \sum_{k=0}^{\infty} (k+1) x^k = 1 + 2x + 3x^2 + \dots$$



Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$



Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\ln(1+x) = \sum_{k=0}^{\infty} \frac{(-1)^{k+1} x^k}{k} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$



Counting and Generating Functions

- **Problem 1** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers with $2 \leq x_1 \leq 5$,
 $3 \leq x_2 \leq 6$, $4 \leq x_3 \leq 7$.



Counting and Generating Functions

- **Problem 1** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers with $2 \leq x_1 \leq 5$,
 $3 \leq x_2 \leq 6$, $4 \leq x_3 \leq 7$.

Using *generating functions*, the number is the **coefficient** of x^{17} in the expansion of

$$(x^2 + x^3 + x^4 + x^5)(x^3 + x^4 + x^5 + x^6)(x^4 + x^5 + x^6 + x^7)$$



Counting and Generating Functions

- **Problem 2** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?



Counting and Generating Functions

- **Problem 2** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?

The coefficient of x^8 in the expansion

$$(x^2 + x^3 + x^4)^3$$



Counting and Generating Functions

- **Problem 3** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?



Counting and Generating Functions

- **Problem 3** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?

This is **equivalent** to the problem of k -combinations from a set with n elements when **repetition is allowed**.



Counting and Generating Functions

- **Problem 3** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where x_1, x_2, x_3 are **nonnegative** integers?

This is **equivalent** to the problem of k -combinations from a set with n elements when **repetition is allowed**.

$$C(n + k - 1, k) = C(19, 17) = C(19, 2)$$



r -Combinations from a Set

- **Definition** A *k -combination* with **repetition allowed**, or a *multiset of size k* , chosen from a set of n elements, is an unordered selection of elements with repetition allowed.



r -Combinations from a Set

- **Definition** A *k -combination* with **repetition allowed**, or a *multiset of size k* , chosen from a set of n elements, is an unordered selection of elements with repetition allowed.

Example Find $\#$ multisets of size 17 from the set $\{1, 2, 3\}$.



r -Combinations from a Set

- **Definition** A *k -combination* with **repetition allowed**, or a *multiset of size k* , chosen from a set of n elements, is an unordered selection of elements with repetition allowed.

Example Find $\#$ multisets of size 17 from the set $\{1, 2, 3\}$.

This is **equivalent** to finding the $\#$ nonnegative solutions to $x_1 + x_2 + x_3 = 17$.



r -Combinations from a Set

- **Definition** A k -combination with repetition allowed, or a *multiset of size k* , chosen from a set of n elements, is an unordered selection of elements with repetition allowed.

Example Find $\#$ multisets of size 17 from the set $\{1, 2, 3\}$.

This is equivalent to finding the $\#$ nonnegative solutions to $x_1 + x_2 + x_3 = 17$.

Q: Use generating functions to find the number of k -combinations from a set with n elements when repetition of elements is allowed.



r -Combinations from a Set

- **Definition** A k -combination with repetition allowed, or a *multiset of size k* , chosen from a set of n elements, is an unordered selection of elements with repetition allowed.

Example Find $\#$ multisets of size 17 from the set $\{1, 2, 3\}$.

This is equivalent to finding the $\#$ nonnegative solutions to $x_1 + x_2 + x_3 = 17$.

Q: Use generating functions to find the number of k -combinations from a set with n elements when repetition of elements is allowed.

Read more on pp. 537-548.



Counting and Generating Functions

- **Problem 4** Use generating functions to find the number of k -combinations of a set with n elements, $C(n, k)$.



Counting and Generating Functions

- **Problem 4** Use **generating functions** to find the number of **k -combinations of a set with n elements**, $C(n, k)$.

Each of the n elements in the set contributes the term $(1 + x)$ to the generating function $f(x) = \sum_{k=0}^n a^k x^k$.
Hence, $f(x) = (1 + x)^n$.

Then by the **binomial theorem**, we have $a_k = \binom{n}{k}$.



Cartesian Product

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$



Cartesian Product

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$

Cartesian product defines a set of all **ordered** arrangements of elements in the two sets.

Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

We use the notation $a R b$ to denote $(a, b) \in R$, and $a \not R b$ to denote $(a, b) \notin R$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

We use the notation $a R b$ to denote $(a, b) \in R$, and $a \not R b$ to denote $(a, b) \notin R$.

Example: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3\}$

- ◇ Is $R = \{(a, 1), (b, 2), (c, 2)\}$ a relation from A to B ?
- ◇ Is $Q = \{(1, a), (2, b)\}$ a relation from A to B ?
- ◇ Is $P = \{(a, a), (b, c), (b, a)\}$ a relation from A to A ?



Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$

Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$


Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and
 $R = \{(0, u), (0, v), (1, v), (2, u)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, v), (2, u)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, u), (2, v)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, u), (2, v)\}$. ($R \subseteq A \times B$)

R	u	v
0	×	×
1	×	
2		×


Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .




Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .



Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .



What is the **difference** between a **relation** and a **function** from A to B ?

Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.



Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$




Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$




Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



R	1	2	3	4
1	×	×	×	×
2		×		×
3			×	
4				×

Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.

Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.

R is a binary relation on A if $R \subseteq A \times A$ (R is subset)



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.

R is a binary relation on A if $R \subseteq A \times A$ (R is subset)

The number of subsets of a set with k elements is 2^k



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. $(1, 1), (2, 2), (3, 3), (4, 4) \in R_{div}$



Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

$$MR_{div} = \begin{matrix} & 1 & 2 & 3 & 4 \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{matrix} 1 \\ 1 \\ 0 \\ 0 \end{matrix} & \begin{matrix} 1 \\ 1 \\ 0 \\ 0 \end{matrix} & \begin{matrix} 1 \\ 0 \\ 1 \\ 0 \end{matrix} & \begin{matrix} 1 \\ 1 \\ 0 \\ 1 \end{matrix} \end{matrix}$$



Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

$$MR_{div} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix}$$

A relation R is reflexive if and only if MR has 1 in every position on its main diagonal.

Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R reflexive?



Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R reflexive?

No. $(1, 1) \notin R$

Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Yes. $(1, 1), (2, 2), (3, 3), (4, 4) \notin R_{\neq}$

Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

A relation R is irreflexive if and only if MR has 0 in every position on its **main diagonal**.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} symmetric?



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *symmetric*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *symmetric*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

No. $(1, 2) \in R_{div}$ but $(2, 1) \notin R$

Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} symmetric?

Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} symmetric?

Yes. If $(a, b) \in R_{\neq}$ then $(b, a) \in R_{\neq}$.



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

A relation R is symmetric if and only if **MR** is symmetric.

Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R antisymmetric?



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R antisymmetric?

Yes.



Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

$$MR = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

$$MR = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

A relation R is antisymmetric if and only if $m_{ij} = 1$ implies $m_{ji} = 0$ for $i \neq j$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|a$, then $a = b$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|a$, then $a = b$.

$$MR_{div} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix}$$

Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for all $a, b, c \in A$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|c$, then $a|c$.



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?

Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?

No. $(1, 2), (2, 1) \in R_{\neq}$ but $(1, 1) \notin R_{\neq}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?

Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?

Yes.

Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.

Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.

Set operations: **union, intersection, difference, etc.**



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?

We may also combine relations by **matrix operations**.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$

$$S \circ R = \{(1, b), (3, a), (3, b)\}$$



Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C
 $S \circ R = \{(1, b), (1, a), (2, a)\}$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$$S \circ R = \{(1, b), (1, a), (2, a)\}$$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix}$$

Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 & 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$

$$M_R \odot M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 & 2 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^k = ? \text{ for } k > 3$$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.

“only if” part: by induction.

Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?

How many subsets on $n(n-1)$ elements are there?



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.

Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.

Transitive Relation: A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



Next Lecture

- relation II...

