# Partitioned Matrices (分块矩阵的乘法)

Lecture 4+

Dept. of Math., SUSTech

2023.9

### Introduction

Often it is useful to think of a matrix as being composed of a number of submatrices. A matrix *C* can be partitioned into smaller matrices by drawing horizontal lines between the rows and vertical lines between the columns. The smaller matrices are often referred to as blocks.

Y. Chen (Dept. of Math.) Partitioned Manness 2023.9

## Example

Example. 
$$C = \begin{bmatrix} 1 & -2 & 4 & 1 & 3 \\ 2 & 1 & 1 & 1 & 1 \\ \hline 3 & 3 & 2 & -1 & 2 \\ 4 & 6 & 2 & 2 & 4 \end{bmatrix}$$
. If lines are drawn between the

3/13

second and third rows and between the third and fourth columns, then C will be divided into four submatrices

$$C = \left[ \begin{array}{c|c} C_{11} & C_{12} \\ \hline C_{21} & C_{22} \end{array} \right]$$

where

$$C_{11} = \begin{bmatrix} 1 & -2 & 4 \\ 2 & 1 & 1 \end{bmatrix}, C_{12} = \begin{bmatrix} 1 & 3 \\ 1 & 1 \end{bmatrix}, C_{21} = \begin{bmatrix} 3 & 3 & 2 \\ 4 & 6 & 2 \end{bmatrix}, C_{22} = \begin{bmatrix} -1 & 2 \\ 2 & 4 \end{bmatrix}.$$

Y. Chen (Dept. of Math.) 2023.9

## **Block Multiplication I**

Block Multiplication I. If A is an  $m \times n$  matrix and B is an  $n \times r$  matrix that has been partitioned into columns  $(b_1, b_2, \dots, b_r)$ , then the block multiplication of A times B is given by

$$AB = (Ab_1, Ab_2, \cdots, Ab_r)$$

Y. Chen (Dept. of Math.) Fartilioned Mathieur 2023.9

## **Block Multiplication II**

Block Multiplication II. Let A be an  $m \times n$  matrix. If we partition A into rows, and if B is an  $n \times r$  matrix, the ith row of the product AB is determined by multiplying the ith row of A times B, then

A = 
$$\begin{bmatrix} \vec{a}_1 \\ \vec{a}_2 \\ \cdot \\ \cdot \\ \cdot \\ \vec{a}_m \end{bmatrix}$$
,  $AB = \begin{bmatrix} \vec{a}_1B \\ \vec{a}_2B \\ \cdot \\ \cdot \\ \cdot \\ \vec{a}_mB \end{bmatrix}$ .

Y. Chen (Dept. of Math.) Parlitioned Mathies 2023.9

## **Block Multiplication III**

Let A be an  $m \times n$  matrix and B and  $n \times r$  matrix. It is often useful to partition A and B and express the product in terms of the submatrices of A and B.

Case 1. 
$$A(B_1 B_2) = (AB_1 AB_2)$$

In this case,  $B_1$  is an  $n \times t$  matrix and  $B_2$  is an  $n \times (r-t)$  matrix.

Case 2. 
$$\begin{bmatrix} A_1 \\ A_2 \end{bmatrix} B = \begin{bmatrix} A_1 B \\ A_2 B \end{bmatrix}$$

Here  $A_1$  is a  $k \times n$  matrix and  $A_2$  is an  $(m-k) \times n$  matrix.

Y. Chen (Dept. of Math.) Parlitioned Mathieus 2023.9

# **Block Multiplication IV**

Case 3. 
$$(A_1 A_2) \begin{pmatrix} B_1 \\ B_2 \end{pmatrix} = A_1 B_1 + A_2 B_2$$

Case
$$4. \begin{bmatrix}
A_{11} & A_{12} \\
A_{21} & A_{22}
\end{bmatrix}
\begin{bmatrix}
B_{11} & B_{12} \\
B_{21} & B_{22}
\end{bmatrix} = \begin{bmatrix}
A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\
A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22}
\end{bmatrix}$$

Can you figure out the sizes of all the above submatrices?

In general, if the blocks have the proper dimensions, the block multiplication can be carried out in the same manner as ordinary matrix multiplication.

Y. Chen (Dept. of Math.) Partitioned Matrices 2023.9

### Examples

Example 1 
$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$
, and  $B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$ . Partition  $A$ 

and *B* into four blocks and perform the block multiplication.

Example 2 Let A be an  $n \times n$  matrix of the form

$$\left[\begin{array}{cc} A_{11} & O \\ O & A_{22} \end{array}\right]$$

where  $A_{11}$  is a  $k \times k$  matrix(k < n). Show that A is nonsingular if and only if  $A_{11}$  and  $A_{22}$  are nonsingular.

Y. Chen (Dept. of Math.) Parlitioned Mathies 2023.9

#### Inner Product

Given two vectors x and y in  $\mathbb{R}^n$ , it is possible to perform a matrix multiplication of the vectors if we transpose one of the vectors first. The matrix product  $x^Ty$  is the product of a row vector (a  $1 \times n$  matrix) and a column vector (an  $n \times 1$  matrix). The result is a  $1 \times 1$  matrix, or simply a scalar. This type of product is referred to as a scalar product or an inner product.

Y. Chen (Dept. of Math.) Parlitioned Mathieus 2023.9

## Outer product

The outer product has special structure in that each of its rows is a multiple of  $y^T$  and each of its column vectors is a multiple of x.

Y. Chen (Dept. of Math.) Parillones Metrices 2023.9

## **Outer Product Expansion**

It is also useful to consider a column vector times a row vector. The matrix product  $xy^T$  is a full  $n \times n$  matrix. The product  $xy^T$  is referred to as the outer product of x and y. The outer product matrix has special structure in that each of its rows is a multiple of  $y^T$  and each of its column vectors is a multiple of x.

Example Compute 
$$\left(\begin{bmatrix} 1\\2\\3 \end{bmatrix}\begin{bmatrix} 1&2&3 \end{bmatrix}\right)^{2023}$$
.

Y. Chen (Dept. of Math.) Southerness 2023.9

## One more example

#### **Example Given**

$$X = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix} \text{ and } Y = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 1 \end{bmatrix}$$

compute the outer product expansion of  $XY^T$ .

## Homework Assignment 5(Suggested)

1.4: 45, 46, 47, 51, 52, 53, 55, 57, 58.

Y. Chen (Dept. of Math.) Parlitioned Manues 2023.9