

An introduction to some concepts in several complex variables

Shijie BAO (包诗界)

Date: September 3, 2025

Abstract

This is a manuscript for giving an introduction to some concepts in several complex variables.

Contents

1	Preliminaries: Basic Knowledge in Several Complex Variables	3
1.1	Holomorphic function	3
1.2	Analytic set	6
1.3	Plurisubharmonic function	8
1.4	Pseudoconvexity	9
1.5	Pluripolar set	11
2	L^2 Methods and Multiplier Ideal Sheaf	13
2.1	L^2 estimate of $\bar{\partial}$ -equation	13
2.2	L^2 extension theorem	17
2.2.1	Applications of the L^2 extension theorem	21
2.3	Multiplier ideal sheaf	31
2.4	Openness and strong openness	33
2.5	Skoda's L^2 division theorem	36
3	Pluripotential Theory	41
3.1	Preliminaries: Distribution	41
3.2	Current	43
3.3	Complex Monge-Ampère operator	46
3.3.1	Definition: Case of locally bounded psh function	46
3.3.2	Definition: Case of unbounded psh function	48
3.4	Lelong Number	49
3.4.1	Demailly's generalized Lelong number	49
3.5	Siu's theorems	52
3.5.1	Siu's slicing theorem	52
3.5.2	Siu's semi-continuity theorem	52
3.5.3	Siu's decomposition theorem	53
3.6	Cegrell's class	54
3.7	Extremal plurisubharmonic function	55
3.7.1	Maximal weight and relative type	55

3.7.2	Pluricomplex Green function	57
3.7.3	Capacity function	58
4	Some applications	59
4.1	Asymptotic behavior of Bergman kernel	59
4.2	Spectrum of Toeplitz operator	61
4.3	Riemann–Roch; Holomorphic Morse Inequality	63
5	Some open problems	65
5.1	Zero mass conjecture	65
5.2	Sublevel sets of pluricomplex Green function	65

1 Preliminaries: Basic Knowledge in Several Complex Variables

Notations

- The disc $D(a_0, r) := \{z \in \mathbb{C} : |z - a_0| < r\}$ for $a_0 \in \mathbb{C}$ and $r > 0$.
- The polydisc $D(a, R) := D(a_1, R_1) \times \cdots \times D(a_n, R_n)$ for $a = (a_1, \dots, a_n) \in \mathbb{C}^n$ and $R = (R_1, \dots, R_n) \in \mathbb{R}_{>0}^n$.
- $\Delta(r) := D(0, r) \subset \mathbb{C}$ for $r > 0$, and $\Delta^n(r) := \Delta(r) \times \cdots \times \Delta(r) \subset \mathbb{C}^n$ for $r > 0$. In particular, $\Delta^n := \Delta^n(1)$.
- The ball $B(a, R) := \{|z - a| < R\} \subset \mathbb{C}^n$ for $a \in \mathbb{C}^n$ and $R > 0$.
- $\wedge^{p,q}(\Omega)$: the set of (p, q) forms in Ω .
- $\mathcal{D}^{p,q}(\Omega)$: the set of smooth (p, q) forms with compact support in Ω .
- $\delta_\Omega(z)$: the distance from z to the boundary of Ω , i.e. $\delta_\Omega(z) = \text{dist}(z, \partial\Omega)$. We also denote $\delta(z) = \delta_\Omega(z)$ for simplicity.

We recall some basic knowledge from several complex variables in this section.

1.1 Holomorphic function

Let $\Omega \subset \mathbb{C}^n$ be an open subset.

DEFINITION 1.1. A function $f: \Omega \rightarrow \mathbb{C}$ is called a holomorphic function, if f is continuous and for each j ,

$$z_j \mapsto f(\dots, z_j, \dots)$$

is holomorphic when $z_1, \dots, z_{j-1}, z_{j+1}, \dots, z_n$ are fixed.

REMARK 1.2. 1. (Hartogs) The continuity condition can be dropped.

2. We usually denote the ring of holomorphic functions on Ω by $\mathcal{O}(\Omega)$ or $H(\Omega)$.

THEOREM 1.3 (Cauchy integral formula on polydiscs). Let $z_0 = (z_{0,1}, \dots, z_{0,n}) \in \mathbb{C}^n$ and $R = (R_1, \dots, R_n) \in \mathbb{R}_{>0}^n$. If $f \in \mathcal{O}(\Omega)$ and $\overline{D(z_0, R)} \subset \Omega$, then for all $w \in D(z_0, R)$, we have

$$f(w) = \frac{1}{(2\pi i)^n} \int_{\Gamma(z_0, R)} \frac{f(z_1, \dots, z_n)}{(z_1 - w_1) \cdots (z_n - w_n)} dz_1 \cdots dz_n,$$

where $\Gamma(z_0, R) = \partial D(z_{0,1}, R_1) \times \cdots \times \partial D(z_{0,n}, R_n)$.

Proof. See [6, p. 22]. ■

注意这里 $\Gamma(z_0, R)$ 是所谓“特征边界”而不是实质上的边界.

COROLLARY 1.4 (Cauchy inequality). Suppose $f \in \mathcal{O}(\Omega)$ and $\overline{D(z_0, R)} \subset \Omega$. Then

$$|f^{(\alpha)}(z_0)| \leq \frac{\alpha!}{R^\alpha} \sup_{\Gamma(z_0, R)} |f|, \quad \forall \alpha \in \mathbb{Z}_{\geq 0}^n.$$

COROLLARY 1.5. We can write

$$f(z) = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} \frac{f^{(\alpha)}(z_0)}{\alpha!} (z - z_0)^\alpha, \quad \forall z \in D(z_0, R),$$

when $f \in \mathcal{O}(D(z_0, R))$, where the expansion is compactly (absolutely) convergent.

这也常作为全纯函数 (复解析函数) 的定义. 例如在 Ohsawa 的书 [14] 中.

COROLLARY 1.6 (Montel's theorem). Every locally uniformly bounded sequence (f_j) in $\mathcal{O}(\Omega)$ has a convergent subsequence.

这表明 $\mathcal{O}(\Omega)$ 为一个 Fréchet 空间.

Recall

$$\frac{\partial}{\partial z_k} = \frac{1}{2} \left(\frac{\partial}{\partial x_k} - i \frac{\partial}{\partial y_k} \right), \quad \frac{\partial}{\partial \bar{z}_k} = \frac{1}{2} \left(\frac{\partial}{\partial x_k} + i \frac{\partial}{\partial y_k} \right), \quad 1 \leq k \leq n,$$

and

$$\begin{aligned} \partial u &= \sum_{I,J} \sum_{1 \leq k \leq n} \frac{\partial u_{I,J}}{\partial z_k} dz_k \wedge dz_I \wedge d\bar{z}_J, \\ \bar{\partial} u &= \sum_{I,J} \sum_{1 \leq k \leq n} \frac{\partial u_{I,J}}{\partial \bar{z}_k} d\bar{z}_k \wedge dz_I \wedge d\bar{z}_J, \\ du &= \partial u + \bar{\partial} u, \end{aligned}$$

for the (p, q) form

$$u = \sum_{|I|=p, |J|=q} u_{I,J} dz_I \wedge d\bar{z}_J, \quad u_{I,J} \in C^s(\Omega, \mathbb{C}).$$

Then

$$d^2 = \partial^2 = \bar{\partial}^2 = 0, \quad \partial \bar{\partial} + \bar{\partial} \partial = 0,$$

and

$$\begin{aligned} f: \Omega \rightarrow \mathbb{C} \text{ holomorphic} &\Leftrightarrow \frac{\partial f}{\partial \bar{z}_k} = 0 \quad \forall 1 \leq k \leq n, \quad \Leftrightarrow \bar{\partial} f = 0 \\ &\Rightarrow \partial \bar{\partial} f = 0 \quad \Rightarrow \quad i \partial \bar{\partial} \log |f| = 0 \text{ on } \Omega \setminus f^{-1}(0). \end{aligned}$$

LEMMA 1.7 (Dolbeault-Grothendieck lemma). Let Ω be a neighborhood of 0 in \mathbb{C}^n and u be a (p, q) form on Ω of C^s such that $\bar{\partial} v = 0$, where $1 \leq s \leq \infty$.

1. If $q = 0$, then $v = \sum_{|I|=p} v_I(z) dz_I$ is a holomorphic p -form, i.e. a form whose coefficients are holomorphic functions.
2. If $q \geq 1$, there exists a neighborhood $\omega \subset \Omega$ of 0 and a $(p, q-1)$ form u of C^s such that $\bar{\partial} u = v$ on ω .

Proof. See [6, p. 28].

Use the **Koppelman formula**:

$$v(z) = \int_{\partial \omega} K_{\text{BM}}^{p,q}(z, \zeta) \wedge v(\zeta) + \bar{\partial}_z \int_{\omega} K_{\text{BM}}^{p,q-1}(z, \zeta) \wedge v(\zeta) + \int_{\omega} K_{\text{BM}}^{p,q}(z, \zeta) \wedge \bar{\partial} v(\zeta),$$

for ω with piecewise C^1 boundary and (p, q) form v of class C^1 on $\bar{\omega}$, where $K_{\text{BM}}^{p,q}(z, \zeta)$ is the component of $K_{\text{BM}}(z, \zeta)$ of type (p, q) in z and $(n - p, n - q - 1)$ in ζ , $K_{\text{BM}}(z, \zeta)$ is the pull back of k_{BM} by the map $\pi: \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}^n$, $(z, \zeta) \mapsto z - \zeta$, and

$$k_{\text{BM}}(z) = c_n \sum_{1 \leq j \leq n} (-1)^j \frac{\bar{z}_j dz_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_1 \wedge \dots \wedge \widehat{d\bar{z}_j} \wedge \dots \wedge d\bar{z}_n}{|z|^{2n}}$$

is the Bochner-Martinelli kernel,

$$c_n = (-1)^{\frac{n(n-1)}{2}} \frac{(n-1)!}{(2\pi i)^n}.$$

■

这个引理说明了 $\bar{\partial}$ - 方程总是局部可解的.

THEOREM 1.8 (Weierstrass preparation theorem). *Let g be a holomorphic function on a neighborhood of 0 in \mathbb{C}^n , such that $\frac{g(0, z_n)}{z_n^s}$ has a not zero finite limit at $z_n = 0$. For $r' = (r'_1, \dots, r'_{n-1})$ and r_n small enough, one can write*

$$g(u) = u(z)P(z', z_n),$$

where u is an invertible holomorphic function in a neighborhood of $\overline{\Delta^n(r', r_n)}$, and P is a Weierstrass polynomial in z_n , that is, a polynomial of the form

$$P(z', z_n) = z_n^s + a_1(z')z_n^{s-1} + \dots + a_s(z'), \quad a_k(0) = 0,$$

with holomorphic coefficients $a_k(z')$ on a neighborhood of $\overline{\Delta^{n-1}(r')}$ in \mathbb{C}^{n-1} .

Proof. See [6, p. 79].

■

COROLLARY 1.9 (Weierstrass division theorem). *Every bounded holomorphic function f on $D = \Delta(r', r_n)$ can be represented in the form*

$$f(z) = g(z)q(z) + R(z', z_n),$$

where q and R are holomorphic in D , $R(z', z_n)$ is a polynomial of degree $\leq s - 1$ in z_n , and

$$\sup_D |q| \leq C \sup |f|, \quad \sup_D |R| \leq C \sup |f|$$

for some constant $C \geq 0$ independent of f . The representation is unique.

Proof. See [6, p. 80].

■

COROLLARY 1.10. *The local ring*

$$\mathcal{O}_n = \mathcal{O}_{\mathbb{C}^n, 0} = \mathbb{C}\{z_1, \dots, z_n\}$$

of convergent power series is a Noetherian (i.e. every ideal \mathcal{I} of \mathcal{O}_n is finitely generated) UFD.

Proof. See [6, pp. 81–83].

■

1.2 Analytic set

DEFINITION 1.11. A subset $A \subset \Omega$ is said to be an analytic subset of Ω if A is **closed** and if for every point $x_0 \in A$ there exist a neighborhood U of x_0 and holomorphic functions $g_1, \dots, g_N \in \mathcal{O}(U)$ such that

$$A \cap U = \{z \in U : g_1(z) = \dots = g_N(z) = 0\}.$$

REMARK 1.12. 1. We can define the germ (A, x) by modulus of the equivalence: $A \sim B$ iff $A \cap V = B \cap V$ for $x \in V$ open neighborhood.

2. Let $\mathcal{I} = (g_1, \dots, g_N)$ be an ideal of \mathcal{O}_x . Then we can denote by $(V(\mathcal{I}), x)$ the zero variety by

$$V(\mathcal{I}) = \{z \in U : g_1(z) = \dots = g_N(z) = 0\}.$$

3. Let (A, x) be a germ of analytic set at x . Denote

$$\mathcal{I}_{A,x} = \{f \in \mathcal{O}_x : f(z) = 0, \forall z \in (A, x)\}.$$

Then $\mathcal{I}_{A,x}$ is an ideal of \mathcal{O}_x , and $\mathcal{I}_{V(\mathcal{I}),x} \supset \mathcal{I}$.

THEOREM 1.13 (Cartan 50'). For any analytic set A , the sheaf of ideals \mathcal{I}_A is a coherent analytic sheaf.

Proof. See [6, pp. 99–100].

Omit. Use the following local parametrization theorem. ■

THEOREM 1.14 (Local parametrization theorem). Let \mathcal{I} be a prime ideal of \mathcal{O} and let $A = V(\mathcal{I})$. Then we can choose the coordinates

$$(z'; z'') = (z_1, \dots, z_d; z_{d+1}, \dots, z_n)$$

such that:

The ring \mathcal{O}_n/I is a finite integral extension of \mathcal{O}_d ; let q be the degree of the extension and let $\delta(z') \in \mathcal{O}_d$ be the discriminant of the irreducible polynomial of a primitive element $u(z'') = \sum_{k>d} c_k z_k$. If Δ', Δ'' are polydiscs of sufficiently small radii r', r'' and if $r' \leq r''/C$ with C large, the projection map $\pi: A \cap (\Delta' \times \Delta'') \rightarrow \Delta'$ is a ramified covering with q sheets, whose ramification locus is contained in $S = \{z' \in \Delta' : \delta(z') = 0\}$. This means that

1. the open subset $A_S = A \cap ((\Delta' \setminus S) \times \Delta'')$ is a smooth d -dimensional manifold, dense in $A \cap (\Delta' \times \Delta'')$;
2. $\pi: A_S \rightarrow \Delta' \setminus S$ is a covering;
3. the fibers $\pi^{-1}(z')$ have exactly q elements if $z' \notin S$ and at most q if $z' \in S$.

Moreover, A_S is a connected covering of $\Delta' \setminus S$, and $A \cap (\Delta' \times \Delta'')$ is contained in a cone $\{|z''| \leq C|z'|\}$.

Proof. See [6, pp. 95–97]. ■

这个定理可以看作是解析版本的 Noether 正规化定理.

THEOREM 1.15 (Hilbert's Nullstellensatz). *For every ideal $\mathcal{I} \subset \mathcal{O}_n$,*

$$\mathcal{I}_{V(I),0} = \sqrt{\mathcal{I}},$$

where \sqrt{I} is the radical of \mathcal{I} , i.e. the set of germs $f \in \mathcal{O}_n$ s.t. some power $f^k \in \mathcal{I}$.

Proof. See [6, p. 97]. ■

DEFINITION 1.16. Let $A \subset \Omega$ be an analytic subset and $x \in A$. Then $x \in A$ is called a regular point of A if $A \cap U$ is a \mathbb{C} -analytic submanifold of U for some neighborhood U of x . Otherwise, x is said to be singular.

The corresponding subsets of A are denoted by A_{reg} and A_{sing} respectively.

DEFINITION 1.17. For an irreducible germ of analytic set (A, x) , define the dimension by

$$\dim(A, x) := \dim(A_{\text{reg}}, x).$$

If (A, x) has several irreducible components (A_l, x) , then

$$\dim(A, x) := \max\{\dim(A_l, x)\}.$$

If all the irreducible components of (A, x) have the same dimension, then (A, x) is said to be of pure dimension. Set $\text{codim}(A, x) := n - \dim(A, x)$.

THEOREM 1.18. A_{sing} is an analytic subset of A .

Proof. See [6, pp. 100–101].

(A Irreducible:) Let g_1, \dots, g_N be local generators of A in a neighborhood U of regular point. Then $A_{\text{sing}} \cap U$ is defined by

$$\det \left(\frac{\partial g_j}{\partial z_k} \right)_{j \in J, k \in K} = 0, \quad J \subset \{1, \dots, N\}, \quad K \subset \{1, \dots, n\}, \quad |J| = |K| = n - d,$$

where $d = \dim A$. ■

EXAMPLE 1.19. Let

$$A = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : z_1 z_2^2 = z_2 z_3^3 = 0\}.$$

Then $A = A_1 \cup A_2$, where

$$A_1 = \{z_1 = z_3 = 0\}, \quad A_2 = \{z_2 = 0\}.$$

Then $\dim A_1 = 1$, $\dim A_2 = 2 \Rightarrow \dim A = 2$. We also can see $A_{\text{sing}} = \{0\}$, and

$$\mathcal{I}_{A,0} = (z_1 z_2, z_2 z_3) = \sqrt{(z_1 z_2^2, z_2 z_3^3)}.$$

1.3 Plurisubharmonic function

DEFINITION 1.20. A function $u: \Omega \rightarrow [-\infty, +\infty)$ is said to be plurisubharmonic (psh for short) if

1. u is upper semicontinuous;
2. for every complex line $L \subset \mathbb{C}^n$, $u|_{\Omega \cap L}$ is subharmonic on $\Omega \cap L$.

We denote the set of psh functions on Ω by $\text{Psh}(\Omega)$.

PROPOSITION 1.21. • If $u_k \in \text{Psh}(\Omega)$ is decreasing, then $u = \lim u_k \in \text{Psh}(\Omega)$.

- If $u_1, \dots, u_p \in \text{Psh}(\Omega)$, and $\chi: \mathbb{R}^p \rightarrow \mathbb{R}$ is convex function s.t. $\chi(t_1, \dots, t_p)$ is non-decreasing in each t_j , then $\chi(u_1, \dots, u_p) \in \text{Psh}(\Omega)$. In particular,

$$u_1 + \dots + u_p, \max\{u_1, \dots, u_p\}, \log(e^{u_1} + \dots + e^{u_p}) \in \text{Psh}(\Omega).$$

- Let $u \in \text{Psh}(\Omega)$, and $u \not\equiv -\infty$ on each component of Ω . If (ρ_ε) is a family of smoothing kernels, then $u \star \rho_\varepsilon \in C^\infty \cap \text{Psh}(\Omega_\varepsilon)$, the family $(u \star \rho_\varepsilon)$ is non-decreasing in ε , and $\lim_{\varepsilon \rightarrow 0} u \star \rho_\varepsilon = u$ (from above).

Proof. See [6, p. 39]. ■

PROPOSITION 1.22. Let $(u_\alpha)_{\alpha \in I}$ be a family of upper semicontinuous functions $\Omega \rightarrow [-\infty, +\infty)$ which is locally uniformly bounded above, and $u := \sup_\alpha u_\alpha$.

The upper semicontinuous regularization of u is defined by:

$$u^*(z) := \lim_{\varepsilon \rightarrow 0} \sup_{B(z, \varepsilon)} u.$$

(**Choquet's lemma**) Then the family (u_α) has a countable subfamily $(v_j) = (u_{\alpha(j)})$ s.t. $v = \sup_\alpha v_j$ satisfies

$$v \leq u \leq u^* \leq u^* = v^*.$$

If all $u_\alpha \in \text{Psh}(\Omega)$, then $u^* \in \text{Psh}(\Omega)$ and $u = u^*$ a.e.

Proof. See [6, p. 38]. ■

EXAMPLE 1.23. If $f_1, \dots, f_q \in \mathcal{O}(\Omega)$ and $\alpha_1, \dots, \alpha_q \geq 0$, then

$$\log(|f_1|^{\alpha_1} + \dots + |f_q|^{\alpha_q}) \in \text{Psh}(\Omega).$$

$u \in \text{Psh}(\Omega)$ is called as having analytic singularities near $z \in \Omega$ if there exists a constant $c > 0$, holomorphic functions f_1, \dots, f_q near z , and a bounded function v near z , such that

$$u = \frac{c}{2} \log(|f_1|^2 + \dots + |f_q|^2) + v \quad \text{near } z.$$

REMARK 1.24. If $u \in C^2(\Omega, \mathbb{R})$, then

$$u \in \text{Psh}(\Omega) \iff \left(\frac{\partial^2 u}{\partial z_j \partial \bar{z}_k} \right)_{1 \leq j, k \leq n} \text{ is semi-positive.}$$

If the matrix is (strictly) positive definite at every point, then u is called strictly psh and denote by $u \in \text{SPsh}(\Omega)$.

PROPOSITION 1.25. *If $\Omega = \omega + i\omega'$, where ω, ω' are open subsets of \mathbb{R}^n , and $u \in \text{Psh}(\Omega)$ which depends only on $x = \text{Re } z$, then $\omega \ni x \mapsto u(x)$ is convex.*

As a corollary, if $u \in \text{Psh}(D(a, R)) \subset \mathbb{C}^n$, then

$$\begin{aligned}\mu(u; r_1, \dots, r_n) &= \frac{1}{(2\pi)^n} \int_0^{2\pi} \cdots \int_0^{2\pi} u(a_1 + r_1 e^{i\theta_1}, \dots, a_n + r_n e^{i\theta_n}) d\theta_1 \cdots d\theta_n, \\ m(u; r_1, \dots, r_n) &= \sup_{z \in D(a, r)} u(z_1, \dots, z_n),\end{aligned}$$

are convex functions of $(\log r_1, \dots, \log r_n)$ that are non-decreasing in each variable.

Proof. See [6, p. 41].

Let

$$\begin{aligned}\tilde{\mu}(u; z_1, \dots, z_n) &= \frac{1}{(2\pi)^n} \int_0^{2\pi} \cdots \int_0^{2\pi} u(a_1 + e^{z_1} e^{i\theta_1}, \dots, a_n + e^{z_n} e^{i\theta_n}) d\theta_1 \cdots d\theta_n, \\ \tilde{m}(u; z_1, \dots, z_n) &= \sup_{|w_j| \leq 1} u(a_1 + e^{z_1} w_1, \dots, a_n + e^{z_n} w_n).\end{aligned}$$

■

1.4 Pseudoconvexity

DEFINITION 1.26. Let Ω be an domain of \mathbb{C}^n . Ω is said to be

- a domain of holomorphy, if every domain Ω' with $\Omega \subsetneq \Omega'$, there exists $F \in \mathcal{O}(\Omega)$, which can not be extended to be a holomorphic function on Ω' ;
- holomorphically convex, if $\hat{K}_\Omega \Subset \Omega$ for every compact subset $K \subset X$, where the holomorphic hull

$$\hat{K}_\Omega := \{z \in \Omega: |f(z)| \leq \sup_K |f|, \forall f \in \mathcal{O}(\Omega)\}.$$

如果 Ω 是全纯凸域, 那我们可以取 Ω 的紧穷竭:

$$K_1 \Subset K_2 \Subset \dots \Subset K_j \Subset \dots \Subset \Omega, \quad \bigcup_{j=1}^{\infty} K_j = \Omega,$$

并使得每个 K_j 满足 $\widehat{(K_j)}_\Omega = K_j$.

REMARK 1.27. • Any domain in \mathbb{C} is a domain of holomorphy.

- (Hartogs phenomenon) There exists a domain in \mathbb{C}^n which is not a domain of holomorphy when $n > 1$.

$$\Omega = \left\{ (z_1, z_2): |z_1| < 1, \frac{1}{2} < |z_2| < 1 \right\} \cup \left\{ (z_1, z_2): \frac{1}{2} < |z_1| < 1, |z_2| < 1 \right\}.$$

In fact, when $n > 1$, every holomorphic function $f \in \mathcal{O}(\Omega \setminus K)$, can be extended to be an element in $\mathcal{O}(\Omega)$, where K is a compact subset of Ω s.t. $\Omega \setminus K$ is connected (Hartogs).

THEOREM 1.28 (Cartan-Thullen theorem). *TFAE:*

1. Ω is a domain of holomorphy;
2. Ω is holomorphically convex;
3. There exists a function $F \in \mathcal{O}(\Omega)$ which is unbounded on any neighborhood of any point of $\partial\Omega$.

Proof. See [6, pp. 48–49]. ■

DEFINITION 1.29. $\psi: \Omega \rightarrow [-\infty, +\infty)$ is said to be an exhaustion if every sublevel set

$$\Omega_c := \{z \in \Omega: \psi(z) < c\} \Subset \Omega, \quad \forall c \in \mathbb{R}.$$

Ω is said to be

1. weakly pseudoconvex if there exists a smooth psh exhaustion function $\psi \in C^\infty(\Omega) \cap \text{Psh}(\Omega)$;
2. strongly pseudoconvex if there exists a smooth strictly psh exhaustion function $\psi \in C^\infty(\Omega) \cap \text{SPsh}(\Omega)$.

THEOREM 1.30. *TFAE:*

1. Ω is weakly pseudoconvex;
2. Ω is strongly pseudoconvex;
3. Ω has a psh exhaustion function ψ ;
4. $-\log d(z, \Omega^c)$ is psh on Ω .

Proof. See [6, pp. 54–55]. ■

THEOREM 1.31. $\Omega \subset \mathbb{C}^n$ domain. Then

$$\Omega \text{ holomorphically convex} \iff \Omega \text{ pseudoconvex}.$$

Proof. See [6, pp. 49–50] and [6, Chapter VIII] (by the theory of L^2 estimates for $\bar{\partial}$). ■

拟凸推全纯凸即是经典的 Levi 问题 (Oka, Norguet, Bremermann). 这些等价性对一般的复流形并不对, 不过全纯凸推弱拟凸是直接的, 而强拟凸实际上等价于为 Stein 流形 (全纯凸 + 局部可分离), 其中强拟凸推 Stein 也被称为 Levi 问题 (由 Grauert 完全解决).

EXAMPLE 1.32. Let $\Omega = \Delta^* \times \Delta \subset \mathbb{C}^2$, where $\Delta^* = \Delta \setminus \{0\}$. Then

$$\psi(z) = \max \{ -\log |z_1|, -\log(1 - |z_1|^2), -\log(1 - |z_2|^2) \} \in \text{Psh}(\Omega),$$

and $\psi(z) \rightarrow +\infty$ as $z \rightarrow \partial\Omega$.

THEOREM 1.33 (Kiselman's Minimum Principle). *Let $\Omega \subset \mathbb{C}^p \times \mathbb{C}^n$ be a pseudoconvex open set s.t. each slice*

$$\Omega_\zeta = \{z \in \mathbb{C}^n : (\zeta, z) \in \Omega\}, \quad \zeta \in \mathbb{C}^p,$$

is a convex tube $\omega_\zeta + i\mathbb{R}^n$, $\omega_\zeta \subset \mathbb{R}^n$. If $v(\zeta, z) \in \text{Psh}(\Omega)$ does not depend on $\text{Im } z$, then

$$u(\zeta) = \inf_{z \in \Omega_\zeta} v(\zeta, z) \in \text{Psh}(\Omega'), \quad \Omega' := \text{pr}_{\mathbb{C}^p}(\Omega),$$

or locally $\equiv -\infty$.

Proof. See [6, p. 57]. ■

如果没有 v 与虚部无关的条件, 这个结论是不对的.

1.5 Pluripolar set

DEFINITION 1.34. • A set $A \subset \Omega$ is said to be (locally) pluripolar if for every $z \in \Omega$, there exists a neighborhood U of x and $u \in \text{Psh}(U)$, $u \not\equiv -\infty$, such that $A \cap U \subset \{x \in U : u(x) = -\infty\}$;

- A set $A \subset \Omega$ is said to be complete pluripolar if for every $z \in \Omega$, there exists a neighborhood U of x and $u \in \text{Psh}(\Omega) \cap L^1_{\text{loc}}(U)$, such that $A \cap U = \{x \in U : u(x) = -\infty\}$;
- $A \subset \mathbb{C}^n$ is globally pluripolar if $A \subset u^{-1}(-\infty)$ for some $u \in \text{Psh}(\mathbb{C}^n)$;
- $A \subset \Omega$ is negligible if there is a family of psh functions $\{u_\alpha\}$ on Ω locally uniformly bounded from above such that $A \subset \{u < u^*\}$, where $u = \sup_\alpha u_\alpha$.

REMARK 1.35. • *Any analytic subset is a (complete) pluripolar set. If we take $A \cap U = \{g_1 = \dots = g_N = 0\}$, then $A \cap U = u^{-1}(-\infty)$, where $u = \log(|g_1|^2 + \dots + |g_N|^2)$.*

注意 analytic set 一定是 pluripolar set, 但反过来不一定, 实际上 analytic set 一定是无处稠密的, 但 pluripolar set 却可能是稠密的.

EXAMPLE 1.36. Let $\Omega = \Delta \subset \mathbb{C}$, and $\psi \in \text{Psh}(\Delta)$ as

$$\psi(z) = \sum_{k=1}^{\infty} \frac{1}{k^2} \log \left| z - \frac{1}{2^k} \right|.$$

Then $A = \psi^{-1}(-\infty) = \{0\} \cup \{1/2^k : k \geq 1\}$ is a (complete) pluripolar set, but A is not an analytic subset since 0 is a cluster point in A .

If we let

$$\phi(z) = \sum_{k=1}^{\infty} \frac{1}{k^3} \log \left| z - \frac{1}{2^k} \right|,$$

then $A = \phi^{-1}(-\infty) = \{1/2^k : k \geq 1\}$ is a pluripolar set which is not closed.

PROPOSITION 1.37. *Let $A \subset \Omega$ be a closed pluripolar subset. Then:*

1. *Every $v \in \text{Psh}(\Omega \setminus A)$ that is locally bounded above near A extends uniquely into a function $\tilde{v} \in \text{Psh}(\Omega)$;*
2. *Every $f \in \mathcal{O}(X \setminus A)$ that is locally bounded near A extends to a function $f \in \mathcal{O}(\Omega)$.*

Proof. See [6, pp. 44–45].

May assume $u \leq 0$ on W where $A \cap W \subseteq u^{-1}(-\infty)$. Let $v_\varepsilon = v + \varepsilon u$ and $v_\varepsilon = -\infty$ on $A \cap W$. Then $\tilde{v} = (\sup v_\varepsilon)^*$.

Apply (1) to $\pm \operatorname{Re} f$ and $\pm \operatorname{Im} f$. ■

THEOREM 1.38. *Let $A \subset \mathbb{C}^n$. Then A is*

$$\text{locally pluripolar} \iff \text{globally pluripolar} \iff \text{negligible}.$$

Proof. See [3, Chapter 3.1]. ■

2 L^2 Methods and Multiplier Ideal Sheaf

2.1 L^2 estimate of $\bar{\partial}$ -equation

We only focus on $(n, 0)$ and $(n, 1)$ forms in this section.

Let Ω be a domain in \mathbb{C}^n , and $\phi \in C^2 \cap \text{Psh}(\Omega)$. We can see that $(L, h) := (\Omega \times \mathbb{C}, e^{-\phi})$ is a trivial line bundle with a singular hermitian metric $h := e^{-\phi}$ on Ω . Let $\omega = \frac{i}{2} \sum_i dz_i \wedge d\bar{z}_i$ and $dV = \frac{\omega^n}{n!}$. Then

$$L_{(2)}^{n,q}(\Omega, h) := \left\{ u \in \wedge^{n,q}(\Omega) : \int_{\Omega} |u|_h^2 dV < +\infty \right\}$$

is a Hilbert space, where

$$|u|_h^2 dV := 2^q \left(\sum_{|J|=q} |u_J|^2 e^{-\phi} \right) dz \wedge d\bar{z},$$

for

$$u(z) = \sum_{|J|=q} u_J dz \wedge d\bar{z}_J, \quad \text{where} \quad dz = dz_1 \wedge \cdots \wedge dz_n.$$

我们也可以将其视为光滑紧支形式在 (加权) L^2 范数下的完备化.

It can be seen that the operator:

$$\bar{\partial}: L_{(2)}^{n,q-1}(\Omega, h) \supset \mathcal{D}^{n,q-1}(\Omega) \longrightarrow \mathcal{D}^{n,q}(\Omega) \subset L_{(2)}^{n,q}(\Omega, h)$$

is a densely defined closed operator, which makes it possible to extend $\bar{\partial}$, and define its adjoint. Let $\text{Dom}(\bar{\partial}) \subset L_{(2)}^{n,q-1}(\Omega, h)$ be its domain and $\bar{\partial}_h^*$ be its adjoint operator.

注意如果考虑的是 (n, q) 形式, 所以 ∂ 算子作用上去都是 0. 我们计算一下 $\bar{\partial}_h^*$ 的表达式: ($q = 1$ 情形) 设

$$U = u dz, \quad V = \sum_j v^j d\bar{z}^j \wedge dz,$$

则

$$\bar{\partial}U = \sum_j u_{\bar{j}} d\bar{z}^j \wedge dz.$$

可得

$$(\bar{\partial}U, V) = 2 \int_{\Omega} \sum_j u_{\bar{j}} \bar{v}^j e^{-\phi} = -2 \int_{\Omega} \sum_j u \left(\bar{v}^j e^{-\phi} \right)_{\bar{j}} = -2 \int_{\Omega} \sum_j u \left(\bar{v}^j_{\bar{j}} e^{-\phi} - \bar{v}^j \phi_{\bar{j}} e^{-\phi} \right),$$

\Rightarrow

$$\bar{\partial}_h^* V = \sum_j (\phi_j v^j - v^j_{\bar{j}}) dz = \sum_j \left(\frac{\partial \phi}{\partial z_j} v^j - \frac{\partial v^j}{\partial z_j} \right) dz.$$

THEOREM 2.1 (Bochner-Kodaira-Nakano inequality). For $u \in \mathcal{D}^{n,q-1}(\Omega)$, one has

$$\|\bar{\partial}u\|^2 + \|\bar{\partial}_h^* u\|^2 \geq (i\Theta_h \Lambda u, u),$$

where Θ_h is the curvature form $i\partial\bar{\partial}\phi$, and $\Lambda = \frac{-i}{2} \sum_j (d\bar{z}_j)^*(dz_j)^*$:

$$(i\Theta_h \Lambda u, u) = 2^q \int_{\Omega} \sum_{|K|=q-1} \sum_{1 \leq i, j \leq n} \phi_{i\bar{j}} u_{K,i} \overline{u_{K,j}} e^{-\phi} dz \wedge d\bar{z}, \quad \phi_{i\bar{j}} = \frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_i}.$$

Proof. See [6, pp. 329–332]. ■

特别地, 如果 $q = 1$:

$$(\mathrm{i}\Theta_h \Lambda u, u) = 2 \int_{\Omega} \sum_{1 \leq i, j \leq n} \phi_{i\bar{j}} u_i \bar{u}_j e^{-\phi} dz \wedge d\bar{z}.$$

在一般的 Kähler 流形上的向量丛 (F, h) 上, 对任意 F - 值的光滑紧支 (p, q) 形式, 有 Bochner-Kodaira-Nakano 等式:

$$\bar{\square} = \square + [\mathrm{i}\Theta_{F,h}, \Lambda].$$

并由此可得

$$\|\bar{\partial}u\|^2 + \|\bar{\partial}_h^* u\|^2 - \|\partial^* u\|^2 - \|\partial_h u\|^2 = ([\mathrm{i}\Theta_h, \Lambda]u, u).$$

为了计算方便, 我们再考虑一个更简单的情形: $n = 1$. 这时若 $V = v d\bar{z} \wedge dz$,

$$\bar{\square}V = \bar{\partial}\bar{\partial}_h^* V = \bar{\partial}((\phi_z v - v_z) dz) = (\phi_{z\bar{z}} v + \phi_z v_{\bar{z}} - v_{z\bar{z}}) d\bar{z} \wedge dz.$$

\Rightarrow

$$\|\bar{\partial}_h^* V\|^2 = (\bar{\square}V, V) = 2 \int_{\Omega} (\phi_{z\bar{z}} v \bar{v} + \phi_z v_{\bar{z}} \bar{v} - v_{z\bar{z}} \bar{v}) e^{-\phi} dz \wedge d\bar{z},$$

其中

$$\begin{aligned} \int_{\Omega} -v_{z\bar{z}} \bar{v} e^{-\phi} dz \wedge d\bar{z} &= \int_{\Omega} v_{\bar{z}} (\bar{v} e^{-\phi})_z dz \wedge d\bar{z} \\ &= \int_{\Omega} v_{\bar{z}} (\bar{v}_{\bar{z}} - \phi_z \bar{v}) e^{-\phi} dz \wedge d\bar{z} \\ &= \|v_{\bar{z}}\|^2 - \int_{\Omega} v_{\bar{z}} \phi_z \bar{v} e^{-\phi} dz \wedge d\bar{z}. \end{aligned}$$

从而

$$\|\bar{\partial}_h^* V\|^2 = 2 \int_{\Omega} \phi_{z\bar{z}} v \bar{v} e^{-\phi} dz \wedge d\bar{z} + \|v_{\bar{z}}\|^2 = 2([\mathrm{i}\Theta_h, \Lambda]V, V) + \|\partial^* V\|^2.$$

值得注意的是, 以上计算过程实际上并不要求 $\mathrm{i}\partial\bar{\partial}\phi$ 的正性.

LEMMA 2.2. *If Ω is pseudoconvex, then $\mathcal{D}^{n,q}(\Omega)$ is dense in $\mathrm{Dom}(\bar{\partial}^*)$ for the graph norm*

$$u \longmapsto \|u\| + \|\bar{\partial}_h^* u\|.$$

Proof. See [6, pp. 368–370]. ■

Consider two closed and densely defined operators T, S among Hilbert spaces:

$$\mathcal{H}_1 \xrightarrow{T} \mathcal{H}_2 \xrightarrow{S} \mathcal{H}_2$$

with $S \circ T = 0$.

LEMMA 2.3. *In order that $\mathrm{Im} T = \mathrm{Ker} S$, it suffices that*

$$\|T^*x\|_1^2 + \|Sx\|_3^2 \geq C\|x\|_2^2, \quad \forall x \in \mathrm{Dom} S \cap \mathrm{Dom} T^*,$$

for some constant $C > 0$. In that case, for every $v \in \mathcal{H}_2$ s.t. $Sv = 0$, there exists $u \in \mathcal{H}_1$ s.t. $Tu = v$ and

$$\|u\|_1^2 \leq \frac{1}{C} \|v\|_2^2.$$

Proof. See [6, pp. 364–365].

Write $x \in \text{Dom } T^*$ as $x = x' + x''$ with $x' \in \text{Ker } S$ and $x'' \in (\text{Ker } S)^\perp \subset \text{Ker } T^*$. If $Sv = 0$, then $\langle v, x'' \rangle_2 = 0$. Note that $Sx' = T^*x'' = 0$. We can get

$$|\langle v, x \rangle_2|^2 = |\langle v, x' \rangle_2|^2 \leq \|v\|_2^2 \|x'\|_2^2 \leq \frac{1}{C} \|v\|_2^2 \|T^*x'\|_2^2 = \frac{1}{C} \|v\|_2^2 \|T^*x\|_1^2.$$

Finally, use the Hahn-Banach theorem and Riesz representation theorem to the operator:

$$\begin{aligned} \mathcal{L}: \text{Im}(T^*) &\longrightarrow \mathbb{C} \\ y = T^*x &\longmapsto \langle x, v \rangle_2. \end{aligned}$$

■

COROLLARY 2.4. Let $\Omega \subseteq \mathbb{C}^n$ be a pseudoconvex domain. If the operator $A := i\Theta_h \Lambda$ is positive definite everywhere on $\wedge^{n,q}(\Omega)$, $q \geq 1$, then for every form $v \in L_{(2)}^{n,q}(\Omega, h)$ satisfying $\bar{\partial}v = 0$ and $\int_\Omega (A^{-1}v, v) dV < +\infty$, there exists $u \in L_{(2)}^{n,q-1}(\Omega, h)$ such that $\bar{\partial}u = v$ and

$$\int_\Omega |u|_h^2 dV \leq \int_\Omega (A^{-1}v, v) dV.$$

Proof. See [6, p. 371].

■

特别地, 如果 $q = 1$:

$$(A^{-1}v, v) = 2 \int_\Omega \sum_{1 \leq i, j \leq n} \phi^{i\bar{j}} v_j \bar{v}_j e^{-\phi} dz \wedge d\bar{z},$$

其中 $(\phi^{i\bar{j}}) = (\phi_{i\bar{j}})^{-1}$ 是 ϕ 的 Hessian 矩阵的逆矩阵.

如果 $i\partial\bar{\partial}\phi \geq \varepsilon\omega$ 对某个 $\varepsilon > 0$ 成立, 也就是 $\phi - \frac{\varepsilon}{2}|z|^2 \in \text{Psh}(\Omega)$ 或者说

$$\sum_{i,j} \phi_{i\bar{j}} v_j \bar{v}_j \geq \varepsilon \sum_j |v_j|^2,$$

则

$$\int_\Omega |u|_h^2 dV \leq \frac{1}{q\varepsilon} \int_\Omega |v|_h^2 dV.$$

COROLLARY 2.5. Let $\Omega \subseteq \mathbb{C}^n$ be a bounded pseudoconvex domain, and $\varphi \in L_{\text{loc}}^1(\Omega \rightarrow [-\infty, +\infty))$ satisfying $i\partial\bar{\partial}\varphi \geq -A\omega$ on Ω for some constant $A > 0$. Then for every $(0, 1)$ form $v = \sum_{j=1}^n v_j d\bar{z}_j$ satisfying $\bar{\partial}v = 0$ and

$$\int_\Omega \left(\sum_{j=1}^n |v_j|^2 \right) e^{-\varphi} d\lambda < +\infty,$$

there exists a function $u \in \text{Dom}(\bar{\partial})$ on Ω such that $\bar{\partial}u = v$ and

$$\int_\Omega |u|^2 e^{-\varphi} d\lambda \leq C_{A,\Omega} \int_\Omega \left(\sum_{j=1}^n |v_j|^2 \right) e^{-\varphi} d\lambda,$$

where $C_{A,\Omega} > 0$ only depends on A and the diameter of Ω .

If v is smooth, then u can be chosen as smooth function.

Proof. Take $h = e^{-\varphi - A'|z|^2}$ for $A' > A$, and transform (n, q) forms into $(0, q)$ forms. \blacksquare

THEOREM 2.6. *Let $\varphi \in \text{Psh}(\Omega)$, where $\Omega \subset \mathbb{C}^n$ is bounded and pseudoconvex. If $z_0 \in \Omega$ and $e^{-\varphi}$ is integrable near z_0 (e.g. $\varphi(z_0) > -\infty$), then there exists $f \in \mathcal{O}(\Omega)$ such that $f(z_0) = 1$ and*

$$\int_{\Omega} |f|^2 e^{-\varphi} d\lambda < +\infty.$$

Proof. Let θ be a cut-off function such that $\theta \equiv 1$ and supported near z_0 . Let $v = \bar{\partial}(1 - \theta)$, and since $e^{-\varphi}$ is integral near z_0 we can take θ such that

$$\int_{\Omega} |v|^2 e^{-\varphi - 2n \log |z - z_0| - |z|^2} d\lambda < +\infty.$$

Then we can get a solution u of $\bar{\partial}u = v = -\bar{\partial}\theta$ with $(\Omega$ is bounded)

$$\int_{\Omega} |u|^2 e^{-\varphi - 2n \log |z - z_0|} d\lambda < +\infty.$$

Let $f = u + \theta$. Then we have $f \in \mathcal{O}(\Omega)$, $u(z_0) = 0 \Rightarrow f(z_0) = \theta(z_0) = 1$, and $\int_{\Omega} |f|^2 e^{-\varphi} d\lambda < +\infty$. \blacksquare

COROLLARY 2.7. *Let $\varphi \in \text{Psh}(\Omega)$. Then*

$$V(\mathcal{I}(\varphi)) := \{z \in \Omega : e^{-\varphi} \text{ is integrable near } z\}$$

is an analytic subset.

Proof. See [6, p. 384].

For every $x \in \Omega$, let U be a small bounded pseudoconvex neighborhood of x in Ω . Denote $A^2(U, e^{-\varphi}) := L^2(U, e^{-\varphi}) \cap \mathcal{O}(U)$. Then

$$V(\mathcal{I}(\varphi)) = \bigcap_{f \in A^2(U, e^{-\varphi})} f^{-1}(0).$$

Then the result follows from the coherence of $\mathcal{I}(\varphi)$ or the fact that *the intersection of a family of analytic subsets is an analytic subset* (Noetherian). \blacksquare

COROLLARY 2.8 (Levi's problem). *A bounded pseudoconvex domain $\Omega \subseteq \mathbb{C}^n$ is a domain of holomorphy.*

Proof. See [6, pp. 373–374]

Since Ω is pseudoconvex, we have $-\log \delta(z) \in \text{Psh}(\Omega)$.

For every $z_0 \in \partial\Omega$, select a sequence of pts $\{z_j\} \subset \Omega$ s.t. $z_j \rightarrow z_0$. For each j , let θ_j be a cut-off function with $\text{Supp } \theta_j \Subset V_j$, where V_j is a sufficiently small neighborhood of z_j and disjoint with each other.

Let

$$\psi(z) = \chi\left(-\log \delta(z) + |z|^2\right) + \sum_{j=1}^{\infty} 2n\theta_j \log |z - z_j| \in \text{Psh}(\Omega), \quad (2.1)$$

where $\delta(z) = \text{dist}(z, \Omega^c)$, and χ is an increasing convex function undetermined.

Let

$$v = \bar{\partial} \left(\sum_{j=1}^{\infty} j \theta_j \right).$$

Suppose

$$\int_{\Omega} |v|^2 e^{-\psi} d\lambda < +\infty. \quad (2.2)$$

We solve $\bar{\partial}u = v$ with L^2 estimates

$$\int_{\Omega} |u|^2 e^{-\psi} d\lambda < +\infty,$$

and let $f = -u + \sum_{j=1}^{\infty} j \theta_j \in \mathcal{O}(\Omega)$. Then $u(z_j) = 0 \Rightarrow f(z_j) = j \Rightarrow f$ can not be extended across z_0 .

Then we only need to find χ s.t. (2.1) and (2.2) hold. Note that $\theta_j \equiv 1$ near z_j and $i\bar{\partial}\bar{\partial}(\theta_j \log |z - z_j|)$ is bounded outside $\{\theta_j \equiv 1\}$. If we take χ increasing so fast, then (2.1) is OK. It also works on (2.2). \blacksquare

2.2 L^2 extension theorem

Cartan B theorem tells us that the extension of a holomorphic function from an analytic subset to a Stein manifold always exists.

THEOREM 2.9. *Let Ω be a pseudoconvex domain in \mathbb{C}^n s.t. $\sup_{z \in \Omega} |z_n| \leq 1$, let $\varphi \in \text{Psh}(\Omega)$, and let $H = \{z_n = 0\}$. Then for every $f \in \mathcal{O}(\Omega \cap H)$ with*

$$\int_{H \cap \Omega} |f|^2 e^{-\varphi} d\lambda_H < +\infty,$$

one can find a holomorphic extension F of f to Ω satisfying

$$\int_{\Omega} |F|^2 e^{-\varphi} d\lambda \leq \mathbf{C} \pi \int_{\Omega \cap H} |f|^2 e^{-\varphi} d\lambda_H.$$

Here \mathbf{C} can be chosen as: $\mathbf{C} = 1620$ (Takegoshi-Ohsawa), ... , $\mathbf{C} = 1$ (Blocki, Guan-Zhou).

THEOREM 2.10 (twisted BKN identity, Ohsawa–Takegoshi 87'). *For $u \in \mathcal{D}^{n,q}(\Omega)$,*

$$\|\sqrt{\eta} \bar{\partial} u\|^2 + \|\sqrt{\eta + c^{-1}} \bar{\partial}_h^* u\|^2 \geq \left(i(-\bar{\partial} \bar{\partial} \eta - c \partial \eta \wedge \bar{\partial} \eta + \eta \Theta_h) \Lambda u, u \right),$$

where $0 < \eta \in C^\infty(\Omega)$ and $0 < c \in C(\Omega)$ are bounded.

Therefore, if we assume the operator

$$\mathbf{B} := i(-\bar{\partial} \bar{\partial} \eta - c_1 \partial \eta \wedge \bar{\partial} \eta + \eta \Theta_h) \Lambda$$

is positive definite everywhere on $\wedge^{n,q}(\Omega)$, Then, for every $v = \mathcal{D}^{n,q}(\Omega)$ with $\bar{\partial} v = 0$ and

$$\left(\mathbf{B}^{-1} v, v \right) < +\infty,$$

we can get a solution of $\bar{\partial} w = v$ satisfying

$$\left\| (\eta + c_1^{-1})^{-1/2} w \right\|^2 \leq \left(\mathbf{B}^{-1} v, v \right).$$

Proof. See [15, pp. 115–117].

我们来算一下 $n = q = 1$ 情形: 令 $V = v d\bar{z} \wedge dz \in \mathcal{D}^{1,1}(\Omega)$, 其中 $\Omega \subset \mathbb{C}$, 回忆 $\bar{\partial}_h^* V = \phi_z v - v_z$ 在这种简单情形下, 则

$$\|\sqrt{\eta} \bar{\partial}_h^* V\|^2 = (\bar{\partial}(\eta \bar{\partial}_h^* V), V)_h = 2 \int_{\Omega} \left(\eta_{\bar{z}} \phi_z v \bar{v} + \eta \phi_{z\bar{z}} v \bar{v} + \eta \phi_z v_{\bar{z}} \bar{v} - \eta_{\bar{z}} v_z \bar{v} - \eta v_{z\bar{z}} \bar{v} \right) e^{-\phi},$$

其中

$$\begin{aligned} \int_{\Omega} -\eta v_{z\bar{z}} \bar{v} e^{-\phi} dz \wedge d\bar{z} &= \int_{\Omega} \left(v_{\bar{z}} \eta_z \bar{v} + \eta v_{\bar{z}} \overline{(v_{\bar{z}})} - v_{\bar{z}} \eta \phi_z \bar{v} \right) e^{-\phi} \\ &= \int_{\Omega} \left(-\eta \phi_z v_{\bar{z}} \bar{v} + \eta_z v_{\bar{z}} \bar{v} \right) e^{-\phi} + \|\sqrt{\eta} \partial^* V\|^2, \end{aligned}$$

以及

$$\int_{\Omega} \eta_z v_{\bar{z}} \bar{v} e^{-\phi} = \int_{\Omega} \left(-\eta_{z\bar{z}} v \bar{v} - \eta_z v \overline{(v_z)} + \eta_z \phi_{\bar{z}} v \bar{v} \right) e^{-\phi}.$$

从而

$$\begin{aligned} \|\eta \bar{\partial}_h^* V\|^2 &\geq 2 \int_{\Omega} \left(\eta \phi_{z\bar{z}} v \bar{v} - \eta_{z\bar{z}} v \bar{v} \right) e^{-\phi} \\ &\quad + 2 \int_{\Omega} \left(\eta_{\bar{z}} \phi_z v \bar{v} + \eta_z \phi_{\bar{z}} v \bar{v} - \eta_{\bar{z}} v_z \bar{v} - \eta_z v \overline{(v_z)} \right) e^{-\phi} \\ &= \dots + 4 \cdot \operatorname{Re} \int_{\Omega} \left(\eta_{\bar{z}} \bar{v} (\phi_z v - v_z) \right) e^{-\phi} \\ &= \dots + 4 \cdot \operatorname{Re} \left(\bar{\partial}_h^* V, \partial \eta \wedge (\Delta V) \right)_h \\ &\geq \dots - \|\sqrt{c^{-1}} \bar{\partial}_h^* V\|^2 - \|\sqrt{c} \partial \eta \wedge (\Delta V)\|^2, \end{aligned}$$

这里用到了很简单的不等式:

$$2\operatorname{Re}(AB) \geq -\frac{1}{C}|A|^2 - C|B|^2, \quad C > 0, \quad A, B \in \mathbb{C}.$$

最后我们得到

$$\|\sqrt{\eta + c^{-1}} \bar{\partial}_h^* V\|^2 \geq \left(i(\eta \Theta_h - \partial \bar{\partial} \eta - c \partial \eta \wedge \bar{\partial} \eta) \Delta V, V \right).$$

■

Proof of Theorem 2.9. We may assume Ω is bounded and φ is smooth up to the boundary of D first (This is because we can approximate Ω with relatively compact pseudoconvex Ω_j and approximate φ with smooth psh φ_k).

Let $\tilde{f} = \operatorname{pr}^*(f dz_1 \wedge \dots \wedge dz_{n-1})$ be the constant extension, and θ a cut-off function such that $\theta = 1$ near $\Omega \cap H$. For example, $\theta = \alpha(\log |z_n|^2)$, where (fix $t < 0$ first)

$$\alpha(x) = \begin{cases} 1 & x \in (-\infty, -t-1), \\ 0 & x \in (-t, 0). \end{cases}$$

Let $v = \bar{\partial}(-\theta \tilde{f}) = -\tilde{f} \bar{\partial} \theta$ and let $\eta, c > 0$ and $\phi \in \operatorname{Psh}(\Omega)$ with

$$\eta(z) = s(-\beta(\psi)), \quad \phi(z) = u(-\beta(\psi)), \quad c(z) = r(-\beta(\psi)),$$

where $\psi = \log |z_n|^2$, such that

1. β is smooth, increasing and convex, $\beta(x) = x$ for x far from $-\infty$, and $\beta(x) = \text{constant}$ near $-\infty$, that is

$$\beta(x) = \begin{cases} x & x \in (-t, 0), \\ \text{constant} & x \in (-\infty, -t-1). \end{cases}$$

For example

$$\beta(x) = \int_0^x (1 - \alpha(y)) dy;$$

2. and $(h = e^{-\psi-\phi-\varphi})$ here need not be positive definite) for some μ ,

$$i(\eta\Theta_h - \partial\bar{\partial}\eta - c\partial\eta \wedge \bar{\partial}\eta) \geq ic_2\partial\mu \wedge \bar{\partial}\mu > 0.$$

Then we can solve $\bar{\partial}w = v = \bar{\partial}(-\theta\tilde{f})$ with L^2 estimates and let $\tilde{F} = w + \theta\tilde{f} \in \mathcal{O}(\Omega)$.

Direct calculation: ($\Delta = \beta(\psi)$ for simplicity)

$$\partial\bar{\partial}\eta = -s'(-\Delta)\partial\bar{\partial}(\Delta) + s''(-\Delta)\partial(\Delta) \wedge \bar{\partial}(\Delta),$$

and

$$\partial\bar{\partial}\phi = -u'(-\Delta)\partial\bar{\partial}(\Delta) + u''(-\Delta)\partial(\Delta) \wedge \bar{\partial}(\Delta),$$

$$(i\partial\bar{\partial}(\psi + \varphi) \geq 0) \Rightarrow$$

$$\begin{aligned} & i(\eta\Theta_h - \partial\bar{\partial}\eta - c\partial\eta \wedge \bar{\partial}\eta) \\ & \geq i(u''s - s'' - rs'^2)(-\Delta)\partial(\Delta) \wedge \bar{\partial}(\Delta) + i(s' - su')(-\Delta)\partial\bar{\partial}(\Delta) \\ & = \dots + (s' - su')(-\Delta) \left(\beta'(\psi)i\partial\bar{\partial}\psi + \beta''(\psi)i\partial\psi \wedge \bar{\partial}\psi \right). \end{aligned}$$

We hope:

$$(a) \quad u''s - s'' - rs'^2 = 0;$$

$$(b) \quad s' - su' = 1.$$

May assume $\beta'' \approx 1$ and $-\alpha' \approx 1$ on an interval with length 1 (e.g. $\alpha'(x) = -1$ on $x \in (-t-1 + \varepsilon, -t - \varepsilon)$ for ε small enough). We then have $-\partial\theta \approx \bar{\partial}\psi$ and thus (let $\varepsilon \rightarrow 0^+$)

$$\begin{aligned} \int_{\Omega} (\eta + c^{-1})^{-1} |\tilde{F} - \theta\tilde{f}|^2 e^{-\psi-\phi-\varphi} & \leq \int_{\{-t-1 < \psi < -t\}} |\tilde{f}|^2 e^{-\psi-\phi-\varphi} \\ & \leq \left(\sup_{\{-t-1 < \psi < -t\}} e^{-\phi} \right) \int_{\{-t-1 < \psi < -t\}} |\tilde{f}|^2 e^{-\psi-\varphi}. \end{aligned} \tag{2.3}$$

We hope:

$$(c) \quad (\eta + c^{-1})^{-1} e^{-\phi} = e^{\beta(\psi)} \geq e^{\psi} = |z_n|^2, \text{ i.e.}$$

$$(s + r^{-1})^{-1} e^{-u} = e^{-t};$$

(d) $\lim_{t \rightarrow +\infty} u(t) = 0$.

Then do approximations (let $t \rightarrow +\infty$ next) to get

$$\int_{\Omega} |F|^2 e^{-\varphi} \leq \mathbf{C} \liminf_{t \rightarrow +\infty} \int_{\{-t-1 < \psi < -t\}} |\tilde{f}|^2 e^{-\psi-\varphi} = \mathbf{C} \pi \int_{\Omega \cap H} |f|^2 e^{-\varphi} d\lambda_H,$$

where

$$\mathbf{C} := \lim_{t \rightarrow +\infty} \left(\sup_{\{-t-1 < \psi < -t\}} e^{-\phi} \right) = 1.$$

Finally, we solve the ODE system

$$\begin{cases} u''s - s'' - rs'^2 = 0, \\ s' - su' = 0, \\ (s + r^{-1})^{-1} e^{-u} = e^{-t}. \end{cases}$$

In fact:

$$s(t) = \frac{t}{1 - e^{-t}} - 1, \quad u(t) = -\log(1 - e^{-t}), \quad r = \frac{u''s - s''}{s'^2} = \frac{1}{e^{t-u} - s} = \frac{1 - e^{-t}}{e^t - t - 1}.$$

The constant $\mathbf{C} = 1$ is the optimal constant.

实际上严格来说需要很多个逼近步骤: 例如区域的逼近, 权函数的光滑化逼近, ε 的趋于 0, 以及 t 的趋于无穷. 所以 Stein 流形上的证明里出现过四个指标依次取极限, 而弱拟凸 Kähler 流形上甚至五个以上. ■

For general $\psi \in \text{Psh}(\Omega)$ with pole:

THEOREM 2.11 (*L^2 estimate part*). Fix $B > 0$. Let Ω be a pseudoconvex domain in \mathbb{C}^n with $o \in \Omega$, and $\psi \in \text{Psh}^-(\Omega)$ with $\psi(o) = -\infty$. Let $F \in A^2(\{\psi < -t_0\})$, where $t_0 \geq 0$. Then there exists a holomorphic function $\tilde{F} \in \mathcal{O}(\Omega)$, s.t.

$$|\tilde{F} - F|^2 e^{-\psi} \text{ is locally } L^1 \text{ integrable near } o,$$

and

$$\int_{\Omega} \left| \tilde{F} - (1 - b_{t_0, B}(\psi))F \right|^2 \leq \mathbf{C} \int_{\Omega} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \psi < -t_0\}} |F|^2 e^{-\psi}, \quad \mathbf{C} = 1 - e^{-(t_0 + B)},$$

where $\mathbb{I}_{\{-t_0 - B < \psi < -t_0\}}$ is the character function of the set $\{-t_0 - B < \psi < -t_0\}$, and $b_{t_0, B}(t) = \int_{-\infty}^t \frac{1}{B} \mathbb{I}_{\{-t_0 - B < s < -t_0\}} ds$.

Proof. ($B = 1$ case) Replace $\log|z_n|^2$ by ψ in the above proof. We will get

$$\int_{\Omega} |\tilde{F} - (1 - b_{t_0}(\psi))F|^2 e^{-\psi + \beta(\psi)} \leq \left(\sup_{\Omega} e^{-\phi} \right) \int_{\{-t_0 - 1 < -\psi < -t_0\}} |F|^2 e^{-\psi}.$$

Then $|\tilde{F} - F|^2 e^{-\psi}$ is locally L^1 integrable near o , and the estimate holds by $\beta(x) \geq x$. ■

THEOREM 2.12 (A twisted version). *Let D be a pseudoconvex domain in \mathbb{C}^n s.t. $\sup_{z \in D} |z_n| \leq 1$, let $\varphi \in \text{Psh}(D)$, and let $H = \{z_n = 0\}$.*

Suppose $c(t) : [0, +\infty) \rightarrow (0, +\infty)$ is smooth s.t. $c(t)e^{-t}$ is decreasing in t , and $\int_0^{+\infty} c(t)e^{-t} dt < +\infty$.

Then for every $f \in \mathcal{O}(D \cap H)$ with

$$\int_{D \cap H} |f|^2 e^{-\varphi} d\lambda < +\infty,$$

one can find a holomorphic extension F of f to D satisfying

$$\int_D c(-2 \log |z_n|) |F|^2 e^{-\varphi} d\lambda \leq \pi \int_0^{+\infty} c(t) e^{-t} dt \int_{D \cap H} |f|^2 e^{-\varphi} d\lambda_H.$$

Proof. We need

$$(\eta + c^{-1})^{-1} e^{-\phi} = c(\beta(\psi)) e^{\beta(\psi)}, \quad \text{that is} \quad (s + r^{-1})^{-1} e^{-u} = c(t) e^{-t}.$$

See [10]. Solve the ODE system:

$$\begin{cases} \left(s + \frac{s'^2}{u''s - s''} \right) e^{u-t} = \frac{1}{c(t)}, \\ s' - su' = 1. \end{cases}$$

We get

$$u(t) = -\log \left(\int_0^t c(t_1) e^{-t_1} dt_1 \right),$$

and

$$s(t) = \frac{\int_0^t \left(\int_0^{t_2} c(t_1) e^{-t_1} dt_1 \right) dt_2}{\int_0^t c(t_1) e^{-t_1} dt_1}.$$

Also, we let $r = (u''s - s'')/s'^2$.

实际上 c 只需要满足正, 连续, 积分有限, 以及

$$\left(\int_0^t c(t_1) e^{-t_1} dt_1 \right)^2 > c(t) e^{-t} \left(\int_0^t \left(\int_0^{t_2} c(t_1) e^{-t_1} dt_1 \right) dt_2 \right), \quad \forall t > 0.$$

■

2.2.1 Applications of the L^2 extension theorem

Let $D = \{z : \rho(z) < 0\} \subset \mathbb{C}^n$ be a bounded strongly pseudoconvex domain with C^2 boundary, where ρ is C^2 and strictly psh on an open neighborhood Ω of \overline{D} . Let $\delta(z) = \text{dist}(z, \partial D)$ for $z \in D$.

Let $\varphi \in \text{Psh}(D) \cap C^0(\overline{D})$. Let

$$A_{\alpha, \varphi}^2(D) = \left\{ f \in \mathcal{O}(D) : \int_D e^{-\varphi} \delta^\alpha |f|^2 d\lambda < +\infty \right\},$$

and

$$\|f\|_{\alpha,\varphi}^2 = (\alpha + 1) \int_D e^{-\varphi} \delta^\alpha |f|^2 d\lambda.$$

Let

$$A_{-1,\varphi}^2(D) = \left\{ f \in \mathcal{O}(D) : \lim_{\alpha \rightarrow -1^+} (1 + \alpha) \int_D e^{-\varphi} \delta^\alpha |f|^2 d\lambda < +\infty \right\},$$

and

$$\|f\|_{-1,\varphi}^2 = \lim_{\alpha \rightarrow -1^+} (1 + \alpha) \int_D e^{-\varphi} \delta^\alpha |f|^2 d\lambda.$$

Let $H = \{h = 0\}$ be a smooth complex hypersurface in Ω , and $dh|_z \neq 0$ for every $z \in H \cap D$.

Assume ∂D intersects with H transversally ([10] says this condition can be dropped, but I do not think so).

THEOREM 2.13 (Beatrous, Diederich–Herbot, Ohsawa, Guan–Zhou). *The extension operator:*

$$A_{\alpha+1,\varphi}^2(D \cap H) \longrightarrow A_{\alpha,\varphi}^2(D)$$

for every $\alpha > -1$ has a bound $C_0 \max\{C_1^\alpha, C_2^\alpha\}$, where $C_0, C_1, C_2 > 0$ are independent of α .

Consequently, the extension operator from $A_{0,\varphi}^2(D \cap H)$ to $A_{-1,\varphi}^2(D)$ is bounded.

Proof. See [10, p. 60] or [8].

Since $\rho \in C^2(\Omega) \cap \text{SPsh}(\Omega)$, and ∂D is C^2 , $-\rho/\delta$ has uniform positive upper and lower bound on D .

Replace δ by $-\rho$. Let

$$\psi = -\log \left(-\frac{\rho}{\varepsilon_0 |h|^2} + 1 \right) < 0,$$

where $\varepsilon_0 > 0$ is sufficiently small s.t. $\rho - \varepsilon_0 |h|^2$ is strictly psh in a neighborhood of D . Since $-\log(-t)$ is increasing and convex for $t < 0$, we have $-\log(-\rho + \varepsilon_0 |h|^2) \in \text{Psh}(D) \Rightarrow \psi \in \text{Psh}(D)$.

Let

$$c(t) = \begin{cases} t^\alpha & 0 < t < 1, \\ 1 & t \geq 1. \end{cases}$$

Then $\int_0^{+\infty} c(t) e^{-t} dt < \frac{1}{1+\alpha} + 1$. Let

$$\phi = -\alpha \log(-\rho + \varepsilon_0 |h|^2) \Rightarrow \phi + \psi \in \text{Psh}(D).$$

For this case we can still use the twisted L^2 extension theorem. We may see h as z_n in the above theorem. The functions

$$e^{-\psi} \approx \begin{cases} (\varepsilon_0 |h|^2)^{-1} & z \rightarrow H \setminus \partial D, \\ -\rho/\varepsilon_0 |h|^2 & z \rightarrow \partial D, \end{cases}$$

and

$$e^{-\phi} = (-\rho + \varepsilon_0 |h|^2)^\alpha \approx \begin{cases} (\varepsilon_0 |h|^2)^\alpha & z \rightarrow H \setminus \partial D, \\ (-\rho)^\alpha & z \rightarrow \partial D \setminus H. \end{cases}$$

If ∂D intersects with H transversally, then we are able to omit the areas near $\partial D \cap H$.

We will have the following estimate of the extension $F|_{D \cap H} = f$ (denote a local holomorphic extension of f by f itself):

$$\begin{aligned} \int_D c(-\psi) |F|^2 e^{-\phi-\varphi} d\lambda &\leq C_{D,H} \frac{2+\alpha}{1+\alpha} \liminf_{t \rightarrow +\infty} \int_{\{-t-1 < \psi < -t\}} |f|^2 e^{-\psi-\phi-\varphi} d\lambda \\ &\leq C_{D,H} C_0^\alpha \frac{1}{1+\alpha} \int_{D \cap H} e^{-\varphi} (-\rho)^{\alpha+1} |f|^2 d\lambda_H. \end{aligned}$$

Finally, note that

$$\begin{aligned} \frac{c(-\psi) e^{-\phi}}{(-\rho)^\alpha} \Big|_{\{\psi > -1\}} &= \left(\log \left(1 + \frac{-\rho}{\varepsilon_0 |h|^2} \right) \right)^\alpha \cdot \left(1 + \frac{\varepsilon_0 |h|^2}{-\rho} \right)^\alpha \\ &\geq \begin{cases} C_3^{-\alpha} & z \rightarrow \partial D \setminus H, \\ C_4^{-\alpha} & z \rightarrow \partial D \cap H, \end{cases} \\ &\geq \min\{C_3^{-\alpha}, C_4^{-\alpha}\}, \end{aligned}$$

Thus, we have

$$\int_D e^{-\varphi} (-\rho)^\alpha |F|^2 d\lambda \leq C \max\{C_1^\alpha, C_2^\alpha\} \frac{1}{1+\alpha} \int_{D \cap H} e^{-\varphi} (-\rho)^{\alpha+1} |f|^2 d\lambda_H.$$

Finally, we can replace $-\rho$ by δ . ■

Let $\Gamma \subseteq \mathbb{C}$ be a discrete set of points such that

$$\inf\{|\gamma - \gamma'| : \gamma, \gamma' \in \Gamma, \gamma \neq \gamma'\} > 0. \quad (2.4)$$

We say that the interpolation problem is solvable for Γ if for any collection $\{a_\gamma \in \mathbb{C} : \gamma \in \Gamma\}$ of complex numbers such that

$$\sum_{\gamma \in \Gamma} |a_\gamma|^2 e^{-|\gamma|^2} < +\infty,$$

there is an entire function f such that

$$f(\gamma) = a_\gamma \quad \text{and} \quad \int_{\mathbb{C}} |f(z)|^2 e^{-|z|^2} d\lambda(z) < +\infty.$$

THEOREM 2.14 (Seip). *The followings are equivalent.*

1. *The discrete set Γ satisfies (2.4), and*

$$D(\Gamma) := \limsup_{R \rightarrow +\infty} \sup_{z \in \mathbb{C}} \frac{\#\{\gamma \in \Gamma : |z - \gamma| < R\}}{R^2} < 1.$$

2. *The interpolation problem is solvable for Γ .*

Proof. (Berndtsson–Ortega Cerdà) We prove $1 \Rightarrow 2$. Let $R > 0$, and

$$\psi(z) = V_R(z) := \sum_{\gamma \in \Gamma} \left(\log |z - \gamma|^2 - \frac{1}{\pi R^2} \int_{\{|\zeta - z| < R\}} \log |\zeta - \gamma|^2 d\lambda(\zeta) \right).$$

Then

1. V_R is well-defined and negative;
2. In the sense of currents,

$$i\partial\bar{\partial}V_R(z) \geq -\frac{\#\{\gamma \in \Gamma: |z - \gamma| < R\}}{R^2} d\lambda(z);$$

3. e^{-V_R} is not locally integrable at each $\gamma \in \Gamma$;
4. There is a constant $C_{R,\varepsilon} > 0$ such that if $\gamma \in \Gamma$ and $\varepsilon/2 < |z - \gamma| < \varepsilon$ then $V_R(z) \geq -C_{R,\varepsilon}$.

■

THEOREM 2.15 (Guan 19'). *Let Ω be a pseudoconvex domain in \mathbb{C}^n with $o \in \Omega$, and $\psi \in \text{Psh}^-(\Omega)$ with $\psi(o) = -\infty$. Assume the holomorphic germ $(F, o) \in \mathcal{O}_o \setminus \mathcal{I}(\psi)_o$. The minimal L^2 integral on any domain $D \ni o$ is defined as*

$$C_{F,\psi}(D) = C_{F,\mathcal{I}(\psi)_o}(D) := \left\{ \int_D |\tilde{F}|^2 : (\tilde{F} - F, o) \in \mathcal{I}(\psi)_o \text{ \& } \tilde{F} \in \mathcal{O}(D) \right\}.$$

Then the function

$$(0, +\infty) \ni t \longrightarrow G(t) := C_{F,\psi}(\Omega \cap \{\psi < -t\})$$

satisfies that $G(-\log r)$ is concave in $r \in [0, 1]$.

Proof. See [7].

Use only mathematical analysis and real analysis. However, the computation is a bit complicated, so we introduce Berndtsson's log-psh of fiberwise Bergman kernels, which can be directly implied by the optimal L^2 extension theorem, fewer calculations are used, and usually have similar efforts. ■

COROLLARY 2.16 (Berndtsson 06'). *Let Ω be a pseudoconvex domain in \mathbb{C}^{n+m} over a pseudoconvex domain $D \subset \mathbb{C}^m$, and pr is the projection. Let $\varphi \in \text{Psh}(\Omega)$. Denote the Bergman kernel on each fiber $\Omega_w = \text{pr}^{-1}(w) \cap \Omega$ w.r.t. the weight $\varphi|_{\Omega_w}$ by*

$$K_\varphi(z, w) := K_{\Omega_w, e^{-\varphi}|_{\Omega_w}}(z) = \sup_{f \in \mathcal{O}(\Omega_w)} \frac{|f(z)|^2}{\int_{\Omega_w} |f|^2 e^{-\varphi}|_{\Omega_w}}.$$

Then $\log K_\varphi(z, w) \in \text{Psh}(\Omega)$.

Proof. See [15, pp. 191–192].

(Guan–Zhou method) We only prove the mean value inequality on the direction of w . Assume $D = \Delta$ the unit disc in \mathbb{C} . Let $f \in \mathcal{O}(\Omega_0)$ such that

$$K_\varphi(z_0, 0) = \frac{|f(z_0)|^2}{\int_{\Omega_0} |f(z)|^2 e^{-\varphi(z)}}.$$

Using the optimal L^2 extension theorem, we can find some $F(z, w) \in \mathcal{O}(\Omega)$, such that $F(z, 0) = f(z)$ and

$$\int_{\Omega} |F|^2 e^{-\varphi} \leq \pi \int_{\Omega_0} |f|^2 e^{-\varphi}.$$

Then apply Jensen's inequality to the increasing convex function $y = \log x$:

$$\begin{aligned}
 \log \left(\int_{\Omega_0} |f|^2 e^{-\varphi} \right) &\geq \log \left(\frac{1}{\pi} \int_{\Omega} |F|^2 e^{-\varphi} \right) \\
 &= \log \left(\frac{1}{\pi} \int_{w \in \Delta} \left(\int_{\Omega_w} |F_w|^2 e^{-\varphi} \right) d\lambda_w \right) \\
 &\geq \frac{1}{\pi} \int_{w \in \Delta} \log \left(\int_{\Omega_w} |F_w|^2 e^{-\varphi} \right) d\lambda_w \\
 &\geq \frac{1}{\pi} \int_{w \in \Delta} \left(\log |F_w(z_0)|^2 - \log K_{\varphi}(z_0, w) \right) d\lambda_w \\
 &= \frac{1}{\pi} \int_{w \in \Delta} \log |F(z_0, w)|^2 - \frac{1}{\pi} \int_{w \in \Delta} \log K_{\varphi}(z_0, w) \\
 &\geq \log |F(z_0, 0)|^2 - \frac{1}{\pi} \int_{w \in \Delta} \log K_{\varphi}(z_0, w) \\
 &= \log |f(z_0)|^2 - \frac{1}{\pi} \int_{w \in \Delta} \log K_{\varphi}(z_0, w),
 \end{aligned}$$

where $F_w(\cdot) = F(\cdot, w)$, and $\log |F_w(z_0)|^2$ is subharmonic in w . Thus,

$$\log K_{\varphi}(z_0, 0) \leq \frac{1}{\pi} \int_{w \in \Delta} \log K_{\varphi}(z_0, w).$$

Berndtsson 最初的证明方法是先在光滑情形直接使用再生核公式计算二阶导, 最后结合 Hörmander L^2 估计. 之后这套方法发展为了他所提出的复 Brunn–Minkowski 理论, 并在 direct image of relative canonical bundle 的正性上有重要应用. \blacksquare

Let D be a domain in the complex plane containing the origin. Let $G(z) := G(z, 0)$ be the Green function on D with pole at 0. Then $2G(z) = \log |z|^2 - h(z)$, where h is a harmonic function such that G vanishes on the boundary of D . Then $c_D := h(0)$ is the Robin constant at 0, and

$$c_{\beta}^2(0) = e^{-c_D} := \exp \lim_{z \rightarrow 0} \left(2G(z) - \log |z|^2 \right),$$

where c_{β} is called the logarithm capacity.

COROLLARY 2.17 (Blocki, Guan–Zhou). *The inequality part of Suita's conjecture says that*

$$\pi K(0) \geq e^{-c_D},$$

where $K(0)$ is the Bergman kernel at 0.

Proof. See [1, p. 2].

(Berndtsson–Lempert) Let $D_t = \{z \in D : 2G(z) < -t\}$, and

$$\Omega := \{(z, w) \in D \times [0, +\infty) + i\mathbb{R} : 2G(z) + \operatorname{Re} w < 0\} \subseteq \mathbb{C}^2,$$

which is a pseudoconvex domain. Then $\log K(0, w)$ is subharmonic in w and only depends on $\operatorname{Re} w$ $\Rightarrow \log K_{D_t}(0) = \log K(0, t + i \cdot)$ is convex in $t \in [0, +\infty)$. Using the definition of c_D , we can find

$$\Delta_{r_1} \subseteq D_t \subseteq \Delta_{r_2}$$

for $r_1 = e^{(-t+c_D-\varepsilon)/2}$ and $r_2 = e^{(-t+c_D+\varepsilon)/2}$ if t is sufficiently large. It follows that

$$K_t \sim \frac{e^{t-c_D}}{\pi}, \quad t \rightarrow +\infty.$$

Thus, $\log K_{D_t}(0) - t$ is convex and bounded from above in $t \in [0, +\infty)$, which implies the function is decreasing \Rightarrow

$$K_D(0) \geq \lim_{t \rightarrow +\infty} e^{-t} K_{D_t}(0) + t = \frac{e^{-c_D}}{\pi}.$$

Suita 猜想在开黎曼面上也是成立的, 并且等号成立当且仅当 D 全纯同构于 Δ 去掉某个容量为 0 的集合 (Guan–Zhou). 这里等号成立条件是更困难得多的, 需要用到后面提到的 twisted version 的 optimal L^2 延拓定理. 而更广范围的 optimal L^2 延拓定理的等号成立条件也是后续利用极小 L^2 积分凹性研究的主要内容之一. \blacksquare

Let $D \subseteq \mathbb{C}$ be a planar regular region bounded with n boundary components which are analytic Jordan curves. Let $H_2^{(c)}(D)$ denote the analytic Hardy class on D as the set of all analytic functions $f(z)$ on D s.t.

$$|f(z)|^2 \leq U(z) \text{ on } D, \quad \exists U \text{ harmonic.}$$

Then each $f \in H_2^{(c)}(D)$ has Fatou's nontangential boundary value a.e. on ∂D belonging to $L^2(\partial D)$.

The conjugate Hardy H^2 kernel on D is the unique kernel function $\widehat{R}_t(z, \bar{w})$ satisfying

- for any fixed $(w, t) \in D \times D$, as a function of z , $\widehat{R}_t(z, \bar{w}) \in H_2^{(c)}(D)$;
- for any $f \in H_2^{(c)}(D)$,

$$f(w) = \frac{1}{2\pi} \int_{\partial D} f(z) \overline{\widehat{R}_t(z, \bar{w})} \left(\frac{\partial G_D(z, t)}{\partial \nu_z} \right)^{-1} d|z|, \quad \forall (z, w, t) \in D \times D \times D,$$

where $G_D(z, t)$ is the Green function on D and $\frac{\partial}{\partial \nu_z}$ denotes the derivative along the outer normal unit vector ν_z .

Or equivalently, $M(z) := \frac{\widehat{R}_t(z, \bar{w})}{\widehat{R}_t(w, \bar{w})} \in H_2^{(c)}(D)$ is the unique solution of the following extremal problem:

$$\inf \left\{ \int_{\partial D} |f(z)|^2 \left(\frac{\partial G_D(z, t)}{\partial \nu_z} \right)^{-1} d|z| : f \in H_2^{(c)}(D), f(w) = 1 \right\}.$$

Let

$$\widehat{R}(z, \bar{w}) := \widehat{R}_w(z, \bar{w}), \quad \text{and} \quad \widehat{R}(z) := \widehat{R}(z, \bar{z}).$$

Let $B(z, \bar{w})$ denote the Bergman kernel on D and $B(z) := B(z, \bar{z})$.

THEOREM 2.18 (Saitoh's conjecture; Guan 19'). *If $n > 1$, then*

$$\widehat{R}(z) > \pi B(z).$$

Proof. Verify for every $f \in H_2^{(2)}(D)$, and $w \in D$,

$$\lim_{t \rightarrow 0^+} \frac{\int_{\{2G_D(\cdot, w) \geq -t\}} |f|^2}{t} = \int_{\partial D} |f(z)|^2 \left(\frac{\partial 2G_D(z, w)}{\partial \nu_z} \right)^{-1} d|z|,$$

which implies

$$\lim_{t \rightarrow 0^+} \frac{B_D^{-1}(w) - B_{\{2G_D(\cdot, w) < -t\}}^{-1}(w)}{t} \geq \frac{\pi}{\widehat{R}(w)}, \quad \forall t > 0.$$

Since the function

$$\log B(t) - t := \log B_{\{2G_D(\cdot, w) < -t\}}(w) - t$$

is convex and decreasing in $t \in (0, +\infty)$, we have

$$-1 + B(0) \lim_{t \rightarrow 0^+} \frac{B^{-1}(0) - B^{-1}(t)}{t} = \lim_{t \rightarrow 0^+} \frac{\log B(t) - t - \log B(0)}{t} \leq 0,$$

and thus

$$\widehat{R}(w) \geq \pi B(z).$$

For $n > 1$, the equality will not hold: if the equality hold, then

$$\frac{d}{dt}(\log B(t) - t) \Big|_{t=0^+} = 0,$$

so $\log B(t) - t$ is a constant function, and then it follows from the equality part of Suita's conjecture. \blacksquare

THEOREM 2.19 (Guan 19'). Let $D \subseteq \mathbb{C}^n$ be a pseudoconvex domain containing the origin o , and $\varphi \in \text{Psh}^-(D)$ with $\varphi(o) = -\infty$. If $\int_D e^{-\varphi} d\lambda < +\infty$, then for every $p > 1$ satisfying

$$\frac{p}{p-1} > K_D(o) \int_D e^{-\varphi} d\lambda,$$

we have $e^{-p\varphi}$ is locally L^1 integrable near o .

Proof. (An optimized version of Berndtsson's proof of the openness conjecture) Let $\Phi = q\varphi$ for some $q > 1$ and $D_t = \{z \in D : \Phi(z) < -t\}$ for every $t \geq 0$. Using the same arguments in the above proof, we have that $\log K_{D_t}(o) - t$ is convex in t , and we can use Theorem 2.11 (only a rough version and very few calculations are needed) to get that $\log K_{D_t}(o) - t$ is also upper bounded if $(1, o) \notin \mathcal{J}(\Phi/2)_o$. Thus, $\log K_{D_t}(o) - t$ is decreasing in t , that is

$$e^{-t} K_{D_t}(o) \leq K_D(o) \quad \Rightarrow \quad K_{D_t}^{-1}(o) \geq e^{-t} K_D^{-1}(o), \quad \forall t > 0.$$

Using Fubini's theorem, when $(1, o) \notin \mathcal{J}(\Phi/2)_o$, i.e. $e^{-q\varphi}$ is not L^1 integrable near o , we have

$$\begin{aligned} \int_D e^{-\varphi} d\lambda &= \int_D e^{-\frac{\Phi}{q}} d\lambda = \int_{-\infty}^{+\infty} \left(\int_{\{\Phi < -qt\}} |1|^2 \right) e^t dt \\ &\geq \lambda(D) + \int_0^{+\infty} K_{D_{qt}}^{-1}(o) dt \\ &\geq K_D^{-1}(o) + K_D^{-1} \int_0^{+\infty} e^{-qt} dt \\ &= \frac{q}{q-1} K_D^{-1}(o). \end{aligned}$$

Consequently, if $q > 1$ satisfies

$$\frac{q}{q-1} > K_D(o) \int_D e^{-\varphi} d\lambda,$$

then we must have that $e^{-q\varphi}$ is locally L^1 integrable near o (note that $\int_D e^{-\varphi} > \lambda(D) \geq K_D^{-1}(o)$).

(Bao–Guan 22'; 一些记号参见后面 “Multiplier ideal sheaf” 小节) 如果条件是 $\int_D |F|^2 e^{-\varphi} < +\infty$ 对某个 $F \in \mathcal{O}(D)$, 我们考虑:

$$\xi \in \ell_1^{(n)} := \left\{ (\xi_\alpha)_{\alpha \in \mathbb{Z}_{\geq 0}^n} : \sum_\alpha |\xi_\alpha| \rho^{|\alpha|} < +\infty \text{ for every } \rho > 0 \right\},$$

以及推广的 Bergman 核

$$K_{\xi, D}(z) := \sup_{f \in \mathcal{O}(D)} \frac{|(\xi \cdot f)(z)|^2}{\int_D |f|^2},$$

其中

$$(\xi \cdot f)(z) := \sum_\alpha \xi_\alpha \frac{f^{(\alpha)}(z)}{\alpha!}.$$

则我们能得到类似的 fiberwise 的多次调和性, 并同样推出 $k(t) := \log K_{\xi, D_t}(o) - t$ 的凸性. 同样地, 由定理 2.11 可以得到如果

$$\xi \in \ell_{\mathcal{J}(\Phi)_o}^{(n)} := \left\{ \eta \in \ell_1^{(n)} : (\eta \cdot f)(o) = 0, \forall (f, o) \in \mathcal{J}(\Phi/2)_o \right\},$$

则 $k(t)$ 上有界, 并因此推出 $k(t)$ 的递减性. 同样我们最终会得到如果 $p > 1$ 满足

$$\frac{p}{p-1} > \frac{\int_D |F|^2 e^{-\varphi}}{B(F, \mathcal{J}_+(c_o^F(\varphi)\varphi)_o, D)},$$

其中

$$B(F, \mathcal{J}_+(c_o^F(\varphi)\varphi)_o, D) := \sup_{\xi \in \ell_{\mathcal{J}_+(c_o^F(\varphi)\varphi)_o}^{(n)}} \frac{|(\xi \cdot F)(o)|^2}{K_{\xi, D}(o)},$$

则必有 $|F|^2 e^{-p\varphi}$ 在 o 附近 L^1 可积. 最后为了保证良定性, 我们只需证明 $B(F, \mathcal{J}_+(c_o^F(\varphi)\varphi)_o, D) > 0$, 即, 若 $(F, o) \in \mathcal{O}_o \setminus I$, 其中 I 为 \mathcal{O}_o 的一个理想, 则必存在 $\xi \in \ell_I$ 使得 $(\xi \cdot F)(o) \neq 0$ 即可 (e.g. 用泛函分析中的分离性定理). 更进一步地, 实际上我们可以证明

$$B(F, \mathcal{J}_+(c_o^F(\varphi)\varphi)_o, D) = C_{F, \mathcal{J}_+(c_o^F(\varphi)\varphi)_o}(D) > 0.$$

最终我们就会得到强开性猜想的一个证明, 并且包含一个 sharp 的有效性结果. ■

THEOREM 2.20. *Suppose $p \in (0, 2]$. Let Ω be a pseudoconvex domain in \mathbb{C}^n s.t. $\sup_{z \in \Omega} |z_n| \leq 1$, let $\varphi \in \text{Psh}(\Omega)$, and let $H = \{z_n = 0\}$. Then for every $f \in \mathcal{O}(D \cap H)$ with*

$$\int_{H \cap \Omega} |f|^p e^{-\varphi} d\lambda < +\infty,$$

one can find a holomorphic extension F of f to Ω satisfying

$$\int_\Omega |F|^p e^{-\varphi} d\lambda \leq C\pi \int_{\Omega \cap H} |f|^p e^{-\varphi} d\lambda.$$

Here C can be chosen as the C in the L^2 extension theorem.

Proof. Berndtsson's iteration technique. See [10, pp. 61–62].

May assume Ω is bounded and φ is smooth up to boundary of Ω . We first take a holomorphic extension F_1 of f satisfying a rough estimate:

$$A_1 := \int_{\Omega} |F_1|^p e^{-\varphi} d\lambda < +\infty.$$

Let

$$\varphi_1 := \varphi + (2-p) \log |F_1| \in \text{Psh}(\Omega).$$

Then the optimal L^2 extension theorem gives some $F_2 \in \mathcal{O}(\Omega)$ s.t. $F_2|_{D \cap H} = F_1|_{D \cap H} = f$ and

$$\int_{\Omega} |F_2|^2 e^{-\varphi_1} \leq \pi \int_{D \cap H} |f|^2 e^{-\varphi_1} = \pi \int_{D \cap H} |f|^p e^{-\varphi} =: A.$$

Note that

$$\left(\int_{\Omega} \frac{|F_2|^2}{|F_1|^{2-p}} e^{-\varphi} \right)^{\frac{p}{2}} \cdot \left(\int_{\Omega} |F_1|^p e^{-\varphi} \right)^{1-\frac{p}{2}} \geq \int_{\Omega} |F_2|^p e^{-\varphi} =: A_2,$$

that is

$$A_2 \leq A_1^{1-p/2} (\pi A)^{p/2}.$$

By iteration, we will finally get an extension F_{∞} with $A_{\infty} \leq \pi A$. ■

THEOREM 2.21 (Demailly's approximation theorem). *Let Ω be a bounded pseudoconvex domain in \mathbb{C}^n , and let $\varphi \in \text{Psh}(\Omega)$. For each $m \geq 1$, the **Demailly approximation** φ_m of φ is defined as*

$$\varphi_m(z) := \frac{1}{2m} \log K_{2m\varphi}(z) := \frac{1}{2m} \log \sup_{\substack{f \in A^2(\Omega, e^{-2m\varphi}), \\ \|f\|_{2m\varphi} = 1}} |f(z)|^2, \quad \forall z \in \Omega, \quad (2.5)$$

where $K_{m\varphi}$ denotes the Bergman kernel for the weight $e^{-2m\varphi}$ on Ω .

Then there are constants $C_1, C_2 > 0$ independent of m such that

$$\varphi(z) - \frac{C_1}{m} \leq \varphi_m(z) \leq \sup_{|\zeta - z| < r} \varphi(\zeta) + \frac{1}{m} \log \frac{C_2}{r^n}$$

for every $z \in \Omega$ and $r < \text{dist}(z, \partial\Omega)$. In particular:

1. φ_m converges to φ pointwise and in L^1_{loc} topology on Ω when $m \rightarrow +\infty$.
2. For every $z \in \Omega$,

$$\nu(\varphi, z) - \frac{n}{m} \leq \nu(\varphi_m, z) \leq \nu(\varphi, z).$$

Proof. See [5, pp. 136–138].

First, applying the mean value inequality to $|f|^2$ for every $f \in A^2(\Omega, e^{-2m\varphi})$, we get

$$\begin{aligned} |f(z)|^2 &\leq \frac{1}{\text{Vol}(B(z, r))} \int_{B(z, r)} |f|^2 d\lambda \\ &\leq \frac{n!}{\pi^n r^{2n}} \left(\sup_{\zeta \in B(z, r)} e^{2m\varphi(\zeta)} \right) \int_{B(z, r)} |f|^2 e^{-2m\varphi} d\lambda \\ &\leq \frac{n!}{\pi^n r^{2n}} \left(e^{2m \sup_{|\zeta - z| < r} \varphi(\zeta)} \right) \int_{\Omega} |f|^2 e^{-2m\varphi} d\lambda, \end{aligned}$$

for every $z \in \Omega$ and $r < \text{dist}(z, \partial\Omega)$, which implies

$$\varphi_m(z) = \frac{1}{2m} \log K_{2m\varphi}(z) \leq \sup_{|\zeta-z|<r} \varphi(\zeta) + \frac{1}{m} \log \frac{C_2}{r^n}.$$

Second, according to the Ohsawa–Takegoshi L^2 extension theorem, there exists a constant C'_1 , which only depends on n and $\text{diam}\Omega$, such that for every $m \geq 1$ and $z \in \Omega$, there exists a holomorphic function F on Ω , satisfying $F(z) = 1$ and

$$\int_{\Omega} |F|^2 e^{-2m\varphi} d\lambda \leq C'_1 e^{-2m\varphi(z)}.$$

It follows that

$$\varphi_m(z) = \frac{1}{2m} \log K_{2m\varphi}(z) \geq \varphi(z) - \frac{\log C'_1}{2m} = \varphi(z) - \frac{C_1}{m}.$$

Third, we choose $r = \frac{1}{m}$ for m sufficiently large, and then we will find that φ_m converges to φ by the upper semi-continuity of φ and $\log(C_2 m^n)/m \rightarrow 0$ as $m \rightarrow +\infty$.

Finally, for the Lelong numbers, from the inequality we get

$$\frac{\sup_{|z-z_0|<2r} \varphi(z) + \frac{1}{m} \log \frac{C_2}{r^n}}{\log r} \leq \frac{\sup_{|z-z_0|<r} \varphi_m(z)}{\log r} \leq \frac{\sup_{|z-z_0|<r} \varphi(z) - \frac{C_1}{m}}{\log r}.$$

By letting $r \rightarrow 0^+$, we get

$$\nu(\varphi, z_0) - \frac{n}{m} \leq \nu(\varphi_m, z_0) \leq \nu(\varphi, z_0), \quad \forall z_0 \in \Omega.$$

■

Note that every φ_m has analytic singularities.

For a psh function φ near z_0 , the Lelong number of φ at z_0 is defined to be

$$\nu(\varphi, z_0) = \liminf_{z \rightarrow z_0} \frac{\varphi(z)}{\log |z - z_0|} = \lim_{r \rightarrow 0^+} \frac{\sup_{B(z_0, r)} \varphi}{\log r}.$$

In particular, $\varphi = \log |f|$, with f holomorphic near z_0 , then

$$\nu(\varphi, z_0) = \text{ord}_{z_0}(f) = \sup\{k: D^\alpha f(z_0) = 0, \forall |\alpha| < k\}.$$

COROLLARY 2.22 (Siu's semi-continuity). *Let $\varphi \in \text{Psh}(\Omega)$. Then*

$$E_c(\varphi) = \{z \in \Omega: \nu(\varphi, z) \geq c\}$$

is an analytic subset of Ω for every $c > 0$.

Proof. See [5, p. 138].

Let φ_m be the Demailly approximation of φ . Then by the above theorem, we have

$$E_c(\varphi) = \bigcap_{m \geq m_0} E_{c-n/m}(\varphi_m)$$

for some $m_0 \gg 1$. For each $E_c(\varphi_m)$, we can see

$$\varphi_m = \left\{ z \in \Omega : \sigma_l^{(\alpha)}(z) = 0, \forall l \text{ \& } |\alpha| < mc \right\},$$

where (σ_l) is a complete orthogonal basis of $A^2(\Omega, e^{-2m\varphi})$. Then every $E_{c-n/m}(\varphi_m)$ is an analytic set, so $E_c(\varphi)$ is an analytic set. \blacksquare

THEOREM 2.23 (Skoda 72'). *If $\nu(\varphi, z) < 1$, then $e^{-2\varphi}$ is locally integrable near z .*

Proof. This can be proved by dynamically using L^2 extension and a simple lemma in complex analysis.

(Guan–Zhou) Assume $\varphi \in \text{Psh}^-(\Delta^n)$, $\nu(\varphi, o) < 1$, and $e^{-2\varphi}$ is not locally integrable near o .

By the definition of the Lelong number, we have that there exists a sequence of points $\{z_j\} \subseteq \Delta^n$ such that

$$z_j \rightarrow o \text{ and } |z_j|^2 e^{-2\varphi(z_j)} \rightarrow 0 \text{ when } j \rightarrow +\infty.$$

For every z_j , using the Ohsawa-Takegoshi L^2 extension theorem, we can find some $F_j \in \mathcal{O}(\Delta^n)$ with $F_j(z_j) = 1$ and

$$\int_{\mathcal{L}_j \cap \Delta^n} |F_j|^2 \leq C \int_{\Delta^n} |F_j|^2 \leq C \int_{\Delta^n} |F_j|^2 e^{-2\varphi} \leq C_1 e^{-2\varphi(z_j)},$$

where \mathcal{L}_j is a 1-dimension plane across z_j and o , and C_1 only depends on the diameter of Δ^n . Since $e^{-2\varphi}$ is not locally integrable near o , we have $F_j(o) = 0$.

A simple lemma For every $h \in \mathcal{O}(\Delta^n)$ with $h(0) = 0$ and $h(z) = 1$, where $z \in \Delta_{1/2}$, there is a constant $C_2 > 0$ independent of h and z , such that

$$\int_{\Delta} |h|^2 d\lambda \geq \frac{C_2}{|z|^2}.$$

It follows that

$$|z_j|^2 e^{-2\varphi(z_j)} \geq \frac{C_2}{C_1} > 0, \quad \forall j \gg 1.$$

Contradiction! \blacksquare

2.3 Multiplier ideal sheaf

DEFINITION 2.24. Let $\varphi \in \text{Psh}(\Omega)$. At each $z \in \Omega$, define the multiplier ideal

$$\mathcal{I}(\varphi)_z := \{f \in \mathcal{O}_{\Omega, z} : |f|^2 e^{-2\varphi} \text{ is locally integrable near } z\}.$$

Then $\mathcal{I}(\varphi)$ can be defined as an ideal sheaf of \mathcal{O}_{Ω} .

DEFINITION 2.25. Let \mathcal{A} be a sheaf of rings on a topological space X and \mathcal{S} be a sheaf of modules over \mathcal{A} . Then \mathcal{S} is said to be locally finitely generated if for every $x_0 \in X$, one can find a neighborhood U and sections $F_1, \dots, F_q \in \mathcal{S}(U)$ such that for every $x \in U$ the stalks \mathcal{S}_x is generated by the germs $F_{1,x}, \dots, F_{q,x}$ as an \mathcal{A}_x -module.

DEFINITION 2.26. Let U be an open subset of X , and $F_1, \dots, F_q \in \mathcal{I}(U)$. The kernel of the sheaf homeomorphism

$$\begin{aligned} F: (\mathcal{A}|_U)^{\otimes q} &\longrightarrow \mathcal{I}|_U \\ (g^1, \dots, g^q) &\longmapsto \sum_{1 \leq j \leq q} g^j F_{j,x}, \quad x \in U \end{aligned}$$

is a subsheaf $\mathcal{R}(F_1, \dots, F_q)$ of $(\mathcal{A}|_U)^{\otimes q}$, called the sheaf of relations between F_1, \dots, F_q .

A sheaf \mathcal{I} of \mathcal{A} -module on X is said to be coherent if

1. \mathcal{I} is locally finitely generated;
2. for any open subset U of X and any $F_1, \dots, F_q \in \mathcal{I}(U)$, the sheaf of relations $\mathcal{R}(F_1, \dots, F_q)$ is locally finitely generated.

PROPOSITION 2.27 (Strong Noetherian property). Let \mathcal{F} be a coherent analytic sheaf on a complex manifold M (i.e. sheaf of modules over \mathcal{O}_M), and let $\mathcal{F}_1 \subset \mathcal{F}_2 \subset \dots$ be an increasing sequence of coherent subsheaves of \mathcal{F} . Then the sequence (\mathcal{F}_k) is stationary on every compact subset of M .

Proof. See [6, p. 90]. ■

PROPOSITION 2.28 (Nadel 89'). $\mathcal{I}(\varphi)$ is a coherent sheaf on Ω .

Proof. See [5, p. 37].

Since the structure sheaf \mathcal{O}_Ω is coherent (Oka), it suffices to prove that $\mathcal{I}(\varphi)$ is a locally finitely generated sheaf. For each $x \in \Omega$, let $U \Subset \Omega$ which is a bounded pseudoconvex domain containing x , and by the strong Noetherian property of coherent sheaves, the family of sheaves generated by finite subsets of $A^2(U, e^{-2\varphi})$ has a maximal element on each relatively compact subset of U . Hence, $A^2(U, e^{-2\varphi})$ generates a coherent ideal sheaf $\mathcal{I} \subset \mathcal{O}_U$. It is clear that $\mathcal{I} \subset \mathcal{I}(\varphi)$ on U .

In view of:

Krull's lemma Let (R, \mathfrak{m}) be a Noetherian local ring, and F be a finitely generated R -module. Then

$$\bigcap_{k \geq 0} (E + \mathfrak{m}^k) = E.$$

We only need to prove at any $x \in U$, we have

$$\mathcal{I}_x + \mathcal{I}(\varphi)_x \cap \mathfrak{m}_{U,x}^{s+1} = \mathcal{I}(\varphi)_x, \quad \forall s \in \mathbb{Z}_{\geq 0}.$$

Then the proof is very like what we did before. Let $f \in \mathcal{I}(\varphi)_x$, and let

$$\tilde{\varphi}(z) = \varphi(z) + (n+s) \log |z-x| + |z|^2.$$

We will solve $\bar{\partial}u = \partial(\theta f)$ with the metric $e^{-2\tilde{\varphi}}$ and let $F = -u + \theta f$, which is holomorphic and

$$f_x - F_x = u_x \in \mathcal{I}(\varphi)_x \cap \mathfrak{m}_{U,x}^{s+1}.$$

This proves the coherence. ■

EXAMPLE 2.29. If φ is a psh function near $0 \in \mathbb{C}$, then

$$\mathcal{I}(\varphi)_0 = (z^k)_0$$

for some $k \in \mathbb{N}$, and k can be determined by $\nu(\varphi, 0) \in [k, k+1)$.

PROPOSITION 2.30 (Restriction formula). *Let H be a complex hyperplane, $Y = H \cap D$, and $\varphi \in \text{Psh}(D)$. Then*

$$\mathcal{I}(\varphi|_Y) \subset \mathcal{I}(\varphi)|_Y.$$

Proof. Direct consequence of O-T L^2 extension theorem. ■

PROPOSITION 2.31 (Subadditive property).

- Let $\varphi_1 \in \text{Psh}(D_1)$, $\varphi_2 \in \text{Psh}(D_2)$, and $\pi_i: D_1 \times D_2 \rightarrow D_i$ be the projections. Then

$$\mathcal{I}(\varphi_1 \circ \pi_1 + \varphi_2 \circ \pi_2) = \pi_1^* \mathcal{I}(\varphi_1) \cdot \pi_2^* \mathcal{I}(\varphi_2).$$

- Let $\varphi, \psi \in \text{Psh}(D)$. Then

$$\mathcal{I}(\varphi + \psi) \subset \mathcal{I}(\varphi) \cdot \mathcal{I}(\psi).$$

Proof. See [5, p. 159]. ■

2.4 Openness and strong openness

Let

$$\mathcal{I}_+(\varphi) := \bigcup_{\varepsilon > 0} \mathcal{I}((1 + \varepsilon)\varphi).$$

Then $\mathcal{I}_+(\varphi)|_K = \mathcal{I}((1 + \varepsilon_K)\varphi)|_K$ for some $\varepsilon_K > 0$ on compact K by the coherence (or simply by the Noetherian of \mathcal{O}_x , we have $\mathcal{I}_+(\varphi)_x = \mathcal{I}((1 + \varepsilon_0)\varphi)_x$ for some $\varepsilon_0 > 0$). Demailly's strong openness conjecture (proved by Guan-Zhou) is:

$$\mathcal{I}_+(\varphi) = \mathcal{I}(\varphi).$$

If $\mathcal{I}(\varphi) = \mathcal{O}$, this is the openness conjecture (proved by Berndtsson).

COROLLARY 2.32. *If $c_o^F(\varphi) < +\infty$, then $(F, o) \notin \mathcal{I}(c_o^F(\varphi)\varphi)_o$.*

THEOREM 2.33. *Let φ_j be sequence of psh function near o such that φ_j increasingly converges to a psh function φ near o in Lebesgue measure. Then*

$$\mathcal{I}(\varphi)_o = \bigcup_{j \geq 1} \mathcal{I}(\varphi_j)_o.$$

Proof. Omit. See [11]. ■

COROLLARY 2.34 (Fornæss). *The strong openness property is also true for every $p > 0$.*

Proof. Let $k \in \mathbb{Z}$ with $2k \geq p$. Observe

$$|F|^p e^{-2\varphi} = |F^k|^2 e^{-(2\varphi + (2k-p)\log|F|)},$$

and for every $q > 1$,

$$|F^k|^2 e^{-q(2\varphi + (2k-p)\log|F|)} = |F|^{2k-2kq+pq} e^{-2q\varphi} \gtrsim |F|^p e^{-2q\varphi},$$

due to $2k - 2kq + pq \leq p \Leftrightarrow q(2k - p) \geq 2k - p \Leftrightarrow q \geq 1$. ■

COROLLARY 2.35. *If φ, ϕ are psh functions near o , and $\nu(\phi, o) = 0$, then*

$$\mathcal{I}(\varphi + \phi)_o = \mathcal{I}(\varphi)_o.$$

Proof. Use Skoda's result.

Since $\nu(\phi, o) = 0$, we have that $e^{-2q\phi}$ is integrable near o for every $q > 0$. Hölder's inequality implies

$$\int |F|^2 e^{-2\varphi - 2\phi} \leq \left(\int |F|^{2(1+\varepsilon)} e^{-2(1+\varepsilon)\varphi} \right)^{1/(1+\varepsilon)} \cdot \left(\int e^{-2\frac{1+\varepsilon}{\varepsilon}\phi} \right)^{\varepsilon/(1+\varepsilon)}.$$

Take $\varepsilon > 0$ such that $\int |F|^2 e^{-2(1+\varepsilon)\varphi} < +\infty$. ■

COROLLARY 2.36. *For psh function φ near o , there exists a psh function φ_A with analytic singularities such that*

$$\mathcal{I}(\varphi)_o = \mathcal{I}(\varphi_A)_o.$$

Proof. See [9].

For every $j \geq 1$, set

$$I_j = |\mathcal{I}(j\varphi)_o|.$$

Claim

$$\mathcal{I}(\varphi)_o = \mathcal{I} \left(\max \left\{ \varphi, \frac{1}{j-1} \log |I_j| \right\} \right)_o, \quad \forall j > 1.$$

The claim follows from, for every positive continuous function χ near o , we have

$$\begin{aligned} 0 &\leq \int_V \chi e^{-2\varphi} - \int_V \chi e^{-2 \max \{ \varphi, \frac{1}{j-1} \log |I_j| \}} \\ &= \int_{V \cap \{ \varphi \leq \frac{1}{j-1} \log |I_j| \}} \chi \left(e^{-2\varphi} - e^{-2 \frac{1}{j-1} \log |I_j|} \right) \\ &\leq \int_{V \cap \{ \varphi \leq \frac{1}{j-1} \log |I_j| \}} \chi e^{-2j\varphi + 2(j-1)\varphi} \\ &\leq \int_V \chi |I_j|^2 e^{-2j\varphi} < +\infty. \end{aligned}$$

Demailly's approximation theorem gives

$$\frac{1}{j} \log |I_j| \geq \varphi + O(1), \quad \forall j \geq 1.$$

According to the strong openness property, we can take some $\varepsilon_0 > 0$ such that

$$\mathcal{I}(\varphi)_o = \mathcal{I}((1 + \varepsilon_0)\varphi)_o.$$

Thus,

$$\begin{aligned} \mathcal{I} \left(\max \left\{ \varphi, \frac{1}{j-1} \log |I_j| \right\} \right)_o &= \mathcal{I}(\varphi)_o = \mathcal{I}((1 + \varepsilon_0)\varphi)_o \\ &\subseteq \mathcal{I} \left(\frac{1 + \varepsilon_0}{j} \log |I_j| \right)_o \subseteq \mathcal{I} \left(\max \left\{ \varphi, \frac{1 + \varepsilon_0}{j} \log |I_j| \right\} \right)_o. \end{aligned}$$

If we take j such that

$$\frac{1}{j-1} \leq \frac{1 + \varepsilon_0}{j} \iff j \geq 1 + \frac{1}{\varepsilon_0},$$

then we have

$$\mathcal{I} \left(\max \left\{ \varphi, \frac{1}{j-1} \log |I_j| \right\} \right)_o \supseteq \mathcal{I} \left(\max \left\{ \varphi, \frac{1 + \varepsilon_0}{j} \log |I_j| \right\} \right)_o,$$

which implies that all the multiplier ideals are equal. Consequently, $\mathcal{I}(\varphi)_o = \mathcal{I}(\varphi_A)_o$, for

$$\varphi_A = \frac{1}{j-1} \log |I_j|, \quad \text{where } j \gg 1 \text{ s.t. } \mathcal{I}(\varphi)_o = \mathcal{I} \left(\frac{j}{j-1} \varphi \right)_o.$$

■

The following is a sharp effectiveness result of the strong openness property, where we have proved the special case for the openness property.

THEOREM 2.37 (Guan 19'). *Let D be a pseudoconvex domain in \mathbb{C}^n containing the origin o , and φ a negative psh function on D such that $\varphi(o) = -\infty$. Suppose $F \in \mathcal{O}(D)$ and $\int_D |F|^2 e^{-2\varphi} < +\infty$. Then for any $p > 1$ satisfying*

$$\frac{p}{p-1} > \frac{\int_D |F|^2 e^{-2\varphi}}{C_{F, \mathcal{I}_+(c_o^F(\varphi)\varphi)_o}(D)},$$

we have $(F, o) \in \mathcal{I}(p\varphi)_o$. Here

$$c_o^F(\varphi) = \sup\{c \geq 0: |F|^2 e^{-2c\varphi} \text{ is locally integrable near } o\},$$

and

$$C_{F,I}(D) = \inf \left\{ \int_D |\tilde{F}|^2: \tilde{F} \in \mathcal{O}(D), (\tilde{F} - F, o) \in I \right\}.$$

2.5 Skoda's L^2 division theorem

THEOREM 2.38 (Skoda, 72'). *Let $\Omega \subseteq \mathbb{C}^n$ be a complete Kähler (\Leftarrow pseudoconvex) domain and $\varphi \in \text{Psh}(\Omega)$. Let $g = (g_1, \dots, g_r)$ be an r -tuple of holomorphic function on Ω s.t. $g^{-1}(0) = \emptyset$. Set $m = \min\{n, r-1\}$. Then for every holomorphy function f on Ω s.t.*

$$I = \int_{\Omega} |f|^2 |g|^{-2(m+1+\varepsilon)} e^{-\varphi} d\lambda < +\infty,$$

there exists holomorphic functions (h_1, \dots, h_r) on Ω s.t. $f = \sum g_j h_j$ and

$$\int_{\Omega} |h|^2 |g|^{-2(m+\varepsilon)} e^{-\varphi} d\lambda \leq \left(1 + \frac{m}{\varepsilon}\right) I.$$

Proof. See [6, pp. 388–392].

Consider the diagram of Hilbert spaces and linear operators:

$$\begin{array}{ccc} H_0 & \xrightarrow{T_1} & H_1 \\ \downarrow T_2 & & \\ H_2 & & \end{array}$$

where T_1 is continuous and T_2 is closed and densely defined. Let G_1 be a closed subspace of H_1 . Suppose $T_1(\text{Ker } T_2) \subset G_1$. We need to following lemma in functional analysis:

LEMMA 2.39. *If there exists some $c > 0$ such that*

$$\|T_1^* x_1 + T_2^* x_2\|_0 \geq c \|x_1\|_1$$

for every $x_1 \in G_1$ and $x_2 \in \text{Dom } T_2^$, then we have $T_1(\text{Ker } T_2) = G_1$.*

Moreover, for every $x_1 \in G_1$, there exists $x_0 \in \text{Ker } T_2$ s.t. $T_1 x_0 = x_1$ and

$$\|x_0\|_0 \leq \frac{1}{c} \|x_1\|_1.$$

Proof. Let $G_0 = \text{Ker } T_2$, and G_0^\perp be the orthogonal complement of G_0 in H_0 . Then $\text{Im } T_2^*$ is dense in G_0^\perp . Note that $G_0 = H_0/G_0^\perp$ be seen as the quotient space equipped with the quotient norm. Let $\widetilde{T}_1^*: H_1 \rightarrow G_0$ be the induced map of $T_1^*: H_1 \rightarrow H_0$. Then for every $u_1 \in H_1$, with respect to the quotient norm, we have

$$\|\widetilde{T}_1^* u_1\|_{G_0}^2 = \inf_{v_0 \in G_0^\perp} \|T_1^* u_1 + v_0\|_0^2 = \inf_{v'_0 \in \text{Im } T_2^*} \|T_1^* u_1 + v'_0\|_0^2 = \inf_{v_2 \in \text{Dom } T_2} \|T_1^* u_1 + T_2^* v_2\|_0^2 \geq c^2 \|u_1\|_1^2.$$

This implies for the given $x_1 \in G_1$,

$$\|\langle x_1, u_1 \rangle_1\|^2 \leq \|x_1\|_1^2 \|u_1\|_1^2 \leq \frac{1}{c^2} \|x_1\|^2 \|\widetilde{T}_1^* u_1\|_{G_0}^2,$$

which shows that the functional

$$\begin{aligned} \mathcal{L}: G_0 &\longrightarrow \mathbb{C} \\ w_1 = \widetilde{T}_1^* u_1 &\longmapsto \langle x_1, u_1 \rangle_1 \end{aligned}$$

is bounded. Thus, by the Riesz representation theorem, we can find some $x_0 \in G_0$ such that

$$\langle x_1, u_1 \rangle_1 = \mathcal{L}(\widetilde{T_1^* u_1}) = \left\langle x_0, \widetilde{T_1^* u_1} \right\rangle_{G_0} = \langle x_0, T_1^* u_1 + G_0^\perp \rangle_0 = \langle T_1 x_0, u_1 \rangle, \quad \forall u_1 \in H_1.$$

Consequently, we have $x_1 = T_1 x_0$, and the norm control comes from

$$\|x_0\|_0 \leq \|\mathcal{L}\|_{G_0 \rightarrow \mathbb{C}} \leq \frac{1}{c} \|x_1\|_1.$$

■

Let

$$\begin{array}{ccc} L_{(2)}^{n,0}(\Omega, e^{-\phi_0})^{\oplus r} = H_0 & \xrightarrow{T_1 := g} & H_1 = L_{(2)}^{n,0}(\Omega, e^{-\phi_1}) \\ \downarrow T_2 := \bar{\partial} & & \\ H_2 = \text{Ker} \left(L_{(2)}^{n,1}(\Omega, e^{-\phi_0})^{\oplus r} \xrightarrow{\bar{\partial}} L_{(2)}^{n,2}(\Omega, e^{-\phi_0})^{\oplus r} \right) & & \end{array}$$

where $\phi_0 = 2(m + \varepsilon) \log |g| + \varphi$, and $\phi_1 = 2(m + 1 + \varepsilon) \log |g| + \varphi$. Let $G_1 = H_1 \cap \text{Ker } \bar{\partial}$. We want to prove $G_1 = T_1(\text{Ker } T_2)$.

First, we can directly compute by the definition:

$$(T_1 s, u)_{H_0} = (s, T_1^* u)_{H_1}, \quad s \in H_0, \quad u \in H_1,$$

to get

$$T_1^* u = u \bar{g} e^{-\tau} = (u \bar{g}_1 e^{-\tau}, \dots, u \bar{g}_r e^{-\tau}), \quad \tau = \log |g|^2.$$

Thus, for our case, $\|T_1^* x_1\|_0^2 = \|x_1\|_1^2$ for every $x_1 \in H_1$, and especially when $x_1 \in G_1$ we have

$$T_2 T_1^* x_1 = \bar{\partial}(x_1 \bar{g} e^{-\tau}) = x_1 \bar{\partial}(\bar{g} e^\tau) =: x_1 \beta, \quad \text{where } \beta = \overline{\partial(g e^{-\tau})} \in \text{Ker } \bar{\partial}.$$

Then, for any real number $\alpha > 1$, and $x_1 \in G_1$, $x_2 \in \text{Dom}(T_2^*) \subset H_2$, we have

$$\begin{aligned} \|T_1^* x_1 + T_2^* x_2\|_0^2 &= \|T_1^* x_1\|_0^2 + \|T_2^* x_2\|_0^2 + 2\text{Re} \langle T_2 T_1^* x_1, x_2 \rangle_2 \\ &= \|x_1\|_1^2 + \|T_2^* x_2\|_0^2 + 2\text{Re} \langle x_1 \beta, x_2 \rangle_2 \\ &\geq \|x_1\|_1^2 + \|T_2^* x_2\|_0^2 - \alpha \int_\Omega |g|^2 \beta \cdot \bar{x}_2 e^{-\phi_0} dV - \frac{1}{\alpha} \int_\Omega |g|^{-2} |x_1|^2 e^{-\phi_0} dV \\ &= \left(1 - \frac{1}{\alpha}\right) \|x_1\|_1^2 + \|T_2^* x_2\|_0^2 - \alpha \left\| |g| \bar{\beta} \cdot x_2 \right\|_2^2. \end{aligned}$$

We need to estimate:

$$\|T_2^* u\|_0^2 - \alpha \left\| |g| \bar{\beta} \cdot u \right\|_2^2 = \|\bar{\partial}^* u\|_{H_0}^2 - \alpha \left\| |g| \partial(g e^{-\tau}) \cdot u \right\|_{H_2}^2 = \sum_{j=1}^r \|\bar{\partial}^* u_j\|^2 - \alpha \left\| |g| \partial(g e^{-\tau}) \cdot u \right\|^2$$

for every $u = (u_1, \dots, u_r) \in \text{Dom}(T_2^*) = \text{Dom}(\bar{\partial}^*) \cap H_2$. In fact, by the Bochner–Kodaira–Nakano identity and the approximation result, we have

$$\|\bar{\partial}^* u_j\|^2 = \|\bar{\partial}^* u_j\|^2 + \|\bar{\partial} u_j\|^2 \geq (i\Theta_{e^{-\phi_0}} \Lambda u_j, u_j), \quad j = 1, \dots, r,$$

for every $u \in \text{Dom}(\bar{\partial}^*) \cap H_2$. We also need the following

Skoda's basic inequality

$$m \sum_{j=1}^r (\mathrm{i} \partial \bar{\partial} \tau)(v_j, v_j) \geq |g|^2 |\partial(g e^{-\tau}) \cdot v|^2,$$

for every $v = (v_1, \dots, v_r) \in (T_\Omega)^{\oplus r}$ at any point in Ω . In other words, for every $\eta = (\eta_j)_{1 \leq j \leq r} \in \mathbb{C}^r$, we have

$$m \sum_{j=1}^r \sum_{\mu, \nu=1}^n \frac{\partial^2 \log |g|^2}{\partial z_\mu \partial \bar{z}_\nu} \eta_{j\mu} \bar{\eta}_{j\nu} \geq |g|^2 \left| \sum_{j=1}^r \sum_{\mu=1}^n \frac{\partial(g_j e^{-\tau})}{\partial \bar{z}_{j\mu}} \eta_\mu \right|^2.$$

Some calculations: On the one hand,

$$\partial \tau = \frac{\langle \partial g, g \rangle}{|g|^2}, \quad \bar{\partial} \tau = \frac{\langle g, \partial g \rangle}{|g|^2}.$$

\Rightarrow

$$\mathrm{i} \partial \bar{\partial} \tau = \mathrm{i} \frac{\langle \partial g, \partial g \rangle}{|g|^2} - \mathrm{i} \frac{\langle \partial g, g \rangle \wedge \langle g, \partial g \rangle}{|g|^4},$$

\Rightarrow

$$\mathrm{i} \partial \bar{\partial} \tau(\xi, \xi) = \frac{|g|^2 |\partial g \cdot \xi|^2 - |\langle \partial g \cdot \xi, g \rangle|^2}{|g|^4} = |g|^{-4} \sum_{1 \leq j < l \leq r} \sum_{\nu=1}^n \left| (g_j \partial_\nu g_l - g_l \partial_\nu g_j) \xi_\nu \right|^2,$$

for every $\xi = \sum_{\nu=1}^n \xi_\nu \frac{\partial}{\partial z_\nu} \in T_\Omega$, by Lagrange's equality. On the other hand,

$$|g|^2 |\partial(g e^{-\tau}) \cdot \eta|^2 = |g|^{-2} \left| \left(|g|^2 \partial g - g \langle \partial g, g \rangle \right) \cdot \eta \right|^2 = |g|^{-6} \left| \sum_{j,k=1}^r \sum_{\nu=1}^n \bar{g}_j (g_j \partial_\nu g_k - g_k \partial_\nu g_j) \eta_{k,\nu} \right|^2.$$

LEMMA 2.40. *Let $n, r \in \mathbb{Z}_{>0}$, and $(a_k), (b_k^\nu), (c_\nu^k)$ be complex numbers for $1 \leq k \leq r, 1 \leq \nu \leq n$. Then, with $m = \min\{n, r-1\}$, and $|a|^2 = \sum_k |a_k|^2$, we have*

$$\left| \sum_{j,k,\nu} \bar{a}_j (a_j b_k^\nu - a_k b_j^\nu) c_\nu^k \right|^2 \leq m |a|^2 \sum_{k,\nu} \sum_{j < l} \left| (a_j b_l^\nu - a_l b_j^\nu) c_\nu^k \right|^2.$$

Proof. The case $m = r-1$ is a direct consequence of the Cauchy-Schwarz inequality, but the case $n < r-1$ is a bit complicated. See [12, pp. 282–284]. \blacksquare

Now by Bochner-Kodaira-Nakano identity and Skoda's basic inequality, we have

$$\sum_{j=1}^r \|\bar{\partial}^* u_j\|^2 \geq (m + \varepsilon) \sum_{j=1}^r (\mathrm{i} \partial \bar{\partial} \tau)(u_j, u_j) \geq \left(1 + \frac{\varepsilon}{m}\right) \left| |g| \partial(g e^{-\tau}) \cdot u \right|^2.$$

Take $\alpha = 1 + \frac{\varepsilon}{m}$. Then we have $1 - \frac{1}{\alpha} = \frac{\varepsilon}{m + \varepsilon}$, and

$$\|T_1^* x_1 + T_2^* x_2\|_0^2 \geq \frac{\varepsilon}{m + \varepsilon} \|x_1\|_1^2, \quad x_1 \in G_1, \quad x_2 \in \mathrm{Dom}(T_2^*).$$

Thus, for every $x_1 \in G_1 = H_1 \cap \text{Ker } \bar{\partial}$, there exists $x_0 \in \text{Ker } T_2$ s.t. $T_1 x_0 = x_1$ and

$$\|x_0\|_0^2 \leq \frac{m+\varepsilon}{\varepsilon} \|x_1\|_1^2,$$

which is the desired result exactly ($f = x_1$ holomorphic, and $h = x_0$ holomorphic r -tuple, with $g \cdot h = f$ and the L^2 estimate). \blacksquare

实际上不要求 $g^{-1}(0) = \emptyset$. 如果我们进一步假设 $|g|$ 在 Ω 上有上界以及正的下界, 那么可以得到 $A^2(\Omega, e^{-\varphi})$ 中的元素都可以被 g 做除法, 并且被除之后的都仍然落在 $A^2(\Omega, e^{-\varphi})$ 中 (即 \mathbb{C}^n 中的弱拟凸域上的关于 psh 函数的加权 Bergman 空间中的 Corona 问题均是可解的).

DEFINITION 2.41. Let $k \in \mathbb{R}_+$, and $\mathcal{J} = (g_1, \dots, g_r) \neq (0)$ be an ideal of $\mathcal{O}_{n,o}$. Define:

- the ideal

$$\overline{\mathcal{J}}^{(k)} := \{f \in \mathcal{O}_{n,o} : |u| \leq C|g|^k \text{ near } o, \exists C > 0\},$$

where $|g|^2 = |g_1|^2 + \dots + |g_r|^2$;

- the ideal

$$\widehat{\mathcal{J}}^{(k)} := \left\{ \int_V |f|^2 |g|^{-2(k+\varepsilon)} d\lambda < +\infty, \exists V \ni o \text{ \& } \varepsilon > 0 \right\}.$$

In fact, we know $\varepsilon = 0$ is enough (e.g. by strong openness property), that is, $\widehat{\mathcal{J}}^{(k)} = \mathcal{J}(k \log |g|)_o$.

COROLLARY 2.42 (Briangon–Skoda, 74’). If set $p = \min\{n-1, r-1\}$, we have

1. $\widehat{\mathcal{J}}^{(k+1)} = \mathcal{J} \widehat{\mathcal{J}}^{(k)} = \overline{\mathcal{J}} \widehat{\mathcal{J}}^{(k)}$, for $k \geq p$;
2. $\overline{\mathcal{J}}^{(k+p)} \subset \widehat{\mathcal{J}}^{(k+p)} \subset \mathcal{J}^k$, for all $k \in \mathbb{Z}_+$.

Proof. Obviously, $\mathcal{J} \widehat{\mathcal{J}}^{(k)} \subset \overline{\mathcal{J}} \widehat{\mathcal{J}}^{(k)} \subset \widehat{\mathcal{J}}^{(k+1)}$. We only need to prove $\widehat{\mathcal{J}}^{(k+1)} \subset \mathcal{J} \widehat{\mathcal{J}}^{(k)}$.

If $r \leq n$, it immediately follows from the L^2 division theorem. If $r > n$, we need the following lemma:

LEMMA 2.43. If $\mathcal{J} = (g_1, \dots, g_r)$ and $r > n$, we can find elements $\widetilde{g}_1, \dots, \widetilde{g}_n$ in \mathcal{J} which are linear combinations of g_i :

$$\widetilde{g}_j = a_j \cdot g = \sum_{1 \leq k \leq r} a_{jk} g_k, \quad a_j \in \mathbb{C}^r \setminus \{0\},$$

such that $C^{-1}|g| \leq |\widetilde{g}| \leq C|g|$ near o . In particular, the ideal $J = (\widetilde{g}_1, \dots, \widetilde{g}_n) \subset \mathcal{J}$ satisfies $\overline{J}^{(k)} = \overline{\mathcal{J}}^{(k)}$ and $\widehat{J}^{(k)} = \widehat{\mathcal{J}}^{(k)}$ for all k .

Proof. See [5, pp. 119–120]. \blacksquare

Let $J \subset \mathcal{J}$ be as in the lemma. Then

$$\widehat{\mathcal{J}}^{(k+1)} = \widehat{J}^{(k+1)} \subset J \widehat{J}^{(k)} \subset \mathcal{J} \widehat{\mathcal{J}}^{(k)}, \quad \forall k \geq n-1.$$

The first result implies

$$\widehat{\mathcal{J}}^{(k+p)} = \mathcal{J}^k \widehat{\mathcal{J}}^{(p)}, \quad \forall k \in \mathbb{Z}_+,$$

by induction, which gives the second result. \blacksquare

COROLLARY 2.44. *The ideal $\overline{\mathcal{J}}$ is the integral closure of \mathcal{J} , i.e. the set of germs $f \in \mathcal{O}_{n,o}$ which satisfies*

$$f^d + a_1 f^{d-1} + \cdots + a_d = 0, \quad a_s \in \mathcal{J}^s, \quad 1 \leq s \leq d.$$

Proof. See [5, pp. 120–121].

If f is in the integral closure of \mathcal{J} , using $|a_s| \leq C|g|^s$ and

$$|\text{roots}| \leq 2 \max_{1 \leq s \leq d} |a_s|^{1/s},$$

we can get $|f| \leq C|g|$.

Let $f \in \overline{\mathcal{J}}$. Let v_1, \dots, v_N be generators of $\widehat{\mathcal{J}}^{(p)}$. Since $f v_j \in \overline{\mathcal{J}} \widehat{\mathcal{J}}^{(p)} = \mathcal{J} \widehat{\mathcal{J}}^{(p)}$, there exists elements $b_{jk} \in \mathcal{J}$ such that

$$f v_j = \sum_{1 \leq k \leq N} b_{jk} v_k.$$

Thus, f satisfies the equation

$$\det(f \delta_{jk} - b_{jk}) = 0.$$

■

The L^2 division theorem can also be used to prove Levi's problem.

Proof of Levi's problem. Let Ω be a bounded pseudoconvex domain, and let $a = (a_1, \dots, a_n) \in \partial\Omega$ be any boundary point. We prove that there exists a holomorphic function in Ω which can not be analytically extended across a .

Let $\varphi(z) = -2 \log \delta(z) \in \text{Psh}(\Omega)$. Then $\varphi(z) \geq -2 \log |z - a|$, and thus for every $\varepsilon > 0$,

$$\int_{\Omega} |z - a|^{-2(n+\varepsilon)} e^{-\varphi(z)} d\lambda < +\infty.$$

In addition, $z_1 - a_1, \dots, z_n - a_n$ have no common zeros in Ω . So we can divide 1 by $z_1 - a_1, \dots, z_n - a_n$, which gives $1 = \sum_{j=1}^n (z_j - a_j) h_j(z)$ with $h_j \in \mathcal{O}(\Omega)$ for every j . Then we can see that there must be one of h_j which can not be analytically extended across a . ■

3 Pluripotential Theory

The “weakly convergence” in this section are in fact “weak*-convergent” what we usually called.

3.1 Preliminaries: Distribution

Let Ω be an open subset of \mathbb{R}^n . Denote by $\mathcal{D}(\Omega)$ the set of smooth real functions on Ω compactly supported in Ω and by \mathcal{D}_K the functions in $\mathcal{D}(\Omega)$ supported in K . Then for every $s \in \mathbb{Z}_{\geq 0}$, we can define a seminorm:

$$p^s(u) := \sup_{x \in \Omega} \max_{|\alpha| \leq s} |u^{(\alpha)}(x)|, \quad \forall u \in \mathcal{D}(\Omega).$$

Besides, for every compact subset K of Ω , and $s \in \mathbb{Z}_{\geq 0}$, we may also define

$$p_K^s(u) := \sup_{x \in K} \max_{|\alpha| \leq s} |u^{(\alpha)}(x)|, \quad \forall u \in \mathcal{D}_K.$$

We can equip $\mathcal{D}(\Omega)$ with the topology defined by the family of seminorms $(p^s)_s$, or equivalently $(p_K^s)_{K,s}$, which makes $\mathcal{D}(\Omega)$ to be a locally convex space.

DEFINITION 3.1. A distribution Λ on Ω , denoted by $\Lambda \in \mathcal{D}'(\Omega)$, is a continuous linear functional on $\mathcal{D}(\Omega)$ (with respect to the above topology), i.e. $\Lambda: \mathcal{D}(\Omega) \rightarrow \mathbb{R}$ is linear and for every compact subset K of Ω , there exists $N \in \mathbb{Z}_{\geq 0}$ and $C > 0$ such that

$$|\Lambda(u)| \leq Cp_K^N(u), \quad \forall u \in \mathcal{D}_K.$$

EXAMPLE 3.2. • Let $p \in \Omega$, and

$$\delta_p(u) := u(p), \quad \forall u \in \mathcal{D}(\Omega).$$

Then the Dirac function (measure) $\delta_p \in \mathcal{D}'(\Omega)$.

- Suppose $f \in L_{\text{loc}}^1(\Omega)$. Define

$$\Lambda_f(u) := \int_{\Omega} u(x)f(x) \, dx, \quad \forall u \in \mathcal{D}(\Omega).$$

Then $\Lambda_f \in \mathcal{D}'(\Omega)$.

- If μ is a positive measure on Ω with $\mu(K) < +\infty$ for every compact $K \subset \Omega$, then

$$\Lambda_{\mu}(u) = \int_{\Omega} u \, d\mu, \quad \forall u \in \mathcal{D}(\Omega)$$

defines a distribution $\Lambda_{\mu} \in \mathcal{D}'(\Omega)$.

REMARK 3.3. • Let $\Lambda \in \mathcal{D}'(\Omega)$ and $\alpha \in \mathbb{Z}_{\geq 0}^n$. Then

$$(D^{\alpha}\Lambda)(u) = (-1)^{|\alpha|}\Lambda(D^{\alpha}u), \quad \forall u \in \mathcal{D}(\Omega)$$

defines a distribution $D^{\alpha}\Lambda \in \mathcal{D}'(\Omega)$.

- Suppose $\Lambda \in \mathcal{D}'(\Omega)$ and $f \in C^\infty(\Omega)$. Then

$$(f\Lambda)(u) = \Lambda(fu), \quad \forall u \in \mathcal{D}(\Omega)$$

defines a distribution $f\Lambda \in \mathcal{D}'(\Omega)$. The Leibniz rule can also be applied.

- Suppose $\Lambda_i \in \mathcal{D}'(\Omega)$, $i = 1, 2, \dots$, and

$$\Lambda u = \lim_{i \rightarrow \infty} \Lambda_i u, \quad \forall u \in \mathcal{D}(\Omega).$$

Then $\Lambda \in \mathcal{D}'(\Omega)$.

DEFINITION 3.4. The order of $\Lambda \in \mathcal{D}'(\Omega)$ is the smallest $N \in \mathbb{Z}_{\geq 0}$ such that there exists $C > 0$ such that

$$|\Lambda(u)| \leq Cp^N(u), \quad \forall u \in \mathcal{D}(\Omega).$$

The order of a distribution can be $+\infty$.

EXAMPLE 3.5. For every $p \in \Omega$ and $\alpha \in \mathbb{Z}_{\geq 0}^n$, the distribution $D^\alpha \delta_p$ is of order $N = |\alpha|$.

DEFINITION 3.6. The support of a distribution $\Lambda \in \mathcal{D}'(\Omega)$ (denoted by $\text{Supp } \Lambda$) is the smallest closed subset A of Ω , such that $\Lambda(u) = 0$ for all $u \in \mathcal{D}(\Omega \setminus A)$.

PROPOSITION 3.7. If $\Lambda \in \mathcal{D}'(\Omega)$ has compact support, then Λ has finite order.

Proof. See [17, pp. 164–165]. Take a cut-off function θ of the support, and notice $\theta\Lambda = \Lambda$:

$$|\Lambda u| = |(\theta\Lambda)u| = |\Lambda(\theta u)| \leq Cp_{K'}^N(\theta u) \leq C'p^N(u).$$

■

EXAMPLE 3.8. • $\text{Supp } \delta_p = \text{Supp } D^\alpha \delta_p = \{p\}$.

- Suppose $\Lambda \in \mathcal{D}'(\Omega)$, $p \in \Omega$, $\text{Supp } \Lambda = \{p\}$, and Λ has order N . Then there are constants c_α such that

$$\Lambda = \sum_{|\alpha| \leq N} c_\alpha D^\alpha \delta_p.$$

THEOREM 3.9. Suppose $\Lambda \in \mathcal{D}'(\Omega)$, and $K \subset \Omega$ compact. Then there exist a continuous function f on Ω and $\alpha \in \mathbb{Z}_{\geq 0}^n$ s.t.

$$\Lambda u = (-1)^{|\alpha|} \int_{\Omega} f(x) (D^\alpha u)(x) dx, \quad \forall u \in \mathcal{D}_K.$$

In particular, if Λ has compact support and of order N , then there exist finitely many compactly supported continuous functions f_β s.t.

$$\Lambda u = \sum_{\beta} (-1)^{|\beta|} \int_{\Omega} f_\beta(x) (D^\beta u)(x), \quad \forall u \in \mathcal{D}(\Omega).$$

PROPOSITION 3.10 (Approximation of distributions). Let $\Lambda \in \mathcal{D}'(\Omega)$, $\varepsilon > 0$ and $\rho_\varepsilon \in \mathcal{D}(\mathbb{R}^n)$ the smoothing kernel with support in $B(o, \varepsilon)$. Then

$$(\Lambda * \rho_\varepsilon)(x) := \Lambda(\rho_\varepsilon(x - \cdot))$$

defines a **smooth function** $\Lambda * \rho_\varepsilon$ on Ω_ε s.t. $\Lambda * \rho_\varepsilon$ converges to Λ in $\mathcal{D}'(\Omega)$ as $\varepsilon \rightarrow 0$. As current,

$$(\Lambda * \rho_\varepsilon)(u) = \Lambda(\rho_\varepsilon * u), \quad \forall u \in \mathcal{D}_K, \quad (\text{somehow nontrivial}).$$

Proof. See [17, p. 173]. ■

EXAMPLE 3.11. If $\Lambda = \delta_o$, then $(\delta_o * \rho_\varepsilon)(x) = \rho_\varepsilon(x - \cdot)(o) = \rho_\varepsilon(x)$.

EXAMPLE 3.12. The following is an extremely wrong calculation, which shows that it is not always appropriate to define the products of distributions (so for currents): Let $\varepsilon \geq 0$ and $H_\varepsilon \in C^\infty(\mathbb{R})$ be increasing with

$$H_\varepsilon(x) = \begin{cases} 1 & x \geq 0, \\ 0 & x \leq -\varepsilon. \end{cases}$$

Then $H_\varepsilon \rightarrow H = \chi_{[0,+\infty)}$ and $dH_\varepsilon \rightarrow \delta_0$ as distributions. We have (really?) $(H_\varepsilon \delta_0)(u) = \delta_0(H_\varepsilon u) = \delta_0(u) \Rightarrow H \delta_0 = \delta_0$ by letting $\varepsilon \downarrow 0$, while $\delta_0 \leftarrow d(H_\varepsilon^2) = 2H_\varepsilon dH_\varepsilon \rightarrow 2H \delta_0 \Rightarrow \delta_0 = H \delta_0 = 0$.

3.2 Current

DEFINITION 3.13. A current on a domain Ω in \mathbb{C}^n of bidegree $(n-p, n-q)$ (or bidimension (p, q)) is a differential form

$$T = \sum_{|I|=n-p, |J|=n-q} T_{IJ} dz_I \wedge d\bar{z}_J$$

whose coefficients T_{IJ} are distributions. The space of bidegree $(n-p, n-q)$ currents on Ω is denoted by $\mathcal{D}'_{p,q}(\Omega)$ or $\mathcal{D}'^{n-p, n-q}(\Omega)$.

Equivalently, a current of bidegree $(n-p, n-q)$ on Ω is a continuous linear functional on the space $\mathcal{D}^{p,q}(\Omega)$ of smooth (p, q) forms with compact support.

PROPOSITION 3.14 (Approximation of currents). Let $T \in \mathcal{D}'_{p,q}(\Omega)$ and $\rho_\varepsilon \in \mathcal{D}(\mathbb{R}^n)$ the smoothing kernel. Then

$$T * \rho_\varepsilon := \sum_{I,J} (T_{I,J} * \rho_\varepsilon) dz_I \wedge d\bar{z}_J$$

is a **smooth $(n-p, n-q)$ form** Ω_ε s.t. $T * \rho_\varepsilon$ converges to T in $\mathcal{D}'_{p,q}(\Omega)$ as $\varepsilon \rightarrow 0$.

DEFINITION 3.15. A current $T \in \mathcal{D}'_{p,p}(\Omega)$ is positive ($T \geq 0$) if $\langle T, u \rangle \geq 0$ for all test forms $u \in \mathcal{D}^{p,p}(\Omega)$ that are **strongly positive** at each point, i.e.,

$$u(x) = \sum \gamma_s i\alpha_{s,1} \wedge \bar{\alpha}_{s,1} \wedge \dots \wedge i\alpha_{s,p} \wedge \bar{\alpha}_{s,p}$$

where $\alpha_{s,j} \in T_{\Omega,x}^*$ and $\gamma_s \geq 0$. Denote by $T \in \mathcal{D}'_{p,p}^+(X)$.

REMARK 3.16. • Positive (p, p) forms are forms with L^1_{loc} coefficients that are positive as (p, p) currents.

- A form $u = i \sum_{j,k} u_{j,k} dz_j \wedge d\bar{z}_k$ of bidegree $(1, 1)$ is positive if and only if (u_{jk}) is semi-positive definite.
- The notions of positive and strongly positive (p, p) -forms coincide for $p = 0, 1, n-1, n$. But this is not true for other cases in general.
- If u_1, \dots, u_s are positive forms, and strongly positive except one, then $u_1 \wedge \dots \wedge u_s$ is positive.
- There is example shows: α, β positive forms but $\alpha \wedge \beta < 0$.

- $T \in \mathcal{D}'_{p,p}(\Omega)$ is positive iff $T \wedge u \in \mathcal{D}'_{(0,0)}(\Omega)$ is a positive measure for all strongly positive smooth (p,p) forms u on Ω (constant coefficients strongly positive forms are enough).

PROPOSITION 3.17. Every positive current T is real and of order 0, i.e. if

$$T = i^{(n-p)^2} \sum T_{I,J} dz_I \wedge d\bar{z}_J,$$

then the coefficients $T_{I,J}$ are complex measures and $\overline{T_{I,J}} = T_{J,I}$.

DEFINITION 3.18. If T is order 0, define the mass measure of T by

$$\|T\| := \sum_{I,J} |T_{I,J}|.$$

(This depends on the choice of coordinates).

PROPOSITION 3.19. If $T \in \mathcal{D}'_{p,p}(X)$ and $v \in C_c^{s,s}(X)$ is strongly positive, then $T \wedge v$ is a positive current.

Proof. See [6, pp. 133–134] or [2, p. 3]. ■

DEFINITION 3.20. A current $T \in \mathcal{D}'_{p,p}(\Omega)$ is closed if it is d-closed, i.e. $dT = 0$, where

$$\langle dT, u \rangle := (-1)^{q+1} \langle T, du \rangle, \quad \forall u \in \mathcal{D}^{(p-1,p-1)}(\Omega).$$

EXAMPLE 3.21. Let $\phi \in \text{Psh} \cap L^1_{\text{loc}}(\Omega)$. Then

$$T = i\partial\bar{\partial}\phi = i \sum_{1 \leq j,k \leq n} \frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} dz_j \wedge d\bar{z}_k$$

is a closed positive current of bidegree $(1,1)$.

Using Dolbeault-Grothendieck's lemma, conversely we have

PROPOSITION 3.22. If $\Theta \in \mathcal{D}'_{n-1,n-1}(\Omega)$ is a closed positive current of bidegree $(1,1)$, then for every point $z_0 \in \Omega$ there exists a neighborhood D of z_0 and $\phi \in \text{Psh}(D)$ s.t. $\Theta = i\partial\bar{\partial}\phi$.

Proof. See [6, p. 135]. ■

EXAMPLE 3.23 (see [5, p. 10], [6, p. 135]). Let $A \subset \Omega$ be a closed analytic subset of pure dimension p . Define

$$\langle [A], \alpha \rangle := \int_{A_{\text{reg}}} \alpha, \quad \alpha \in \mathcal{D}^{p,p}(\Omega).$$

The current $[A] \in \mathcal{D}'_{p,p}(\Omega)$ is a closed positive current (Lelong 57'; see [6, p. 140]).

若 A 是子流形, 这个 current 的良好性是直接的, 一般情形可以通过局部参数化定理来得到. 闭性在正则部分由 Stokes 定理得到, 奇异部分要用到 Skoda–El Mir 延拓定理.

DEFINITION 3.24. For every $T \in \mathcal{D}'_{p,p}(\Omega)$, the trace measure of T w.r.t. ω is the positive measure

$$\sigma_T = \frac{1}{2^p} T \wedge \omega^p.$$

When $\omega = \frac{i}{2} \sum_j dz_j \wedge d\bar{z}_j$, and $T = i^{(n-p)^2} \sum_{I,J} T_{IJ} dz_I \wedge d\bar{z}_J$,

$$\sigma_T = 2^{-p} \left(\sum_{|I|=n-p} T_{I,I} \right) i dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge i dz_n \wedge d\bar{z}_n.$$

EXAMPLE 3.25. Let Z be a p -dimensional complex analytic submanifold of Ω . Then

$$\sigma_{[Z]} = \text{Riemannian volume measure on } Z.$$

Proof. See [6, pp. 136–137]. ■

若没有 complex analytic 条件, 则 $\sigma_{[Z]} = \alpha dV_Z$ 且 $|\alpha| \leq 1$ 以及等号成立当且仅当 complex analytic. 这说明解析子流形是极小子流形. 实际上 current theory 非常普遍应用在几何测度论的研究中.

PROPOSITION 3.26. For positive current $T \in \mathcal{D}'_{p,p}(\Omega)$, the mass measure is dominated by its trace measure, that is

$$\|T\| \leq C\sigma_T, \quad C \text{ is a constant.}$$

Proof. See [6, pp. 133–134]. ■

REMARK 3.27. With the weak topology defined by seminorms $T \mapsto |\langle T, f_\nu \rangle|$, where (f_ν) dense in $\mathcal{D}_{p,p}(\Omega)$, $\mathcal{D}'_{p,p}(\Omega)$ is a metrizable subspace of $\mathcal{D}'_{p,p}(\Omega)$.

THEOREM 3.28 (Skoda-El Mir). Let $E \subset \Omega$ be a closed complete pluripolar set and $T \in \mathcal{D}'_{p,p}(\Omega, E)$ a closed positive current. Assume that T has finite mass in a neighborhood of every point in E . Then the trivial extension $\tilde{T} \in \mathcal{D}'_{p,p}(\Omega)$ (obtained by extending measure T_{IJ} by 0 on E) is closed on Ω .

Proof. See [6, pp. 138–140]. ■

DEFINITION 3.29. A current Θ is said to normal if Θ and $d\Theta$ are currents of order 0.

实际上我们这里不关心 normal current, 只关心其中的 closed positive current, 但因为这个概念太有名了于是提一下.

PROPOSITION 3.30. Let $\Theta \in \mathcal{D}'_{p,p}(\Omega)$ be a normal current. If $\text{Supp } \Theta$ is contained in an analytic subset A of dimension $< p$, then $\Theta = 0$.

Proof. See [6, p. 141].

On A_{reg} , observe that $\bar{\partial}g_k \wedge \Theta = \bar{\partial}(g_k \Theta) - g_k \bar{\partial}\Theta = 0$ for every generator g_k of A . Use dimensional assumption to deduce each $\Theta_{IJ} = 0$. Then $\text{Supp } \Theta \subset A_{\text{sing}}$ and conclude by induction on $\dim A$. ■

PROPOSITION 3.31. Let A be an analytic subset of Ω with global irreducible components A_j of pure dimension p . Then any closed current $\Theta \in \mathcal{D}'_{p,p}(\Omega)$ of order 0 with support in A is of the form

$$\Theta = \sum_j \lambda_j [A_j], \quad \lambda_j \in \mathbb{C}.$$

Moreover, Θ is positive iff all $\lambda_j \geq 0$.

这地方就类似于除子了. 但是除子我们可不会让系数取复数值, 这也说明 current 作为一个更灵活的工具可以作为应用.

Proof. See [6, p. 143]. ■

THEOREM 3.32 (Lelong-Poincaré equation). *Let $f \in \mathcal{M}(\Omega)$ be a meromorphic function, and let $\sum m_j Z_j$ be the divisor of f . Then*

$$\frac{i}{\pi} \partial \bar{\partial} \log |f| = \sum m_j [Z_j]$$

in the space $\mathcal{D}'_{n-1, n-1}(X)$.

Proof. See [6, p. 143].

Let $Z = \bigcup Z_j$ be its support. Near $a \in Z_j \cap Z_{\text{reg}}$, using Weierstrass' preparation theorem, write $f(w) = u(w)w_1^{m_j}$. Note that $i\partial\bar{\partial} \log |w_1|^2 = \pi[\{w_1 = 0\}]$. At last, use Proposition 3.30. ■

3.3 Complex Monge-Ampère operator

3.3.1 Definition: Case of locally bounded psh function

Denote

$$d = \partial + \bar{\partial}, \quad d^c = \frac{1}{2\pi i}(\partial - \bar{\partial}).$$

Then $dd^c = \frac{i}{\pi} \partial \bar{\partial}$.

PROPOSITION 3.33. *Let Ω be a smoothly bounded open set and let f, g be forms of class C^2 on $\bar{\Omega}$ of pure bidegree (p, p) and (q, q) with $p + q = n - 1$. Then*

$$\int_{\Omega} f \wedge dd^c g - dd^c f \wedge g = \int_{\partial\Omega} f \wedge d^c g - d^c f \wedge g.$$

Proof. See [6, p. 144]. Using Stokes' formula. ■

Let u be a **locally bounded** psh function and T a closed positive current.

DEFINITION 3.34 (Bedford-Taylor 82'). Then uT is well-defined: u is locally Borel function and T has measure coefficients. Define

$$dd^c u \wedge T := dd^c(uT).$$

REMARK 3.35. *Generally, we can not define the product of two currents (Kiselman 83').*

PROPOSITION 3.36. $dd^c u \wedge T$ is a closed positive current.

Proof. See [6, p. 145].

Let smooth $u_k \rightarrow u$ decreasing convergent. Note that dd^c is weakly continuous. ■

DEFINITION 3.37. Given locally bounded psh functions u_1, \dots, u_q , define inductively

$$dd^c u_1 \wedge dd^c u_2 \wedge \dots \wedge dd^c u_q \wedge T := dd^c (u_1 dd^c u_2 \wedge \dots \wedge dd^c u_q \wedge T).$$

In particular, for locally bounded psh function u , the bidegree (n, n) current $(dd^c u)^n$ is well-defined and is a positive measure.

REMARK 3.38. If u is of class C^2 , then

$$(\mathrm{dd}^c u)^n = \det \left(\frac{\partial^2 u}{\partial z_j \partial \bar{z}_k} \right) \cdot \frac{n!}{\pi^n} \mathrm{id}z_1 \wedge \mathrm{d}\bar{z}_1 \wedge \dots \wedge \mathrm{id}z_n \wedge \mathrm{d}\bar{z}_n.$$

All operators $(\mathrm{dd}^c)^q$ are called Monge-Ampère operators.

THEOREM 3.39 (Chern–Levine–Nirenberg inequality, 69'). For all compact subsets K, L of Ω with $L \subset K^\circ$, there exists a constant $C_{K,L} \geq 0$ s.t.

$$\|\mathrm{dd}^c u_1 \wedge \mathrm{dd}^c u_2 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T\|_L \leq C_{K,L} \|u_1\|_{L^\infty(K)} \cdots \|u_q\|_{L^\infty(K)} \|T\|_K,$$

where $\|\Theta\|_K = \|\Theta\|(K) = \int_K \sum_{I,J} |T_{I,J}|$.

Proof. See [6, p. 146].

Note that $\|T\|$ and σ_T are dominated by each other when T is positive. ■

COROLLARY 3.40. Let u_1, \dots, u_q be *continuous* psh functions, and u_1^k, \dots, u_q^k be sequences of psh functions s.t. $u_j^k \rightarrow u_j$ *locally uniformly* for each j . If a sequence of closed positive currents $T_k \rightarrow T$ weakly, then

1. $u_1^k \mathrm{dd}^c u_2^k \wedge \dots \wedge \mathrm{dd}^c u_q^k \wedge T_k \longrightarrow u_1 \mathrm{dd}^c u_2 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ weakly;
2. $\mathrm{dd}^c u_1^k \wedge \dots \wedge \mathrm{dd}^c u_q^k \wedge T_k \longrightarrow \mathrm{dd}^c u_k \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ weakly.

Proof. See [6, p. 147].

(2) implies (1) due to the weak continuity of dd^c .

By induction, only need to prove (1) when $q = 1$. Let u_ε be regularization of u , and write

$$u^k T_k - uT = (u^k - u)T_k + (u - u_\varepsilon)T_k + u_\varepsilon(T_k - T).$$

Then apply the Chern–Levine–Nirenberg inequality to the first two terms and $u_\varepsilon(T_k - T)$ converges weakly to 0 since u_ε is smooth. ■

THEOREM 3.41 (Bedford–Taylor 82'). Let u_1, \dots, u_q be *locally bounded* psh functions, and u_1^k, \dots, u_q^k be *decreasing* sequences of psh functions s.t. $u_j^k \rightarrow u_j$ *pointwise* for each j . Then

1. $u_1^k \mathrm{dd}^c u_2^k \wedge \dots \wedge \mathrm{dd}^c u_q^k \wedge T \longrightarrow u_1 \mathrm{dd}^c u_2 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ weakly;
2. $\mathrm{dd}^c u_1^k \wedge \dots \wedge \mathrm{dd}^c u_q^k \wedge T \longrightarrow \mathrm{dd}^c u_k \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ weakly.

Proof. See [6, pp. 147–149]. ■

COROLLARY 3.42 ([6, p. 149]). The product $\mathrm{dd}^c u_1 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ is symmetric w.r.t. u_1, \dots, u_q .

3.3.2 Definition: Case of unbounded psh function

Assume that Ω is pseudoconvex. Let $u \in \text{Psh}(\Omega)$ and $T \in \mathcal{D}'_{p,p}(\Omega)$ a closed positive current on Ω of bidimension (p, p) .

DEFINITION 3.43. The unbounded locus $L(u)$ is defined as:

$$L(u) := \{x \in \Omega : u \text{ is unbounded in every neighborhood of } x\} \subseteq u^{-1}(-\infty).$$

PROPOSITION 3.44. *If*

1. T has nonzero bidimension (p, p) (i.e. degree of $T < 2n$),
2. the boundary $\partial\Omega \cap L(u) \cap \text{Supp } T = \emptyset$,

then the current uT has locally finite mass in Ω .

Proof. See [6, pp. 150–152].

(1) is necessary: $T = \delta_0$, $u = \log |z|$. ■

DEFINITION 3.45. Let $u_1, \dots, u_q \in \text{Psh}(\Omega)$, and $\partial\Omega \cap L(u_j) \cap \text{Supp } T = \emptyset$. Define inductively

$$\text{dd}^c u_1 \wedge \text{dd}^c u_2 \wedge \dots \wedge \text{dd}^c u_q \wedge T := \text{dd}^c (u_1 \text{dd}^c u_2 \wedge \dots \wedge \text{dd}^c u_1 \wedge T).$$

If u_1^k, \dots, u_q^k are decreasing sequence of psh functions with $u_j^k \rightarrow u_j$ pointwise, then Theorem 3.41 (1,2) hold.

Proof. See [6, p. 152]. ■

REMARK 3.46. [6, Proposition 4.4, p. 152] Moreover, we can define $V \text{dd}^c u_1 \wedge \dots \wedge \text{dd}^c u_q$ for $V \in \text{Psh}(\Omega)$, when $\partial\Omega \cap L(u_j) = \emptyset$ (V can be unbounded near $\partial\Omega$, but it is OK due to V is L^1_{loc}).

REMARK 3.47. [6, Theorem 4.5, pp. 152–156] The currents $u_1 \text{dd}^c u_2 \wedge \dots \wedge \text{dd}^c u_q \wedge T$ and $\text{dd}^c u_1 \wedge \dots \wedge \text{dd}^c u_1 \wedge T$ are also well-defined and has locally finite mass in Ω if

1. $q \leq p$;
2. and the Hausdorff measure

$$\mathcal{H}_{2p-2m+1}(L(u_{j_1}) \cap \dots \cap L(u_{j_m}) \cap \text{Supp } T) = 0$$

for all choices of indices $j_1 < \dots < j_m$ in $\{1, \dots, q\}$. Moreover, the analogue of the monotone convergence Theorem 3.41 (1,2) hold.

REMARK 3.48. [6, Proposition 4.12, pp. 156–157] Let $u_j = \log |f_j|$ for $f_j \in \mathcal{O}^*(\Omega)$, and $[Z_j] = \text{dd}^c u_j$, $1 \leq j \leq q$. If the supports satisfy

$$\text{codim}(|Z_{j_1}| \cap \dots \cap |Z_{j_m}|) = m, \quad \forall m,$$

then it follows from the above remark that $[Z_1] \wedge \dots \wedge [Z_q]$ is well-defined.

Moreover, if $(C_k)_{k \geq 1}$ is the irreducible components of the point set intersection $|Z_1| \cap \dots \cap |Z_q|$, then there exists integers $m_k > 0$ s.t.

$$[Z_1] \wedge \dots \wedge [Z_q] = \sum m_k [C_k].$$

The integer m_k is the multiplicity of intersection of Z_1, \dots, Z_q along the component C_k .

3.4 Lelong Number

3.4.1 Demailly's generalized Lelong number

Let $\Omega \subset \mathbb{C}^n$ be a bounded pseudoconvex domain, and $\varphi: \Omega \rightarrow [-\infty, +\infty)$ a continuous (here means e^φ is continuous) psh function. Set

$$\begin{aligned} S(r) &= \{x \in \Omega: \varphi(x) = r\}, \\ B(r) &= \{x \in \Omega: \varphi(x) < r\}, \\ \overline{B}(r) &= \{x \in \Omega: \varphi(x) \leq r\}. \end{aligned}$$

DEFINITION 3.49. We say φ is semi-exhaustive if $\exists R$ s.t. $B(R) \Subset \Omega$; semi-exhaustive on $A \subset \Omega$ closed if $A \cap B(R) \Subset \Omega$, $\exists R$.

We now let $T \in \mathcal{D}'_{p,p}(\Omega)$, and φ semi-exhaustive on Ω . Assume $B(R) \cap \text{Supp } T \Subset \Omega$.

DEFINITION 3.50. Set for $r \in (-\infty, R)$,

$$\begin{aligned} \nu(T, \varphi, r) &= \int_{B(r)} T \wedge (\text{dd}^c \varphi)^p, \\ \nu(T, \varphi) &= \int_{S(-\infty)} T \wedge (\text{dd}^c \varphi)^p = \lim_{r \rightarrow -\infty} \nu(T, \varphi, r). \end{aligned}$$

PROPOSITION 3.51. For convex increasing $\chi: \mathbb{R} \rightarrow \mathbb{R}$,

$$\int_{B(r)} T \wedge (\text{dd}^c \chi \circ \varphi)^p = \chi'(r - 0) \nu(T, \varphi, r).$$

Proof. See [6, p. 158]. ■

EXAMPLE 3.52. • Let $\chi(r) = e^{2r}$:

$$\nu(T, \varphi, r) = e^{-2pr} \int_{B(r)} T \wedge (\text{dd}^c e^{2\varphi})^p.$$

- Let $\varphi(z) = \log |z - a|$:

$$\mu(T, \varphi, \log r) = \frac{1}{r^{2p}} \int_{|z-a| < r} T \wedge (\text{dd}^c |z|^2)^p = \frac{\sigma_T(B(a, r))}{\pi^p r^{2p}/p!}.$$

- Denote $\nu(T, a) = \nu(T, \log |z - a|)$, which is the (ordinary) Lelong number.
- When $T = 1 = [\Omega]$, $\varphi = \log |z - a|$, we have

$$\int_{B(a, r)} (\text{dd}^c \log |z - a|)^n = \nu(T, \varphi) = \frac{|B(a, r)|}{\pi^n r^{2n}/n!} = 1.$$

Thus, $(\text{dd}^c \log |z - a|)^n = \delta_a$.

- When $T = [A]$, and A is an analytic subset of pure dimension p ,

$$\nu([A], x) = \begin{cases} 0 & x \notin A, \\ 1 & x \in A \text{ is a regular point.} \end{cases}$$

Since $\sigma_{[A]}$ is the volume measure on A near regular points.

PROPOSITION 3.53. *For every $c > 0$, the set $E_c = \{x \in \Omega : \nu(T, x) \geq c\}$ is a closed set of locally finite \mathcal{H}_{2p} Hausdorff measure in Ω .*

Proof. See [6, p. 160]. ■

REMARK 3.54. • $T_k \rightarrow T$ weakly \Rightarrow

$$\limsup_{k \rightarrow \infty} \nu(T_k, \varphi) \leq \nu(T, \varphi).$$

- $e^{\varphi_k} \rightarrow e^{\varphi}$ compactly uniform \Rightarrow

$$\limsup_{k \rightarrow \infty} \nu(T, \varphi_k) \leq \nu(T, \varphi).$$

Set $\varphi_{\geq r} = \max\{\varphi, r\}$. Then

$$(\mathrm{dd}^c \varphi_{\geq r})^n = \begin{cases} (\mathrm{dd}^c \varphi)^n & \text{on } \Omega \setminus \overline{B}(r), \\ 0 & \text{on } B(r). \end{cases}$$

DEFINITION 3.55. Define the measure

$$\mu_r = (\mathrm{dd}^c \varphi_{\geq r})^n - \mathbb{I}_{\Omega \setminus B(r)} (\mathrm{dd}^c \varphi)^n, \quad r \in (-\infty, R).$$

Then the measure μ_r is supported on $S(r)$ and $\mu_r(S(r)) = \mu_r(B(s)) = \int_{B(r)} (\mathrm{dd}^c \varphi)^n$ for $s \geq r$.

THEOREM 3.56 (Jensen-Lelong formula). *Let $V \in \mathrm{Psh}(\Omega)$. Then*

$$\mu_r(V) - \int_{B(r)} V(\mathrm{dd}^c \varphi)^n = \int_{-\infty}^r \nu(\mathrm{dd}^c V, \varphi, t) dt.$$

Proof. See [6, pp. 163–164]. ■

For $r < r_0 < R$, we then have

$$\mu_r(V) - \mu_{r_0}(V) + \int_{B(r_0) \setminus B(r)} V(\mathrm{dd}^c \varphi)^n = \int_{r_0}^r \nu(\mathrm{dd}^c V, \varphi, t) dt.$$

COROLLARY 3.57. *If $(\mathrm{dd}^c \varphi)^n = 0$ on $\Omega \setminus S(-\infty)$, then $r \mapsto \mu_r(V)$ is convex increasing, and*

$$\mu(\mathrm{dd}^c V, \varphi) = \lim_{r \rightarrow -\infty} \frac{\mu_r(V)}{r}.$$

Proof. See [6, p. 164].

Because $\nu(\mathrm{dd}^c V, \varphi, t)$ is increasing and non-negative. ■

EXAMPLE 3.58. If $\varphi(z) = \log |z - a|$, then

$$\nu(\text{dd}^c V, \log |z - a|, \log t) = \frac{(n-1)!}{\pi^{n-1} t^{2n-2}} \int_{B(a,t)} \frac{1}{2\pi} \Delta V.$$

Thus,

$$\nu(\text{dd}^c V, a) = \lim_{r \rightarrow 0} \frac{\int_{S(a,r)} V \frac{dS}{\lambda(S(a,r))}}{\log r} = \lim_{r \rightarrow 0} \frac{\sup_{S(a,r)} V}{\log r} = \sup\{\gamma : V \leq \gamma \log |z - a| + O(1) \text{ near } a\},$$

by Harnack's inequality and the convexity of $\log r \mapsto \sup_{S(a,r)} V$.

THEOREM 3.59. Let $\varphi, \psi \in C(\Omega) \cap \text{Psh}(\Omega)$, and semi-exhaustive on $\text{Supp } T$. If

$$\ell := \limsup \frac{\psi(x)}{\varphi(x)} < +\infty, \quad \text{as } x \in \text{Supp } T \text{ and } \varphi(x) \rightarrow -\infty,$$

then $\nu(T, \psi) \leq \ell \nu(T, \varphi)$, and the equality holds if $\ell = \lim \psi/\varphi$.

Proof. See [6, p. 166].

Set $u_c = \max\{\psi - c, \varphi\}$ for c large enough. ■

REMARK 3.60. • This shows the Lelong numbers $\nu(T, x)$ are independent of the choice of local coordinates.

- For $\lambda_1, \dots, \lambda_n > 0$, it implies

$$\left(\text{ddc} \log \max_{1 \leq j \leq n} |z_j|^{\lambda_j} \right)^n = \left(\text{ddc} \log \sum_{1 \leq j \leq n} |z_j|^{\lambda_j} \right)^n = \lambda_1 \cdots \lambda_n \delta_0.$$

- The Kiselman number is

$$\nu(T, x, \lambda) := \nu \left(T, \log \max_{1 \leq j \leq n} |z_j - x_j|^{\lambda_j} \right),$$

and $\nu(T, x) = \nu(T, x, (1, \dots, 1))$.

THEOREM 3.61 (Thie 67'). Let A be an analytic set of dimension p in \mathbb{C}^n . The **Local parametrization theorem** shows we can take coordinates $(z'; z'') = (z_1, \dots, z_d; z_{d+1}, \dots, z_n)$ near x such that (A, x) is contained in $|z''| \leq C|z'|$, and the projection map $\pi : A \cap (\Delta' \times \Delta'') \rightarrow \Delta'$ is a ramified covering with m sheets. The $m \in \mathbb{Z}_{\geq 0}$ is called the multiplicity of A at x . Then

$$\nu([A], x) = m.$$

Proof. See [6, p. 168]. ■

THEOREM 3.62. Let u_1, \dots, u_q and v_1, \dots, v_q be psh functions s.t. each q -tuple satisfies the conditions in Definition 3.45 or Remark 3.47. Assume $u_j = -\infty$ on $\text{Supp } T \cap \varphi^{-1}(-\infty)$ and

$$\ell_j := \limsup \frac{v_j(z)}{u_j(z)} \quad \text{when } z \in \text{Supp } T \setminus u_j^{-1}(-\infty), \varphi(z) \rightarrow -\infty.$$

Then

$$\nu(\text{dd}^c v_1 \wedge \dots \wedge \text{dd}^c v_q \wedge T, \varphi) \leq \ell_1 \cdots \ell_q \nu(\text{dd}^c u_1 \wedge \dots \wedge \text{dd}^c u_q \wedge T, \varphi).$$

Proof. See [6, p. 169].

Set $w_{j,c} = \max\{v_j - c, u_j\}$ for c large. ■

COROLLARY 3.63. *If $\mathrm{dd}^c u_1 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T$ is well-defined, then at each $x \in \Omega$,*

$$\nu(\mathrm{dd}^c u_1 \wedge \dots \wedge \mathrm{dd}^c u_q \wedge T, x) \geq \nu(\mathrm{dd}^c u_1, x) \cdots \nu(\mathrm{dd}^c u_q, x) \nu(T, x).$$

Proof. See [6, p. 169]. Take $\varphi(z) = v_1(z) = \dots = v_q(z) = \log |z - x|$. ■

3.5 Siu's theorems

3.5.1 Siu's slicing theorem

THEOREM 3.64 (Siu's slicing theorem). *For almost $S \in G(q, n)$ with $q \geq n - p$, the slice $T|_S$ satisfies $\nu(T|_S, 0) = \nu(T, 0)$, where $G(q, n)$ is the Grassmannian of q -dimensional subspace in \mathbb{C}^n .*

Proof. See [6, pp. 172–173].

The slice of current is somehow a little complicated, but we can think about the psh functions. One important ingredient in the proof is **Crofton's formula**:

$$\int_{S \in G(q, n)} [S] \, dv(S) = (\mathrm{dd}^c \log |z|)^{n-q},$$

where dv is the unique $U(n)$ -invariant measure of mass 1 on $G(q, n)$.

Think that ($p = n - 1, q = 1$ and $T = \mathrm{dd}^c \psi$ smooth)

$$\nu(\psi|_S, 0, r) = \int_{S \cap B(0, r)} T|_S = \int_{B(0, r)} T \wedge S,$$

so

$$\int_{S \in \mathbb{CP}^{n-1}} \nu(\psi|_S, 0, r) \, dv(S) = \int_S \int_{B(0, r)} T \wedge [S] = \int_{B(0, r)} T \wedge (\mathrm{dd}^c \log |z|)^{n-1} = \nu(\psi, 0, r).$$

In this special case, we also have $\nu(\psi|_S, 0) \leq \nu(\psi, 0)$ for every S . ■

EXAMPLE 3.65. Let $\psi = \log \max_{i=1,2} |z_i|^{\lambda_i}$, where $0 < \lambda_1 < \lambda_2$. Then $\nu(\psi|_S, 0) = \lambda_2$ except for $S = \{z_2 = 0\}$.

3.5.2 Siu's semi-continuity theorem

Let $\varphi: \Omega \times D \rightarrow [-\infty, +\infty)$ be a continuous psh function, and $T \in \mathcal{D}_{p,p}^+(\Omega)$. Assume for every compact subset $K \subset D$, there exists $R = R(K) < 0$ s.t.

$$\{(x, y) \in \mathrm{Supp} T \times K : \varphi(x, y) \leq R\} \Subset \Omega \times D.$$

For every $y \in D$, let $\varphi_y(x) = \varphi(x, y)$. Assume that φ is locally Hölder continuous w.r.t. y , i.e. locally we have

$$|\varphi(x, y_1) - \varphi(x, y_2)| \leq M |y_1 - y_2|^\gamma, \quad \gamma \in (0, 1].$$

THEOREM 3.66 (Demailly 87'). *The upperlevel sets*

$$E_c(T, \varphi) = \{y \in D : \nu(T, \varphi_y) \geq c\}$$

are analytic subsets of Y .

Proof. See [6, pp. 173–180].

The proof is an application of Kiselman's minimum principle and Corollary 2.7. The proof is complicated, but we will give a proof for the semi-continuity of relative types (another generalization of Lelong number) with the similar methods later. ■

COROLLARY 3.67 (Siu's semi-continuity theorem). *If $T \in \mathcal{D}_{p,p}^+(\Omega)$, then*

$$E_c(T) = \{x \in \Omega : \nu(T, x) \geq c\}$$

of Lelong numbers are analytic subsets of dimension $\leq p$.

Proof. Let $\varphi(x, y) = \log |x - y|$. ■

3.5.3 Siu's decomposition theorem

THEOREM 3.68. *If $T \in \mathcal{D}_{p,p}^+(\Omega)$, there is a unique decomposition of T as a (finite or infinite) weakly convergent series*

$$T = \sum_{j \geq 1} \lambda_j [A_j] + R, \quad \lambda_j > 0,$$

where A_j are irreducible p -dimensional analytic subset, and R is a closed positive current with the property that $\dim E_c(R) < p$ for every $c > 0$.

Proof. See [6, p. 181].

Existence: Let A_j be the p -dimensional irreducible components occuring in one of $E_c(T)$, $c \in \mathbb{Q}_+$, and $\lambda_j > 0$ be the generic Lelong number

$$\lambda_j := \inf\{\nu(T, x) : x \in A_j\}.$$

The rest is to show $R_N = T - \sum_{1 \leq j \leq N} \lambda_j [A_j]$ is positive. ■

EXAMPLE 3.69. Let $\psi \in \text{Psh}^-(\Delta)$ in \mathbb{C} , and $\nu(\psi, 0) = c \in (0, +\infty)$. Then $0 \in E_c(\psi)$. Since the 0-dimensional irreducible analytic subset must be a point, by Siu's decomposition, we have

$$0 \leq R := T - c[0] = \text{dd}^c \psi - c \text{dd}^c \log |z| = \text{dd}^c(\psi - c \log |z|),$$

that is, $\phi := \psi - c \log |z| \in \text{Psh}(\Delta)$. Note that $\phi \leq 0$ near $\partial\Delta$, so $\phi \in \text{Psh}^{-1}(\Delta)$. Thus, $\psi \leq c \log |z|$ on Δ . From Siu's decomposition (or by definition), we can also find that $\nu(\phi, 0) = 0$.

Now if $c \in [k, k+1)$ for $k \in \mathbb{Z}_{\geq 0}$, then

$$\int_{\Delta(r)} |f|^2 e^{-2\psi} \approx \int_{\Delta(r)} |z|^{2m} e^{-2c \log |z| - 2\phi} = \int_{\Delta(r)} |z|^{2m-2c} e^{-2\phi}, \quad m = \text{ord}_0(f).$$

Since $\nu(\phi, 0) = 0$, by Skoda's result, $e^{-2\phi}$ is locally integrable. We can then conclude that $\mathcal{J}(\psi)_0 = (z^k)_0$.

3.6 Cegrell's class

DEFINITION 3.70. A domain $\Omega \subset \mathbb{C}^n$ is said to be hyperconvex if there exists a bounded psh exhaustion function, i.e. $\exists \varphi \in \text{Psh}^-(\Omega)$, s.t. $\{\varphi < -t\} \Subset \Omega$ for any $t < 0$.

In this “subsection”, we will always assume that Ω is a bounded hyperconvex domain in \mathbb{C}^n .

PROPOSITION 3.71. Suppose $u \in \text{Psh}^-(\Omega)$. Then there is a decreasing sequence of functions $u_j \in \text{Psh}(\Omega) \cap C(\overline{\Omega})$ with $u_j|_{\partial\Omega} = 0$ for every $j \in \mathbb{Z}_{>0}$, $\lim_{j \rightarrow \infty} u_j(z) = u(z)$ for every $z \in \Omega$, and $\int_{\Omega} (\text{dd}^c u_j)^n < +\infty$.

Proof. See [4, p. 3]. ■

DEFINITION 3.72. $\mathcal{E}_0(\Omega)$ is the convex cone of bounded psh function u with $\lim_{z \rightarrow \xi} u(z) = 0$ for any $z \in \partial\Omega$, and $\int_{\Omega} (\text{dd}^c u)^n < +\infty$.

REMARK 3.73. For $u \in \text{Psh}^-(\Omega) \setminus \{0\}$ with $\lim_{z \rightarrow \xi} u(z) = 0$ for any $z \in \partial\Omega$, and T a closed positive current on Ω , the current $\text{dd}^c u \wedge T$ is well-defined.

DEFINITION 3.74. Let $u \in \text{Psh}^-(\Omega)$. We say $u \in \mathcal{E}(\Omega)$ if for every $z_0 \in \Omega$ there is a neighborhood ω of z_0 in Ω and a decreasing sequence $h_j \in \mathcal{E}_0(\Omega)$ s.t. $h_j \rightarrow u$ on ω and $\sup_j \int_{\Omega} (\text{dd}^c h_j)^n < +\infty$.

For $u_1, \dots, u_n \in \mathcal{E}(\Omega)$, define $\text{dd}^c u_1 \wedge \text{dd}^c u_2 \wedge \dots \wedge \text{dd}^c u_n$ to be the (weakly convergence) limit measure obtained by the decreasing sequences (independent of the choice in fact, see [4, pp. 7–8]).

DEFINITION 3.75 (Cegrell's class). We denote by $\mathcal{F}(\Omega)$ the subclass of functions u in $\mathcal{E}(\Omega)$ s.t. there exists a decreasing sequence $u_j \in \mathcal{E}_0(\Omega)$ s.t. $u_j \rightarrow u$ on Ω and $\sup_j \int_{\Omega} (\text{dd}^c u_j)^n < +\infty$.

REMARK 3.76. • The integration by parts is allowed in $\mathcal{F}(\Omega)$.

• Every $u \in \mathcal{E}(\Omega)$ is locally in $\mathcal{F}(\Omega)$.

• In the unit ball \mathbb{B} of \mathbb{C}^2 , $u = \log |z_2| \notin \mathcal{E}(\mathbb{B})$. Moreover, $-(-\log |z_2|)^{\alpha} \in \mathcal{E}(\mathbb{B})$ iff $0 < \alpha < 1/2$.

THEOREM 3.77. Suppose $u_1, \dots, u_n \in \mathcal{F}(\Omega)$ and $h \in \mathcal{E}_0(\Omega)$. Then

$$\int_{\Omega} -h \text{dd}^c u_1 \wedge \dots \wedge \text{dd}^c u_n \leq \prod_{j=1}^n \left(\int_{\Omega} -h (\text{dd}^c u_j)^n \right)^{1/n}.$$

Especially, we have

$$\int_{\Omega} \text{dd}^c u_1 \wedge \dots \wedge \text{dd}^c u_n \leq \prod_{j=1}^n \left(\int_{\Omega} (\text{dd}^c u_j)^n \right)^{1/n},$$

and

$$\int_{\Omega} (\text{dd}^c u_1)^j \wedge (\text{dd}^c u_2)^{n-j} \leq \left(\int_{\Omega} (\text{dd}^c u_1)^n \right)^{j/n} \cdot \left(\int_{\Omega} (\text{dd}^c u_2)^n \right)^{1-j/n}.$$

Proof. See [4, pp. 169–170]. ■

COROLLARY 3.78. Suppose $u, v \in \mathcal{F}(\Omega)$. Then

$$\left(\int_{\Omega} (\text{dd}^c(u+v))^n \right)^{1/n} \leq \left(\int_{\Omega} (\text{dd}^c u)^n \right)^{1/n} + \left(\int_{\Omega} (\text{dd}^c v)^n \right)^{1/n}.$$

Proof. Actually we need to show $u + v \in \mathcal{F}(\Omega)$, but directly we compute

$$\begin{aligned}
 \text{MA}(u + v)^{1/n} &= \left(\int_{\Omega} (\text{dd}^c u + \text{dd}^c v)^n \right)^{1/n} \\
 &= \left(\int_{\Omega} \sum_{j=1}^n (\text{dd}^c u)^j \wedge (\text{dd}^c v)^{n-j} \right)^{1/n} \\
 &\leq \left(\sum_{j=1}^n \int_{\Omega} \text{MA}(u)^{j/n} \cdot \text{MA}(v)^{1-j/n} \right)^{1/n} \\
 &= \left((\text{MA}(u)^{1/n} + \text{MA}(v)^{1/n})^n \right)^{1/n} = \text{MA}(u)^{1/n} + \text{MA}(v)^{1/n},
 \end{aligned}$$

where $\text{MA}(u) = \text{MA}_{\Omega}(u) = \int_{\Omega} (\text{dd}^c u)^n$. ■

COROLLARY 3.79. Suppose $u \in \mathcal{E}(\Omega)$, and $0 \in \Omega$. Then the (higher) Lelong numbers

$$e_j(u) = \int_{\{0\}} (\text{dd}^c u)^j \wedge (\text{dd}^c \log |z|)^{n-j},$$

satisfy

1. $e_j(u)^2 \leq e_{j-1}(u)e_{j+1}(u)$, for $1 \leq j \leq n-1$;
2. $e_j(u) \geq e_1(u)^j$, for $0 \leq j \leq n$;
3. $e_k(\varphi) \leq e_j(u)^{(l-k)/(l-j)} e_l(u)^{(k-j)/(l-j)}$, for $0 \leq j < k < l \leq n$.

THEOREM 3.80. Let $u, v \in L^{\infty} \cap \text{Psh}(\Omega)$, and $\liminf_{z \rightarrow \partial\Omega} (u(z) - v(z)) \geq 0$. Then

$$\int_{\{u < v\}} (\text{dd}^c v)^n \leq \int_{\{u < v\}} (\text{dd}^c u)^n.$$

Epecially, if $(\text{dd}^c u)^n \leq (\text{dd}^c v)^n$, then $v \leq u$ in Ω .

Proof. See [2, p. 10].

For $u, v \in C(\overline{\Omega})$ this case, take $v_{\varepsilon} = \max\{v, u + \varepsilon\}$. ■

3.7 Extremal plurisubharmonic function

We continue letting Ω bounded hyperconvex.

3.7.1 Maximal weight and relative type

Assume $0 \in \Omega$. Let $\varphi \in L_{\text{loc}}^{\infty}(\Omega \setminus \{0\}) \cap \text{Psh}^{-}(\Omega)$ which satisfies $\varphi(0) = -\infty$ and $(\text{dd}^c \varphi)^n = 0$ on $\Omega \setminus \{0\}$, i.e. φ is maximal on $\Omega \setminus \{0\}$. We call φ a *maximal weight* near 0.

DEFINITION 3.81 (Rashkovskii). For every psh germ u at 0, the relative type of u to φ is defined by

$$\sigma(u, \varphi) := \liminf_{z \rightarrow 0} \frac{u(z)}{\varphi(z)} = \sup\{c \geq 0: u \leq c\varphi + O(1) \text{ near } 0.\}$$

PROPOSITION 3.82. Assume $u \in \text{Psh}(\Omega)$, and $u(0) = -\infty$. Then

$$\Lambda(u, \varphi, r) := \sup_{z \in B(r)} u(z)$$

is convex and decreasing in $\log r$, where $B(r) = \{\varphi < r\}$.

Proof. See [16, p. 8]. ■

COROLLARY 3.83. We have

$$u \leq \sigma(u, \varphi)\varphi + O(1)$$

near 0.

Proof. See [16, p. 9]. ■

THEOREM 3.84. Let $\Phi: \Omega \times \Omega \rightarrow [-\infty, 0)$ be a negative psh function s.t.

1. $\{x: \Phi(x, \zeta) = -\infty\} = \{\zeta\}$;
2. for any $\zeta \in \Omega$ and $r < 0$, there exists a neighborhood U of ζ s.t.

$$\{(x, \zeta): \Phi(x, \zeta) < r, \zeta \in U\} \Subset \Omega \times \Omega;$$

3. $(\text{dd}^c \Phi_\zeta)^n = 0$ on $\Omega \setminus \{\zeta\}$ for every fixed ζ , where $\Phi_\zeta(x) := \Phi(x, \zeta)$;
4. $e^{\Phi(x, \zeta)}$ is locally Hölder continuous in ζ , i.e. $\exists \beta > 0$ s.t.

$$\left| e^{\Phi(x, \zeta_1)} - e^{\Phi(x, \zeta_2)} \right| \leq C |\zeta_1 - \zeta_2|^\beta.$$

Let $u \in \text{Psh}(\Omega)$, and denote

$$\mathcal{S}_c(u, \Phi; \Omega) := \{\zeta \in \Omega: \sigma(u, \Phi_\zeta) \geq c\}, \quad c > 0.$$

Then $\mathcal{S}_c(u, \Phi; \Omega)$ is an analytic subset of Ω for each $c > 0$.

Proof. See [16, pp. 22–23]. ■

COROLLARY 3.85. Let φ be a maximal weight near 0 which is Hölder continuous. Then for every $u \in \text{Psh}(\Omega)$ and $c > 0$,

$$E_c(u; \varphi) := \{z \in \Omega: \sigma_z(u, \tau_z \varphi) \geq c\}$$

is an analytic set.

Proof. Let $R > 0$ such that $\varphi \in \text{Psh}^-(B(0, 2R))$, and let $\Phi(x, \zeta) = \varphi(x - \zeta)$, which is a negative psh function on $B(0, R) \times B(0, R)$. ■

3.7.2 Pluricomplex Green function

DEFINITION 3.86. A locally bounded psh function u is maximal if $(dd^c u)^n = 0$.

In fact, it is the maximal psh function among those have same boundary values.

DEFINITION 3.87. For every $z_0 \in \Omega$, the function

$$G_\Omega(z, z_0) = \sup \{v(z) : v \in \text{Psh}^-(\Omega), v(z) \leq \log |z - z_0| + O(1) \text{ near } z_0\}$$

is the solution to the Dirichlet problem

$$\begin{cases} u \in \text{Psh}(\Omega) \cap C(\overline{\Omega}) \\ (dd^c u)^n = 0 \text{ on } \Omega \setminus \{z_0\} \\ u|_{\partial\Omega} = 0 \\ u(z) = \log |z - z_0| + O(1) \text{ near } z_0, \end{cases}$$

where $u \in C(\overline{\Omega})$ means $e^u \in C(\overline{\Omega})$. We call $G_\Omega(\cdot, z_0)$ the pluricomplex Green function on Ω with a logarithm pole at z_0 .

REMARK 3.88. • Actually we only need Ω bounded or hyperconvex to define the pluricomplex Green function, but we may have $G_\Omega \equiv -\infty$ when unbounded.

- $G_\Omega(z, w)$ is continuous on $\overline{\Omega} \times \Omega \setminus \{z = w\}$.
- If Ω is strongly pseudoconvex with $C^{2,1}$ boundary, then $G_\Omega(\cdot, z_0)$ is $C^{1,1}$ in $\overline{\Omega} \setminus \{z_0\}$.
- Even Ω is strongly pseudoconvex with C^∞ boundary, the pluricomplex Green function can be not C^2 up to boundary.
- When $n \geq 2$, in general, $G_\Omega(z, w)$ is not symmetric in z and w .

DEFINITION 3.89. A bounded domain $D \subset \mathbb{C}^n$ is said to be strictly hyperconvex if there exists a bounded domain Ω and a function $\varrho: \Omega \rightarrow (-\infty, 1)$ such that $\varrho \in C(\Omega) \cap \text{Psh}(\Omega)$, $D = \{z \in \Omega: \varrho(z) < 0\}$, ϱ is exhaustive for Ω and for any real number $c \in [0, 1]$, the open set $\{z \in \Omega: \varrho(z) < c\}$ is connected.

We have the following approximation theorem of pluricomplex Green function:

THEOREM 3.90. Let D be a bounded strictly hyperconvex domain and $0 \in D$. For any $m \in \mathbb{N}_+$, define

$$\mathcal{E}_m(D) := \{f \in \mathcal{O}(D) : \sup_{z \in D} |f(z)| \leq 1, \text{ord}_0(f) \geq m - 1\},$$

and

$$\phi_m(z) := \sup_{f \in \mathcal{E}_m(D)} \frac{1}{m} \log |f(z)|, \quad \forall z \in D.$$

Then ϕ_m converges to $G_\Omega(\cdot, 0)$ pointwise.

Proof. See [13].

Use Demailly's approximation theorem. ■

COROLLARY 3.91. If $0 \in D$ is a bounded strictly hyperconvex domain, then

$$G_\Omega(w, 0) = \sup \left\{ \frac{1}{\nu(\log |f|, 0)} \log |f(w)| : f \in \mathcal{O}(D), \sup_D |f| \leq 1, f(0) = 0, f \not\equiv 0 \right\}.$$

3.7.3 Capacity function

4 Some applications

4.1 Asymptotic behavior of Bergman kernel

Suppose Ω is an open set in \mathbb{C}^n , and $\phi: \Omega \rightarrow \mathbb{R}$ is a C^2 psh function. Consider the weighted Bergman space $\mathcal{H}_k := A^2(\Omega, e^{-k\phi})$ and Let B_k be the Bergman kernel on the diagonal for \mathcal{H}_k .

LEMMA 4.1. *We have that*

$$\limsup_{k \rightarrow \infty} \frac{1}{k^n} B_k(z) e^{-k\phi(z)} d\lambda(z) \leq \frac{1}{\pi^n n!} (i\partial\bar{\partial}\phi)^n(z).$$

Proof. Let $h \in \mathcal{H}_k$ with

$$\int_{\Omega} |h|^2 e^{-k\phi} \leq 1.$$

WLOG, say $z = 0$. We first assume ϕ has the Taylor expansion

$$\phi(\zeta) = \phi(0) + \sum \gamma_j |\zeta_j|^2 + o(|\zeta|^2),$$

where each $\gamma_j > 0$. Then for every $A < +\infty$,

$$1 \geq \int_{\sum \gamma_j |\zeta_j|^2 < A/k} |h|^2 e^{-k\phi} d\lambda \geq (1 - \varepsilon_k) e^{-k\phi(0)} \int_{\gamma_j |\zeta_j|^2 < A/k} |h|^2 e^{-k \sum \gamma_j |\zeta_j|^2} d\lambda,$$

where $\varepsilon_k \rightarrow 0$ as k tends to ∞ . By the mean value inequality,

$$\begin{aligned} \int_{\gamma_j |\zeta_j|^2 < A/k} |h|^2 e^{-k \sum \gamma_j |\zeta_j|^2} d\lambda &\geq |h(0)|^2 \int_{\gamma_j |\zeta_j|^2 < A/k} e^{-k \sum \gamma_j |\zeta_j|^2} d\lambda \\ &= |h(0)|^2 (\gamma_1 \cdots \gamma_n)^{-1} \int_{B(0, A/k)} e^{-k|\zeta|^2} d\lambda \\ &= |h(0)|^2 (\gamma_1 \cdots \gamma_n k^n)^{-1} \pi^n (1 - \delta_A), \end{aligned}$$

where $\delta_A \rightarrow 0$ as A tends to ∞ . Hence,

$$|h(0)|^2 e^{-k\phi(0)} / k^n \leq \pi^{-n} \gamma_1 \cdots \gamma_n (1 + 2\delta_A) (1 + 2\varepsilon_k)$$

for A and k large. Taking the supremum over h of norm at most 1 we then get

$$B_k(0) e^{-k\phi(0)} / k^n \leq \pi^{-n} \gamma_1 \cdots \gamma_n (1 + 2\delta_A) (1 + 2\varepsilon_k).$$

It follows that (e.g. taking $A = C\sqrt{k}$)

$$\limsup_{k \rightarrow \infty} \frac{1}{k^n} B_k(0) e^{-k\phi(0)} d\lambda \leq \frac{1}{\pi^n} \gamma_1 \cdots \gamma_n = \frac{1}{\pi^n n!} (i\partial\bar{\partial}\phi)^n(0).$$

It is easy to see that this also holds when some of γ_j 's are 0. A general ϕ has the Taylor expansion

$$\phi(\zeta) = \phi(0) + 2\operatorname{Re}(p(\zeta)) + q(\zeta, \bar{\zeta}) + o(|\zeta|^2).$$

We can reduce this case by change of variables and the substitution $h \mapsto h e^{-p}$. ■

THEOREM 4.2. *Additionally assume that D is bounded pseudoconvex. We have that*

$$\frac{1}{k^n} B_k(z) e^{-k\phi(z)} d\lambda(z) \rightarrow \frac{1}{\pi^n n!} (i\partial\bar{\partial}\phi)^n(z)$$

pointwise.

Proof. We only need to show the reverse estimate. We may assume $z = 0$, and near 0,

$$\phi(\zeta) = \phi(0) + |\zeta|^2 + o(|\zeta|^2).$$

Let $\chi \in C_c^\infty$ be compactly supported in $B(0, 1)$ and $\chi \equiv 1$ in $B(0, 1/2)$ and δ_k be a sequence of positive numbers s.t. $\delta_k \rightarrow 0$ and $\tau_k := \sqrt{k}\delta_k \rightarrow \infty$. Set

$$g_k(\zeta) = k^{n/2} e^{k\phi(0)/2} \chi(\tau_k \zeta).$$

Then

$$\begin{aligned} \int_{\Omega} |g_k|^2 e^{-k\phi} d\lambda &\leq \int_{\Omega} k^n e^{k\phi(0)} |\chi(\tau_k \zeta)|^2 e^{-k\phi} d\lambda \\ &\leq k^n e^{k\phi(0)} \int_{|\zeta|^2 < \tau_k^{-2}} e^{-k\phi} d\lambda \\ &\leq e^{k\phi(0)} \delta_k^{-2n} \int_{B(0,1)} e^{-k\phi(\zeta/\tau_k)} d\lambda \\ &\leq (1 + \epsilon_k) e^{k\phi(0)} \delta_k^{-2n} e^{-k\phi(0)} \int_{B(0,1)} e^{-|\zeta/\delta_k|^2} d\lambda \\ &\leq (1 + \epsilon_k) \int_{B(0, \delta_k^{-1})} e^{-|\zeta|^2} d\lambda \rightarrow \pi^n \quad (k \rightarrow \infty). \end{aligned}$$

We can also see

$$|\bar{\partial} g_k(z)|^2 \leq C k^n e^{k\phi(0)} \tau_k^2,$$

and $\bar{\partial} g_k = 0$ except when $\tau_k^{-1}/2 < |\zeta| < \tau_k^{-1}$. So we can use Hörmander's L^2 estimate to solve $\bar{\partial} u_k = \bar{\partial} g_k$ with

$$\begin{aligned} \int_{\Omega} |u_k(z)|^2 e^{-k\phi} d\lambda &\leq C \frac{1}{k} \int_{\tau_k^{-1}/2 < |\zeta| < \tau_k^{-1}} k^n e^{k\phi(0)} \tau_k^2 e^{-k\phi} d\lambda \\ &\leq C k^n \delta_k^2 \int_{\tau_k^{-1}/2 < |\zeta| < \tau_k^{-1}} e^{-k|\zeta|^2} d\lambda \\ &\leq C \delta_k^2 \int_{\delta_k^{-1}/2 < |\zeta| < \delta_k^{-1}} e^{-|\zeta|^2} d\lambda \rightarrow 0 \quad (k \rightarrow \infty). \end{aligned}$$

Set $F_k = g_k - u_k$, which is holomorphic. Then $F_k(0) = g_k(0) = k^{n/2} e^{k\phi(0)/2}$, and

$$\int_{\Omega} |F_k|^2 e^{-k\phi} d\lambda \leq \pi^n + o(1).$$

Thus,

$$\liminf_{k \rightarrow \infty} \frac{1}{k^n} B_k(z) e^{-k\phi(0)} \geq \liminf_{k \rightarrow \infty} \frac{1}{k^n} \cdot \frac{k^n e^{k\phi(0)}}{\pi^n} e^{-k\phi(0)} = \frac{1}{\pi^n}.$$

The rest is similar as above. ■

4.2 Spectrum of Toeplitz operator

Let Ω be a bounded pseudoconvex subset in \mathbb{C}^n and φ a C^2 psh function on Ω . Let χ be a compactly supported bounded function on Ω . Define the Toeplitz concentration operator $T_{\chi,\varphi}$ by

$$T_{\chi,\varphi}f(z) = \int_{\Omega} \chi(\zeta)f(\zeta)B_{\varphi}(z,\bar{\zeta})e^{-\varphi(\zeta)},$$

where $B_{\varphi}(z,\bar{\zeta})$ is the off-diagonal Bergman kernel of $A^2(\Omega, e^{-\varphi})$. It can be seen that $T_{\chi,\varphi}$ is of trace class, where

$$\text{Tr}(T_{\chi,\varphi}) = \int_D \chi(z)B_{\varphi}(z,\bar{z})e^{-\varphi(z)}$$

and

$$\text{Tr}(T_{\chi,\varphi}^2) = \int_{\Omega} \int_{\Omega} |\chi(\zeta)B_{\varphi}(z,\bar{\zeta})|^2 e^{-\varphi(z)}e^{-\varphi(\zeta)} = \int_{\Omega} |\chi(z)|^2 B_{\varphi}(z,\bar{z})e^{-\varphi(z)}.$$

If f is an eigenfunction of $T_{\chi,\varphi}$ with eigenvalue λ , then

$$\lambda \|f\|_{\varphi}^2 = (T_{\chi,\varphi}f, f) = (\chi f, P_{\varphi}f) = (\chi f, f) = \int_{\Omega} \chi(z)|f(z)|^2 e^{-\varphi(z)}, \quad (4.6)$$

where P_{φ} is the Bergman projection.

Now let χ be a characteristic of a relatively compact open subset K , and denote the eigenvalues of $T_{\chi,\varphi}$ be $\lambda_m(D, \varphi) = \lambda_m(\chi, \varphi)$, ordered in a non-increasing sequence (counted with multiplicity)

$$\lambda_0(D, \varphi) \geq \lambda_1(D, \varphi) \geq \lambda_2(D, \varphi) \geq \cdots$$

It is well-known [Weyl–Courant lemma]

$$\begin{aligned} \lambda_m(\chi, \varphi) &= \min_{\dim E = m-1} \max_{\substack{f \perp E \\ \|f\|_{\varphi}=1}} (T_{\chi,\varphi}f, f) \\ &= \max_{\dim F = m} \min_{\substack{f \in F \\ \|f\|_{\varphi}=1}} (T_{\chi,\varphi}f, f), \end{aligned}$$

where E, F are subspaces of $A^2(\Omega, e^{-\varphi})$.

THEOREM 4.3. *Let D be a relatively compact smooth open subset of Ω . Then for any $\gamma \in (0, 1)$ we have*

$$\lim_{k \rightarrow \infty} k^{-n} \#\{m \in \mathbb{Z}_{\geq 0} : \lambda_m(D, k\varphi) > \gamma\} = \frac{1}{\pi^n n!} \int_D (i\partial\bar{\partial}\varphi)^n.$$

Proof. According to Theorem 4.2, we have

$$\frac{1}{k^n} \chi(z) B_{k\varphi}(z) e^{-k\varphi(z)} d\lambda(z) \rightarrow \frac{1}{\pi^n n!} \chi(z) (i\partial\bar{\partial}\varphi)^n(z)$$

pointwise, which implies $(T_k = T_{\chi,k\varphi})$

$$\lim_{k \rightarrow \infty} k^{-n} \text{Tr}(T_k) = \lim_{k \rightarrow \infty} k^{-n} \text{Tr}(T_k^2) = \frac{1}{\pi^n n!} \int_D (i\partial\bar{\partial}\varphi)^n. \quad (4.7)$$

By definition, we have $(\lambda_m(k) = \lambda_m(T_{\chi, k\varphi}))$

$$\text{Tr}(T_k) = \sum_m \lambda_m(k), \quad \text{Tr}(T_k) = \sum_m \lambda_m(k)^2.$$

It follows from (4.6) that $\lambda_m(k) \leq 1$. Note that

$$\lambda_1(k) = \sup \frac{(T_k f, f)}{\|f\|_{k\varphi}^2} \geq \frac{(T_k 1, 1)}{\|1\|_{k\varphi}^2} > 0.$$

Thus, we have

$$0 < \sum_m \lambda_m(k)^2 \leq \sum_m \lambda_m(k).$$

By (4.7), for any $\delta > 0$, there exists k_δ s.t. when $k \geq k_\delta$,

$$1 - \delta \leq \frac{\sum_m \lambda_m(k)}{\sum_m \lambda_m(k)} \leq 1.$$

Now fix k_δ and for every $\beta \in (0, 1)$ denote

$$S_\beta = \frac{\sum_{\lambda_m \leq \beta} \lambda_m(k)}{\sum_m \lambda_m(k)}.$$

Then

$$\begin{aligned} (1 - \delta) \sum_m \lambda_m(k) &\leq \sum_{\lambda_m > \gamma} \lambda_m(k)^2 + \sum_{\lambda_m \leq \gamma} \lambda_m(k)^2 \\ &\leq \sum_{\lambda_m > \gamma} \lambda_m(k) + \gamma \cdot \sum_{\lambda_m \leq \gamma} \lambda_m(k) \\ &\leq (1 - S_\gamma) \sum_m \lambda_m(k) + \gamma S_\gamma \sum_m \lambda_m(k), \end{aligned}$$

which shows

$$S_\gamma \leq \frac{\delta}{1 - \gamma}.$$

Hence,

$$\#\{m: \lambda_m(k) > \gamma\} \geq \sum_{\lambda_m > \gamma} \lambda_m(k) \geq \left(1 - \frac{\delta}{1 - \gamma}\right) \sum_m \lambda_m(k). \quad (4.8)$$

Note that for $\gamma < \gamma' < 1$ we have

$$\begin{aligned} \#\{m: \lambda_m(k) > \gamma\} &= \#\{m: \lambda_m(k) > \gamma'\} + \#\{m: \gamma' \geq \lambda_m(k) > \gamma\} \\ &\leq \frac{1}{\gamma'} \sum_{\lambda_m > \gamma'} \lambda_m(k) + \frac{1}{\gamma} \sum_{\gamma' \geq \lambda_m > \gamma} \lambda_m(k) \\ &\leq \frac{1}{\gamma'} \sum_m \lambda_m(k) + \frac{1}{\gamma} S_{\gamma'} \sum_m \lambda_m(k). \end{aligned} \quad (4.9)$$

Now combining (4.7), (4.8) and (4.9), we get

$$\liminf_{k \rightarrow \infty} \#\{m: \lambda_m(k) > \gamma\} \geq \left(1 - \frac{\delta}{1 - \gamma}\right) \frac{1}{\pi^n n!} \int_D (\mathrm{i}\partial\bar{\partial}\varphi)^n,$$

and

$$\limsup_{k \rightarrow \infty} \#\{m: \lambda_m(k) > \gamma\} \leq \left(\frac{1}{\gamma'} + \frac{\delta}{\gamma(1 - \gamma')}\right) \frac{1}{\pi^n n!} \int_D (\mathrm{i}\partial\bar{\partial}\varphi)^n.$$

The result follows since $\delta > 0$ and $\gamma' \in (\gamma, 1)$ are arbitrary. \blacksquare

4.3 Riemann–Roch; Holomorphic Morse Inequality

LEMMA 4.4. *Let \mathcal{H} be a subspace of $L^2(X, \mu)$ consisting of continuous functions, and denote by $K(x, y)$ the reproducing kernel of \mathcal{H} . Then*

$$\dim \mathcal{H} = \int_X K(x, x) \, \mathrm{d}\mu.$$

Proof. Let (h_j) be an orthonormal basis for \mathcal{H} . Then

$$\dim \mathcal{H} = \int_X \sum |h_j|^2 \, \mathrm{d}\mu = \int_X K(x, x) \, \mathrm{d}\mu. \quad \blacksquare$$

PROPOSITION 4.5 (Bouche, Tian, ...). *Let $L \rightarrow X$ be a hermitian holomorphic line bundle over a compact complex manifold. Assume $h = e^{-\phi}$ is a smooth metric on L with $\mathrm{i}\partial\bar{\partial}\phi \geq 0$ and positive somewhere. Let X_0 be the open subset where $\mathrm{i}\partial\bar{\partial}\phi$ is positive. Then*

$$\lim_{k \rightarrow \infty} \frac{1}{k^n} h^0(X, kL) = \frac{1}{\pi^n n!} \int_{X_0} (\mathrm{i}\partial\bar{\partial}\phi)^n.$$

Proof. Since h is smooth, by the above lemma,

$$h^0(X, kL) = \int_X B_k e^{-k\phi}.$$

The proposition then follows from Theorem 4.2. \blacksquare

Now let X be a compact complex manifold, E a holomorphic vector bundle of rank r and L a line bundle over X . If L is equipped with a smooth metric h of curvature form $\Theta_{L,h}$, we define the q -index set of L to be the open subset $X(L, h, q)$ where $\mathrm{i}\Theta_{L,h}$ has exactly q negative eigenvalues and $(n - q)$ positive eigenvalues for $0 \leq q \leq n$. Denote

$$\Delta = \{x \in X: \det(\Theta_{L,h}(x)) = 0\}.$$

Then $X = \Delta \cup \bigcup_q X(L, h, q)$. Set

$$X(L, h, \leq q) = \bigcup_{0 \leq j \leq q} X(L, h, j).$$

THEOREM 4.6 (Demailly, 85'; Holomorphic Morse Inequalities). *The cohomology groups $H^q(X, E \otimes \mathcal{O}(kL))$ satisfy the following asymptotic inequalities:*

1. *Weak Morse inequalities*

$$h^q(X, \mathcal{O}(E) \otimes \mathcal{O}(kL)) \leq r \frac{k^n}{n!} \int_{X(L, h, q)} (-1)^q \left(\frac{i}{2\pi} \Theta_{L, h} \right)^n + o(k^n).$$

2. *Strong Morse inequalities*

$$\sum_{0 \leq j \leq q} (-1)^{q-j} h^j(X, \mathcal{O}(E) \otimes \mathcal{O}(kL)) \leq r \frac{k^n}{n!} \int_{X(L, h, \leq q)} (-1)^q \left(\frac{i}{2\pi} \Theta_{L, h} \right)^n + o(k^n).$$

5 Some open problems

5.1 Zero mass conjecture

5.2 Sublevel sets of pluricomplex Green function

References

- [1] Bo Berndtsson and László Lempert. “A Proof of the Ohsawa–Takegoshi Theorem with Sharp Estimates”. In: *Journal of the Mathematical Society of Japan* 68.4 (Oct. 2016), pp. 1461–1472. DOI: [10.2969/jmsj/06841461](https://doi.org/10.2969/jmsj/06841461).
- [2] Zbigniew Blocki. “Minicourse on Pluripotential Theory”. 2012. URL: <https://gamma.im.uj.edu.pl/~blocki/publ/ln/ln-vienna.pdf>.
- [3] Zbigniew Blocki. “The Complex Monge-Ampère Operator in Pluripotential Theory”. 1997. URL: <https://gamma.im.uj.edu.pl/~blocki/publ/ln/wykl.pdf>.
- [4] Urban Cegrell. “The General Definition of the Complex Monge-Ampère Operator”. In: *Annales de l’institut Fourier* 54.1 (2004), pp. 159–179. ISSN: 0373-0956. DOI: [10.5802/aif.2014](https://doi.org/10.5802/aif.2014). (Visited on 12/26/2024).
- [5] Jean-Pierre Demailly. *Analytic Methods in Algebraic Geometry*. 2nd ed. Surveys of Modern Mathematics 1. Somerville: International press, 2012. ISBN: 978-1-57146-234-3. URL: https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/analmeth_book.pdf.
- [6] Jean-Pierre Demailly. *Complex Analytic and Differential Geometry*. URL: <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [7] Qi’an Guan. “A Sharp Effectiveness Result of Demailly’s Strong Openness Conjecture”. In: *Advances in Mathematics* 348 (2019/05/25/), pp. 51–80. DOI: [10.1016/j.aim.2019.03.017](https://doi.org/10.1016/j.aim.2019.03.017).
- [8] Qi’an Guan and Zhitong Mi. “Concavity of Minimal L^2 Integrals Related to Multiplier Ideal Sheaves on Weakly Pseudoconvex Kähler Manifolds”. In: *Science China Mathematics* 65.5 (May 2022), pp. 887–932. ISSN: 1674-7283, 1869-1862. DOI: [10.1007/s11425-021-1930-2](https://doi.org/10.1007/s11425-021-1930-2).
- [9] Qi’an Guan and Xiangyu Zhou. “A Proof of Demailly’s Strong Openness Conjecture”. In: *Annals of Mathematics* 182.2 (2015), pp. 605–616. ISSN: 0003-486X, 1939-8980. DOI: [10.4007/annals.2015.182.2.5](https://doi.org/10.4007/annals.2015.182.2.5).
- [10] Qi’an Guan and Xiangyu Zhou. “A Solution of an L^2 Extension Problem with an Optimal Estimate and Applications”. In: *Annals of Mathematics* 181.3 (2015), pp. 1139–1208. ISSN: 0003-486X, 1939-8980. DOI: [10.4007/annals.2015.181.3.6](https://doi.org/10.4007/annals.2015.181.3.6).
- [11] Qi’an Guan and Xiangyu Zhou. “Effectiveness of Demailly’s Strong Openness Conjecture and Related Problems”. In: *Inventiones mathematicae* 202.2 (Nov. 2015), pp. 635–676. ISSN: 0020-9910, 1432-1297. DOI: [10.1007/s00222-014-0575-3](https://doi.org/10.1007/s00222-014-0575-3).
- [12] Jeffery D. McNeal and Mircea Mustață. *Analytic and Algebraic Geometry: Common Problems, Different Methods*. IAS/Park City Mathematics Series Volume 17. Providence (R.I.): American mathematical society, 2010. ISBN: 978-0-8218-4908-8.
- [13] Stéphanie Nivoche. “The Pluricomplex Green Function, Capacitative Notions, and Approximation Problems in \mathbb{C}^n ”. In: *Indiana University Mathematics Journal* 44.2 (1995), pp. 489–510. ISSN: 0022-2518, 1943-5258. DOI: [10.1512/iumj.1995.44.1998](https://doi.org/10.1512/iumj.1995.44.1998).
- [14] Takeo Ohsawa. *Analysis of Several Complex Variables*. Trans. by Shu Gilbert Nakamura. Translations of Mathematical Monographs Volume 211. Providence, Rhode Island: American Mathematical Society, 2002. ISBN: 978-0-8218-2098-8 978-1-4704-4636-9.
- [15] Takeo Ohsawa. *L^2 Approaches in Several Complex Variables: Towards the Oka–Cartan Theory with Precise Bounds*. Springer Monographs in Mathematics. Tokyo: Springer Japan, 2018. ISBN: 978-4-431-56851-3 978-4-431-56852-0. DOI: [10.1007/978-4-431-56852-0](https://doi.org/10.1007/978-4-431-56852-0).

- [16] Alexander Rashkovskii. “Relative Types and Extremal Problems for Plurisubharmonic Functions”. In: *International Mathematics Research Notices* (Jan. 2006), p. 76283. ISSN: 1073-7928, 1687-0247. DOI: [10.1155/IMRN/2006/76283](https://doi.org/10.1155/IMRN/2006/76283).
- [17] Walter Rudin. *Functional Analysis*. 2. ed., [Nachdr.] International Series in Pure and Applied Mathematics. Boston, Mass.: McGraw-Hill, 2007. ISBN: 978-0-07-054236-5.