

L^2 Extension and Effectiveness of L^p Strong Openness Property

Shi Jie BAO Qi An GUAN¹⁾

School of Mathematical Sciences, Peking University, Beijing 100871, P. R. China

E-mail: bsjie@pku.edu.cn guanqian@math.pku.edu.cn

Abstract In this note, we present an L^2 extension approach to the effectiveness result of L^p strong openness property of multiplier ideal sheaves.

Keywords Optimal L^2 extension, multiplier ideal sheaf, L^p strong openness property

MR(2010) Subject Classification 32D15, 32E10, 32L10, 32U05, 32W0

1 Introduction

The multiplier ideal sheaf associated to a plurisubharmonic function was widely discussed in several complex variables, complex geometry and algebraic geometry (see e.g. [10, 20, 25, 26]). We recall the definition of multiplier ideal sheaf as follows. Let φ be a plurisubharmonic function (see [7]) on a complex manifold X . The multiplier ideal sheaf $\mathcal{I}(\varphi)$ is the sheaf on X whose germs are the holomorphic functions F such that $|F|^2 e^{-\varphi}$ is locally integrable.

The following strong openness property was conjectured by Demailly [8, 9] (the well-known strong openness conjecture), and established by Guan–Zhou [15].

Strong openness property:

$$\mathcal{I}_+(\varphi) = \mathcal{I}(\varphi),$$

where $\mathcal{I}_+(\varphi) := \bigcup_{p>1} \mathcal{I}(p\varphi)$.

When $\mathcal{I}(\varphi) = \mathcal{O}$, the strong openness conjecture is called the openness conjecture, which was posed by Demailly–Kollár [10] and proved by Berndtsson [3].

The effectiveness of openness conjecture was established by Berndtsson [3], which implies the openness conjecture. In the proof of the strong openness conjecture, Ohsawa–Takegoshi L^2 extension was used by Guan–Zhou [15]. After that, Guan–Zhou [17] established an effectiveness result of the strong openness property by solving the $\bar{\partial}$ equations with L^2 estimates. In [1], authors of the present article gave an L^2 extension approach to an effectiveness result of the strong openness property.

In [11], Fornæss established the following L^p strong openness property by using the strong openness property [15]:

Let F be a holomorphic function on a domain $D \subset \mathbb{C}^n$ containing the origin o , φ a plurisubharmonic function on D and $p \in (0, +\infty)$. If $|F|^p e^{-\varphi}$ is L^1 on a neighborhood of o , then there exists $q > 1$ such that $|F|^p e^{-q\varphi}$ is L^1 on a neighborhood of o .

Received August 3, 2021, accepted December 31, 2021

The second author was supported by NSFC (Grant Nos. 11825101, 11522101 and 11431013)

1) Corresponding author

In [14], Guan–Yuan gave an effectiveness result of the L^p strong openness property by using the general concavity of minimal L^2 integrals related to multiplier ideal sheaves in [13]. Following [1], it is natural to ask

Question 1.1 Can one obtain an L^2 extension approach to the effectiveness result of the L^p strong openness property for any $p > 0$?

In the present note, we give an affirmative answer to the above question.

Let D be a bounded psuedoconvex domain in \mathbb{C}^n with $o \in D$. Let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$. Let F be a holomorphic function on D . For any $p > 0$, denote that

$$c_{o,p}^F(\varphi) := \sup\{c \geq 0 : |F|^p e^{-c\varphi} \text{ is locally integrable near } o\}.$$

Let I be a proper ideal of \mathcal{O}_o , and ψ be a plurisubharmonic function on D . Denote that

$$B^\psi(F, I, D) := \sup_{\xi \in \ell_I} \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)}$$

if $I \not\supseteq \mathcal{I}(\psi)_o$. Here $\xi \in \ell_1 := \{\xi = (\xi_\alpha)_{\alpha \in \mathbb{N}^n} : \sum_{\alpha \in \mathbb{N}^n} |\xi_\alpha| \rho^{|\alpha|} < +\infty, \text{ for any } \rho > 0\}$, and $(\xi \cdot F)(z) := \sum_{\alpha \in \mathbb{N}^n} \xi_\alpha F^{(\alpha)}(z)/\alpha!$. $\xi \in \ell_I$ means that $\xi \neq (0, \dots, 0, \dots)$ and $(\xi \cdot f)(o) = 0$ for any $(f, o) \in I$. The weighted Bergman kernel

$$K_{\xi,D}^\psi(z) := \sup_{f \in A^2(D, e^{-\psi})} \frac{|(\xi \cdot f)(z)|^2}{\int_D |f|^2 e^{-\psi}},$$

where $A^2(D, e^{-\psi}) = \{f \in \mathcal{O}(D) : |f|^2 e^{-\psi} \text{ is integrable on } D\}$.

Denote that $F_1 = F^{\lceil \frac{p}{2} \rceil}$, $\varphi_1 = (2\lceil \frac{p}{2} \rceil - p) \log |F|$, and here $\lceil m \rceil := \min\{n \in \mathbb{Z} : n \geq m\}$. We will prove the following effectiveness result of the L^p strong openness theorem.

Theorem 1.2 Let C_1 and C_2 be two positive constants. If there exists $a \geq 0$, such that

- (1) $\int_D |F|^p e^{-(1+a)\varphi} \leq C_1$;
- (2) $B^{\varphi_1+a\varphi}(F_1, \mathcal{I}_+(\varphi_1 + c_{o,p}^F(\varphi)\varphi)_o, D) \geq C_2$.

Then for any $q > a + 1$ satisfying

$$\tilde{\theta}_a(q) > \frac{C_1}{C_2},$$

$|F|^p e^{-q\varphi}$ is locally L^1 integrable near o , where $\tilde{\theta}_a(q) = \frac{q-a}{q-a-1}$.

Remark 1.3 Let $D = \Delta$, $F = z$ and $\varphi = \frac{p+2}{q} \log |z|$, then $c_{o,p}^F(\varphi) = q$. We can find that $\int_D |F|^p e^{-(1+a)\varphi} = \frac{2q\pi}{(q-a-1)(p+2)}$, and $B^{\varphi_1+a\varphi}(F_1, \mathcal{I}_+(\varphi_1 + c_{o,p}^F(\varphi)\varphi)_o, D) = \frac{2q\pi}{(q-a)(p+2)}$ by straightforward calculations, then $\frac{\int_D |F|^p e^{-(1+a)\varphi}}{B^{\varphi_1+a\varphi}(F_1, \mathcal{I}_+(\varphi_1 + c_{o,p}^F(\varphi)\varphi)_o, D)} = \frac{q-a}{q-a-1}$. It means that Theorem 1.2 is a sharp effectiveness result of the L^p strong openness theorem.

When $p = 2$, the above Theorem 1.2 induces the following corollary.

Corollary 1.4 Let D be a bounded psuedoconvex domain in \mathbb{C}^n with $o \in D$. Let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$. Let F be a holomorphic function on D . Let C_1 and C_2 be two positive constants. If there exists $a \geq 0$, such that

- (1) $\int_D |F|^2 e^{-(1+a)\varphi} \leq C_1$;

$$(2) B^{a\varphi}(F, \mathcal{I}_+(c_o^F(\varphi)\varphi)_o, D) \geq C_2.$$

Then for any $q > a + 1$ satisfying

$$\tilde{\theta}_a(q) > \frac{C_1}{C_2},$$

$|F|^2 e^{-q\varphi}$ is locally L^1 integrable near o , where $\tilde{\theta}_a(q) = \frac{q-a}{q-a-1}$.

Here $c_o^F(\varphi)$ is the jumping number

$$c_o^F(\varphi) := \sup\{c \geq 0 : |F|^2 e^{-c\varphi} \text{ is locally integrable near } o\}.$$

In particular, let $a = 0$. Then Corollary 1.4 induces Theorem 1.6 in [1].

In addition, when $F \equiv 1$, note that

$$B^{a\varphi}(1, \mathcal{I}_+(c_o^F(\varphi)\varphi)_o, D) \geq K_{D,a\varphi}^{-1}(o).$$

Here $K_{D,a\varphi}$ is the Bergman kernel on D with respect to the weight $e^{-a\varphi}$. Then Corollary 1.4 induces the following effectiveness result of the openness property.

Corollary 1.5 *Let D be a bounded psuedoconvex domain in \mathbb{C}^n with $o \in D$. Let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$. Let C_1 and C_2 be two positive constants. If there exists $a \geq 0$, such that*

$$(1) \int_D e^{-(1+a)\varphi} \leq C_1;$$

$$(2) (K_{D,a\varphi})^{-1}(o) \geq C_2.$$

Then for any $q > a + 1$ satisfying

$$\tilde{\theta}_a(q) > \frac{C_1}{C_2},$$

$e^{-q\varphi}$ is locally L^1 integrable near o , where $\tilde{\theta}_a(q) = \frac{q-a}{q-a-1}$.

2 Definition and Basic Properties of the Bergman Kernel

Firstly, we recall a linear space of sequences of complex numbers (see [1]),

$$\ell_1 := \left\{ \xi = (\xi_\alpha)_{\alpha \in \mathbb{N}^n} : \sum_{\alpha \in \mathbb{N}^n} |\xi_\alpha| \rho^{|\alpha|} < +\infty, \text{ for any } \rho > 0 \right\}.$$

Any element in ℓ_1 can be a linear functional over \mathcal{O}_{z_0} for any $z_0 \in \mathbb{C}^n$ as follows.

For any $F(z) \in \mathcal{O}_{z_0}$, we can write that $F(z) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha (z - z_0)^\alpha$ near z_0 . Then we define the value that ξ acts on $F(z)$ as

$$(\xi \cdot F)(z_0) := \sum_{\alpha \in \mathbb{N}^n} \xi_\alpha \frac{F^{(\alpha)}(z_0)}{\alpha!} = \sum_{\alpha \in \mathbb{N}^n} \xi_\alpha a_\alpha,$$

for any $\xi \in \ell_1$.

We have showed that $(\xi \cdot F)(z_0)$ is well defined in [1].

Let ψ be a plurisubharmonic function on D . For any $z \in D$, $\xi \in \ell_1$, we denote the weighted Bergman kernel by

$$K_{\xi,D}^\psi(z) := \sup_{F \in A^2(D, e^{-\psi})} \frac{|(\xi \cdot F)(z)|^2}{\int_D |F|^2 e^{-\psi}},$$

where D is a bounded domain in \mathbb{C}^n , and

$$A^2(D, e^{-\psi}) = \{f \in \mathcal{O}(D) : |f|^2 e^{-\psi} \text{ is integrable on } D\}.$$

If $A^2(D, e^{-\psi}) = \{0\}$, we denote that $K_{\xi, D}^\psi(z) = 0$.

Secondly, we list some lemmas. Lemmas 2.1, 2.2 and 2.3 were proved in [1].

Lemma 2.1 ([1]) *Let D be a bounded domain in \mathbb{C}^n , and let F be a holomorphic function on D . Then $(\xi \cdot F)(z)$ is also a holomorphic function on D , where $\xi \in \ell_1$.*

Lemma 2.2 ([1]) *Let D be a bounded domain in \mathbb{C}^n , and let $\{F_j\}$ be a sequence of holomorphic functions on D uniformly converging to F on every compact subset of D . Then for any $z \in D$, and $\xi \in \ell_1$, $\{(\xi \cdot F_j)(z)\}$ converges to $(\xi \cdot F)(z)$ uniformly on every compact subset of D .*

Lemma 2.3 ([1]) *Let D be a bounded domain in \mathbb{C}^n , and let $\{z_j\}$ be a sequence of points in D such that $\lim_{j \rightarrow +\infty} z_j = z \in D$. Let $\{F_j\}$ be a sequence of holomorphic functions on D uniformly converging to F on every compact subset of D . Then for any $\xi \in \ell_1$, $\lim_{j \rightarrow +\infty} (\xi \cdot F_j)(z_j) = (\xi \cdot F)(z)$.*

The following lemma shows that the weighted Bergman kernel is finite.

Lemma 2.4 *Let D be a bounded domain in \mathbb{C}^n , and let ψ be a plurisubharmonic function on D . Then $K_{\xi, D}^\psi(z_0) < +\infty$ for any $z_0 \in D$, $\xi \in \ell_1$.*

In fact, we can prove the following stronger result.

Lemma 2.5 *Let D be a bounded domain in \mathbb{C}^n , ψ be a plurisubharmonic function on D , and let $\xi \in \ell_1$. Then for any compact subset $K \subseteq D$, there is a finite constant $C > 0$ such that*

$$|(\xi \cdot F)(z)|^2 \leq C \int_D |F|^2 e^{-\psi},$$

for any $F \in A^2(D, e^{-\psi})$, and any $z \in K$.

Proof It is trivial when $\xi = 0$ or $A^2(D, e^{-\psi}) = \{0\}$. Now we assume $\xi \neq 0$ and $A^2(D, e^{-\psi}) \neq \{0\}$. For the compact subset K , we are able to find some $R > 0$ such that $\bigcup_{w \in K} \Delta_{w, R}^n \subset\subset D$. Then there exists a finite constant a such that $\psi(z) < a$ for any $z \in \bigcup_{w \in K} \Delta_{w, R}^n$. And for any nonzero holomorphic function $F(z)$ on D , if we write that $F(z) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha (z - z_0)^\alpha$ in $\Delta_{z_0, R}^n$ for any $z_0 \in K$, then

$$\begin{aligned} \int_D |F|^2 e^{-\psi} &\geq \int_{\Delta_{z_0, R}^n} |F|^2 e^{-\psi} \\ &\geq \int_{\Delta_{z_0, R}^n} |F|^2 e^{-a} \\ &= e^{-a} \sum_{\alpha \in \mathbb{N}^n} \frac{\pi^n |a_\alpha|^2}{(\alpha_1 + 1) \cdots (\alpha_n + 1)} R^{2(|\alpha| + n)}. \end{aligned}$$

By Cauchy–Schwarz’s inequality,

$$|(\xi \cdot F)(z_0)|^2 \leq \left(\sum_{\alpha \in \mathbb{N}^n} \frac{(\alpha_1 + 1) \cdots (\alpha_n + 1)}{\pi^n R^{2(|\alpha| + n)}} |\xi_\alpha|^2 \right) \left(\sum_{\alpha \in \mathbb{N}^n} \frac{\pi^n |a_\alpha|^2}{(\alpha_1 + 1) \cdots (\alpha_n + 1)} R^{2(|\alpha| + n)} \right),$$

and it implies that

$$\frac{|(\xi \cdot F)(z_0)|^2}{\int_D |F|^2 e^{-\psi}} \leq e^a \sum_{\alpha \in \mathbb{N}^n} \frac{(\alpha_1 + 1) \cdots (\alpha_n + 1)}{\pi^n R^{2(|\alpha| + n)}} |\xi_\alpha|^2.$$

Since $\xi \in \ell_1$, we can choose some ρ with $\rho > 1/R$ such that

$$|\xi_\alpha| \rho^{|\alpha|} < M, \quad \forall \alpha \in \mathbb{N}^n,$$

for some positive constant M . Hence

$$\begin{aligned} \frac{|(\xi \cdot F)(z_0)|^2}{\int_D |F|^2 e^{-\psi}} &\leq e^a \sum_{\alpha \in \mathbb{N}^n} \frac{(\alpha_1 + 1) \cdots (\alpha_n + 1)}{\pi^n R^{2(|\alpha| + n)}} |\xi_\alpha|^2 \\ &\leq e^a \sum_{\alpha \in \mathbb{N}^n} \frac{(\alpha_1 + 1) \cdots (\alpha_n + 1)}{\pi^n R^{2(|\alpha| + n)}} \cdot \frac{M^2}{\rho^{2|\alpha|}} \\ &= \frac{e^a M^2}{\pi^n R^{2n}} \sum_{\alpha \in \mathbb{N}^n} (\alpha_1 + 1) \cdots (\alpha_n + 1) \frac{1}{(R\rho)^{2|\alpha|}} < +\infty. \end{aligned}$$

Now we choose

$$C = \frac{e^a M^2}{\pi^n R^{2n}} \sum_{\alpha \in \mathbb{N}^n} (\alpha_1 + 1) \cdots (\alpha_n + 1) \frac{1}{(R\rho)^{2|\alpha|}},$$

which is independent of the choice of $z_0 \in K$, and get the result. \square

Now let $K = \{z_0\}$, then Lemma 2.4 can be induced by Lemma 2.5.

We prove the following lemma using Lemma 2.3.

Lemma 2.6 *Let D be a bounded domain in \mathbb{C}^n , ψ be a plurisubharmonic function on D with $A^2(D, e^{-\psi}) \neq \{0\}$, and let $z \in D$. Then for any $\xi \in \ell_1$, there exists a holomorphic function F_0 on D such that*

$$K_{\xi, D}^\psi(z) = \frac{|(\xi \cdot F_0)(z)|^2}{\int_D |F_0|^2 e^{-\psi}}.$$

Proof It is trivial when $\xi = 0$. Now we assume $\xi \neq 0$.

By the definition of Bergman kernel, there exists a sequence of holomorphic functions $\{F_j\}$ on D such that $\int_D |F_j|^2 e^{-\psi} = 1$, and $\lim_{j \rightarrow +\infty} |(\xi \cdot F_j)(z)|^2 = K_{\xi, D}^\psi(z)$. Then by Montel's theorem (see [22, Theorem 1.5]), there is a subsequence of $\{F_j\}$ which is uniformly convergent on every compact subset of D . Denote the limit of the subsequence by F_0 . Then Fatou's lemma and Lemma 2.2 imply that $\int_D |F_0|^2 e^{-\psi} \leq 1$, and $|(\xi \cdot F_0)(z)|^2 = K_{\xi, D}^\psi(z)$. It means that $\frac{|(\xi \cdot F_0)(z)|^2}{\int_D |F_0|^2 e^{-\psi}} \geq K_{\xi, D}^\psi(z)$. Then by the definition of $K_{\xi, D}^\psi(z)$, we get $\frac{|(\xi \cdot F_0)(z)|^2}{\int_D |F_0|^2 e^{-\psi}} = K_{\xi, D}^\psi(z)$. \square

The following lemma shows that the weighted Bergman kernel is continuous.

Lemma 2.7 *Let D be a bounded domain in \mathbb{C}^n , ψ be a plurisubharmonic function on D . Then for any $\xi \in \ell_1$, $K_{\xi, D}^\psi(z)$ is a continuous function on D .*

Proof It is trivial when $\xi = 0$ or $A^2(D, e^{-\psi}) = \{0\}$. Now we assume $\xi \neq 0$ and $A^2(D, e^{-\psi}) \neq \{0\}$. By the definition of $K_{\xi, D}^\psi(z)$,

$$K_{\xi, D}^\psi(z) = \sup \left\{ |(\xi \cdot F)(z)|^2 : \int_D |F|^2 e^{-\psi} = 1, F \in \mathcal{O}(D) \right\}.$$

Combining with Lemma 2.1, we know that $K_{\xi, D}^\psi(z)$ is lower semicontinuous.

Next we prove that $K_{\xi, D}^\psi(z)$ is also upper continuous. Let $\{z_j\}$ be a sequence of points in D such that $\lim_{j \rightarrow +\infty} z_j = z_0 \in D$. And we may assume that $\{z_{k_j}\}$ is the subsequence of $\{z_j\}$ such that

$$\lim_{j \rightarrow +\infty} K_{\xi, D}^\psi(z_{k_j}) = \limsup_{j \rightarrow +\infty} K_{\xi, D}^\psi(z_j).$$

Using Lemma 2.6, we can get a sequence of holomorphic function $\{F_j\}$ on D , such that $\int_D |F_j|^2 e^{-\psi} = 1$, and $|(\xi \cdot F_j)(z_j)|^2 = K_{\xi, D}^\psi(z_j)$, for any $j \geq 1$. Then using Montel's theorem

and the diagonal method, we can select a subsequence of $\{F_{k_j}\}$ which is uniformly convergent on every compact subset of D . We may denote the subsequence by $\{F_{k_j}\}$ itself, and denote the limit function by F_0 . It follows from Fatou's lemma and Lemma 2.3 that $\int_D |F_0|^2 e^{-\psi} \leq 1$, $\lim_{j \rightarrow +\infty} (\xi \cdot F_{k_j})(z_{k_j}) = (\xi \cdot F_0)(z_0)$. Then

$$\begin{aligned} K_{\xi,D}^\psi(z_0) &\geq \frac{|(\xi \cdot F_0)(z_0)|^2}{\int_D |F_0|^2 e^{-\psi}} \\ &\geq |(\xi \cdot F_0)(z_0)|^2 \\ &= \lim_{j \rightarrow +\infty} |(\xi \cdot F_{k_j})(z_{k_j})|^2 \\ &= \lim_{j \rightarrow +\infty} K_{\xi,D}^\psi(z_{k_j}) \\ &= \limsup_{j \rightarrow +\infty} K_{\xi,D}^\psi(z_j). \end{aligned}$$

We get that $K_{\xi,D}^\psi(z)$ is upper semicontinuous.

It is known that $K_{\xi,D}^\psi(z)$ is lower semicontinuous, which implies that $K_{\xi,D}^\psi(z)$ is continuous. \square

The weighted Bergman kernel is also log-plurisubharmonic.

Lemma 2.8 *Let D be a bounded domain in \mathbb{C}^n , ψ be a plurisubharmonic function on D , and let $\xi \in \ell_1$. Then $\log K_{\xi,D}^\psi(z)$ is plurisubharmonic on D .*

Proof There is $\log K_{\xi,D}^\psi(z) \equiv -\infty$ when $\xi = 0$ or $A^2(D, e^{-\psi}) = \{0\}$. Now we assume that $\xi \neq 0$ and $A^2(D, e^{-\psi}) \neq \{0\}$. By the definition, we have

$$\log K_{\xi,D}^\psi(z) = \sup \left\{ 2 \log |(\xi \cdot F)(z)| : \int_D |F|^2 e^{-\psi} = 1, F \in \mathcal{O}(D) \right\}.$$

Lemma 2.1 shows that $(\xi \cdot F)(z)$ is holomorphic on D , when F is holomorphic on D . Then $\log |(\xi \cdot F)(z)|$ is plurisubharmonic. As $\log K_{\xi,D}^\psi(z)$ is upper semicontinuous according to Lemma 2.7, $\log K_{\xi,D}^\psi(z)$ is plurisubharmonic on D . \square

3 Optimal L^2 Extension and Guan–Zhou Method

In this section, we will recall the Guan–Zhou Method, i.e., an optimal L^2 extension approach to a log-convexity property of the fibrewised Bergman kernel.

Let Ω be a pseudoconvex domain in \mathbb{C}^{n+1} with coordinate (z, t) , where $z \in \mathbb{C}^n$, $t \in \mathbb{C}$. Let p, q be the natural projections $p(z, t) = t$, $q(z, t) = z$ on Ω , $p(\Omega) = D$. For any $t \in D$, suppose that $\Omega_t = p^{-1}(t) \subseteq \Omega$ is bounded in \mathbb{C}^n . Let ψ be a plurisubharmonic function on Ω , and let $K_{\xi,t}^\psi(z) = K_{\xi,\Omega_t}^\psi(z)$ be the Bergman kernels of the domains Ω_t defined as which in the above section with respect to some fixed $\xi \in \ell_1$.

We will use the following version of the optimal L^2 extension theorem.

Lemma 3.1 (Optimal L^2 extension theorem ([5], see [23, 24])) *Let $D = \Delta_{t_0,r}$ be the disk in the complex plane centered on t_0 with radius r . Then for any f in $A^2(\Omega_{t_0}, e^{-\psi})$, there exists a holomorphic function F on Ω , such that $F|_{\Omega_{t_0}} = f$, and*

$$\frac{1}{\pi r^2} \int_\Omega |F|^2 e^{-\psi} \leq \int_{\Omega_{t_0}} |f|^2 e^{-\psi}.$$

Guan–Zhou Method shows that Lemma 3.1 implies the following

Proposition 3.2 (see [1, 2, 18, 19]) $\log K_{\xi,t}^\psi(z)$ is a plurisubharmonic function with respect to (z, t) , for any $\xi \in \ell_1$.

Proof Firstly, we prove that $\log K_{\xi,t}^\psi(z)$ is upper semicontinuous on Ω . Let (z_j, t_j) be a sequence of points in Ω , such that $(z_j, t_j) \rightarrow (z_0, t_0) \in \Omega$, $j \rightarrow +\infty$. Since Ω is a domain in \mathbb{C}^{n+1} , we know that for any compact subset of $q(\Omega_{t_0})$ denoted by K , there exists $j_K \geq 1$, such that $K \subseteq q(\Omega_{t_j})$ in the sense of domains in \mathbb{C}^n , for any $j \geq j_K$.

We denote $q(\Omega_{t_j})$ by Ω_j . Then we may assume that $\{(z_{k_j}, t_{k_j})\}$ is the subsequence of $\{(z_j, t_j)\}$ such that

$$\lim_{j \rightarrow +\infty} K_{\xi, \Omega_{k_j}}^\psi(z_{k_j}) = \limsup_{j \rightarrow +\infty} K_{\xi, \Omega_j}^\psi(z_j),$$

and $z_0 \in \Omega_{j_0}$ for some $j_0 \in \mathbb{N}_+$. Using Lemma 2.6, we can get a sequence of holomorphic function $\{F_j\}$ on Ω_j , such that $\int_{\Omega_j} |F_j|^2 e^{-\psi} = 1$, and $|(\xi \cdot F_j)(z_j)|^2 = K_{\xi, \Omega_j}^\psi(z_j)$, for any $j \geq j_0$. Since for any compact subset K of $q(\Omega_{t_0})$, there exists $j_K \geq 1$, such that $K \subseteq \Omega_j$ for any $j \geq j_K$, then we can use Montel's theorem and the diagonal method to get a subsequence of $\{F_{k_j}\}$ which is uniformly convergent on every compact subset of $q(\Omega_{t_0})$. We may denote the subsequence by $\{F_{k_j}\}$ itself, and denote the limit function by F_0 . Fatou's lemma and Lemma 2.3 imply that $\int_{q(\Omega_{t_0})} |F_0|^2 e^{-\psi} \leq 1$, and $\lim_{j \rightarrow +\infty} (\xi \cdot F_{k_j})(z_{k_j}) = (\xi \cdot F_0)(z_0)$. It follows that

$$\begin{aligned} K_{\xi, \Omega_{t_0}}^\psi(z_0) &\geq \frac{|(\xi \cdot F_0)(z_0)|^2}{\int_{q(\Omega_{t_0})} |F_0|^2 e^{-\psi}} \\ &\geq |(\xi \cdot F_0)(z_0)|^2 \\ &= \lim_{j \rightarrow +\infty} |(\xi \cdot F_{k_j})(z_{k_j})|^2 \\ &= \lim_{j \rightarrow +\infty} K_{\xi, \Omega_{k_j}}^\psi(z_{k_j}) \\ &= \limsup_{j \rightarrow +\infty} K_{\xi, \Omega_{t_j}}^\psi(z_j), \end{aligned}$$

and

$$\limsup_{j \rightarrow +\infty} \log K_{\xi, t_j}^\psi(z_j) \leq \log K_{\xi, t_0}^\psi(z_0).$$

Then we obtain that $\log K_{\xi,t}^\psi(z)$ is upper semicontinuous on Ω .

Secondly we need to check that for any complex line L , $\log K_{\xi,t}^\psi(z)|_L$ is subharmonic. If the complex line lies on some Ω_t for some fixed t , we know that $\log K_{\xi,t}^\psi(z)|_L$ is subharmonic using Lemma 2.8. Then without loss of generality, we assume that L is the complex line on $\{t|(z, t)\}$ (meaning that L lies on $q^{-1}(z)$ for some $z \in \mathbb{C}^n$) and $D = \Delta_{t_0,r} = L$.

If $\log K_{\xi,t_0}^\psi(z) = -\infty$, we are done. Then we assume that there exists $f \in A^2(\Omega_{t_0}, e^{-\psi})$ such that

$$K_{\xi, t_0}^\psi(z) = \frac{|(\xi \cdot f)(z)|^2}{\int_{\Omega_{t_0}} |f|^2 e^{-\psi}}.$$

Using the optimal L^2 extension theorem (Lemma 3.1), we can get a holomorphic function F on Ω such that $F(z, t_0) = f(z)$ and

$$\frac{1}{\pi r^2} \int_{\Omega} |F|^2 e^{-\psi} \leq \int_{\Omega_{t_0}} |f|^2 e^{-\psi}.$$

Denote that $F_t(z) = F(z, t) = F|_{\Omega_t}$. Note that the function $y = \log x$ is concave, and by Jensen's inequality, it follows from Guan-Zhou Method ([18], see also [23, 24]) that

$$\begin{aligned}
 \log \left(\int_{\Omega_{t_0}} |f|^2 e^{-\psi} \right) &\geq \log \left(\frac{1}{\pi r^2} \int_{\Omega} |F|^2 e^{-\psi} \right) \\
 &= \log \left(\frac{1}{\pi r^2} \int_{\Delta_{t_0, r}} \int_{\Omega_t} |F_t|^2 e^{-\psi} \right) \\
 &\geq \frac{1}{\pi r^2} \int_{\Delta_{t_0, r}} \log \left(\int_{\Omega_t} |F_t|^2 e^{-\psi} \right) \\
 &\geq \frac{1}{\pi r^2} \int_{\Delta_{t_0, r}} (\log |(\xi \cdot F_t)(z)|^2 - \log K_{\xi, t}^{\psi}(z)) \\
 &\geq \log |(\xi \cdot f)(z)|^2 - \frac{1}{\pi r^2} \int_{\Delta_{t_0, r}} \log K_{\xi, t}^{\psi}(z).
 \end{aligned}$$

The last inequality above holds, since we can prove that $\log |(\xi \cdot F_t)(z)|^2$ is subharmonic with respect to t . In fact, if we write

$$F_t(w) = \sum_{\alpha \in \mathbb{N}^n} a_{\alpha}(t)(w - z)^{\alpha},$$

then all the

$$a_{\alpha}(t) = \frac{1}{\alpha!} \frac{\partial^{\alpha} F(w, t)}{\partial w^{\alpha}}(z, t)$$

are holomorphic with respect to t . In addition, $(\xi \cdot F_t)(z) = \sum_{\alpha \in \mathbb{N}^n} \xi_{\alpha} a_{\alpha}(t)$ is absolutely and uniformly convergent on every compact subset of $\Delta_{t_0, r}$. Since for any compact subset of $\Delta_{t_0, r}$, denoted by K , we can find some $R > 0$ such that $\Delta_{z, R}^n \subseteq q(\Omega_t)$ for any $t \in K$. Combining with $\xi \in \ell_1$, we get that

$$|\xi_{\alpha} a_{\alpha}(t)| \leq \frac{MM'}{(\rho R)^{|\alpha|}}, \quad \forall \alpha \in \mathbb{N}^n$$

for some $M, M' > 0$, $\rho > 1/R$, and any $t \in K$. This means that $\sum_{\alpha \in \mathbb{N}^n} \xi_{\alpha} a_{\alpha}(t)$ is absolutely and uniformly convergent on K . Then $(\xi \cdot F_t)(z)$ is holomorphic with respect to t for any fixed z , which implies that $\log |(\xi \cdot F_t)(z)|^2$ is subharmonic with respect to t . Then

$$\log K_{\xi, t_0}^{\psi}(z) \leq \frac{1}{\pi r^2} \int_{t \in \Delta_{t_0, r}} \log K_{\xi, t}^{\psi}(z),$$

which implies that $\log K_{\xi, t}^{\psi}(z)$ is plurisubharmonic with respect to (z, t) . □

4 Effectiveness Result of L^p Strong Openness Property

In this section, we show the L^2 extension approach to the L^p effectiveness of the strong openness property.

Let I be an ideal of \mathcal{O}_o such that $I \neq \mathcal{O}_o$. We consider about a subset $\ell_I \subseteq \ell_1$ such as

$$\ell_I = \{0 \neq \xi \in \ell_1 : (\xi \cdot F)(o) = 0, \forall F \in I\}.$$

It is obvious that ℓ_I is always nonempty since $\xi = (1, 0, \dots, 0, \dots) \in \ell_I$ when $I \neq \mathcal{O}_o$.

Especially, we denote $\ell_{\mathcal{I}(\varphi)_o}$ by ℓ_{φ} .

Let φ be a negative plurisubharmonic function on a bounded pseudoconvex domain $D \subseteq \mathbb{C}^n$, and let ψ be a plurisubharmonic function on D . Denote that

$$D_t := \{z \in D : \varphi(z) < -t\},$$

and

$$K_\xi^\psi(t) := K_{\xi, D_t}^\psi(o),$$

for $t \in [0, +\infty)$. We need the following lemma.

Lemma 4.1 ([7], Theorem 5.13, Chapter I) *Let $\Omega = I + i\mathbb{R}$ be a domain in \mathbb{C} with the coordinate $z = x + iy$, where I is an interval in \mathbb{R} . Let $\phi(z)$ be a subharmonic function on Ω which is independent of y . Then $\phi(x) := \phi(x + i\mathbb{R})$ is a convex function with respect to $x \in I$.*

This result is also used in [2, 4] and [19].

Note that the domain

$$\{(z, \tau) : \varphi(z) - \operatorname{Re} \tau < 0\}$$

is pseudoconvex in \mathbb{C}^{n+1} , then according to Proposition 3.2, $\log K_{\xi, \tau}^\psi(o)$ is subharmonic for $\tau \in [0, +\infty) + i\mathbb{R}$, and independent of $\operatorname{Im} \tau$. Lemma 4.1 shows that $\log K_\xi^\psi(t)$ is convex for $t \in [0, +\infty)$. This implies that $-\log K_\xi^\psi(t) + t$ is concave, which will be increasing if it has a lower bound. We state the following result.

Lemma 4.2 *Let D be a bounded pseudoconvex domain in \mathbb{C}^n such that $o \in D$, and let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$, and ψ be a plurisubharmonic function on D such that $\mathcal{I}(\varphi + \psi)_o \neq \mathcal{I}(\psi)_o$. Then for any fixed $\xi \in \ell_{\varphi+\psi}$,*

$$-\log K_\xi^\psi(t) + t \geq -\log K_\xi^\psi(0), \quad \forall t \in [0, +\infty).$$

Lemma 4.2 can be proved by the following lemma.

Lemma 4.3 ([21]) *Let D be a bounded pseudoconvex domain in \mathbb{C}^n such that $o \in D$, and let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$, and ψ be a plurisubharmonic function on D . Let F be an holomorphic function on $\{\varphi < -t_0\}$ such that $\int_{\{\varphi < -t_0\}} |F|^2 e^{-\psi} < +\infty$. Then there exists a holomorphic function F_{t_0} on D , such that*

$$(F_{t_0} - F, o) \in \mathcal{I}(\varphi + \psi)_o$$

and

$$\int_D |F_{t_0} - (1 - b_{t_0}(\varphi))F|^2 e^{-\psi} \leq C_D \int_{\{-t_0-1 < \varphi < -t_0\}} |F|^2 e^{-\varphi-\psi},$$

where $b_{t_0}(t) = \int_{-\infty}^t \mathbb{I}_{\{-t_0-1 < s < -t_0\}} ds$ and $t_0 \geq 0$, and here C_D is a positive constant only dependent of D .

Lemma 4.3 can also be referred to [12, 16–18].

Proof of Lemma 4.2 We only need to prove that there is a lower bound of $-\log K_\xi^\psi(t) + t$.

Lemma 2.6 shows that for any $t \in [0, +\infty)$, there exists $F \in \mathcal{O}(\{\varphi < -t\})$, such that

$$K_\xi^\psi(t) = \frac{|(\xi \cdot F)(o)|^2}{\int_{\{\varphi < -t\}} |F|^2 e^{-\psi}}.$$

It follows from Lemma 4.3 that there exists a holomorphic function F_t on D such that

$$(F_t - F, o) \in \mathcal{I}(\varphi + \psi)_o$$

and

$$\int_D |F_t - (1 - b_t(\varphi))F|^2 e^{-\psi} \leq C_D \int_{\{-t-1 < \varphi < -t\}} |F|^2 e^{-\varphi-\psi}.$$

Note that $\xi \in \ell_{\varphi+\psi}$, then $(F_t - F, o) \in \mathcal{I}(\varphi + \psi)_o$ induces that $(\xi \cdot F_t)(o) = (\xi \cdot F)(o)$. On the one hand,

$$\begin{aligned} & \left(\int_D |F_t - (1 - b_t(\varphi))F|^2 e^{-\psi} \right)^{\frac{1}{2}} \\ & \geq \left(\int_D |F_t|^2 e^{-\psi} \right)^{\frac{1}{2}} - \left(\int_D |(1 - b_t(\varphi))F|^2 e^{-\psi} \right)^{\frac{1}{2}} \\ & \geq (K_{\xi, D}^\psi(o))^{-\frac{1}{2}} |(\xi \cdot F_t)(o)| - \left(\int_{\{\varphi < -t\}} |F|^2 e^{-\psi} \right)^{\frac{1}{2}} \\ & = (K_{\xi, D}^\psi(o))^{-\frac{1}{2}} |(\xi \cdot F)(o)| - \left(\int_{\{\varphi < -t\}} |F|^2 e^{-\psi} \right)^{\frac{1}{2}}. \end{aligned}$$

On the other hand, we have

$$\int_{\{-t-1 < \varphi < -t\}} |F|^2 e^{-\varphi-\psi} \leq e^{t+1} \int_{\{-t-1 < \varphi < -t\}} |F|^2 e^{-\psi} \leq e^{t+1} \int_{\{\varphi < -t\}} |F|^2 e^{-\psi}.$$

Then

$$(C_D^{\frac{1}{2}} e^{\frac{t+1}{2}} + 1)^2 \int_{\{\varphi < -t\}} |F|^2 e^{-\psi} \geq (K_{\xi, D}^\psi(o))^{-1} |(\xi \cdot F)(o)|^2,$$

which means

$$\int_{\{\varphi < -t\}} |F|^2 e^{-\psi} \geq C e^{-t} |(\xi \cdot F)(o)|^2,$$

where $C = (2(e+1)K_{\xi, D}^\psi(o) \max\{C_D, 1\})^{-1}$ is a positive constant independent of the choices of F and t .

Then we get that

$$-\log K_\xi^\psi(t) + t \geq \log C$$

for any $t \in [0, +\infty)$. Then $-\log K_\xi^\psi(t) + t$ has a lower bound, inducing that it is increasing, and

$$-\log K_\xi^\psi(t) + t \geq -\log K_\xi^\psi(0). \quad \square$$

By Lemma 4.2, we get the following proposition.

Proposition 4.4 *Let D be a bounded pseudoconvex domain in \mathbb{C}^n such that $o \in D$, and let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$, and ψ be a plurisubharmonic function on D such that $\mathcal{I}(\varphi + \psi)_o \neq \mathcal{I}(\psi)_o$. Let F be a holomorphic function on D . Then for any $q > 1$, and any $\xi \in \ell_{\varphi+\psi}$,*

$$\int_D |F|^2 e^{-\frac{\varphi}{q}-\psi} \geq \frac{q}{q-1} \cdot \frac{|(\xi \cdot F)(o)|^2}{K_{\xi, D}^\psi(o)}.$$

Proof Using Lemma 4.2, we have that $K_\xi^\psi(t) \leq e^t K_\xi^\psi(0)$ for any $t \in [0, +\infty)$ if $\xi \in \ell_{\varphi+\psi}$. And It is known that

$$\int_D |F|^2 e^{-\frac{\varphi}{q}-\psi} = \int_{-\infty}^{+\infty} \left(\int_{\{\frac{\varphi}{q} < -t\}} |F|^2 e^{-\psi} \right) e^t dt,$$

(this equality can be referred in [12]), which implies that

$$\begin{aligned} \int_0^{+\infty} \left(\int_{\{\varphi < -qt\}} |F|^2 e^{-\psi} \right) e^t dt &\geq \int_0^{+\infty} \frac{|(\xi \cdot F)(o)|^2}{K_\xi^\psi(qt)} e^t dt \\ &\geq \frac{|(\xi \cdot F)(o)|^2}{K_\xi^\psi(0)} \int_0^{+\infty} e^{(1-q)t} dt \\ &= \frac{1}{q-1} \cdot \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)}, \end{aligned}$$

and

$$\begin{aligned} \int_{-\infty}^0 \left(\int_{\{\varphi < -qt\}} |F|^2 e^{-\psi} \right) e^t dt &\geq \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)} \int_{-\infty}^0 e^t dt = \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)}, \\ \int_D |F|^2 e^{-\frac{\varphi}{q} - \psi} &\geq \frac{q}{q-1} \cdot \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)}. \end{aligned} \quad \square$$

Let D be a bounded domain in \mathbb{C}^n , and let I be an ideal of \mathcal{O}_o such that $I \not\supseteq \mathcal{I}(\psi)_o$. Let ψ be a plurisubharmonic function on D . Let (F, o) be an element in \mathcal{O}_o . Then we denote that

$$B^\psi(F, I, D) = \sup_{\xi \in \ell_I} \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)}.$$

Then we get the following corollary of Proposition 4.4.

Corollary 4.5 *Let D be a bounded pseudoconvex domain in \mathbb{C}^n such that $o \in D$, and let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$, and ψ be a plurisubharmonic function on D such that $\mathcal{I}(\varphi + \psi)_o \neq \mathcal{I}(\psi)_o$. Let F be a holomorphic function on D . Then for any $q > 1$,*

$$\int_D |F|^2 e^{-\frac{\varphi}{q} - \psi} \geq \frac{q}{q-1} B^\psi(F, \mathcal{I}(\varphi + \psi)_o, D).$$

The following proposition shows that $B^\psi(F, I, D) > 0$ if $(F, o) \notin I$ and $A^2(D, e^{-\psi}) \neq \{0\}$.

Proposition 4.6 ([1]) *Let D be a bounded domain in \mathbb{C}^n such that $o \in D$, and let I be an ideal of \mathcal{O}_o such that $I \neq \mathcal{O}_o$. Let $(F, o) \in \mathcal{O}_o$ such that $(F, o) \notin I$. Then we are able to find some $\xi \in \ell_I$ such that $(\xi \cdot F)(o) \neq 0$.*

Moreover, if $F \in A^2(D, e^{-\psi})$, then $B^\psi(F, I, D) \leq \int_D |F|^2 e^{-\psi}$, since

$$\begin{aligned} B^\psi(F, I, D) &= \sup_{\xi \in \ell_I, (\xi \cdot F)(o) \neq 0} \frac{|(\xi \cdot F)(o)|^2}{K_{\xi,D}^\psi(o)} \\ &\leq \sup_{\xi \in \ell_I, (\xi \cdot F)(o) \neq 0} \frac{|(\xi \cdot F)(o)|^2}{|(\xi \cdot F)(o)|^2 / \int_D |F|^2 e^{-\psi}} \\ &= \int_D |F|^2 e^{-\psi}. \end{aligned}$$

It is clear that if there are two ideals I_1 and I_2 of \mathcal{O}_o such that $I_1 \subseteq I_2 \neq \mathcal{O}_o$ and $I_2 \not\supseteq \mathcal{I}(\psi)_o$, then $B^\psi(F, I_1, D) \geq B^\psi(F, I_2, D)$.

Now we can prove the following theorem.

Theorem 4.7 *Let D be a bounded pseudoconvex domain in \mathbb{C}^n such that $o \in D$, and let φ be a negative plurisubharmonic function on D such that $\varphi(o) = -\infty$, and ψ be a plurisubharmonic*

function on D . Let F be a holomorphic function on D with $\int_D |F|^2 e^{-\varphi-\psi} < +\infty$. Then for any $q > 1$ satisfying

$$\frac{q}{q-1} > \frac{\int_D |F|^2 e^{-\varphi-\psi}}{B^\psi(F, \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o, D)},$$

$|F|^2 e^{-q\varphi-\psi}$ is locally integrable near o .

Here $c_o^F(\varphi, \psi) := \sup\{c \geq 0 : |F|^2 e^{-c\varphi-\psi} \text{ is locally integrable near } o\}$.

Proof For $q > c_o^F(\varphi, \psi)$, $|F|^2 e^{-q\varphi-\psi}$ is not integrable near o , and $B^\psi(F, \mathcal{I}(\psi + q\varphi)_o, D) \geq B^\psi(F, \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o, D)$. Then Corollary 4.5 shows that

$$\int_D |F|^2 e^{-\varphi-\psi} \geq \frac{q}{q-1} B^\psi(F, \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o, D).$$

Let $q \rightarrow c_o^F(\varphi, \psi)^+$. Then the above inequality also holds for $q \geq c_o^F(\varphi, \psi)$. Then if $q > 1$ satisfying

$$\int_D |F|^2 e^{-\varphi-\psi} < \frac{q}{q-1} B^\psi(F, \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o, D),$$

we get $q < c_o^F(\varphi, \psi)$, which means that $|F|^2 e^{-q\varphi-\psi}$ is integrable near o .

Since $F \notin \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o$, we know that

$$0 < B^\psi(F, \mathcal{I}_+(\psi + c_o^F(\varphi, \psi)\varphi)_o, D) \leq \int_D |F|^2 e^{-\psi} < \int_D |F|^2 e^{-\varphi-\psi}.$$

Then the proof is done. \square

Then we give the proof of Theorem 1.2.

Proof of Theorem 1.2 In Theorem 4.7, let $\psi = \varphi_1 + a\varphi$. Then $\int_D |F_1|^2 e^{-\varphi-\psi} = \int_D |F|^p e^{-(1+a)\varphi}$, and $\mathcal{I}_+(\psi + c_o^{F_1}(\varphi, \psi)\varphi)_o = \mathcal{I}_+(\varphi_1 + c_{o,p}^F(\varphi)\varphi)_o$. Thus for any $q' > 1$ satisfying

$$\frac{q'}{q'-1} > \frac{\int_D |F|^p e^{-(1+a)\varphi}}{B^{\varphi_1+a\varphi}(F_1, \mathcal{I}_+(\varphi_1 + c_{o,p}^F(\varphi)\varphi)_o, D)},$$

$|F|^p e^{-(q'+a)\varphi}$ is locally integrable near o . Let $q = q' + a$. Then the proof is done. \square

Acknowledgements We would like to thank Zheng Yuan and Zhitong Mi for checking this article. And we thank the referees for their time and comments.

References

- [1] Bao, S. J., Guan, Q. A.: L^2 extension and effectiveness of strong openness property. submitted (2021)
- [2] Berndtsson, B.: Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains. *Ann. Inst. Fourier (Grenoble)*, **56**(6), 1633–1662 (2006)
- [3] Berndtsson, B.: The openness conjecture for plurisubharmonic functions. arXiv:1305.5781, (2013)
- [4] Berndtsson, B., Lempert, L.: A proof of the Ohsawa-Takegoshi theorem with sharp estimates. *J. Math. Soc. Japan*, **68**(4), 1461–1472 (2016)
- [5] Blocki, Z.: Suita conjecture and the Ohsawa–Takegoshi extension theorem. *Invent. Math.*, **193**, 149–158 (2013)
- [6] Demailly, J. -P.: Transcendental proof of a generalized Kawamata–Viehweg vanishing theorem. In: Geometrical and Algebraical Aspects in Several Complex Variables (Cetraro, 1989), Vol. 8 of Sem. Conf., 81–94 (1991)
- [7] Demailly, J. -P.: Complex analytic and differential geometry. Electronically accessible at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>

- [8] Demailly, J. -P.: Multiplier ideal sheaves and analytic methods in algebraic geometry. In: School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000), volume 6 of ICTP Lect. Notes, 1–148 (2001)
- [9] Demailly, J. -P.: Analytic Methods in Algebraic Geometry. volume 1 of Surveys of Modern Mathematics. International Press, Somerville, MA; Higher Education Press, Beijing (2012)
- [10] Demailly, J. -P., Kollár, J.: Semi-continuity of complex singularity exponents and Kähler–Einstein metrics on Fano orbifolds. *Ann. Sci. École Norm. Sup. (4)*, **34**(4), 525–556 (2001)
- [11] Fornæss, J. E.: Several complex variables. arXiv:1507.00562 (2015)
- [12] Guan, Q. A.: A sharp effectiveness result of Demailly’s strong openness conjecture. *Adv. Math.*, **348**, 51–80 (2019)
- [13] Guan, Q. A.: General concavity of minimal L^2 integrals related to multiplier sheaves. arXiv:1811.03261v4 (2019)
- [14] Guan, Q. A., Yuan, Z.: Effectiveness of strong openness property in L^p . arXiv:2106.03552v3 (2021)
- [15] Guan, Q. A., Zhou, X. Y.: A proof of Demailly’s strong openness conjecture. *Ann. of Math. (2)*, **182**(2), 605–616 (2015)
- [16] Guan, Q. A., Zhou, X. Y.: Strong openness conjecture and related problems for plurisubharmonic functions. arXiv:1401.7158 (2014)
- [17] Guan, Q. A., Zhou, X. Y.: Effectiveness of Demailly’s strong openness conjecture and related problems. *Invent. Math.*, **202**(2), 635–676 (2015)
- [18] Guan, Q. A., Zhou, X. Y.: A solution of an L^2 extension problem with an optimal estimate and applications. *Ann. of Math. (2)*, **181**(3), 1139–1208 (2015)
- [19] Maitani, F., Yamaguchi, H.: Variation of Bergman metrics on Riemann surfaces. *Math. Ann.*, **330**(3), 477–489 (2004)
- [20] Nadel, A. M.: Multiplier ideal sheaves and Kähler–Einstein metrics of positive scalar curvature. *Ann. of Math. (2)*, **132**(3), 549–596 (1990)
- [21] Ohsawa, T.: On the extension of L^2 holomorphic functions. V. Effects of generalization. *Nagoya Math. J.*, **161**, 1–21 (2001)
- [22] Ohsawa, T.: Analysis of Several Complex Variables. Translations of Mathematical Monographs, Vol. 211, American Mathematical Society, Providence, RI, 2002. Translated from the Japanese by Shu Gilbert Nakamura, Iwanami Series in Modern Mathematics
- [23] Ohsawa, T.: L^2 Approaches in Several Complex Variables. Development of Oka–Cartan Theory by L^2 Estimates for the $\bar{\partial}$ operator. Springer Monographs in Mathematics, Springer, Tokyo (2015) ix+196 pp. ISBN: 978-4-431-55746-3; 978-4-431-55747-0
- [24] Ohsawa, T.: L^2 Approaches in Several Complex Variables. Towards the Oka–Cartan Theory with Precise Bounds. Second edition of [23]. Springer Monographs in Mathematics, Springer, Tokyo (2018) xi+258 pp. ISBN: 978-4-431-56851-3; 978-4-431-56852-0
- [25] Siu, Y. T.: Multiplier ideal sheaves in complex and algebraic geometry. *Sci. China Ser. A*, **48**, 1–31 (2005)
- [26] Tian, G.: On Kähler–Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$. *Invent. Math.*, **89**(2), 225–246 (1987)