

# Number Theory: Notes on Xie Junyi's classes

BAO

*Date: December 24, 2025*

## Abstract

These are the notes on Prof. Xie Junyi's classes.

## Contents

<b>1</b>	<b>Algebraic Integers</b>	<b>3</b>
1.1	Note on 20250922 . . . . .	3
1.1.1	Finding the ring of integers . . . . .	5
1.2	Note on 20250924 . . . . .	6
1.2.1	General strategy . . . . .	6
<b>2</b>	<b>Dedekind Domains and Factorization</b>	<b>8</b>
2.1	Note on 20250924 . . . . .	8
2.1.1	Discrete valuation rings . . . . .	8
2.1.2	Dedekind domains . . . . .	9
2.2	Note on 20250929 . . . . .	10
2.2.1	Unique factorization of ideals . . . . .	10
2.3	Note on 20251020 . . . . .	13
2.3.1	Ideal class group . . . . .	13
<b>3</b>	<b>Discrete valuations</b>	<b>14</b>
3.1	Note on 20251022 . . . . .	14
<b>4</b>	<b>Factorization in extensions</b>	<b>17</b>
4.1	Note on 20251027 . . . . .	17
4.1.1	The primes that ramify . . . . .	18
4.2	Note on 20251103 . . . . .	20
<b>5</b>	<b>The finiteness of the class number</b>	<b>21</b>
5.1	Note on 20251103 . . . . .	21
5.1.1	Norms of ideals . . . . .	21
5.2	Note on 20251105 . . . . .	23
5.2.1	Lattices . . . . .	25
5.3	Note on 20251110 . . . . .	27
5.3.1	Finiteness of the class number . . . . .	28
5.3.2	Binary quadratic forms . . . . .	29
5.4	Note on 20251117 . . . . .	31

<b>6</b>	<b>The Unit Theorem</b>	<b>32</b>
6.0.1	Proof of that $U_K$ is finitely generated	32
6.1	Note on 20251119	35
6.1.1	$S$ -units	35
6.1.2	Example: CM fields	35
6.1.3	Cyclotomic extensions	36
6.2	Note on 20251126	39
6.3	Note on 20251201	42
<b>7</b>	<b>Absolute values</b>	<b>45</b>
7.0.1	NA absolute value	45
7.1	Note on 20251203	47
7.1.1	Places of a number field	49
7.2	Note on 20251215	51
7.2.1	The weak approximation theorem	51
<b>8</b>	<b>Completions</b>	<b>53</b>
8.0.1	Completion for discrete valuation field	54
8.0.2	Newton's lemma	55
8.1	Note on 20251217	56
8.1.1	Hensel's lemma	56
8.1.2	Extensions of NA absolute values	57
8.2	Note on 20251222	59
8.2.1	Newton's polygon	59
<b>9</b>	<b>Locally compact field</b>	<b>61</b>
9.0.1	Unramified extensions of a field	61
9.1	Note on 20251224	63
9.1.1	Totally ramified extension of $K$	63
9.1.2	Ramification groups	64
9.1.3	Krasner's lemma	65

# 1 Algebraic Integers

## 1.1 Note on 20250922

**COROLLARY 1.1.** *Let  $L$  be a number field. Then  $\mathcal{O}_L$  is the biggest subring of  $L$  which is finitely generated as a  $\mathbb{Z}$ -module.*

**Proof.**  $\mathcal{O}_L$  is f. g.  $\mathbb{Z}$ -module. Let  $B$  be a subring of  $L$  which is f. g. as a  $\mathbb{Z}$ -module. Then for any  $b \in B$ ,  $b$  is integral over  $\mathbb{Z}$ , so  $B \subset \mathcal{O}_L$ . ■

**EXAMPLE 1.2.** If  $\text{char } p > 0$ , there exists DVR, which is not Japanese, i.e. there exists  $A$  DVR,  $L$  finite field extension of  $\text{Frac}(A)$ , such that the integral closure  $B$  of  $A$  in  $L$  is not a finitely generated  $A$ -module.

For example, let

$$k = \mathbb{F}_p(t_0, t_1, \dots), \quad K = \mathbb{F}_p(t_0^{1/p}, t_1^{1/p}, \dots),$$

and

$$A := \left\{ h \in K[[x]] \mid h = \sum_{i \geq 0} a_i x^i, \ a_i \in K, \ k(a_0, a_1, \dots) \text{ is a finite extension over } L \right\}.$$

Then  $A$  is a DVR. Note

$$A^\times = \{a_0 + \dots \mid a_0 \neq 0\},$$

so

$$f := \sum_{i \geq 0} t_i^{1/p} x^i \notin A.$$

Let  $R := A[f]$ . Only need to show the integral closure  $B$  of  $R$  in  $L := \text{Frac}(R)$  is not f. g. over  $R$ .

For all  $n \geq 0$ , set

$$h_n := \sum_{i \geq n} t_i^{1/p} x^{i-n} = \frac{1}{x^n} \left( f - \sum_{i < n} t_i^{1/p} x^i \right) \in L.$$

Moreover,

$$h_n^p = \sum_{i \geq n} t_i x^{p(i-n)} \in A \implies h_n \in B, \quad \forall n.$$

Hence,  $f \in A + x^n B$  for all  $n$ .

Assume  $B$  is f. g. over  $A$ . Take

$$M := B/A, \quad N := \bigcap_{n \geq 1} x^n M,$$

which are f. g.  $A$ -modules, and  $xN = N$ . By Nakayama's lemma ( $\mathfrak{m} = (x)$ ),  $N = 0$ . So  $f \in A$ , which is a contradiction. Q.E.D.

Let  $C = \sum A\beta_i \subset B$  with  $\beta_i$  a basis for  $L/K$ .

Define

$$\begin{aligned} C^* &:= \{\beta \in L \mid \text{Tr}(\beta \cdot \gamma) \in A, \forall \gamma \in C\} \\ &= \{\beta \in L \mid \text{Tr}(\beta \cdot \beta_i) \in A \text{ } i = 1, \dots, m\} \\ &= \sum A\beta'_i, \end{aligned}$$

where  $\{\beta'_i\}$  is the dual basis of  $\beta_i$ . We have

$$C = \sum A\beta_i \subset B \subset \sum A\beta'_i = C^*.$$

Then, how to find  $C^*$ ?

Assume  $L = \mathbb{Q}[\beta]$  with  $\beta \in \mathcal{O}_L$ . Let  $f(x)$  be the minimal polynomial of  $\beta$  with  $\deg f = m$ . Let

$$C = \mathbb{Z}[\beta] = \bigoplus_{i=0}^{m-1} \beta^i.$$

Want to find  $C^*$ .

**LEMMA 1.3 (Euler).**

$$\text{Tr}(\beta^i / f'(\beta)) = \begin{cases} 0, & 0 \leq i \leq m-2, \\ 1, & i = m-1. \end{cases}$$

**Proof.** Let  $\beta_1 = \beta, \dots, \beta_m$  be the roots of  $f$ . Then

$$\text{Tr}(\beta^i / f'(\beta)) = \sum_{j=1}^m \frac{\beta_j^i}{\prod_{k \neq j} (\beta_j - \beta_k)}.$$

Consider

$$D_j(x) := \prod_{k \neq j} (x - \beta_k) \in \overline{\mathbb{Q}}[x],$$

$\deg D_j = m-1$ , and

$$\frac{D_j(x)}{D_j(\beta_j)} = \begin{cases} 1, & x = \beta_j, \\ 0, & x = \beta_k, \text{ } k \neq j. \end{cases}$$

So any polynomial  $P \in \overline{\mathbb{Q}}[x]$  of  $\deg \leq m-1$ , we have

$$P(x) = \sum_{j=1}^m \frac{D_j(x)}{D_j(\beta_j)} P(\beta_j),$$

so

$$x^l = \sum_{j=1}^m \frac{D_j(x)}{D_j(\beta_j)} \cdot \beta_j^l, \quad l = 0, \dots, m-1.$$

Compare the coefficients of  $x^{n-1}$ . We get

$$\sum_{j=1}^m \frac{\beta_j^l}{D_j(\beta_j)} = \begin{cases} 0, & l < m-1, \\ 1, & l = m-1. \end{cases}$$

■

As

$$\mathbb{Z}[\beta] = \bigoplus_{i=0}^{m-1} \mathbb{Z}\beta^i,$$

Lemma  $\implies$

$$\mathrm{Tr} \left( \beta^l / f'(\beta) \right) \in A, \quad \forall l \geq 0.$$

Moreover,

$$\det \left( \mathrm{Tr} \left( \beta^i \cdot \frac{\beta^j}{f'(\beta)} \right)_{0 \leq i, j \leq m-1} \right) = (-1)^m,$$

which is a unit in  $\mathbb{Z}$ . Hence,

$$\left\{ \frac{\beta^i}{f'(\beta)} \mid i = 0, \dots, m-1 \right\}$$

is a basis of  $C^*$   $\implies$

$$C^* = (f'(\beta))^{-1} A[\beta].$$

### 1.1.1 Finding the ring of integers

Let  $K$  be a field of char 0.

**PROPOSITION 1.4.** *Let  $L = K[\beta]$  for some  $\beta$ , and  $f(x)$  the minimal polynomial of  $\beta$  over  $K$  with  $\deg f = m$ . Suppose  $\beta_1, \dots, \beta_m$  are the roots of  $f$  in  $\bar{K}$ . Then the discriminant of  $f$ :*

$$D(1, \beta, \dots, \beta^{m-1}) = \prod_{1 \leq i < j \leq m} (\beta_i - \beta_j)^2 = (-1)^{\frac{m(m-1)}{2}} \mathrm{Nm}_{L/K} (f'(\beta)).$$

**Proof.**

$$\begin{aligned} D(1, \beta, \dots, \beta^{m-1}) &= \det (\sigma_i(\beta^j))^2 = \det(\beta_i^j)^2 \\ &= \prod_{1 \leq i < j \leq m} (\beta_i - \beta_j)^2 \\ &= (-1)^{\frac{m(m-1)}{2}} \prod_i \prod_{j \neq i} (\beta_i - \beta_j) \\ &= (-1)^{\frac{m(m-1)}{2}} \prod_i f'(\beta_i) \\ &= (-1)^{\frac{m(m-1)}{2}} \mathrm{Nm}_{L/K} (f'(\beta)). \end{aligned}$$

■

**REMARK 1.5.**  $D(1, \beta, \dots, \beta^{m-1}) = 0$  iff  $f$  has multiple roots.

Let  $L$  be a number field.

**PROPOSITION 1.6.** *Let  $\beta_1, \dots, \beta_m$  be a basis of  $L/\mathbb{Q}$ , and  $d := D(\beta_1, \dots, \beta_m) \in \mathbb{Z}$ . Then*

$$\mathbb{Z}\beta_1 + \dots + \mathbb{Z}\beta_m \subset \mathcal{O}_L \subset \mathbb{Z}\frac{\beta_1}{d} + \dots + \mathbb{Z}\frac{\beta_m}{d}.$$

**Proof.** For  $\beta \in \mathcal{O}_L$ , write

$$\beta = \sum x_i \beta_i, \quad x_i \in \mathbb{Q}.$$

Let  $\sigma_1, \dots, \sigma_m$  be the embeddings of  $L$  into  $\overline{\mathbb{Q}}$ . Then

$$\sigma_j(\beta) = \sum_i x_i \sigma_j(\beta_i), \quad j = 1, \dots, m.$$

Solve  $x_i$ , we get

$$x_i = \frac{A_i}{\det(\sigma_j(\beta_k))} \in \frac{\mathcal{O}_L}{\det(\sigma_j(\beta_k))}, \quad i = 1, \dots, m.$$

Note  $\det(\sigma_j(\beta_k))^2 = d$ . Hence, every  $dx_i \in \mathcal{O}_L \cap \mathbb{Q} = \mathbb{Z}$ . So  $\beta \in \mathbb{Z} \frac{\beta_1}{d} + \dots + \mathbb{Z} \frac{\beta_m}{d}$ . ■

Now write  $L = \mathbb{Q}[\alpha]$  with  $\alpha \in \mathcal{O}_L$ . Compute  $d = D(1, \alpha, \dots, \alpha^{m-1})$ . Then

$$\mathbb{Z}[\alpha] \subset \mathcal{O}_L d^{-1} \mathbb{Z}[\alpha].$$

Note

$$(d^{-1} \mathbb{Z}[\alpha] : \mathbb{Z}[\alpha]) = d^m.$$

For every coset  $\beta + \mathbb{Z}[\alpha]$  of  $d^{-1} \mathbb{Z}[\alpha]$  is in  $\mathcal{O}_L$  iff

$$(\beta + \mathbb{Z}[\alpha]) \cap \mathcal{O}_L \neq \emptyset.$$

Let  $\beta_1, \dots, \beta_m \in d^{-1} \mathbb{Z}[\alpha]$  represent all the cosets of  $d^{-1} \mathbb{Z}[\alpha] / \mathbb{Z}[\alpha]$ . Test any  $\beta_i$  whether  $\beta_i \in \mathcal{O}_L$  or not.

## 1.2 Note on 20250924

### 1.2.1 General strategy

Write  $K = \mathbb{Q}[\alpha]$  with  $\alpha \in \mathcal{O}_K$ . Compute  $D(1, \alpha, \dots, \alpha^{m-1})$ . If it is square-free, then  $\{1, \alpha, \dots, \alpha^{m-1}\}$  is automatically an integral basis as

$$D(1, \alpha, \dots, \alpha^{m-1}) = \text{disc}(\mathcal{O}_K/\mathbb{Z})(\mathcal{O}_K : \mathbb{Z}[\alpha])^2.$$

If it is not square-free,  $\{1, \alpha, \dots, \alpha^{m-1}\}$  may still be an integral basis. Sometimes we can show this by [Stickelberger's thm](#) or look at [how prime ramify](#). If  $\{1, \alpha, \dots, \alpha^{m-1}\}$  is not an integral basis, one has to look for algebraic integers outside  $\mathbb{Z}[\alpha]$ .

**PROPOSITION 1.7.** *Let  $K$  be a number field.*

- (a) *The sign of  $\text{disc}(K/\mathbb{Q})$  is  $(-1)^s$ , where  $2s$  is the number for homomorphisms  $K \hookrightarrow \mathbb{C}$  whose image is not in  $\mathbb{R}$ ;*
- (b) *([Stickelberger's thm](#))  $\text{disc}(\mathcal{O}_K/\mathbb{Z}) \equiv 0 \text{ or } 1 \pmod{4}$ .*

**Proof.** (a) Let  $K = \mathbb{Q}[\alpha]$  and  $\alpha_1, \dots, \alpha_r$  be the real conjugates of  $\alpha$ , and  $\alpha_{r+1}, \overline{\alpha_{r+1}}, \dots, \alpha_{r+s}, \overline{\alpha_{r+s}}$  be the complex conjugates of  $\alpha$ . Then

$$\text{sign}\left(D(1, \alpha, \dots, \alpha^{m-1})\right) = \text{sign}\left(\prod_{1 \leq i \leq s} (\alpha_{r+i} - \overline{\alpha_{r+i}})\right)^2 = (-1)^s.$$

(b) Let  $\alpha_1, \dots, \alpha_m$  be an integral basis of  $\mathcal{O}_K$ . Let  $\sigma_1, \dots, \sigma_m$  be the embeddings of  $K \hookrightarrow \overline{\mathbb{Q}}$ . Then

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) = \det(\sigma_i \alpha_j)^2.$$

Let

$$\begin{aligned} P &= \sum_{\substack{i_1 \dots i_m \\ \text{even permutation}}} (\sigma_{i_1} \alpha_1) \cdots (\sigma_{i_m} \alpha_m), \\ N &= \sum_{\substack{i_1 \dots i_m \\ \text{odd permutation}}} (\sigma_{i_1} \alpha_1) \cdots (\sigma_{i_m} \alpha_m). \end{aligned}$$

Then

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) = P^2 - N^2 = (P + N)^2 - 4PN,$$

where  $P + N$  and  $PN$  are integral over  $\mathbb{Z}$ .

For any  $\tau \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ , either  $\tau P = P, \tau N = N$  or  $\tau P = N, \tau N = P$ . So

$$\tau(P + N) = P + N, \quad \tau(PN) = PN,$$

which implies  $P + N, PN \in \mathbb{Q} \implies \in \mathbb{Z}$ . Then

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) \equiv (P + N)^2 \equiv 0 \text{ or } 1 \pmod{4}.$$

■

**EXAMPLE 1.8.** Consider  $K = \mathbb{Q}(\sqrt{m})$  with  $m \in \mathbb{Z}$  square-free.

Case  $m \equiv 2, 3 \pmod{4}$ . Then

$$D(1, \sqrt{m}) = \text{disc}(x^2 - m) = 4m.$$

By Stickelberger's thm,

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) = 4m,$$

hence  $\{1, \sqrt{m}\}$  is an integral basis.

Case  $m \equiv 1 \pmod{4}$ . The element  $\frac{1 + \sqrt{m}}{2}$  is integral, and

$$D\left(1, \frac{1 + \sqrt{m}}{2}\right) = m.$$

Then  $\left\{1, \frac{1 + \sqrt{m}}{2}\right\}$  is an integral basis.

## 2 Dedekind Domains and Factorization

- Definition of Dedekind domains;
- Ideals in Dedekind domains factor uniquely into products of prime ideals;
- Rings of integers in number fields are Dedekind domains.

### 2.1 Note on 20250924

#### 2.1.1 Discrete valuation rings

**DEFINITION 2.1.** A ring  $A$  is called a discrete valuation ring (DVR) if it is a principal ideal domain which has the following equivalent conditions:

- (a)  $A$  has exactly one non-zero prime ideal  $\mathfrak{m}$ ;
- (b) up to a unit, there exists a unique prime element  $\pi \in A$ ;
- (c)  $A$  is a local ring, and is not a field.

**Proof.** (a)  $\Leftrightarrow$  (b), (c)  $\Rightarrow$  (a).

(c)  $\Rightarrow$  (a). There exists  $(\pi)$  nonzero maximal ideal  $\Rightarrow (\pi) \neq (0)$ . If  $(\pi') \subset (\pi)$  is another nonzero prime ideal, then  $\pi' = \pi \cdot h$ . If  $h \in (\pi')$ , then  $h = \pi' \cdot g$ , then  $\pi' = \pi' \pi g$ ,  $\Rightarrow \pi$  is a unit, which is a contradiction. So  $h \notin (\pi')$ , hence  $(\pi') = (\pi)$ . ■

**EXAMPLE 2.2.**

$$\mathbb{Z}_{(p)} := \left\{ \frac{m}{n} \in \mathbb{Q} \mid n \text{ not divisible by } p \right\}$$

is a DVR with the unique maximal ideal  $\mathfrak{m} = (p)$ .

Recall that any  $A$ -module  $M$  and  $m \in M$ , the annihilator of  $m$  is defined as

$$\text{Ann}(m) := \{a \in A \mid am = 0\},$$

which is an ideal of  $A$ , and proper if  $m \neq 0$ .

**PROPOSITION 2.3.** *An integral domain  $A$  is a DVR iff*

- (a)  $A$  is Noetherian,
- (b)  $A$  is integrally closed,
- (c)  $A$  has exactly one non-zero prime ideal.

**Proof.**  $A$  is a DVR  $\Rightarrow$  (a), (b), (c).

(a)+(b)+(c)  $\Rightarrow A$  is a DVR. (c)  $\Rightarrow A$  is a local ring, not a field. Only need to show  $A$  is a PID. Choose  $c \in A$  with  $c \neq 0$ , not a unit. Consider  $M := A/(c)$ . Pick  $m \in M \setminus \{0\}$  s.t.  $\mathfrak{p} := \text{Ann}(m)$  is maximal. Such  $m$  exists as  $M$  is a f. g.  $A$ -module, and  $A$  is Noetherian. Write  $m = b + (c)$ . Then

$$\mathfrak{p} = \{a \in A : c \mid ab\}.$$



**Claim:**  $\mathfrak{p}$  is a prime ideal.

Otherwise,  $\exists x, y \notin \mathfrak{p}$ , s.t.  $xy \in \mathfrak{p}$ . Then  $yb + (c) \in M \setminus \{0\}$  as  $y \notin \mathfrak{p}$ . But,

$$\text{Ann}(yb + (c)) \supset \text{Ann}(ym) \supsetneq \text{Ann}(m),$$

where the last inequality holds as  $x \in \text{Ann}(ym) \setminus \text{Ann}(m)$ . This contradicts the maximality of  $\mathfrak{p}$ .

As  $m \neq 0$   $c \nmid b$ , i.e.  $\frac{b}{c} \notin A$ .

**Claim:**  $\frac{c}{b} \in A$  and  $\mathfrak{p} = (\frac{c}{b})$ .

$\mathfrak{p} \cdot b \subset (c) \implies \frac{b}{c}\mathfrak{p} \subset A$  is an ideal. If  $\frac{b}{c}\mathfrak{p} \subset \mathfrak{p}$ , then  $\frac{b}{c}$  is integral over  $A$ , hence in  $A$  as  $A$  is integrally closed, which is a contradiction. So  $\frac{b}{c}\mathfrak{p} = A$ , i.e.  $\mathfrak{p} = (\frac{c}{b})$ .

Let  $\pi := \frac{c}{b}$ . Then  $\mathfrak{p} = (\pi)$  is the unique nonzero prime ideal of  $A$ . Let  $\mathfrak{a}$  be a proper of  $A$ . Consider

$$\mathfrak{a} \subset \mathfrak{a}\pi^{-1} \subset \mathfrak{a}\pi^{-2} \subset \dots$$

If  $\mathfrak{a}\pi^{-r} = \mathfrak{a}\pi^{-r-1}$  for some  $r \geq 0$ , then  $\pi^{-1}(\mathfrak{a}\pi^{-r}) = \mathfrak{a}\pi^{-r} \implies \pi^{-1} \in A$ , which is a contradiction. Thus, the sequence is strictly increasing. As  $A$  Noetherian, there exists maximal  $m$  s.t.  $\mathfrak{a}\pi^{-m} \subset A$ ,  $\mathfrak{a}\pi^{-m-1} \not\subset A$ . Then  $\mathfrak{a}\pi^{-m} \not\subset \mathfrak{p} \implies \mathfrak{a} \cdot \pi^{-m} = A \implies \mathfrak{a} = (\pi^m)$ . ■

### 2.1.2 Dedekind domains

**DEFINITION 2.4.** A Dedekind domain is an integral domain  $A$  s.t.

- (a)  $A$  is Noetherian,
- (b)  $A$  is integrally closed,
- (c)  $A$  not a field and every nonzero prime ideal  $\mathfrak{p}$  is maximal.

**REMARK 2.5.** Proposition 2.3  $\implies$   $A$  local domain is a Dedekind domain iff it is a DVR.

**PROPOSITION 2.6.** Let  $A$  be a domain, and  $S$  a multiplicative subset of  $A$ .

- (a) If  $A$  is Noetherian, so is  $S^{-1}A$ ;
- (b) If  $A$  is integrally closed, so also is  $S^{-1}A$ .

**Proof.** Omit. ■

**PROPOSITION 2.7.** Let  $A$  be a Noetherian integral domain. Then  $A$  is a Dedekind domain iff for every nonzero prime ideal  $\mathfrak{p}$  in  $A$ , the localization  $A_{\mathfrak{p}}$  is a DVR.

**Proof.** The “only if” part follows from the above Proposition.

For the “if” part: Only to show  $A$  is integrally closed. Let  $x \in \text{Frac}(A)$  be integral over  $A$ . Set

$$\mathfrak{a} := \{a \in A \mid ax \in A\}.$$

If  $\mathfrak{a} \neq A$ , then there exists a nonzero prime ideal  $\mathfrak{p}$  of  $A$  s.t.  $\mathfrak{a} \subset \mathfrak{p}$ . Since  $A_{\mathfrak{p}}$  is integrally closed,  $x \in A_{\mathfrak{p}}$ . Hence, there exists  $s \in A \setminus \mathfrak{p}$  s.t.  $sx \in A \implies s \in \mathfrak{a} \implies s \in \mathfrak{p}$ , which is a contradiction. Thus,  $\mathfrak{a} = A \implies x \in A$ . ■

## 2.2 Note on 20250929

### 2.2.1 Unique factorization of ideals

**THEOREM 2.8.** *Let  $A$  be a Dedekind domain. Then every nonzero ideal  $\mathfrak{a}$  of  $A$  can be written uniquely (up to ordering) as a product of nonzero prime ideals:*

$$\mathfrak{a} = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_n^{r_n}$$

where the  $\mathfrak{p}_i$  are distinct nonzero prime ideals of  $A$  and  $r_i \geq 1$ .

**LEMMA 2.9.** *Let  $A$  be a Noetherian ring. Then any nonzero ideal  $\mathfrak{a}$  in  $A$  contains a product of nonzero prime ideals.*

**Proof.** Suppose not, choose a maximal counterexample  $\mathfrak{a}$ . Then  $\mathfrak{a}$  is not a prime ideal  $\implies \exists x, y \in \mathfrak{a}$  but  $xy \in \mathfrak{a} \implies$  both  $(x) + \mathfrak{a}$  and  $(y) + \mathfrak{a}$  strictly contain  $\mathfrak{a} \implies$  each of them contains a product of prime ideals  $\implies \mathfrak{a} \supset ((x) + \mathfrak{a}) \cdot ((y) + \mathfrak{a})$  contains a product of prime ideals, which is a contradiction. ■

**LEMMA 2.10.** *Let  $A$  be a ring, and  $\mathfrak{a}, \mathfrak{b}$  be relatively prime ideals on  $A$ , i.e.  $\mathfrak{a} + \mathfrak{b} = A$ . Then for any  $m, n \in \mathbb{N}$ ,  $\mathfrak{a}^m$  and  $\mathfrak{b}^n$  are relatively prime.*

**Proof.** Let  $a \in \mathfrak{a}, b \in \mathfrak{b}$ , s.t.  $a + b = 1$ . Then for  $r \geq m + n$ ,

$$1 = (a + b)^r = \sum_{i=0}^r \binom{r}{i} a^i b^{r-i} \in \mathfrak{a}^m + \mathfrak{b}^n.$$

■

**LEMMA 2.11.** *Let  $\mathfrak{p}$  be a maximal ideal of an integral domain  $A$ , and  $\mathfrak{q} = \mathfrak{p}A_{\mathfrak{p}}$  the maximal ideal of  $A_{\mathfrak{p}}$ . Then the map*

$$\mathfrak{a} + \mathfrak{p}^m \mapsto \mathfrak{a} + \mathfrak{q}^m : A/\mathfrak{p}^m \longrightarrow A_{\mathfrak{p}}/\mathfrak{q}^m$$

*is an isomorphism.*

**Proof.** Injectivity.

Only need to show  $\mathfrak{q} \cap A = \mathfrak{p}^m$ . Clearly  $\mathfrak{p}^m \subset \mathfrak{q}^m \cap A$ . Let  $a \in \mathfrak{q}^m \cap A$ . As  $a \in \mathfrak{q}^m$ , there exists  $s \in A \setminus \mathfrak{p}$  s.t.  $as \in \mathfrak{p}^m \implies s$  is invertible in the field  $A/\mathfrak{p} \implies \exists t \in A$  s.t.  $st = 1 - u$  for  $u \in \mathfrak{p} \implies$

$$st' := st(1 + u + \cdots + u^{m-1}) = 1 - u^m,$$

where  $u^m \in \mathfrak{p}^m$ . Hence,  $a - au^m = ast' \in \mathfrak{p}^m \implies a \in \mathfrak{p}^m$ .

Surjectivity.

Let  $a/s \in A_{\mathfrak{p}}$  with  $a \in A, s \in A \setminus \mathfrak{p}$ . As  $\mathfrak{p}$  maximal,  $(s) + \mathfrak{p} = A \implies (s) + \mathfrak{p}^m = A$  by Lemma 2.10  $\implies \exists b \in A, u \in \mathfrak{p}^m$  s.t.  $sb + u = 1 \implies$

$$\frac{a}{s} = \frac{ab}{sb} = \frac{ab}{1 - u} = ab + \frac{uab}{1 - u}.$$

So  $a/s + \mathfrak{q}^m = ab + \mathfrak{q}^m$ . ■

**Proof of Theorem 2.8.** Now  $A$  is a Dedekind domain.

Existence: We prove that any ideal  $\mathfrak{a}$  of  $A$  can be factored into a product of prime ideals.

By Lemma 2.9, there exists an ideal  $\mathfrak{b} \subset A$  s.t.

$$\mathfrak{b} = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_m^{r_m} \subset \mathfrak{a},$$

where  $\mathfrak{p}_i$  are distinct prime ideals and each two of them are relatively prime. Hence, by Lemma 2.11,

$$A/\mathfrak{b} \simeq A/\mathfrak{p}_1^{r_1} \times \cdots \times A/\mathfrak{p}_m^{r_m} \simeq A_{\mathfrak{p}_1}/\mathfrak{q}_1^{r_1} \times \cdots \times A_{\mathfrak{p}_m}/\mathfrak{q}_m^{r_m},$$

where each  $A_{\mathfrak{p}_i}$  is a DVR  $\implies$  any ideal in  $A_{\mathfrak{p}_i}/\mathfrak{q}_i^{r_i}$  takes form  $\mathfrak{q}_i^{s_i}/\mathfrak{q}_i^{r_i}$ ,  $0 \leq s_i \leq r_i \implies \mathfrak{a}/\mathfrak{b}$  corresponds to

$$\mathfrak{q}_1^{s_1}/\mathfrak{q}_1^{r_1} \times \cdots \times \mathfrak{q}_m^{s_m}/\mathfrak{q}_m^{r_m}.$$

$\implies$

$$\frac{\mathfrak{a}}{\mathfrak{b}} = \frac{\mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m}}{\mathfrak{b}} \implies \mathfrak{a} = \mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m}.$$

Uniqueness.

Suppose

$$\mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m} = \mathfrak{a} = \mathfrak{p}_1^{t_1} \cdots \mathfrak{p}_m^{t_m}$$

(some  $s_i, t_j$  may be 0). Assume  $s_i + t_i \geq 1$ . Take

$$\mathfrak{b} = \mathfrak{p}_1^{s_1+t_1} \cdots \mathfrak{p}_m^{s_m+t_m}.$$

Consider

$$A/\mathfrak{b} \simeq A_{\mathfrak{p}_1}/\mathfrak{q}_1^{s_1+t_1} \times \cdots \times A_{\mathfrak{p}_m}/\mathfrak{q}_m^{s_m+t_m}.$$

$\implies s_i = t_i$  for all  $i$ . ■

**REMARK 2.12.**  $s_i > 0 \iff \mathfrak{a}A_{\mathfrak{p}_i} \neq A_{\mathfrak{p}_i} \iff \mathfrak{a} \subset \mathfrak{p}_i$ .

**COROLLARY 2.13.** Let  $\mathfrak{a}, \mathfrak{b}$  be two ideals in  $A$ . Then  $\mathfrak{a} \subset \mathfrak{b} \iff \mathfrak{a}A_{\mathfrak{p}} \subset \mathfrak{b}A_{\mathfrak{p}}$  for every nonzero prime ideal  $\mathfrak{p}$  of  $A$ .

In particular,  $\mathfrak{a} = \mathfrak{b} \iff \mathfrak{a}A_{\mathfrak{p}} = \mathfrak{b}A_{\mathfrak{p}}$  for all  $\mathfrak{p}$ .

**Proof.** “ $\implies$ ” is clear.

“ $\Leftarrow$ ”: Write  $\mathfrak{a} = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_m^{r_m}$  and  $\mathfrak{b} = \mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m}$ , where  $\mathfrak{p}_i$  distinct,  $s_i, r_i \geq 0$ . For any  $i$ ,  $\mathfrak{a}A_{\mathfrak{p}_i} \subset \mathfrak{b}A_{\mathfrak{p}_i} \implies r_i \geq s_i \implies \mathfrak{a} \subset \mathfrak{b}$ . ■

**COROLLARY 2.14.** Let  $A$  be an integral domain with only finitely many prime ideals. Then  $A$  is a Dedekind domain iff it is a principal ideal domain (PID).

**Proof.** “ $\Leftarrow$ ” is clear.

Assume  $A$  is a Dedekind domain. To show  $A$  is a PID, only need to show prime ideals are principle. Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be these prime ideals. Choose  $x_1 \in \mathfrak{p}_1 \setminus \mathfrak{p}_1^2$ . By Chinese Remainder Theorem,  $\exists x$  s.t.

$$\begin{cases} x \equiv x_1 \pmod{\mathfrak{p}_1^2}; \\ x \equiv 1 \pmod{\mathfrak{p}_i}, \quad \forall i > 1. \end{cases}$$

Then  $(x) = \mathfrak{p}_1$ . ■

**COROLLARY 2.15.** Let  $\mathfrak{a} \supset \mathfrak{b} \neq 0$  be two ideals of a Dedekind domain. Then  $\mathfrak{a} = \mathfrak{b} + (c)$  for some  $c \in A$ .

**Proof.** Write  $\mathfrak{a} = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_m^{r_m}$  and  $\mathfrak{b} = \mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m}$ , where  $\mathfrak{p}_i$  distinct,  $s_i, r_i \geq 0$ .  $\mathfrak{a} \supset \mathfrak{b} \implies s_i \leq r_i$ ,  $\forall i$ . For  $i = 1, \dots, m$ , choose  $x_i \in A$  s.t.

$$x_i \in \mathfrak{p}_i^{s_i} \setminus \mathfrak{p}_i^{s_i+1}.$$

By Chinese Remainder Theorem,  $\exists c \in A$  s.t.

$$c \equiv x_i \pmod{\mathfrak{p}_i^{r_i}}, \quad \forall i.$$

Then  $\mathfrak{b} + (c) = \mathfrak{a}$ . ■

**COROLLARY 2.16.** Let  $\mathfrak{a}$  be an ideal of a Dedekind domain. Let  $a \in \mathfrak{a} \setminus \{0\}$ . Then  $\exists b \in \mathfrak{a}$  s.t.  $\mathfrak{a} = (a, b)$ .

**Proof.** Take  $\mathfrak{b} = (a) \subset \mathfrak{a}$  in the above corollary. ■

**COROLLARY 2.17.** Let  $\mathfrak{a}$  be a nonzero ideal in a Dedekind domain  $A$ . Then there exists a nonzero ideal  $\mathfrak{a}^*$  in  $A$  s.t.  $\mathfrak{a} \cdot \mathfrak{a}^*$  is principal.

Moreover,  $\mathfrak{a}^*$  can be chosen (but not both):

1. to be relatively prime any particular ideal  $\mathfrak{c}$ ; and
2. s.t.  $\mathfrak{a} \cdot \mathfrak{a}^* = (a)$  with any given  $a \in \mathfrak{a}$ .

**Proof.** Let  $a \in \mathfrak{a}$ ,  $a \neq 0$ .  $\mathfrak{a} \supset (a) \implies (a) = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_m^{r_m}$  and  $\mathfrak{a} = \mathfrak{p}_1^{s_1} \cdots \mathfrak{p}_m^{s_m}$ ,  $s_i \leq r_i$ . Take

$$\mathfrak{a}^* = \mathfrak{p}_1^{r_1-s_1} \cdots \mathfrak{p}_m^{r_m-s_m}.$$

Then  $\mathfrak{a} \cdot \mathfrak{a}^* = (a)$ .

Now show that  $\mathfrak{a}^*$  can be chosen relatively prime to  $\mathfrak{c}$ . We have  $\mathfrak{a} \supset \mathfrak{a}\mathfrak{c} \implies \exists a \in \mathfrak{a}$  s.t.

$$\mathfrak{a} = \mathfrak{a}\mathfrak{c} + (a).$$

Above argument  $\implies \exists \mathfrak{a}^*$  s.t.  $(a) = \mathfrak{a} \cdot \mathfrak{a}^* \implies$

$$(a) = \mathfrak{a} \cdot \mathfrak{a}^* = \mathfrak{a} \cdot \mathfrak{c} \cdot \mathfrak{a}^* + (a) \cdot \mathfrak{a}^* = (a) \cdot \mathfrak{c} + (a) \cdot \mathfrak{a}^*,$$

$\implies \exists c \in \mathfrak{c}$ ,  $a' \in \mathfrak{a}^*$  s.t.

$$a = a \cdot c + a \cdot a' \implies (1) = \mathfrak{c} + \mathfrak{a}^*.$$
■

**REMARK 2.18.** We know  $PID \implies UFD$ , but the inverse is not true in general. For example,  $k[x, y]$  is UFD but the ideal  $(x, y)$  is not principle.

**PROPOSITION 2.19.** Let  $A$  be a Dedekind domain. Then  $A$  UFD  $\implies A$  PID.

**Proof.** Only need to show every prime ideal  $\mathfrak{p}$  of  $A$  is principle.

Let  $a \in \mathfrak{p} \setminus \{0\}$ . Then

$$a = u\pi_1^{r_1} \cdots \pi_n^{r_n},$$

for  $u$  unit,  $r_i \geq 1$  and  $\pi_i$  irreducible. Thus, there exists  $\pi_i \in \mathfrak{p}$ . Hence,  $(\pi_i)$  prime  $\implies$  maximal  $\implies (\pi_i) = \mathfrak{p}$ . ■

## 2.3 Note on 20251020

### 2.3.1 Ideal class group

### 3 Discrete valuations

#### 3.1 Note on 20251022

Let  $K$  be a field.

**DEFINITION 3.1.** A discrete valuation on  $K$  is a nonzero homomorphism  $v: K^\times \rightarrow \mathbb{Z}$  s.t.

$$v(a+b) \geq \min\{v(a), v(b)\}.$$

$v$  is called normalized if  $v(K^\times) = \mathbb{Z}$ .

Indeed, if  $v$  is not normalized, write  $v(K^\times) = m\mathbb{Z}$  for some  $m \geq 1$ . Then  $\frac{1}{m}v: K^\times \rightarrow \mathbb{Z}$  is a normalized discrete valuation. We extend  $v$  to a map

$$v: K \rightarrow \mathbb{Z} \cup \{+\infty\}$$

sending 0 to  $+\infty$ .

**EXAMPLE 3.2.** •  $A =$  Dedekind domain,  $\mathfrak{p} \subset A$  a prime ideal in  $A$ .  $K := \text{Frac}(A)$ .  $\forall c \in K^\times$ , define

$$v_{\mathfrak{p}}(c) = \max\{n \in \mathbb{Z} \mid c \in \mathfrak{p}^n\}.$$

Then  $v_{\mathfrak{p}}$  is a normalized discrete valuation on  $K$ .

- In particular, when  $A$  is PID,  $\pi$  a prime element of  $A$ , then for every  $c \in K^\times$ , write  $c = \pi^m \frac{a}{b}$  with  $a, b \in A$  not divisible by  $\pi$ . Define  $v(c) = m$ .
- e.g.  $A = \mathbb{Z}$ ,  $\pi = p$  prime number,  $K = \mathbb{Q}$ . Then for every  $\frac{m}{n} \in \mathbb{Q}^\times$ , write  $\frac{m}{n} = p^r \frac{a}{b}$  with  $a, b$  not divisible by  $p$ . Define  $v_p\left(\frac{m}{n}\right) = r$ .
- e.g.  $A = \mathbb{C}[x]$ ,  $\mathfrak{p} = (t - a)$  for some  $a \in \mathbb{C}$ ,  $K = \mathbb{C}(x)$ . Then for every  $f(x)/g(x) \in K^\times$ , write

$$\frac{f(x)}{g(x)} = (x - a)^r \frac{h(x)}{k(x)}$$

with  $h(a), k(a) \neq 0$ . Define  $v_{\mathfrak{p}}\left(\frac{f(x)}{g(x)}\right) = r$ .

- Let  $U$  be a connected open subset of  $\mathbb{C}$ , and

$$\mathcal{M}(U) := \{\text{meromorphic functions on } U\}.$$

Then  $K = \mathcal{M}(U)$  is a field. For any  $p \in U$ ,  $\forall f \in K^\times$ ,  $\text{ord}_p(f) :=$  vanishing order of  $f$  at  $p$ , i.e.  $f = c \cdot (z - p)^{\text{ord}_p f} + o(z^{\text{ord}_p f})$  with  $c \neq 0$ . Then  $\text{ord}_p$  is a discrete valuation on  $K$ .

**REMARK 3.3.**  $\mathcal{O}(U) := \{\text{holomorphic functions on } U\}$ ,  $\mathcal{M}(U) = \text{Frac}(\mathcal{O}(U))$ . However,  $\mathcal{O}(U)$  is not a Dedekind domain in general.

Fact:  $\exists x_n \in U$ , with  $x_n \rightarrow x \in \partial U$ . For any  $m \geq 0$ ,

$$I_m := \{f \in \mathcal{O}(U) \mid f(x_n) = 0, \forall n \geq m\}.$$

Then  $I_m$  is increasing, and  $\exists h \in \mathcal{O}(U)$  s.t.  $h(x_n) = 0, \forall n, \implies I_m \subsetneq I_{m+1}, \forall m$ . So  $\mathcal{O}(U)$  is not Noetherian.

**LEMMA 3.4.** *If  $v(a) > v(b)$ , then  $v(a + b) = v(b)$ .*

**Proof.**  $v(a + b) \geq \min\{v(a), v(b)\} = v(b)$ . Also,

$$v(b) = v(a + b - a) \geq \min\{v(a + b), v(a)\} = v(a + b).$$

■

**REMARK 3.5.**  $v$  induces a map

$$\begin{aligned} K &\longrightarrow \mathbb{R} \\ a &\longmapsto e^{-v(a)} = |a|_v. \end{aligned}$$

We can check that  $|\cdot|_v$  is a non-archimedean absolute value on  $K$ :

$$|a + b| \leq \max\{|a|, |b|\}.$$

**PROPOSITION 3.6.** *Let  $v$  be a discrete valuation on a field  $K$ . Then*

$$A := \{a \in K \mid v(a) \geq 0\}$$

*is a DVR with the unique maximal ideal*

$$\mathfrak{m} = \{a \in K \mid v(a) > 0\}.$$

$\exists \pi \in A$  s.t.  $\mathfrak{m} = (\pi)$ , and every  $a \in K^\times$  can be written uniquely as  $a = u\pi^{v(a)}$ .

**Proof.** Easy. ■

**PROPOSITION 3.7.** *Let  $A$  be a Dedekind domain. Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be distinct primes ideals of  $A$ , and  $x_1, \dots, x_m \in A$ . Then for any  $n \in \mathbb{Z}_+$ , there exists  $x \in A$  s.t.*

$$\text{ord}_{\mathfrak{p}_i}(x - x_i) > n, \quad i = 1, \dots, m.$$

**Proof.** The ideals  $\mathfrak{p}_i^{n+1}$  are relatively prime two by two. By Chinese Remainder Theorem,  $\exists x \in A$  s.t.

$$x \equiv x_i \pmod{\mathfrak{p}_i^{n+1}}, \quad i = 1, \dots, m.$$

$$\implies \text{ord}_{\mathfrak{p}_i}(x - x_i) > n.$$

■

**THEOREM 3.8.** *Let  $A$  be a Dedekind domain, and  $K = \text{Frac}(A)$ . Let  $L/K$  be a finite separable field extension, and  $B$  the integral closure of  $A$  in  $L$ . Then  $B$  is a Dedekind domain.*

**Proof.** We have proved that  $B$  is Noetherian as an  $A$ -mod. Any ideal of  $B$  is a finitely generated  $A$ -mod  $\implies B$  is Noetherian.  $B$  is integrally closed by definition. We only need to show that every nonzero prime ideal  $\mathfrak{q}$  of  $B$  is maximal.

Let  $\beta \in \mathfrak{q} \setminus \{0\}$ . As  $\beta$  is integral over  $A$ , we have

$$\beta^n + a_1\beta^{n-1} + \dots + a_n = 0, \quad a_i \in A,$$

with minimal degree  $n \implies a_n \neq 0$ . Then  $a_n \in \beta B \cap A \implies \mathfrak{q} \cap A \neq 0$ . Let  $\mathfrak{p} = \mathfrak{q} \cap A$ . Then  $\mathfrak{p}$  is a nonzero prime ideal of  $A \implies \mathfrak{p}$  is maximal  $\implies A/\mathfrak{p}$  is a field.  $B/\mathfrak{q}$  is an integral domain and algebraic over  $A/\mathfrak{p} \implies B/\mathfrak{q}$  is a field  $\implies \mathfrak{q}$  is maximal. ■

**REMARK 3.9.** • *This Theorem shows  $\mathcal{O}_K$  is a Dedekind domain for any number field  $K$ .*

- *In fact, we do not need the full strength of separability.*

**LEMMA 3.10.** *Any integral domain  $B$  containing a field  $k$  and algebraic over  $k$  is itself a field.*

**Proof.** Let  $\beta \in B \setminus \{0\}$ . As  $\beta$  is algebraic over  $k$ ,  $\dim_k k[\beta] < \infty$ . Consider the  $k$ -linear map

$$\begin{aligned} L_\beta: k[\beta] &\longrightarrow k[\beta] \\ x &\longmapsto \beta x. \end{aligned}$$

$L_\beta$  is injective  $\implies$  surjective  $\implies \exists \gamma \in k[\beta] \subset B$  s.t.  $L_\beta(\gamma) = 1 \implies \beta\gamma = 1$ . ■



## 4 Factorization in extensions

Let  $A$  be a Dedekind domain,  $K = \text{Frac}(A)$ , and  $L/K$  a finite separable field extension. Let  $B$  be the integral closure of  $A$  in  $L$ . Let  $\mathfrak{p}$  be a nonzero prime ideal of  $A$ . Then

$$\mathfrak{p}B = \beta_1^{e_1} \cdots \beta_g^{e_g},$$

where  $\beta_i$  are distinct prime ideals of  $B$ , and  $e_i \geq 1$ .

Say  $e_i$  is the ramification index. If some  $e_i > 1$ , then  $\mathfrak{p}$  is ramified over in  $B$  (or  $L$ ); if  $e_i = 1$  for all  $i$ , then  $\mathfrak{p}$  is unramified in  $B$  (or  $L$ ).

Say  $\beta$  divides  $\mathfrak{p}$  (written  $\beta \mid \mathfrak{p}$ ) if  $\beta$  occurs in the factorization of  $\mathfrak{p}$  in  $B$ .

$e(\beta/\mathfrak{p}) :=$  ramification index.

$f(\beta/\mathfrak{p}) := [B/\beta : A/\mathfrak{p}]$  (residue class degree).

A prime  $\mathfrak{p}$  is said to be split (or split completely) in  $L$  if  $e_i = f_i = 1$  for all  $i$ , and it said to be inert in  $L$  if  $\mathfrak{p}B$  is a prime ideal ( $g = 1$  and  $e_1 = 1$ ).

**EXAMPLE 4.1.** •  $(2) = (1+i)^2$  in  $\mathbb{Z}[i]$ , so  $(2)$  ramifies with ramification index 2.

- $(3)$  is inert in  $\mathbb{Q}[i]$  as  $\mathbb{Z}[i]/(3) \simeq \mathbb{F}_9$ .
- $(5) = (2+i)(2-i)$  splits completely in  $\mathbb{Q}[i]$ .

**LEMMA 4.2.** A prime ideal  $\beta$  of  $B$  divides  $\mathfrak{p}$  iff  $\mathfrak{p} = \beta \cap K$ .

**Proof.** “only if”:  $\mathfrak{p} \subset \beta \cap K, \beta \cap K \neq A \implies \mathfrak{p} = \beta \cap K$ .

“if”:  $\mathfrak{p} = \beta \cap K \implies \mathfrak{p}B \subset \beta \implies \beta$  occurs in the factorization of  $\mathfrak{p}B$ . ■

### 4.1 Note on 20251027

**THEOREM 4.3.** Let  $m = [L : K]$ . Let  $\beta_1, \dots, \beta_g$  be the prime ideals dividing  $\mathfrak{p}$ . Then

$$\sum_{i=1}^g e_i f_i = m. \quad (4.1)$$

If  $L$  is Galois over  $K$ , then all ramification numbers  $e_i$  and the residue class degree  $f_i$  are equal  $\implies efg = m$ .

**Proof.** To prove (4.1), we shall show each side equals  $[B/\mathfrak{p}B : A/\mathfrak{p}] = \dim_{A/\mathfrak{p}}(B/\mathfrak{p}B)$ .

First, show  $\sum_{i=1}^g e_i f_i = [B/\mathfrak{p}B : A/\mathfrak{p}]$ . By Chinese Remainder Theorem,

$$B/\mathfrak{p}B = B / \prod_{i=1}^g \beta_i^{e_i} \simeq \prod_{i=1}^g B/\beta_i^{e_i}.$$

Only need to show  $[B/\beta_i^{e_i} : A/\mathfrak{p}] = e_i f_i$ . For any  $r \geq 0$ ,  $0 \neq \beta_i^r / \beta_i^{r+1}$  is a  $B/\beta_i$ -module, and there is no non-trivial submodule. So  $\beta_i^r / \beta_i^{r+1}$  is a one-dimensional  $B/\beta_i$ -vector space  $\implies$

$$\dim_{A/\mathfrak{p}}(\beta_i^r / \beta_i^{r+1}) = \dim_{A/\mathfrak{p}} B/\beta_i = f_i.$$

Consider the chain

$$B \supset \beta_i \supset \beta_i^2 \cdots \supset \beta_i^{e_i}.$$

Each quotient  $\beta_i^r/\beta_i^{r+1}$  has dimension  $f_i$  over  $A/\mathfrak{p} \implies$

$$[B/\beta_i^{e_i} : A/\mathfrak{p}] = \sum_{r=0}^{e_i-1} \dim_{A/\mathfrak{p}}(\beta_i^r/\beta_i^{r+1}) = e_i f_i.$$

Then prove  $[B/\mathfrak{p}B : A/\mathfrak{p}] = m$ . Want to replace  $A, B$  by  $A' = A_{\mathfrak{p}}, B' = B_{\mathfrak{p}}$  respectively. Note that  $K = \text{Frac}(A) = \text{Frac}(A')$ ,  $L = \text{Frac}(B) = \text{Frac}(B')$ , and  $B'$  is the integral closure of  $A'$  in  $L$ . Let  $\mathfrak{p}' = \mathfrak{p}A'$ . Then  $\mathfrak{p}'B' = (\mathfrak{p}B)B' = \beta_1^{e_1} \cdots \beta_g^{e_g}$ , where  $\beta'_i = \beta_i B'$ ,  $B'_i/\beta'_i = B_i/\beta_i$  and  $A/\mathfrak{p}' = A/\mathfrak{p} \implies [B'_i/\beta'_i : A'/\mathfrak{p}'] = f_i$ . So  $[B'/\mathfrak{p}'B' : A'/\mathfrak{p}'] = \sum_{i=1}^g e_i f_i$ .

Only need to show  $[B/\mathfrak{p}B : A/\mathfrak{p}] = m$  when  $A$  is a DVR. In this case  $B$  is a free  $A$ -module  $\implies$  there exists an isomorphism of  $A$ -modules  $A^n \rightarrow B$ . Tensoring with  $K$ , we get an isomorphism of  $K$ -vector spaces  $K^n \rightarrow B \otimes_A K = L \implies n = m$ . We can also show  $[B/\mathfrak{p}B : A/\mathfrak{p}] = n$  by tensoring with  $A/\mathfrak{p} \implies [B/\mathfrak{p}B : A/\mathfrak{p}] = m$ .

Now assume  $L/K$  is Galois. Any  $\sigma \in \text{Gal}(L/K)$  induces  $\sigma : B \rightarrow B$  isomorphism. For any prime ideal  $\beta$  of  $B$  dividing  $\mathfrak{p}$ ,  $\sigma(\beta)$  is also a prime ideal of  $B$  dividing  $\mathfrak{p}$ . As  $\sigma$  is invertible, the map  $\beta \mapsto \sigma(\beta)$  is a bijection on the set of prime ideals of  $B$  dividing  $\mathfrak{p} \implies$  the Galois group  $\text{Gal}(L/K)$  acts transitively on the set of prime ideals of  $B$  dividing  $\mathfrak{p}$ .

Suppose both  $\beta, \beta'$  divide  $\mathfrak{p}$ , and  $\beta'$  is not conjugate to  $\beta$ , i.e.  $\beta' \notin \text{Gal}(L/K)\beta$ . By Chinese Remainder Theorem,  $\exists a \in \beta' \setminus \bigcup_{\sigma \in \text{Gal}(L/K)} \sigma(\beta)$ . Take  $b = \text{Nm}(a) = \prod_{\sigma \in \text{Gal}(L/K)} \sigma(a)$ . Then  $a \in \beta' \implies b \in \beta' \cap A = \mathfrak{p}$ . On the other hand, any  $\sigma \in \text{Gal}(L/K)$ ,  $a \notin \sigma^{-1}(\beta)$  i.e.  $\sigma(a) \notin \beta$  for all  $\sigma \in \text{Gal}(L/K) \implies b = \prod_{\sigma \in \text{Gal}(L/K)} \sigma(a) \notin \beta \cap A = \mathfrak{p}$ . Contradiction! ■

#### 4.1.1 The primes that ramify

**THEOREM 4.4.** *Let  $K$  be a number field,  $L/K$  a finite field extension,  $A \subset K$  a Dedekind domain (e.g.  $A = \mathcal{O}_K$ ), and  $B$  the integral closure of  $A$  in  $L$ . Assume that  $B$  is a free  $A$ -mod (true when  $A$  is a PID). Then a prime  $\mathfrak{p}$  ramifies in  $L$  iff  $\mathfrak{p} \mid \text{disc}(B/A)$ .*

*In particular, only finitely many prime ideals ramify.*

**LEMMA 4.5.** *Let  $A$  be a ring and  $B$  admitting a finite basis  $\{e_1, \dots, e_m\}$  as an  $A$ -mod. For any ideal  $\mathfrak{a}$  of  $A$ ,  $\{\overline{e_1}, \dots, \overline{e_m}\}$  is a basis of  $B/\mathfrak{a}B$  as an  $A/\mathfrak{a}$ -mod, and*

$$D(\overline{e_1}, \dots, \overline{e_m}) \equiv D(e_1, \dots, e_m) \pmod{\mathfrak{a}}.$$

**Proof.** Easy. ■

**LEMMA 4.6.** *Let  $A$  be a ring. Let  $B_1, \dots, B_g$  be rings containing  $A$  and free of finite rank as  $A$ -mods. Then*

$$\text{disc}\left(\left(\prod B_i\right)/A\right) = \prod \text{disc}(B_i/A).$$

**Proof.** Direct computation. ■

We say an element  $\alpha$  in a ring is nilpotent if  $\exists m > 0$  s.t.  $\alpha^m = 0$ .

Fact:  $A = \text{ring}$ :

$$\text{Rad}(A) := \{\alpha \in A \mid \alpha \text{ nilpotent}\} = \bigcap_{\mathfrak{p} \subset A \text{ prime}} \mathfrak{p}.$$

A ring is called reduced if  $\text{Rad}(A) = 0$ . e.g.  $A/\text{Rad}(A)$  is reduced.

Recall that a field  $k$  is called perfect if any finite extension  $K/k$  is separable.

Fact:

1. If  $\text{char } k = 0$ , then  $k$  is perfect.
2. If  $\text{char } k = p > 0$ , then  $k$  is perfect iff  $\forall \alpha \in k, \exists \beta \in k$  s.t.  $\beta^p = \alpha$ .

**LEMMA 4.7.** *Let  $k$  be a perfect field, and  $B$  a finite-dimensional reduced  $k$ -algebra. Then  $B$  is reduced iff  $\text{disc}(B/k) \neq 0$ .*

**Proof.** Let  $\beta \neq 0$  be a nilpotent element in  $B$ . Let  $e_1, \dots, e_m$  be a  $k$ -basis of  $B$  with  $e_1 = \beta$ . Then  $e_1 e_j$  are all nilpotent  $\implies \text{Tr}(e_1 e_j) = 0 \implies$  the first column of the matrix  $(\text{Tr}_{B/k}(e_i e_j))_{1 \leq i, j \leq m}$  is zero  $\implies \text{disc}(B/k) = 0$ . ■

...

## 4.2 Note on 20251103

**PROPOSITION 4.8.** *Let  $f(x) \in A[x]$  be an Eisenstein polynomial w.r.t. a prime ideal  $\mathfrak{p}$  of a Dedekind domain  $A$ . Then  $f(x)$  is irreducible, and if  $\alpha$  is a root of  $f(x)$ , then  $\mathfrak{p}$  is totally ramified in  $K[\alpha]$ . In fact,  $\mathfrak{p}B = \beta^m$  with  $\beta = (\mathfrak{p}, \alpha)$  and  $m = \deg f$ .*

**Proof.**  $L := K(\alpha)$ . Then

$$[L : K] \leq m = \deg f.$$

Let  $\beta$  be a prime ideal of  $B$  dividing  $\mathfrak{p}$  with ramification index  $e$ ,  $e \leq m$ . Consider the equation

$$\alpha^m + a_1\alpha^{m-1} + \cdots + a_m = 0.$$

Then

- $\text{ord}_\beta(\alpha^m) = m \cdot \text{ord}_\beta(\alpha)$ ;
- for  $1 \leq i \leq m-1$ ,

$$\begin{aligned} \text{ord}_\beta(a_i\alpha^{m-i}) &= (m-i) \cdot \text{ord}_\beta(\alpha) + \text{ord}_\beta(a_i) \\ &= (m-i) \cdot \text{ord}_\beta(\alpha) + e \cdot \text{ord}_{\mathfrak{p}}(a_i) \geq (m-i) \cdot \text{ord}_\beta(\alpha) + e; \end{aligned}$$

- $\text{ord}_\beta(a_m) = e \cdot \text{ord}_{\mathfrak{p}}(a_m) = e$ .

If  $\text{ord}_\beta(\alpha) = 0$ , then  $0 = \text{ord}_\beta(\alpha^m) < \text{ord}_\beta(\text{other terms})$ , contradiction. So  $\text{ord}_\beta(\alpha) \geq 1$ . For all  $i = 1, \dots, m-1$ ,

$$\text{ord}_\beta(a_i\alpha^{m-i}) \geq (m-i) \cdot \text{ord}_\beta(\alpha) + e > e.$$

$$\implies \text{ord}_\beta(\alpha^m) = m \cdot \text{ord}_\beta(\alpha) = \text{ord}_\beta(a_m) = e \implies e = m \text{ and } \text{ord}_\beta(\alpha) = 1 \text{ since } e \leq m. \quad \blacksquare$$

## 5 The finiteness of the class number

### 5.1 Note on 20251103

#### 5.1.1 Norms of ideals

Let  $A$  be a Dedekind domain,  $K = \text{Frac}(A)$ , and  $L/K$  a finite separable field extension. Let  $B$  be the integral closure of  $A$  in  $L$ . We want to define a homomorphism

$$\text{Nm}_{B/A}: \text{Id}(B) \longrightarrow \text{Id}(A),$$

makes the following diagram commutes:

$$\begin{array}{ccc} L^\times & \xrightarrow{b \mapsto (b)} & \text{Id}(B) \\ \downarrow \text{Nm} & & \downarrow \text{Nm} \\ K^\times & \longrightarrow & \text{Id}(A) \end{array}$$

$\text{Id}(B)$  is the free abelian group on the set of prime ideals. Only need to define  $\text{Nm}(\mathfrak{p})$  for  $\mathfrak{p}$  prime.

Let  $\mathfrak{p}$  be a prime ideal of  $A$ ,

$$\mathfrak{p}B = \prod_i \beta_i^{e_i}.$$

If  $\mathfrak{p}$  is principle, say  $\mathfrak{p} = (\pi)$ , then

$$\text{Nm}(\mathfrak{p}B) = \text{Nm}(\pi B) = \text{Nm}(\pi) \cdot A = (\pi^m) = \mathfrak{p}^m, \quad m = [L : K]. \quad (5.2)$$

Generally,

$$\text{Nm}(\mathfrak{p}B) = \text{Nm}\left(\prod_i \beta_i^{e_i}\right) = \prod_i \text{Nm}(\beta_i)^{e_i}. \quad (5.3)$$

Compare (5.2) and (5.3). Recall  $\sum_i e_i f_i = m$ . So we can define

$$\text{Nm}(\beta_i) = \mathfrak{p}^{f_i}.$$

Then (5.3) holds.

We take this as our definition:

**DEFINITION 5.1.** Let  $\beta$  be a prime ideal of  $B$ . The norm of  $\beta$  is defined as

$$\text{Nm}(\beta) := \mathfrak{p}^{f(\beta/\mathfrak{p})}, \quad \text{where } \mathfrak{p} := \beta \cap A, \quad f(\beta/\mathfrak{p}) = [B/\beta : A/\mathfrak{p}].$$

To avoid confusion, we also use  $\mathcal{N}$  to denote norms of ideals. If we have a tower of fields  $M \supset L \supset K$ , then

$$\mathcal{N}_{L/K}(\mathcal{N}_{M/L}\mathfrak{a}) = \mathcal{N}_{M/K}\mathfrak{a}.$$

**PROPOSITION 5.2.** 1. Any nonzero ideal  $\mathfrak{a}$  of a Dedekind domain  $A$ ,

$$\mathcal{N}_{L/K}(\mathfrak{a}B) = \mathfrak{a}^m, \quad m = [L : K].$$

2. Suppose  $L/K$  is Galois. Let  $\beta$  be a nonzero prime ideal of  $B$  with  $\mathfrak{p} = \beta \cap A$ . Write  $\mathfrak{p}B = (\beta_1 \cdots \beta_g)^e$ . Then

$$\mathcal{N}(\beta) \cdot B = (\beta_1 \cdots \beta_g)^{ef} = \prod_{\sigma \in \text{Gal}(L/K)} \sigma(\beta).$$

3. Any nonzero  $b \in B$ ,

$$\mathcal{N}(bB) = (\text{Nm}_{L/K}(b))A.$$

**Proof.** 1.  $\sum_i e_i f_i = m$ ;

2.  $efg = m$ ;

3. First, treat the case  $L/K$  Galois. Let  $\mathfrak{b} = bB$ . Then the map

$$\begin{aligned} \text{Id}(A) &\longrightarrow \text{Id}(B) \\ \mathfrak{a} &\longmapsto \mathfrak{a}B \end{aligned}$$

is injective (by 1). Only need to show  $\text{Nm}(\mathfrak{b}) \cdot B = \mathcal{N}(\mathfrak{b}) \cdot B$ . By 2,

$$\mathcal{N}(\mathfrak{b}) \cdot B = \prod \sigma(bB) = \prod \sigma(b) \cdot B = \text{Nm}(\mathfrak{b}) \cdot B.$$

In the general case, let  $E/K$  be a finite Galois extension s.t.  $E \supset L$ .  $d = [E : L]$ . Then we have

$$\begin{aligned} \mathcal{N}_{L/K}(bB)^d &= \mathcal{N}_{L/K}(\mathcal{N}_{E/L}(bC)) = \mathcal{N}_{E/K}(bC) = \text{Nm}_{E/K}(b)A \\ &= \text{Nm}_{L/K}(\text{Nm}_{E/L}(b))A = \text{Nm}_{L/K}(b^d)A = (\text{Nm}_{L/K}(b)A)^d. \end{aligned}$$

$$\implies \mathcal{N}_{L/K}(bB) = \text{Nm}_{L/K}(b)A.$$

■

Now assume  $K$  is a number field. Every  $\mathfrak{a}$  nonzero ideal of  $\mathcal{O}_K$  is of finite index in  $\mathcal{O}_K$ , i.e.  $\sharp(\mathcal{O}_K/\mathfrak{a}) < \infty$ .

**DEFINITION 5.3.** Define  $\mathcal{N}\mathfrak{a} := (\mathcal{O}_K : \mathfrak{a}) = \sharp(\mathcal{O}_K/\mathfrak{a}) \in \mathbb{Z}_{\geq 1}$ , and call it numerical norm of  $\mathfrak{a}$ .

**PROPOSITION 5.4.** 1. For any ideal  $\mathfrak{a}$  in  $\mathcal{O}_K$ ,

$$\mathcal{N}_{K/\mathbb{Q}}(\mathfrak{a}) = (\mathcal{N}(\mathfrak{a})),$$

in particular,  $\mathcal{N}(\mathfrak{a}\mathfrak{b}) = \mathcal{N}(\mathfrak{a})\mathcal{N}(\mathfrak{b})$ .

2. Let  $\mathfrak{b} \subset \mathfrak{a}$  be fractional ideals in  $K$ , then

$$(\mathfrak{a} : \mathfrak{b}) = \mathcal{N}(\mathfrak{a}^{-1}\mathfrak{b}).$$

**Proof.** 1. Write  $\mathfrak{a} = \prod \beta_i^{r_i}$ ,  $f_i = f(\beta_i/p_i)$ ,  $(p_i) = \beta_i \cap \mathbb{Z}$ , where  $p_i$  are prime numbers. Then  $\text{Nm}(\beta_i) = (p_i)^{f_i}$ , and

$$\mathcal{O}_K/\mathfrak{a} \simeq \prod \mathcal{O}_K/\beta_i^{r_i},$$

where

$$\sharp(\mathcal{O}_K/\beta_i^{\gamma_i}) = \prod_{0 \leq s \leq \gamma_i - 1} \sharp(\beta_i^s/\beta_i^{s+1}) = \left( \sharp(\mathcal{O}_K/\beta_i) \right)^{r_i},$$

and  $[\mathcal{O}_K/\beta_i : \mathbb{Z}/(p_i)] = f_i \implies \#(\mathcal{O}_K/\beta_i) = p_i^{f_i}$ . Hence,  $\mathcal{N}(\mathfrak{a}) = \#(\mathcal{O}_K/\mathfrak{a}) = \prod p_i^{f_i r_i} \implies$   

$$(\mathcal{N}(\mathfrak{a})) = \prod (p_i)^{f_i r_i} = \mathcal{N}_{K/\mathbb{Q}}(\mathfrak{a}).$$

2. Any  $d \in K^*$ , the map

$$\begin{aligned} (K, +) &\longrightarrow (K, +) \\ x &\longmapsto d \cdot x \end{aligned}$$

is an isomorphism  $\implies (d\mathfrak{a} : a\mathfrak{b}) = (\mathfrak{a} : \mathfrak{b})$ . Since  $\mathfrak{a}^{-1}\mathfrak{b} = (d\mathfrak{a})^{-1}(a\mathfrak{b})$ , may assume  $\mathfrak{a}, \mathfrak{b} \subset \mathcal{O}_K$ .  
 $(\mathfrak{a}^{-1}\mathfrak{b})\mathfrak{a} = \mathfrak{b} \implies \mathcal{N}(\mathfrak{a}^{-1}\mathfrak{b})\mathcal{N}(\mathfrak{a}) = \mathcal{N}(\mathfrak{b})$ . Also,  $(\mathcal{O}_K : \mathfrak{a})(\mathfrak{a} : \mathfrak{b}) = (\mathcal{O}_K : \mathfrak{b})$ . Hence,  
 $(\mathfrak{a} : \mathfrak{b}) = \mathcal{N}(\mathfrak{a}^{-1}\mathfrak{b})$ . ■

**THEOREM 5.5.** *Let  $K$  be a number field with  $[K : \mathbb{Q}] = n$ . Let  $\Delta_K = \text{disc}(K/\mathbb{Q})$ , and*

$$2s := \# \{ \text{non-real complex embedding of } K \}.$$

*Then there exists a set of representatives for the ideal class group of  $K$  consisting of integral ideals  $\mathfrak{a}$  with*

$$\mathcal{N}(\mathfrak{a}) \leq \frac{n!}{n^n} \left( \frac{4}{\pi} \right)^s \sqrt{|\Delta_K|}.$$

The bound is called the Minkowski bound (denoted by  $B_K$ ), and the Minkowski constant

$$C_K := \frac{n!}{n^n} \left( \frac{4}{\pi} \right)^s.$$

Note that  $(2s \leq n)$

$$C_K \approx \sqrt{2\pi n} \frac{1}{e^n} \left( \frac{4}{\pi} \right)^s \leq \sqrt{2\pi n} \left( \frac{2}{e\sqrt{\pi}} \right)^n \rightarrow 0, \quad n \rightarrow \infty.$$

**THEOREM 5.6.** *The class group number of  $K$  is finite.*

**Proof.** Only need to show for any  $M > 0$ , the set

$$\# \{ \mathfrak{a} \mid \text{integral ideal of } \mathcal{O}_K \text{ with } \mathcal{N}(\mathfrak{a}) < M \} < \infty.$$

If  $\mathfrak{a} = \prod \beta_i^{r_i}$ , then

$$\mathcal{N}(\mathfrak{a}) = \prod p_i^{r_i f_i}, \quad (p_i) = \beta_i \cap \mathbb{Z},$$

where  $p_i > 0$  prime.  $\mathcal{N}(\mathfrak{a}) < M \implies p_i < M$  for all  $i \implies$  finitely many  $\beta_i$ ,  $r_i \leq n$ ,  $f_i \leq n$ . ■

## 5.2 Note on 20251105

Let  $S := \{ \text{integral ideals in } K \text{ with norm } < B_K \}$ , which is finite, and  $\text{Cl}(\mathcal{O}_K) = S / \sim$ , where  $\mathfrak{a} \sim \mathfrak{b}$  if  $\mathfrak{a} \cdot \mathfrak{b}^{-1}$  is principal.

To decide whether  $\mathfrak{a} \sim \mathfrak{b}$ , only need to decide whether  $\mathfrak{c} := \mathfrak{a} \cdot \mathfrak{b}^{-1}$  is principal. If  $\mathfrak{c} = (\gamma)$ , then

$$\mathcal{N}(\mathfrak{c}) = |\text{Nm}(\gamma)|,$$

which is a diophantine equation (has algorithm to solve). Fix a  $\mathbb{Z}$ -basis of  $\mathcal{O}_K$ ,  $e_1, \dots, e_m$ , and  $\gamma = \sum x_i e_i$ .

**EXAMPLE 5.7.**  $K = \mathbb{Q}(\sqrt{-5})$ . Then  $\text{Cl}(\mathcal{O}_K)$  is generated by  $\mathfrak{a}$  with  $\mathcal{N}(\mathfrak{a}) \leq 0.63 \times \sqrt{20} < 3$ .  $\Delta_K = 20$ .

For  $\mathfrak{a} \neq \mathcal{O}_K$ , we have  $\mathfrak{a} \mid (2)$ .  $(2) = \mathfrak{p}^2$ , where  $\mathfrak{p} = (2, 1 + \sqrt{-5})$ .  $\mathcal{N}(\mathfrak{p}^2) = \mathcal{N}(2) = 4 \implies \mathcal{N}(\mathfrak{p}) = 2$ . If  $\mathfrak{p}$  is principal, then  $\mathfrak{p} = (\alpha) = (m + n\sqrt{-5})$ ,  $2 = \mathcal{N}(\mathfrak{p}) = \mathcal{N}(\alpha) = m^2 + 5n^2$ . No solution.  $\implies \text{Cl}(\mathcal{O}_K)$  has order 2.

**DEFINITION 5.8.** An extension  $L$  of a number field  $K$  is said to be unramified over  $K$  if no prime ideal of  $\mathcal{O}_K$  ramified in  $\mathcal{O}_L$ .

**THEOREM 5.9.** No unramified extension of  $\mathbb{Q}$ .

**Proof.** Let  $K/\mathbb{Q}$  be a finite extension of  $\mathbb{Q}$ . Since a set of representatives for  $\text{Cl}(K) \geq 1$ , and it has numerical norm  $\geq 1$ . Theorem 5.5  $\implies$

$$|\Delta|^{1/2} \geq \frac{n^n}{n!} \left(\frac{\pi}{4}\right)^s \geq \frac{n^n}{n!} \left(\frac{\pi}{4}\right)^{n/2} =: a_n.$$

Then

- $a_2 > 1$ ;
- $\frac{a_{n+1}}{a_n} = \left(\frac{\pi}{4}\right)^{1/2} \left(1 + \frac{1}{n}\right)^n > 1$ .

$\implies a_n > 1, \forall n \implies |\Delta| > 1 \implies K$  can not be unramified. ■

**COROLLARY 5.10.** No irreducible monic polynomial  $f(x) \in \mathbb{Z}[x]$  of degree  $> 1$  with discriminant  $= 1$ .

**Proof.** Let  $f$  be such polynomial. Let  $\alpha$  be a root of  $f(x)$ . Then  $\text{disc}(\mathbb{Z}[\alpha]/\mathbb{Z}) = \pm 1 \implies \mathbb{Z}[\alpha]$  is the ring of integers of  $K = \mathbb{Q}[\alpha] \implies \text{disc}(\mathcal{O}_K/\mathbb{Z}) = \pm 1$ . Contradiction with Theorem 5.9. ■



### 5.2.1 Lattices

Let  $V$  be a  $\mathbb{Q}$ -vector space,  $\dim V = n$ .

**DEFINITION 5.11.** A lattice  $\Lambda$  in  $V$  is a subgroup of the form

$$\Lambda = \mathbb{Z}e_1 + \cdots + \mathbb{Z}e_r$$

with  $e_1, \dots, e_r$  linearly independent elements on  $V$  (for  $\mathbb{R}$ ).

Fact:  $r \leq n$ .

When  $r = n$ , the lattice is said to be full.

A full lattice in  $V$  is a subgroup  $\lambda$  for  $V$  s.t. the map

$$\begin{aligned} \mathbb{R} \otimes \Lambda &\longrightarrow V \\ \sum r_i \otimes x_i &\longmapsto \sum r_i x_i \end{aligned}$$

is an isomorphism.

**EXAMPLE 5.12.** The group  $\mathbb{Z} + \mathbb{Z}\sqrt{2}$  of  $\mathbb{R}$  is a free abelian group of rank 2, but it is not a lattice in  $\mathbb{R}$ .

**LEMMA 5.13.** *The following conditions on a subgroup  $\Lambda$  of  $V$  are equivalent:*

1.  $\Lambda$  is a discrete group;
2.  $\exists$  an open subset  $U$  of  $V$  s.t.  $U \cap \Lambda = \{o\}$ ;
3.  $\forall$  compact subset  $K$  of  $V$ ,  $\sharp(K \cap \Lambda) < \infty$ ;
4.  $\forall$  bounded subset  $B$  of  $V$ ,  $\sharp(B \cap \Lambda) < \infty$ .

**Proof.** Easy. ■

**PROPOSITION 5.14.** *A subgroup  $\Lambda$  of  $V$  is a lattice if and only if it is discrete.*

**Proof.** “Only if”:  $\Lambda = \sum \mathbb{Z}e_i$ ,  $i = 1, \dots, r$ ,  $e_i$  linearly independent. Extend  $e_1, \dots, e_r$  to a basis  $e_1, \dots, e_n$  of  $V$ . Assume  $r = n$ . Then  $U := \sum_{i=1}^n (-1/2, 1/2)e_i$  is open in  $V$  and  $U \cap \Lambda = \{o\}$ .

“If”: Replace  $V$  by the  $\mathbb{R}$ -subspace generated by  $\Lambda$ . We may assume  $V$  is generated by  $\Lambda$  (over  $\mathbb{R}$ ).  $\exists e_1, \dots, e_n \in \Lambda$  forms a  $\mathbb{R}$ -basis of  $V$ .

$$\Lambda' := \sum_{i=1}^n \mathbb{Z}e_i < \Lambda,$$

and  $V/\Lambda' \simeq (\mathbb{R}/\mathbb{Z})^n$  compact. Let  $K := \sum_{i=1}^n [0, 1)e_i$  bounded, and  $K \rightarrow V/\Lambda'$  bijective. Moreover,  $K \cap \Lambda \rightarrow \Lambda/\Lambda'$  bijection.  $\Lambda$  discrete  $\implies K \cap V$  finite  $\implies \Lambda/\Lambda'$  finite  $\implies \exists N \in \mathbb{Z}_{\geq 1}$  s.t.  $N \cdot \Lambda/\Lambda' = 0 \implies \Lambda \subset \frac{1}{N}\Lambda' \implies \Lambda$  is free abelian of rank  $n$ . Then for  $f_1, \dots, f_n$  a  $\mathbb{Z}$ -basis of  $\Lambda$ , they are linearly independent over  $\mathbb{R}$ . ■

**DEFINITION 5.15.** Let  $V$  be a  $\mathbb{R}$ -vector space,  $\dim V = n$ , and  $\Lambda$  a full lattice in  $V$ . Write  $\Lambda = \sum \mathbb{Z}e_i$ .

For any  $\lambda_0 \in \Lambda$ , let

$$D_{(\lambda_0)} := \{\lambda_0 + \sum a_i e_i \mid 0 \leq a_i < 1\}.$$

Such a set is called a fundamental parallelopiped for  $\Lambda$ .

Its shape depends on the choice of the basis  $(e_i)$ . Fix  $e_i$ ,

$$V = \sum_{\lambda \in \Lambda} D_\lambda.$$

**REMARK 5.16.** 1.  $\forall$  FP  $D$  of  $\Lambda = \mathbb{Z}f_1 + \cdots + \mathbb{Z}f_n$  in  $\mathbb{R}^n$ ,

$$\text{Vol}(D) = |\det(f_1, \dots, f_n)|.$$

If also  $\Lambda = \mathbb{Z}f'_1 + \cdots + \mathbb{Z}f'_n$ , then  $\det(f'_1, \dots, f'_n) = \pm \det(f_1, \dots, f_n)$ . So  $\text{Vol}(D)$  does not depend on the choice of the basis for  $\Lambda$ .

2. When  $\Lambda \supset \Lambda'$  are two full lattices in  $\mathbb{R}^n$ , we can choose the bases  $(e_i)$  and  $(f_i)$  for  $\Lambda$  and  $\Lambda'$  s.t.  $f_i = m_i e_i$ , where  $m_i \in \mathbb{Z}_{\geq 1}$ . With the choice of basis, the FP  $D'$  of  $\Lambda'$  is a disjoint union of  $(\Lambda : \Lambda')$  FP  $D$  of  $\Lambda$ . Hence,

$$\frac{\mu(D')}{\mu(D)} = (\Lambda : \Lambda'). \quad (5.4)$$

The choice of a basis for  $V$  determines an isomorphism  $V \simeq \mathbb{R}^n$ , hence a measure  $\mu$  on  $V$ .  $\mu$  is invariant under translations  $\implies \mu$  is well defined up to multiplication by a nonzero constant.

Thus the ratio of measures of two sets is well defined. The equality (5.4) holds for two full lattices  $\Lambda \supset \Lambda'$  in  $V$ .

**THEOREM 5.17.** Let  $D_0$  be a FP for a full lattice  $V$ , and  $S$  a measurable subset of  $V$ . If  $\mu(S) > \mu(D_0)$ , then  $S$  contains distinct points  $\alpha$  and  $\beta$  s.t.  $\beta - \alpha \in \Lambda$ .

**Proof.**

$$\mu(D_0) < \mu(S) = \sum_{\lambda \in \Lambda} \mu(S \cap D_\lambda) = \sum_{\lambda \in \Lambda} \mu((S - \lambda) \cap D_0),$$

$\implies$  there exists  $\lambda_1, \lambda_2 \in \Lambda$  distinct s.t.

$$[(S - \lambda_0) \cap D_0] \cap [(S - \lambda_2) \cap D_0] \neq \emptyset \implies D_0 \cap [(\lambda_1 - \lambda_2) + S] \neq \emptyset.$$

Pick  $\alpha$  in the above set, then  $\beta = \alpha - (\lambda_1 - \lambda_2) \in S$ ,  $\alpha - \beta \in \Lambda$ . ■

### 5.3 Note on 20251110

Let  $T \subset V$ . We say  $T$  is symmetric in the origin if  $\alpha \in T \implies -\alpha \in T$ .

If  $T$  is convex and symmetric in the origin, we have:

$$\forall \alpha, \beta \in T \implies \frac{\alpha - \beta}{2} \in T. \quad (5.5)$$

**LEMMA 5.18.** Assume  $T$  satisfies (5.5) and  $\mu(T) > 2^n \mu(D)$ . Then  $T \cap (\Lambda \setminus \{0\}) \neq \emptyset$ .

**Proof.** Let  $S = \frac{1}{2}T$ . Then  $\mu(S) > \mu(D)$ . Theorem 5.17  $\implies \exists \alpha, \beta \in S, \alpha \neq \beta$  s.t.  $\alpha - \beta \in \Lambda$ . By (5.5),  $\alpha - \beta = \frac{2\alpha - 2\beta}{2} \in T$ . Hence,  $T \cap (\Lambda \setminus \{0\}) \neq \emptyset$ . ■

**THEOREM 5.19 (Minkowski; 1896).** Let  $T$  be a subset of  $V$ , that is compact, convex, and symmetric in the origin. If  $\mu(T) \geq 2^n \mu(D)$ , then  $T$  contains a point of the lattice other than the origin.

**Proof.** Let  $\epsilon > 0$ . Since  $T$  is compact, for  $T_\epsilon = (1+\epsilon)T$ , we have  $T \subset S$  and  $\mu(T_\epsilon) = (1+\epsilon)^n \mu(T) > 2^n \mu(D)$ . By the previous lemma,  $T_\epsilon \cap (\Lambda \setminus \{0\}) \neq \emptyset \implies T \cap (\Lambda \setminus \{0\}) \neq \emptyset$  by the compactness of  $T$ . ■

**REMARK 5.20.** Theorem 5.5 has many non-trivial consequences. It was the starting point for “geometry of numbers”.

**COROLLARY 5.21.** Any  $n \in \mathbb{Z}_{\geq 0}$  is a sum of four squares.

**Proof.** From the identity

$$\begin{aligned} & (a_1^2 + a_2^2 + a_3^2 + a_4^2)(b_1^2 + b_2^2 + b_3^2 + b_4^2) \\ &= (a_1b_1 - a_2b_2 - a_3b_3 - a_4b_4)^2 + (a_1b_2 + a_2b_1 + a_3b_4 - a_4b_3)^2 \\ & \quad + (a_1b_3 - a_2b_4 + a_3b_1 + a_4b_2)^2 + (a_1b_4 + a_2b_3 - a_3b_2 + a_4b_1)^2, \end{aligned}$$

we only need to show that any prime number  $p$  is a sum of four squares.  $2 = 1^2 + 1^2 + 0^2 + 0^2$ . For odd prime  $p$ :

**Claim:**  $m^2 + n^2 + 1 \equiv 0 \pmod{p}$  has a solution  $(m, n) \in \mathbb{Z}^2$ .

**Proof of the claim.** Consider the sets

$$A = \{m^2 \mid m = 0, 1, \dots, (p-1)/2\}, \quad B = \{-n^2 - 1 \mid n = 0, 1, \dots, (p-1)/2\}.$$

Then  $\#A = \#B = (p+1)/2 \implies A \cap B \neq \emptyset \implies \exists m, n$  s.t.  $m^2 \equiv -n^2 - 1 \pmod{p}$ . ■

Fix a solution  $(m, n)$  of the claim. Consider the lattice  $\Lambda \subset \mathbb{Z}^4$  consisting of points  $(a_1, a_2, a_3, a_4)$  satisfying

$$a_1 \equiv ma_3 + na_4 \pmod{p}, \quad a_2 \equiv na_3 - ma_4 \pmod{p}.$$

Then  $\Lambda$  is of index  $p^2$  in  $\mathbb{Z}^4$ . Let

$$T = \{(a_1, a_2, a_3, a_4) \in \mathbb{R}^4 \mid a_1^2 + a_2^2 + a_3^2 + a_4^2 < r^2\}.$$

Then  $\mu(T) = \pi^2 r^4 / 2$ . A FP  $D$  of  $\Lambda$  has measure  $\mu(D) = p^2$ . Let  $2p > r^2 > 1.9p$ . Then  $\mu(T) > 16\mu(D)$ . Theorem 5.19  $\implies$   $T$  contains a point of  $\Lambda$  other than the origin, say  $(a_1, a_2, a_3, a_4)$ . Then

$$a_1^2 + a_2^2 + a_3^2 + a_4^2 \equiv 0 \pmod{p}, \quad a_1^2 + a_2^2 + a_3^2 + a_4^2 < 2p \implies a_1^2 + a_2^2 + a_3^2 + a_4^2 = p.$$

■

### 5.3.1 Finiteness of the class number

Let  $K$  be a number field,  $[K : \mathbb{Q}] = n$ . Let  $r$  be the number of real embeddings  $\{\sigma_1, \dots, \sigma_r\}$  of  $K$ , and  $2s$  the number of non-real complex embeddings  $\{\sigma_{r+1}, \overline{\sigma_{r+1}}, \dots, \sigma_{r+s}, \overline{\sigma_{r+s}}\}$ . Then  $n = r + 2s$ .

We have an embedding

$$\begin{aligned} \sigma : K &\longrightarrow \mathbb{R}^r \times \mathbb{C}^s \simeq \mathbb{R}^{r+2s} = \mathbb{R}^n = V \\ \alpha &\longmapsto (\sigma_1(\alpha), \dots, \sigma_r(\alpha), \sigma_{r+1}(\alpha), \dots, \sigma_{r+s}(\alpha)). \end{aligned}$$

Metric on  $V$ :

$$\|(x_1, \dots, x_r, z_1, \dots, z_s)\| := |x_1|^2 + \dots + |x_r|^2 + 2(|z_1|^2 + \dots + |z_s|^2).$$

**PROPOSITION 5.22.** *Let  $\mathfrak{a}$  be a nonzero ideal of  $\mathcal{O}_K$ . Then  $\sigma(\mathfrak{a})$  is a full lattice in  $V$  with fundamental parallelepiped  $D$  satisfying*

$$\mu(D) = 2^{-s} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a}).$$

**Proof.** Let  $\alpha_1, \dots, \alpha_n$  be a basis of  $\mathfrak{a}$  as a  $\mathbb{Z}$ -module. To prove  $\sigma(\mathfrak{a})$  is a lattice, we show that the matrix  $A =$

$$\begin{pmatrix} \sigma_1(\alpha_1) & \dots & \sigma_1(\alpha_1) & \operatorname{Re}(\sigma_{r+1}(\alpha_1)) & \dots & \operatorname{Re}(\sigma_{r+1}(\alpha_1)) & \operatorname{Im}(\sigma_{r+1}(\alpha_1)) & \dots & \operatorname{Im}(\sigma_{r+1}(\alpha_1)) \\ \vdots & & \vdots & \vdots & & \vdots & \vdots & & \vdots \\ \sigma_r(\alpha_n) & \dots & \sigma_r(\alpha_n) & \operatorname{Re}(\sigma_{r+s}(\alpha_n)) & \dots & \operatorname{Re}(\sigma_{r+s}(\alpha_n)) & \operatorname{Im}(\sigma_{r+s}(\alpha_n)) & \dots & \operatorname{Im}(\sigma_{r+s}(\alpha_n)) \end{pmatrix}$$

has nonzero determinant. Let the matrix  $B =$

$$\begin{pmatrix} \sigma_1(\alpha_1) & \dots & \sigma_1(\alpha_1) & \sigma_{r+1}(\alpha_1) & \overline{\sigma_{r+1}(\alpha_1)} & \dots & \sigma_{r+1}(\alpha_s) & \overline{\sigma_{r+s}(\alpha_1)} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots & \vdots \\ \sigma_1(\alpha_n) & \dots & \sigma_1(\alpha_n) & \sigma_{r+1}(\alpha_n) & \overline{\sigma_{r+1}(\alpha_n)} & \dots & \sigma_{r+1}(\alpha_n) & \overline{\sigma_{r+s}(\alpha_n)} \end{pmatrix}.$$

Then  $\det(B) = \det(A)(\det(1, 1; i, -i))^s \implies$

$$\det(A) = (-2i)^{-s} \det(B) = \pm(-2i)^{-s} D(\alpha_1, \dots, \alpha_n)^{1/2} \neq 0.$$

Moreover,

$$\begin{aligned} \mu(D) &= |\det(A)| = 2^{-s} |D(\alpha_1, \dots, \alpha_n)|^{1/2} \\ &= 2^{-s} \sqrt{|\Delta_K|} \cdot [\mathcal{O}_K : \mathfrak{a}] \\ &= 2^{-s} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a}). \end{aligned}$$

■

**PROPOSITION 5.23.** *Let  $\mathfrak{a}$  be a nonzero ideal of  $\mathcal{O}_K$ . Then there exists a nonzero element  $\alpha \in \mathfrak{a}$  s.t.*

$$|\mathrm{Nm}_{K/\mathbb{Q}}(\alpha)| \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a}).$$

**Proof.** Let

$$X_t := \{x \in V \mid \|x\| \leq t\},$$

which is compact, convex, and symmetric in the origin. Choose  $t$  s.t.  $\mu(X_t) = 2^n \mu(D)$ . Then

$$\begin{aligned} t^n \frac{\pi^s}{2^{r+s} n!} &= 2^n \cdot 2^{-s} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a}) \\ \implies t &= \left(\frac{4}{\pi}\right)^{s/n} \frac{n!^{1/n}}{n} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a})^{1/n}. \end{aligned}$$

By Theorem 5.19,  $\exists \alpha \in \mathfrak{a}$ ,  $\alpha \neq 0$  s.t.  $\sigma(\alpha) \in X_t$ . Then

$$\begin{aligned} |\mathrm{Nm}_{K/\mathbb{Q}}(\alpha)| &= \prod_{i=1}^r |\sigma_i(\alpha)| \cdot \prod_{j=1}^s |\sigma_{r+j}(\alpha)|^2 \\ &\leq \left(\frac{\|\sigma(\alpha)\|}{n}\right)^n \quad (\text{by AM-GM inequality}) \\ &\leq \left(\frac{t}{n}\right)^n = \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{a}). \end{aligned}$$

■

**THEOREM 5.24.** *There exists a set of representatives for the ideal class group of  $K$  consisting of integral ideals  $\mathfrak{a}$  with*

$$\mathcal{N}(\mathfrak{a}) \leq \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{|\Delta_K|}.$$

**Proof.** Let  $\mathfrak{c}$  be a fractional ideal of  $K$ . We want to show that the class of  $\mathfrak{c}$  in  $\mathrm{Cl}(K)$  is represented by an integral ideal  $\mathfrak{a}$  with  $\mathcal{N}(\mathfrak{a}) \leq B_K$ . Since  $\mathfrak{c}^{-1}$  is a fractional ideal,  $\exists d \in \mathcal{O}_K$ ,  $d \neq 0$  s.t.  $d\mathfrak{c}^{-1} = \mathfrak{b}$  is an integral ideal. By the previous proposition,  $\exists \beta \in \mathfrak{b}$ ,  $\beta \neq 0$  s.t.

$$|\mathrm{Nm}_{K/\mathbb{Q}}(\beta)| \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \sqrt{|\Delta_K|} \cdot \mathcal{N}(\mathfrak{b}).$$

Let  $\mathfrak{a} = (\beta)d^{-1}\mathfrak{c}$ . Then  $\mathfrak{a}$  is an integral ideal in the class of  $\mathfrak{c}$ , and

$$\mathcal{N}(\mathfrak{a}) = |\mathrm{Nm}_{K/\mathbb{Q}}(\beta)| \cdot \mathcal{N}(\mathfrak{b})^{-1} \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \sqrt{|\Delta_K|}.$$

■

### 5.3.2 Binary quadratic forms

Let

$$Q(x, y) = ax^2 + bxy + cy^2.$$

We call it integral if  $Q(m, n) \in \mathbb{Z}$  for all  $m, n \in \mathbb{Z}$ , i.e.  $a, b, c \in \mathbb{Z}$ . Its discriminant is defined as

$$d_Q = b^2 - 4ac.$$

A form is said to be non-degenerate if  $d_Q \neq 0$ .

Two integral binary quadratic forms  $Q$  and  $Q'$  are said to be equivalent if there exists  $M \in \text{SL}_2(\mathbb{Z})$  s.t.

$$Q'(x, y) = Q(px + qy, rx + sy), \quad M = \begin{pmatrix} p & q \\ r & s \end{pmatrix}.$$

The question considered by Gauss was to try to describe the set of equivalence classes of forms with a fixed discriminant.

## 5.4 Note on 20251117

Let  $d \neq 1$  be a square-free integer. Let  $K = \mathbb{Q}(\sqrt{d})$  and  $d_K := \text{disc}(\mathcal{O}_K/\mathbb{Z})$ .

Recall

$$\begin{cases} d_K = 4d & \text{if } d \equiv 2, 3 \pmod{4}, \\ d_K = d & \text{if } d \equiv 1 \pmod{4}. \end{cases}$$

Define norm form  $q_K$  by

$$q_K(x, y) = \text{Nm}_{K/\mathbb{Q}}(x + y\sqrt{d}) = x^2 - dy^2 \quad \text{if } d \equiv 2, 3 \pmod{4},$$

and

$$q_K(x, y) = \text{Nm}_{K/\mathbb{Q}}\left(x + y\frac{1 + \sqrt{d}}{2}\right) = x^2 + xy + \frac{1-d}{4}y^2 \quad \text{if } d \equiv 1 \pmod{4},$$

In general, if  $Q$  is an integral binary quadratic form, then  $d_Q = d_K \cdot f^2$  for some integer  $f$ , where  $K = \mathbb{Q}(\sqrt{d_Q})$ . Moreover, if  $d_Q = d_K$ , then  $Q$  is primitive,  $\gcd(a, b, c) = 1$ .

Fix a field  $K = \mathbb{Q}(\sqrt{d})$  and an embedding  $K \hookrightarrow \mathbb{C}$ . Choose  $\sqrt{d}$  to be positive if  $d > 0$ , and have positive imaginary part if  $d < 0$ .

Write  $\text{Gal}(K/\mathbb{Q}) = \{\text{id}, \sigma\}$ . If  $d < 0$ , define  $\text{Cl}^+(K) := \text{Cl}(K)$ , and if  $d > 0$ , define

$$\text{Cl}^+(K) := \text{Id}(K)/P^+(K),$$

where  $P^+(K)$  is the group of principle ideals of form  $(\alpha)$  with  $\alpha > 0$  under any embedding of  $K$  into  $\mathbb{R}$ .

Let  $\mathfrak{a}$  be a fractional ideal in  $K$ , let  $a_1, a_2$  be a basis of  $\mathfrak{a}$  as  $\mathbb{Z}$ -mod. We know

$$\begin{vmatrix} a_1 & a_2 \\ \sigma(a_1) & \sigma(a_2) \end{vmatrix}^2 = d_K \cdot \mathcal{N}(\mathfrak{a})^2.$$

After possibly reorder of  $a_1, a_2$ , we may ask

$$\begin{vmatrix} a_1 & a_2 \\ \sigma(a_1) & \sigma(a_2) \end{vmatrix} = \sqrt{d_K} \cdot \mathcal{N}(\mathfrak{a}).$$

For such a pair, define

$$Q_{a_1, a_2}(x, y) = \mathcal{N}(\mathfrak{a})^{-1} \text{Nm}_{K/\mathbb{Q}}(a_1x + a_2y).$$

This is an integral binary quadratic form with discriminant  $d_K$ .

**THEOREM 5.25.** *The equivalent class of  $Q_{a_1, a_2}(x, y)$  depends only on the image of  $\mathfrak{a}$  in  $\text{Cl}^+(K)$ . Moreover, the map sending  $\mathfrak{a}$  to the equivalence class of  $Q_{a_1, a_2}$  defines a bijection from  $\text{Cl}^+(K)$  to the set of equivalence classes of integral binary quadratic form with discriminant  $d_K$ .*

## 6 The Unit Theorem

Let  $K$  be a number field,  $r$  be the number of real embeddings of  $K$ , and  $2s$  be the number of non-real complex embeddings.

Thus,

$$K \otimes_{\mathbb{Q}} \mathbb{R} \simeq \mathbb{R}^s \times \mathbb{C}^s$$

and  $r + 2s = [K : \mathbb{Q}]$ .

**THEOREM 6.1.** *The group of units in  $\mathcal{O}_K$  is finitely generated with rank  $= r + s - 1$ .*

Denote  $U_K := \mathcal{O}_K^\times$ . The torsion group of  $U_K$  is  $\mu(K) = \{\text{roots of 1 in } K\}$ .

**EXAMPLE 6.2.** Let  $K$  be a real quadratic field. Then  $\text{rk}(U_K) = 2 + 0 - 1 = 1$ .

Let  $K$  be a complex quadratic field. Then  $\text{rk}(U_K) = 0 + 1 - 1 = 0$ .

A set of units  $u_1, \dots, u_{r+s-1}$  is called a fundamental system of units, if it forms a basis for  $U_K$  modulo torsions i.e. any unit  $u$  can be written uniquely in the form

$$u = \xi \cdot u_1^{m_1} \cdots u_{r+s-1}^{m_{r+s-1}}, \quad \xi \in \mu(K),$$

with  $m_i \in \mathbb{Z}$ .

**LEMMA 6.3.** *An element  $\alpha \in K$  is a unit iff  $\alpha \in \mathcal{O}_K$  and  $\text{Nm}_{K/\mathbb{Q}}(\alpha) = \pm 1$ .*

**Proof.** If  $\alpha \in U_K$ , then  $\exists \beta \in \mathcal{O}_K$  s.t.

$$\alpha \cdot \beta = 1 \implies \text{Nm}(\alpha) \cdot \text{Nm}(\beta) = 1 \implies \text{Nm}(\alpha) = \pm 1.$$

For the converse, fix an embedding  $\sigma_0: K \hookrightarrow \mathbb{C}$  and identify  $K$  with  $\sigma_0(K)$ . Then

$$\pm 1 = \text{Nm}(\alpha) = \prod_{\sigma} \sigma(\alpha) = \alpha \prod_{\sigma \neq \sigma_0} \sigma(\alpha) =: \alpha \cdot \beta.$$

Then  $\alpha \in \mathcal{O}_K \implies \sigma(\alpha)$  are algebraic integers  $\implies \beta$  is an algebraic integer. As  $\beta = \pm \alpha^{-1}$ ,  $\beta \in K \implies \beta \in \mathcal{O}_K \implies \alpha \in U_K$ . ■

### 6.0.1 Proof of that $U_K$ is finitely generated

**PROPOSITION 6.4.** *Given  $m, M > 0$ . The set of all algebraic integers  $\alpha$  s.t.*

1. *the degree of  $\alpha \leq m$ ; and*
2.  *$|\alpha'| \leq M$  for all conjugates  $\alpha'$  of  $\alpha$ ,*

*is finite.*

**Proof.** By (1),  $\alpha$  is a root of a monic irreducible polynomial of  $\deg \leq m$  over  $\mathbb{Z}$ .

By (2), the coefficients of the polynomial are bounded in term of  $M$ . Only finitely many such polynomials. ■

**COROLLARY 6.5.** *An algebraic integer  $\alpha$ , each of whose conjugates, in  $\mathbb{C}$  has absolute value 1, is a root of unity.*

**Proof.** By Proposition 6.4,  $\{1, \alpha, \alpha^2, \dots\}$  is a finite set. ■



**REMARK 6.6.** *It is essential to require  $\alpha$  to be an algebraic integer. For example,  $\alpha = \frac{3+4i}{5}$  is not a root of unity.*

Consider

$$\begin{aligned}\sigma: K &\longrightarrow \mathbb{R}^r \times \mathbb{C}^s \\ \alpha &\longmapsto (\sigma_1\alpha, \dots, \sigma_r\alpha, \sigma_{r+1}\alpha, \dots, \sigma_{r+s}\alpha).\end{aligned}$$

Take logarithm, we consider

$$\begin{aligned}L: K^\times &\longrightarrow \mathbb{R}^{r+s} \\ \alpha &\longmapsto (\log |\sigma_1\alpha|, \dots, \log |\sigma_{r+s}\alpha|),\end{aligned}$$

It is a homomorphism for  $\times$ .

For any  $u \in U_K$ , since  $\text{Nm}_{K/\mathbb{Q}}(u) = \pm 1$ , we have

$$|\sigma_1 u| \cdots |\sigma_r(u)| |\sigma_{r+1} u|^2 \cdots |\sigma_{r+s} u|^2 = 1,$$

$\implies L(u)$  is contained in the hyperplane

$$H: x_1 + \dots + x_r + 2x_{r+1} + \dots + 2x_{r+s} = 0,$$

$$H \simeq \mathbb{R}^{r+s-1}.$$

**PROPOSITION 6.7.** *The image of  $L$  in  $H$  is a lattice in  $H$ .*

$\ker L|_{U_K}$  is a finite group (hence  $= \mu(K)$ ).

**Proof.** Set

$$C = \{x \in H \mid |x_i| \leq M\},$$

for which  $o \in C$  and bounded. Let  $L(u) \in C$ , then  $|\sigma_i u| \leq e^M$  for any  $j$ . By Proposition 6.4, only finitely many such  $u \in U := \sigma(U_K)$ , i.e.  $\sharp(L^{-1}(C) \cap U_K) < \infty \implies L(U_K)$  is a lattice in  $H$ , and  $\ker L|_{U_K}$  is finite.  $\blacksquare$

Consider the exact sequence:

$$0 \longrightarrow \mu(K) \longrightarrow U_K \xrightarrow{L \circ \sigma} L(U_K) \rightarrow 0,$$

where  $\mu_K$  is finite, and  $L(U_K)$  is a lattice in  $H$  hence free of rank  $\leq \dim H = r + s - 1 \implies U_K$  is finitely generated and  $\text{rk}(U_K) \leq r + s - 1$ .

**THEOREM 6.8.** *The image  $L(U_K)$  in  $H$  is a full lattice ( $\implies \text{rk}(U_K) = r + s - 1$ ).*

**Proof.** We work with

$$\sigma: K \hookrightarrow \mathbb{R}^r \times \mathbb{C}^s \simeq \mathbb{R}^{r+2s}.$$

For  $x = (x_1, \dots, x_r, x_{r+1}, \dots) \in \mathbb{R}^r \times \mathbb{C}^s$ , define

$$\text{Nm}(x) = x_1 \dots x_r x_{r+1} \overline{x_{r+1}} \cdots x_{r+s} \overline{x_{r+s}}.$$

$L: K^\times \rightarrow \mathbb{R}^{r+s}$  extends to

$$\begin{aligned}L: \mathbb{R}^r \times \mathbb{C}^s &\longrightarrow (\mathbb{R} \cup \{-\infty\})^{r+s} \\ (x_1, \dots, x_r, x_{r+1}, \dots) &\longmapsto (\log |x_1|, \dots, \log |x_{r+1}|, \dots)\end{aligned}$$

continuous, surjective.

Consider

$$Y := \{x \in \mathbb{R}^r \times \mathbb{C}^s \mid |\mathrm{Nm}(x)| = 1\},$$

which is a group for  $\times$ . Then

- $Y$  is closed;
- $Y = L^{-1}(H)$ .

$\implies L|_Y := Y \rightarrow H$  surjective, continuous, and preserves the multiplication.

**LEMMA 6.9.**  $\exists \Omega \subset Y$  compact containing  $(1, \dots, 1)$  s.t.  $Y = \bigcup_{u \in U} u\Omega$ .

**Proof.** Let  $y \in Y$ , the map

$$\begin{aligned} \mathbb{R}^r \times \mathbb{C}^s &\longrightarrow \mathbb{R}^r \times \mathbb{C}^s \\ x &\longmapsto y \cdot x \end{aligned}$$

has  $|\text{Jacobian}| = |\mathrm{Nm}(y)| = 1$ .  $\implies$  It preserves the volume. Minkowski's thm  $\implies \exists B$  s.t. any compact convex subset  $T \subseteq \mathbb{R}^r \times \mathbb{C}^s$  symmetric in the origin, if  $\mu(T) \geq B$ , then  $T \cap (\sigma(\mathcal{O}_K) \setminus \{0\}) \neq \emptyset$ .

Pick such  $T$ . Any  $y \in Y$ ,  $y^{-1}T$  satisfies the same condition. For example,  $\mu(y^{-1}T) = \mu(T) \geq B \implies \exists, \gamma_y \in \mathcal{O}_K \setminus \{0\}$  s.t.  $\gamma_y \in y^{-1}T \implies \mathcal{N}((\gamma_y)) = \mathrm{Nm}(\gamma_y) \leq \max_{t \in T} \mathrm{Nm}(t) =: B_1$ .

The set of principle ideals  $\mathcal{N}(\gamma) \leq B_1$  is finite. Denote it by  $\{(\gamma_1), \dots, (\gamma_m)\}$ . Then  $(\gamma_y) = (\gamma_i)$  for some  $i = 1, \dots, m$ . Thus,  $\gamma_y = \gamma_i \cdot \varepsilon_y^{-1}$  for some  $\varepsilon_y \in U \implies \gamma_i \cdot \varepsilon_y^{-1} \in y^{-1}T \implies y \in \varepsilon_y \cdot \gamma_i^{-1}T \implies$

$$y \in \varepsilon_y \cdot \left( \bigcup_{i=1}^m \gamma_i^{-1}T \right).$$

As  $y \in Y$ ,  $\varepsilon_y \in Y$ ,  $y \cdot \varepsilon_y^{-1} \in Y \implies$

$$y \cdot \varepsilon_y^{-1} \in Y \cap \left( \bigcup_{i=1}^m \gamma_i^{-1}T \right) =: \Omega$$

compact,  $\implies Y = \bigcup_{\varepsilon \in U} \varepsilon\Omega$ . ■

Lemma  $\implies$

$$H = L(Y) \subseteq \bigcup_{u \in U} (L(u) + L(\Omega))$$

$\implies H = \bigcup_{u \in L(U)} (u + L(\Omega)) \implies L(U)$  is full. Otherwise, there exists a nonzero linear function  $g: H \rightarrow \mathbb{R}$  s.t.  $g(L(U)) = 0$ . Then

$$g(H) = g\left( \bigcup_{u \in L(U)} (u + L(\Omega)) \right) = g(L(\Omega))$$

is bounded. Contradiction! ■

## 6.1 Note on 20251119

### 6.1.1 $S$ -units

Let  $S$  be a finite set of prime ideals of  $\mathcal{O}_K$ .

**DEFINITION 6.10.** The ring of  $S$ -integers is

$$\mathcal{O}_K(S) := \bigcap_{\mathfrak{p} \notin S} \mathcal{O}_{\mathfrak{p}} = \{\alpha \in K \mid \text{ord}_{\mathfrak{p}}(\alpha) \geq 0, \forall \mathfrak{p} \notin S\}.$$

If  $S = \emptyset$ , then  $\mathcal{O}_K(S) = \mathcal{O}_K$ .

**DEFINITION 6.11.** The group of  $S$ -units is

$$U(S) := \mathcal{O}_K(S)^{\times} = \{\alpha \in K \mid \text{ord}_{\mathfrak{p}}(\alpha) = 0, \forall \mathfrak{p} \notin S\}.$$

Clearly, the torsion subgroup of  $U(S)$  is  $\mu(K)$ .

**THEOREM 6.12.** The group of  $S$ -units is finitely generated with rank  $= r + s + \#S - 1$ .

**Proof.** Write  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t\}$ . Consider the homomorphism

$$\begin{aligned} \theta: U(S) &\longrightarrow \mathbb{Z}^t \\ u &\longmapsto (\text{ord}_{\mathfrak{p}_1}(u), \dots, \text{ord}_{\mathfrak{p}_t}(u)), \end{aligned}$$

where  $\ker \theta = U(S)$ . Only need to show  $\text{rk im}(\theta) = t$ . Let  $h = \# \text{Cl}(K) < \infty$ . Then  $\mathfrak{p}_i^h$  is principal for any  $i$ . Write  $\mathfrak{p}_1^h = (\pi_i)$ . Then  $\pi_i$  is an  $S$ -unit with

$$\theta(\pi_i) = (0, \dots, 0, h, 0, \dots, 0),$$

where  $h$  is at the  $i$ -th position.  $\implies \text{im}(\theta) \supset h \cdot \mathbb{Z}^t$ , hence  $\text{rk im}(\theta) = t$ . ■

**EXAMPLE 6.13.**  $K = \mathbb{Q}$ ,  $S = \{(2), (3), (5)\}$ . Then  $U(S) = \{\pm 2^k 3^m 5^n \mid k, m, n \in \mathbb{Z}\}$ .

### 6.1.2 Example: CM fields

**DEFINITION 6.14.** We say a number field  $K$  is totally real if all of its embeddings in  $\mathbb{C}$  lie in  $\mathbb{R}$ , i.e.  $r = n$ ,  $s = 0$ . And it is totally imaginary if non of its embedding in  $\mathbb{C}$  is in  $\mathbb{R}$ , i.e.  $r = 0$ ,  $s = n/2$ .

A CM field  $K$  is a totally imaginary quadratic extension of a totally real field, i.e.

- $K = K^+(\sqrt{\alpha})$ ;
- $K^+$  is totally real;
- $\alpha \in K^+$ , any conjugate of  $\alpha$  is negative.

Such  $K^+$  is unique. Indeed, any  $\sigma: K \hookrightarrow \mathbb{C}$ ,  $K^+ = \sigma^{-1}(\sigma(K) \cap \mathbb{R})$ .

Let  $K$  be a CM field.  $m = [K^+ : \mathbb{Q}] \implies [K : \mathbb{Q}] = 2m$ . So  $\text{rk}(U_K) = m - 1 = \text{rk}(U_{K^+}) \implies [U_K : U_{K^+}] < \infty$ .

**PROPOSITION 6.15.** The index of  $\mu(K) \cdot U_{K^+}$  in  $U_K$  is either 1 or 2.

**Proof.**  $\text{Gal}(K/K^+) = \{\text{Id}, \tau\}$ . For any  $a \in K$ , write  $\tau(a) = \bar{a}$ . For any field embedding  $\rho: K \hookrightarrow \mathbb{C}$ , we have  $\overline{\rho(a)} = \rho(\bar{a}) \implies \forall a \in K$ , any conjugate of  $a/\bar{a}$  in  $\mathbb{C}$  has norm 1  $\implies a/\bar{a} \in \mu(K)$ .

Consider

$$\begin{aligned} \phi: U_K &\longrightarrow \mu(K)/\mu(K)^2 \\ a &\longmapsto \frac{a}{\bar{a}} \pmod{\mu(K)^2}. \end{aligned}$$

Any  $u \in \ker(\phi)$ ,  $u/\bar{u} = \xi^2$  for some  $\xi \in \mu(K)$ . Then

$$\frac{u \cdot \bar{\xi}}{\bar{u} \cdot \xi} = \xi^2 \cdot \frac{\bar{\xi}}{\xi} = 1$$

$$\implies u \cdot \bar{\xi} \in K^+ \implies u \in \mu(K) \cdot U_{K^+}.$$

Conversely, if  $u = \xi \cdot \overline{u^+}$  with  $\xi \in \mu(K)$  and  $u^+ \in U_{K^+}$ , then  $u/\bar{u} = \xi^2 \in \ker(\phi)$ . So  $\ker(\phi) = \mu(K) \cdot U_{K^+}$ . Note that  $\mu(K)$  is a cyclic group  $\implies \#\mu(K)/\mu(K)^2 \leq 2$ .  $\blacksquare$

### 6.1.3 Cyclotomic extensions

Let  $K$  be a field.

**DEFINITION 6.16.**  $\xi \in K$  is said to be a primitive  $n$ -th root of 1 if  $\xi^n = 1$  but  $\xi^d \neq 1$  for all  $d < n$ .

Then the  $n$ -th roots of 1 in  $\mathbb{C}$  are the numbers  $e^{\frac{2\pi im}{n}}$ ,  $0 \leq m \leq n-1$ , which is primitive iff  $(m, n) = 1$ .

**LEMMA 6.17.** Let  $\xi$  be a primitive  $n$ -th primitive root of 1. Then  $\xi^m$  is primitive iff  $(m, n) = 1$ .

Let  $K = \mathbb{Q}[\xi]$ , where  $\xi$  is a primitive  $n$ -th root of 1. Then  $K$  is the splitting field of  $x^n - 1 \implies K$  is Galois over  $\mathbb{Q}$ .

Denote  $\mathcal{G} := \text{Gal}(\mathbb{Q}[\xi]/\mathbb{Q})$ .

It permutes the set of primitive  $n$ -th root of 1 in  $K$ .

For any  $\sigma \in \mathcal{G}$ ,  $\sigma(\xi) = \xi^m$  for some  $m$  with  $(m, n) = 1$ . The map

$$\begin{aligned} \mathcal{G} &\longrightarrow (\mathbb{Z}/n\mathbb{Z})^\times \\ \sigma &\longmapsto [m] \end{aligned}$$

is an isomorphism.

**DEFINITION 6.18.** The cyclotomic polynomial  $\Phi_n$  is defined by

$$\Phi_n = \prod_{m \in (\mathbb{Z}/n\mathbb{Z})^\times} (x - \xi^m) = \prod_{\xi': \text{primitive } n\text{-th root of 1}} (x - \xi')$$

$$\implies x^n - 1 = \prod_{d|n} \Phi_d(x).$$

**PROPOSITION 6.19.** TFAE:

1. The map  $\mathcal{G} \rightarrow (\mathbb{Z}/n\mathbb{Z})^\times$  is an isomorphism;
2.  $[\mathbb{Q}[\xi] : \mathbb{Q}] = \varphi(n)$ ;
3.  $\mathcal{G}$  acts transitively on the set of primitive  $n$ -th roots of 1;
4.  $\Phi_n(x)$  is irreducible ( $\implies$  it is the minimal polynomial of  $\xi$ ).

**Proof.** Easy. ■

We now prove theses statements.

First treat the case  $n = p^r$ , where  $p$  is prime.

**PROPOSITION 6.20.** *Let  $\xi$  be a primitive  $p^r$ -th root of 1 with  $p$  prime. Then*

1.  $[K : \mathbb{Q}] = \varphi(p^r) = p^{r-1}(p-1)$ ;
2.  $\mathcal{O}_K = \mathbb{Z}[\xi]$ ;
3.  $\pi := 1 - \xi$  is a prime element of  $\mathcal{O}_K$  and  $(p) = (\pi)^e$  with  $e = \varphi(p^r)$ ;
4. The discriminant of  $\mathcal{O}_K$  over  $\mathbb{Z}$  is  $\pm p^c$ , where  $c = p^{r-1}(pr - r - 1)$  ( $\implies (p)$  is the only prime to ramify in  $\mathbb{Q}[\xi]$ ).

**Proof.**  $\xi$  integral  $\implies \mathbb{Z}[\xi] \subset \mathcal{O}_K$ . If  $\xi'$  is another primitive  $p^r$ -th root of 1, then  $\xi' = \xi^s$  and  $\xi = \xi'^t$  s.t.  $p \nmid s, p \nmid t$ .  $\implies \mathbb{Z}[\xi] = \mathbb{Z}[\xi']$  and  $\mathbb{Q}[\xi] = \mathbb{Q}[\xi']$ . Moreover,

$$\frac{1 - \xi'}{1 - \xi} = \frac{1 - \xi^s}{1 - \xi} = 1 + \xi + \cdots + \xi^{s-1} \in \mathbb{Z}[\xi].$$

$\frac{1 - \xi}{1 - \xi'} \in \mathbb{Z}[\xi]$  similarly.  $\implies \frac{1 - \xi'}{1 - \xi}$  is a unit of  $\mathbb{Z}[\xi]$ . Set  $t := x^{p^{r-1}}$ . Then

$$\Phi_{p^r}(x) = \frac{x^{p^r} - 1}{x^{p^{r-1}} - 1} = \frac{t^p - 1}{t - 1} = 1 + t + \cdots + t^{p-1},$$

and  $\Phi_{p^r}(1) = p$ . So

$$p = \Phi_{p^r}(1) = \prod (1 - \xi') = \prod \frac{1 - \xi'}{1 - \xi} \cdot (1 - \xi) = u \cdot (1 - \xi)^{\varphi(p^r)} \quad (6.6)$$

for some unit  $u$  in  $\mathbb{Z}[\xi]$ .

So we get  $(p) = (\pi)^e$ , for  $\pi = 1 - \xi$  and  $e = \varphi(p^r)$ .  $\implies (p)$  has  $\geq \varphi(p^r)$  prime factors in  $\mathcal{O}_K$ .

As  $[\mathbb{Q}[\xi] : \mathbb{Q}] \leq \deg \Phi_{p^r} = \varphi(p^r)$ , we get  $= \varphi(p^r)$ .  $\implies (1)$ .

Moreover,  $\pi \cdot \mathcal{O}_K$  is a prime ideal, otherwise  $(p) \cdot \mathcal{O}_K$  has too many prime ideal factors  $\implies (3)$ .

Then in  $\mathcal{O}_K$ ,

$$(p) = \mathfrak{p}^{\varphi(p^r)}, \quad \mathfrak{p} = (\pi) \text{ prime}, \quad f(\mathfrak{p}/(p)) = 1.$$

We next show (up to sign)  $\text{disc}(\mathbb{Z}[\xi]/\mathbb{Z})$  is a power of  $p$ .

Since

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) \cdot (\mathcal{O}_K : \mathbb{Z}[\xi])^2 = \text{disc}(\mathbb{Z}[\xi]/\mathbb{Z}),$$

this will imply

- $\text{disc}(\mathcal{O}_K/\mathbb{Z})$  is a power of  $p$ ;
- $(\mathcal{O}_K : \mathbb{Z}[\xi])$  is a power of  $p \implies$

$$p^M \mathcal{O}_K \subseteq \mathbb{Z}[\xi] \text{ for some } M. \quad (6.7)$$

We have

$$\text{disc}(\mathbb{Z}[\xi]/\mathbb{Z}) = \pm \text{Nm}_{K/\mathbb{Q}}(\Phi'_{p^r}(\xi)),$$

where

$$(x^{p^r-1} - 1) \cdot \Phi_{p^r}(x) = x^{p^r} - 1 \implies \Phi'_{p^r}(\xi)(\xi^{p^r} - 1) = p^r \xi^{p^r-1} \implies \Phi'_{p^r}(\xi) = \frac{p^r \xi^{p^r-1}}{\xi^{p^r-1} - 1},$$

$$\text{Nm}_{K/\mathbb{Q}}(\xi) = \pm 1, \text{Nm}_{K/\mathbb{Q}}(p^r) = (p^r)^{\varphi(p^r)} = p^{r\varphi(p^r)}.$$

**Claim:**  $\text{Nm}_{K/\mathbb{Q}}(1 - \xi^{p^s}) = \pm p^{p^s}, \forall 0 \leq s \leq r.$

**Proof of Claim.** Case  $s = 0$ :

The minimal polynomial of  $(1 - \xi)$  is  $\Phi_{p^r}(1 - x)$ . Then constant term is  $\Phi_{p^r}(1) = p \implies \text{Nm}_{K/\mathbb{Q}}(1 - \xi) = \pm p.$

Case  $1 \leq s \leq r$ :

$\xi^{p^s}$  is a primitive  $p^{r-s}$ -th root of 1. The  $s = 0$  case for  $\mathbb{Q}[\xi^{p^s}]/\mathbb{Q}$  implies  $\text{Nm}_{\mathbb{Q}[\xi^{p^s}]/\mathbb{Q}}(1 - \xi^{p^s}) = \pm p,$   
 $[\mathbb{Q}[\xi^{p^r}] : \mathbb{Q}[\xi^{p^s}]] = \frac{\varphi(p^r)}{\varphi(p^s)} = p^{r-s} \implies$

$$\text{Nm}_{K/\mathbb{Q}}(1 - \xi^{p^s}) = (\pm p)^{[\mathbb{Q}[\xi^{p^r}]:\mathbb{Q}[\xi^{p^s}]]} = (\pm p)^{p^s}.$$

■

Claim  $\implies \text{Nm}_{K/\mathbb{Q}}(\Phi'_{p^r}(\xi)) = \pm p^c$ , where  $c = r(p-1)p^{r-1} - p^{r-1} = p^{r-1}(pr - r - 1) \implies (4).$

Recall  $\mathfrak{p} = (1 - \xi) = (\pi)$  and  $f(\mathfrak{p}/(p)) = 1$ . Then  $\mathbb{Z}/(p) \simeq \mathcal{O}_K/(\pi) \implies$  for any  $m$ ,

$$\mathcal{O}_K = \mathbb{Z} + \pi \mathcal{O}_K = \dots = \mathbb{Z} + \pi \mathbb{Z} + \dots + \pi^{m-1} \mathbb{Z} + \pi^m \mathcal{O}_K \subseteq \mathbb{Z}[\xi] + \pi^m \mathcal{O}_K.$$

Recall  $p = \pi^{\varphi(n)} \times u$  where  $u$  is a unit in  $\mathbb{Z}[\xi]$ , and  $p^M \mathcal{O}_K \subseteq \mathbb{Z}[\xi]$ . Then by taking  $m$  sufficiently large ( $m \gg \varphi(n) \cdot M$ ), we have

$$\mathcal{O}_K \subseteq \mathbb{Z}[\xi] + \pi^m \mathcal{O}_K \subseteq \mathbb{Z}[\xi].$$

$\implies (2).$

■



**LEMMA 6.24.** *Let  $K, L$  be finite field extensions of  $\mathbb{Q}$ , s.t.*

$$[KL : \mathbb{Q}] = [K : \mathbb{Q}][L : \mathbb{Q}],$$

*and let  $d$  be the greatest common divisor of  $\text{disc}(\mathcal{O}_K/\mathbb{Z})$  and  $\text{disc}(\mathcal{O}_L/\mathbb{Z})$ . Then*

$$\mathcal{O}_{KL} \subseteq d^{-1}\mathcal{O}_K\mathcal{O}_L.$$

**Proof.** Let  $\{\alpha_1, \dots, \alpha_m\}$  and  $\{\beta_1, \dots, \beta_n\}$  be integral bases for  $\mathcal{O}_K$  and  $\mathcal{O}_L$  respectively. Then  $\{\alpha_i\beta_j\}$  is a basis for  $KL$  over  $\mathbb{Q}$ . Thus,  $\gamma \in \mathcal{O}_{KL}$  can be written as

$$\gamma = \sum_{i,j} \frac{a_{ij}}{r} \alpha_i \beta_j, \quad r \in \mathbb{Z},$$

with  $\frac{a_{ij}}{r}$  uniquely determined. We may assume  $(r, a_{ij} \forall ij) = 1$ . We have to show  $r \mid d$ .

We identify  $L$  with a subfield of  $\mathbb{C}$ . Any embedding  $\sigma : K \hookrightarrow \mathbb{C}$  extend uniquely to an embedding  $KL \hookrightarrow \mathbb{C}$ .

To see this, write  $K = \mathbb{Q}[\alpha]$ .  $KL = L[\alpha]$ . The hypothesis on the degree  $\implies$  the minimal polynomial of  $\alpha$  does not change when we pass from  $\mathbb{Q}$  to  $L$ . So there exists a unique  $L$ -homomorphism  $L[\alpha] \hookrightarrow \mathbb{C}$  sending  $\alpha$  to  $\sigma(\alpha)$ .

Applying  $\sigma$  to  $\gamma$ , we get

$$\sigma(\gamma) = \sum_{i,j} \frac{a_{ij}}{r} \sigma(\alpha_i) \beta_j.$$

Write  $x_i = \sum_j \frac{a_{ij}}{r} \beta_j$  and let  $\sigma_1, \dots, \sigma_m$  be the distinct embeddings of  $K$  into  $\mathbb{C}$ . We get  $m$  linear equations:

$$\sum_{j=1}^m \sigma_k(\alpha_i) x_i = \sigma_k(\gamma), \quad k = 1, \dots, m.$$

$\implies$

$$(x_1, \dots, x_m)^T = \left( \sigma_k(\alpha_i) \right)^{-1} \cdot (\sigma_1(\gamma), \dots, \sigma_m(\gamma))^T.$$

Denote  $D := \det(\sigma_k(\alpha_i))$ . Then

$$x_i = \frac{D_i}{D}, \quad D_i \in \mathcal{O}_{KL},$$

and  $D^2 = \text{disc}(\mathcal{O}_K/\mathbb{Z}) =: \Delta_K \implies$

$$\sum \frac{\Delta_K a_{ij}}{r} \beta_j = \Delta_K x_i = DD_i \in \mathcal{O}_{KL}.$$

$\implies$

$$\frac{\Delta_K a_{ij}}{r} \in \mathbb{Z}, \quad \forall ij \implies r \mid \Delta_K a_{ij}, \quad \forall ij.$$

$\implies$

$$r \mid (a_{ij} \forall ij) \Delta_K \implies r \mid \Delta_K.$$

Similarly,  $r \mid \Delta_L$ . So  $r \mid (\Delta_K, \Delta_L)$ . ■



Back to prove the theorem.

Induction hypothesis and  $p \nmid m \implies \text{disc}(\mathbb{Q}[\xi_{p^r}])$  and  $\text{disc}(\mathbb{Q}[\xi_m])$  are coprime. Lemma 6.24  $\implies$

$$\mathcal{O}_{\mathbb{Q}[\xi]} = \mathcal{O}_{\mathbb{Q}[\xi_{p^r}]} \mathcal{O}_{\mathbb{Q}[\xi_m]} = \mathbb{Z}[\xi_{p^r}] \mathbb{Z}[\xi_m] = \mathbb{Z}[\xi],$$

$\implies$  (2).

LEMMA 6.25. *Same setting of Lemma 6.24, i.e.*

$$[KL : \mathbb{Q}] = [K : \mathbb{Q}][L : \mathbb{Q}],$$

and assume  $\mathcal{O}_{KL} = \mathcal{O}_K \mathcal{O}_L$ . Then

$$\text{disc}(KL/\mathbb{Q}) = \text{disc}(K/\mathbb{Q})^{[L:\mathbb{Q}]} \text{disc}(L/\mathbb{Q})^{[K:\mathbb{Q}]}.$$

**Proof.** Let  $\{\alpha_1, \dots, \alpha_m\}$  and  $\{\beta_1, \dots, \beta_n\}$  be integral bases for  $\mathcal{O}_K$  and  $\mathcal{O}_L$  respectively. Then  $\{\alpha_i \beta_j\}$  is a basis for  $KL$  over  $\mathbb{Q}$ .

Let  $\{\sigma_1, \dots, \sigma_m\}$  and  $\{\tau_1, \dots, \tau_n\}$  be the embeddings of  $K \hookrightarrow \mathbb{C}$  and  $L \hookrightarrow \mathbb{C}$  respectively.

The proof of Lemma 6.24 shows that there exists a unique embedding  $\delta_{st} : KL \hookrightarrow \mathbb{C}$  s.t.  $\delta_{st}|_K = \sigma_s$  and  $\delta_{st}|_L = \tau_t$ . Those  $\delta_{st}$  are all embeddings  $KL \hookrightarrow \mathbb{C}$ .

$$\det(\delta_{st}(\alpha_i \beta_j)) = \det(\sigma_s(\alpha_i) \tau_t(\beta_j)) = \det(M_{tj}),$$

where

$$M_{tj} = \left( \tau_t(\beta_j) \cdot \sigma_s(\alpha_i) \right)_{s,i} = \tau_t(\beta_j) \cdot \left( \sigma_s(\alpha_i) \right)_{s,i}.$$

Since  $M_{tj}$  commutes with each other,

$$\det(M_{tj}) = \det \left( \det(\tau_t(\beta_j)) \cdot \left( \sigma_s(\alpha_i) \right)^n \right) = \left( \det(\tau_t(\beta_j)) \right)^m \cdot \left( \det(\sigma_s(\alpha_i)) \right)^n,$$

$\implies$

$$\text{disc}(KL/\mathbb{Q}) = \text{disc}(K/\mathbb{Q})^{[L:\mathbb{Q}]} \text{disc}(L/\mathbb{Q})^{[K:\mathbb{Q}]}.$$

■  
■

$\implies$  (3) by induction.

### 6.3 Note on 20251201

**REMARK 6.26.** • Statement (c) of the above Theorem shows that if  $p \mid n$ , then  $p$  ramifies unless  $\varphi(p^r) = 1$ , i.e.  $p^r = 2$ . Thus, if  $p \mid n$ , then  $p$  ramifies in  $\mathbb{Q}[\xi_n]$  except  $p = 2$  and  $n = 2 \times$  (odd number).

- Let  $m \in \mathbb{Z}_{>1}$ . Then  $\varphi(mn) > \varphi(n)$  except  $n$  is odd and  $m = 2 \implies \mu(\mathbb{Q}[\xi_n])$  is cyclic of order  $n$  (generated by  $\xi_n$ ) except when  $n$  is odd, in which case it is cyclic of order  $2n$  (generated by  $-\xi_n$ ).

**THEOREM 6.27 (Kummer).** Let  $p$  be an odd prime. If  $p \nmid \text{Cl}(\mathbb{Q}[\xi_p])$ , then there is no nonzero integer solution  $(x, y, z)$  to  $x^p + y^p = z^p$ .

**REMARK 6.28.** If  $p \nmid \text{Cl}(\mathbb{Q}[\xi_p])$ , call  $p$  a regular prime.

**Proof of Kummer's theorem for the case  $p$  relatively prime to  $xyz$ .**  $p = 3$  case: looking modulo 9.

$p = 5$  case: looking modulo 25.

Now assume  $p > 5$ . We may assume  $(x, y) = 1$ , i.e.  $x, y, z$  relatively prime in pair.

If  $x \equiv y \equiv -z \pmod{p}$ , then  $-2z^p \equiv z^p \pmod{p} \implies 3z^p \equiv 0 \pmod{p} \implies p \mid 3z \implies p \mid z$  contradiction.

Hence, either  $x \not\equiv y$  or  $x \not\equiv z \pmod{p}$ .

After rewriting the equation to  $x^p + (-z)^p = (-y)^p$ , we may assume  $x \not\equiv y \pmod{p}$ .

Set  $\xi := \xi_p$ . The roots of  $x^p + 1 = 0$  are  $-1, -\xi, \dots, -\xi^{p-1}$ . So

$$X^p + 1 = \prod_{i=0}^{p-1} (X + \xi^i) \implies \prod_{i=0}^{p-1} (x + \xi^i y) = z^p.$$

Let  $\mathfrak{p}$  be the unique prime ideal of  $\mathbb{Z}[\xi]$  dividing  $p \implies$

$$\mathfrak{p} = (1 - \xi^i), \quad \forall 1 \leq i \leq p-1.$$

**LEMMA 6.29.** The elements  $x + \xi^i y$  of  $\mathbb{Z}[\xi]$  are relatively prime in pairs.

**Proof.** Assume there exists a prime ideal  $\mathfrak{q}$  dividing  $x + \xi^i y$  and  $x + \xi^j y$ ,  $i \neq j$ ,  $\implies$

$$\mathfrak{q} \mid (\xi^i - \xi^j)y = \mathfrak{p}y$$

and

$$\mathfrak{q} \mid (\xi^j - \xi^i)x = \mathfrak{p}x.$$

Since  $(x, y) = 1$ ,  $\mathfrak{q} \mid \mathfrak{p} \implies \mathfrak{q} = \mathfrak{p}$ . As  $x + y \equiv (x + \xi^i y) + (1 - \xi^i)y$ , we have  $\mathfrak{p} \mid x + y \implies p \mid x + y \implies z^p \equiv x^p + y^p \equiv x^p + (-y)^p \equiv 0 \pmod{p}$ , contradiction. ■

**LEMMA 6.30.** Any  $\alpha \in \mathbb{Z}[\xi]$ ,  $\alpha^p \in \mathbb{Z} + p\mathbb{Z}[\xi]$ .

**Proof.**  $\mathbb{Q}[\xi]/\mathbb{Q}$  totally ramifies at  $\mathfrak{p} \implies \mathbb{Z}/p\mathbb{Z} \simeq \mathbb{Z}[\xi]/\mathfrak{p}$ . So  $\alpha = u + v \in \mathbb{Z} + \mathfrak{p}$ . Thus,

$$\alpha^p = (u + v)^p = u^p + \sum_{i=1}^{p-1} \binom{p}{i} u^i v^{p-i} + v^p,$$

where  $p \mid \binom{p}{i}$  and  $v^p \in \mathfrak{p}^p = (p)$ . Hence,  $\alpha^p \in \mathbb{Z} + p\mathbb{Z}[\xi]$ . ■

**LEMMA 6.31.** *Let*

$$\alpha = a_0 + a_1\xi + \cdots + a_{p-1}\xi^{p-1}$$

*with  $a_i \in \mathbb{Z}$  and  $a_0 \cdots a_{p-1} = 0$ . If  $\alpha \in n\mathbb{Z}[\xi]$  for some  $n \in \mathbb{Z}$ , then  $n \mid a_i, \forall i$ .*

**Proof.** Since  $1 + \xi + \cdots + \xi^{p-1} = 0$ , any subset of  $\{1, \xi, \dots, \xi^{p-1}\}$  with  $p-1$  elements will be a  $\mathbb{Z}$ -basis. This lemma is clear.  $\blacksquare$

View

$$\prod_{i=0}^{p-1} (x + \xi^i y) = (z)^p$$

as an equality in  $\mathbb{Z}[\xi]$ . Then factors in the LHS are relatively prime in pairs. So each one as an ideal is a  $p$ -th power, i.e.

$$(x + \xi^i y) = \mathfrak{a}_i^p$$

for some ideal  $\mathfrak{a}_i$  in  $\mathbb{Z}[\xi]$ .

As  $p \nmid \text{Cl}(\mathbb{Z}[\xi])$ ,  $\mathfrak{a}_i$  is principal. Say  $\mathfrak{a}_i = (\alpha_i)$ . Take  $i = 1$ , we have

$$x + \xi y = u \cdot \alpha_1^p, \quad \text{where } u \in \mathbb{Z}[\xi]^\times.$$

**Claim**  $u = \xi^r \cdot v$ , where  $v = \bar{v}$ .

Then Lemma 6.30 implies there exists  $a \in \mathbb{Z}$  s.t.

$$\alpha_1^p \equiv a \pmod{p} \implies x + \xi y = \xi^r v \alpha_1^p \equiv \xi^r v a \pmod{p}.$$

$\implies$

$$x + \bar{\xi} y = \xi^{-r} v \bar{\alpha}^p \equiv \xi^{-r} v a \pmod{p}.$$

Then we get

$$\xi^{-r}(x + \xi y) \equiv \xi^r(x + \xi^1 y) \pmod{p}.$$

$\implies$

$$x + \xi y - \xi^{2r} x - \xi^{2r-1} y \equiv 0 \pmod{p}.$$

If  $1, \xi, \xi^{2r-1}, \xi^{2r}$  are distinct, then  $p \mid x$ , contradiction. The only remaining possibilities are

1.  $1 = \xi^{2r}$ , then

$$\xi y - \xi^{2r-1} y \equiv 0 \pmod{p} \implies p \mid y,$$

contradiction.

2.  $1 = \xi^{2r-1} \Leftrightarrow \xi = \xi^{2r}$ . Then

$$(x - y) - (x - y)\xi \equiv 0 \pmod{p} \implies p \mid (x - y),$$

contradiction.

3.  $\xi = \xi^{2r-1}$ . Then

$$x - \xi^2 x = x - \xi^{2r} x \equiv 0 \pmod{p} \implies p \mid x,$$

contradiction.

**Proof of Claim.**  $\xi = \xi_n$ ,  $n > 2$ . Set

$$\mathbb{Q}[\xi]^+ := \mathbb{Q}[\xi + \xi^{-1}].$$

Under any embedding  $\rho: \mathbb{Q}[\xi] \hookrightarrow \mathbb{C}$ ,  $\rho(\xi^{-1}) = \overline{\rho(\xi)}$ . So  $\mathbb{Q}[\xi]^+$  is a totally real field. Then  $\mathbb{Q}[\xi]$  is a CM field. Hence, the index of  $\mu(\mathbb{Q}[\xi]) \cdot U_{\mathbb{Q}[\xi]^+}$  in  $U_{\mathbb{Q}[\xi]}$  is 1 or 2.

**LEMMA 6.32.** *If  $n$  is an odd prime power, then any unit  $u \in \mathbb{Q}[\xi]$  can be written as  $u = \xi \cdot v$ , where  $\xi$  is a root of 1, and  $v$  is a unit in  $\mathbb{Q}[\xi]^+$ .*

**Proof.** By contradiction, if the homomorphism ( $\mu = \mu(\mathbb{Q}[\xi])$ )

$$\begin{aligned} U_{\mathbb{Q}[\xi]} &\longrightarrow \mu/\mu^2 \\ u &\longmapsto u/\bar{u} \end{aligned}$$

were surjective, then there exists  $u \in U_{\mathbb{Q}[\xi]}$  s.t.  $\bar{u} = \beta u$  where  $\beta$  is a root of 1 which is not a square. As  $n$  is odd,  $\mu = \{\pm 1\} \cdot \langle \xi \rangle$ . So  $\mu^2 = \langle \xi \rangle \implies \beta = -\xi^m$  for some  $m \in \mathbb{Z}$ . Let

$$u = a_0 + \cdots + a_{\varphi(n)-1} \xi^{\varphi(n)-1}, \quad a_i \in \mathbb{Z}.$$

Then

$$\bar{u} = a_0 + \cdots + a_{\varphi(n)-1} \bar{\xi}^{\varphi(n)-1}.$$

Modulo the prime  $\mathfrak{p} := (1 - \xi) = (1 - \bar{\xi}) \implies$

$$u \equiv a_0 + \cdots + a_{\varphi(n)-1} \equiv \bar{u} \pmod{\mathfrak{p}} \implies u \equiv -\xi^m u \equiv -u \pmod{p},$$

$$\implies 2u \in \mathfrak{p} \implies 2 \in \mathfrak{p}, \text{ contradiction.} \quad \blacksquare$$

Lemma  $\implies$  Claim.  $\blacksquare$

The proof is complete.  $\blacksquare$

## 7 Absolute values

**DEFINITION 7.1.** An absolute value on a field  $K$  is a function

$$\begin{aligned} K &\longrightarrow \mathbb{R}_{\geq 0} \\ x &\longmapsto |x| \end{aligned}$$

s.t.

- (a)  $|x| = 0$  iff  $x = 0$ ;
- (b)  $|xy| = |x| \cdot |y|$ ;
- (c)  $|x + y| \leq |x| + |y|$ .

If the stronger condition

$$(c') \quad |x + y| \leq \max\{|x|, |y|\},$$

holds, the  $|\cdot|$  is called non archimedean. An absolute value is called archimedean if it is not non archimedean.

**REMARK 7.2.**  $(a)(b) \implies |\cdot|: K^\times \rightarrow (\mathbb{R}_{>0}, \times)$  is a homomorphism.  $\mathbb{R}_{>0}$  is torsion free  $\implies |\mu(K)| = 1$ . In particular,  $\forall x \in K, |x| = |-x|$ .

**EXAMPLE 7.3.** 1. Let  $K$  be a field, and  $\sigma: K \hookrightarrow \mathbb{C}$ . Then we get an archimedean absolute value on  $K$  by  $|x| := |\sigma(x)|$ .

- 2. Let  $\text{ord}: K^\times \rightarrow \mathbb{Z}$  be a discrete valuation, then  $|a| := c^{-\text{ord}(a)}$  for  $c > 1$  is a NA absolute value.

For example, any prime number  $p$ , we have the  $p$ -adic absolute value  $|\cdot|_p$  on  $\mathbb{Q}$ :

$$|a|_p := c^{-\text{ord}_p(a)}, \quad c > 1.$$

Usually we normalize this by taking  $c = p$ .

Similarly, any prime ideal  $\mathfrak{p}$  in a number field  $K$ , we have a normalized  $\mathfrak{p}$ -adic absolute value

$$|a|_{\mathfrak{p}} := \left( \frac{1}{\mathcal{N}(\mathfrak{p})} \right)^{\text{ord}_{\mathfrak{p}}(a)}.$$

- 3. On any field  $K$ , we define the trivial absolute value:  $|a| = 1, \forall a \neq 0$ .

### 7.0.1 NA absolute value

Recall (c')  $|x + y| \leq \max\{|x|, |y|\}$ . Then

$$\left| \sum_{\text{finite}} x_i \right| \leq \max\{|x_i|\}.$$

**PROPOSITION 7.4.** An absolute value  $|\cdot|$  is NA iff  $|m \cdot 1|, m \in \mathbb{Z}$  is bounded.

**Proof.** “Only if” trivial.

“If” part: Assume  $|m \cdot 1| \leq N$ ,  $\forall m \in \mathbb{Z}$ . Then

$$|x + y|^n = \left| \sum_{i=0}^n \binom{n}{i} x^i y^{n-i} \right| \leq \sum_{i=0}^n \left\{ \left| \binom{n}{i} \right| \cdot 1 \cdot |x|^i |y|^{n-i} \right\} \leq (n+1)N \cdot \max\{|x|, |y|\}^n$$

for any  $n \geq 1 \implies$

$$|x + y| \leq (n+1)^{1/n} N^{1/n} \max\{|x|, |y|\}$$

$\implies$  (c') by  $n \rightarrow \infty$ . ■

## 7.1 Note on 20251203

**COROLLARY 7.5.** *If  $\text{char } K \neq 0$ , then any absolute value on  $K$  is NA.*

**PROPOSITION 7.6.** *Let  $|\cdot|$  be a non-trivial NA absolute value and put*

$$v(x) := -\log_c |x|, \quad x \neq 0, \quad c > 1,$$

*then  $v: K^\times \rightarrow \mathbb{R}$  satisfies the following conditions:*

- $v(xy) = v(x) + v(y)$ ;
- $v(x + y) \geq \min\{v(x), v(y)\}$ .

*If  $v(K^\times)$  is discrete in  $\mathbb{R}$ , then  $v$  is a multiple of a normalized discrete valuation  $\text{ord}: K^\times \rightarrow \mathbb{Z} \subset \mathbb{R}$ .*

We say  $|\cdot|$  is discrete when  $|K^\times|$  is a discrete subgroup of  $\mathbb{R}_{>0}$ .

**PROPOSITION 7.7.** *Let  $|\cdot|$  be a NA absolute value. Then*

$$A := \{a \in K \mid |a| \leq 1\}$$

*is a subring of  $K$ , with*

$$U := \{a \in K \mid |a| = 1\}$$

*the group of units in  $A$  and*

$$\mathfrak{m} := \{a \in K \mid |a| < 1\}$$

*the unique maximal ideal of  $A$ .*

*In particular,  $|\cdot|$  is discrete iff  $A$  is a DVR.*

**EXAMPLE 7.8.** Let  $R = \mathbb{Q}[t]$  and  $r > 0$ . Then any  $f = \sum_{i \geq 0} a_i t^i \in R$ , define

$$|f|_r := \max\{|a_i| r^i\}$$

where  $|a_i|$  denotes the  $p$ -adic absolute value with  $|p| = p^{-1}$ . One can check

- $|f|_r = 0$  iff  $f = 0$ ;
- $|f_1 f_2|_r = |f_1|_r \cdot |f_2|_r$ .

Hence  $|\cdot|_r$  extends to an absolute value on  $K = \mathbb{Q}(t) = \text{Frac } \mathbb{Q}[t]$ . If  $\log_p r \notin \mathbb{Q}$ , then  $|\cdot|_r$  is not discrete.

An absolute value  $|\cdot|$  defines a metric on  $K$  with

$$d(a, b) := |a - b|,$$

which induces a topology on  $K$ .

**EXAMPLE 7.9.** On  $(\mathbb{Q}, |\cdot|_p)$ , we have  $p^n \rightarrow 0$  as  $n \rightarrow \infty$ .

**PROPOSITION 7.10.** *Let  $|\cdot|_1, |\cdot|_2$  be absolute values on  $K$ , and  $|\cdot|_1$  nontrivial. Then TFAE:*

1.  $|\cdot|_1, |\cdot|_2$  defines the same topology on  $K$ ;
2.  $|\alpha|_1 < 1 \implies |\alpha|_2 < 1$ ;

3.  $|\cdot|_2 = |\cdot|_1^a$  for some  $a > 0$ .

**Proof.** (1)  $\implies$  (2): If  $|\alpha|_1 < 1$ , then  $\alpha^n \rightarrow 0 \implies |\alpha|_2 < 1$ .

(2)  $\implies$  (3):  $|\cdot|_1$  nontrivial  $\implies \exists x \in K$  with  $|x|_1 > 1 \implies |x|_2 > 1$ . Let  $y \in K^\times$ ,  $m, n \in \mathbb{Z} \setminus \{0\}$ . Then

$$\frac{\log |y|_2}{\log |x|_2} < \frac{m}{n} \iff \left| \frac{y^m}{x^n} \right|_2 < 1 \implies \left| \frac{y^m}{x^n} \right|_1 < 1 \iff \frac{\log |y|_1}{\log |x|_1} < \frac{m}{n},$$

$\implies \frac{\log |y|_1}{\log |x|_1} = \frac{\log |y|_2}{\log |x|_2}$  (replacing  $y$  by  $y^{-1}$  for the converse inequality). Set  $a := \frac{\log |x|_2}{\log |x|_1}$ , then  $\log |y|_2 = a \log |y|_1$ .

(3)  $\implies$  (1): Clear. ■

Two absolute values are said to be equivalent if they satisfy the above equivalent conditions.

**THEOREM 7.11 (Ostrowski).** Let  $|\cdot|$  be a nontrivial absolute value on  $\mathbb{Q}$ .

- If  $|\cdot|$  is archimedean, then  $|\cdot|$  is equivalent to  $|\cdot|_\infty$ ;
- If  $|\cdot|$  is NA, then it is equivalent to  $|\cdot|_p$  for exactly one prime  $p$ .

**Proof.** Let  $m, n \in \mathbb{Q}_{>1}$ . Write

$$m = a_0 + a_1 n + \cdots + a_r n^r,$$

with  $0 \leq a_i \leq n - 1$ ,  $a_r \neq 0$ . Then  $r \leq \frac{\log m}{\log n}$ , and

$$|a_i| \leq |1| + \cdots |1| = a_i < n. \quad (7.8)$$

Let  $N := \max\{1, |n|\}$ . Then

$$|m| \leq \sum_{i=0}^r |a_i| |n|^r \leq \sum_{i=0}^r |a_i| N^r \leq (1+r)n \cdot N^r \leq \left(1 + \frac{\log m}{\log n}\right) n \cdot N^{\log m / \log n}, \quad (7.9)$$

and again applying this to  $m^t$ , we get

$$|m|^t \leq \left(1 + t \frac{\log m}{\log n}\right) n \cdot N^{t \cdot \log m / \log n}$$

$\implies$

$$|m| \leq \left(1 + t \frac{\log m}{\log n}\right)^{1/t} n^{1/t} \cdot N^{\log m / \log n}.$$

Letting  $t \rightarrow \infty$ , we get

$$|m| \leq N^{\log m / \log n}. \quad (7.10)$$

**Case (i)**  $|n| > 1$  for any  $n > 1$ .

Then  $N = |n| \implies$

$$|m|^{1/\log m} \leq |n|^{1/\log n}, \quad \forall m, n > 1.$$



Hence  $|n|^{1/\log n} = c$  a constant  $\implies |n| = c^{\log n} = n^{\log c}$ . Let  $a := \log c$ . Then

$$|n| = |n|_{\infty}^a, \quad \forall n \in \mathbb{Z}_{>1}.$$

Note that  $|\cdot|$  and  $|\cdot|_{\infty}$  are homomorphisms  $\mathbb{Q}^{\times} \rightarrow \mathbb{R}$  and  $\{\mathbb{Z}_{>1}, \pm 1\}$  generates  $\mathbb{Q}^{\times} \implies |\cdot| = |\cdot|_{\infty}^a$ .

**Case (ii)**  $\exists n > 1$  s.t.  $|n| \leq 1$ .

Now  $N = 1$ . We then have  $|m| \leq 1$  for any  $m \in \mathbb{Z} \implies |\cdot|$  is NA. Let

$$A := \{x \in \mathbb{Q} \mid |x| \leq 1\}, \quad \mathfrak{m} := \{x \in \mathbb{Q} \mid |x| < 1\}.$$

Then  $\mathfrak{m} \cap \mathbb{Z}$  is a prime ideal of  $\mathbb{Z}$  and  $\mathfrak{m} \cap \mathbb{Z} \neq 0$ . Otherwise,  $|\cdot|$  is trivial. So  $\mathfrak{m} \cap \mathbb{Z} = (p)$  with  $p$  prime.

$$|m| = 1 \quad \text{if } m \in \mathbb{Z}, p \nmid m.$$

So for any  $x \in \mathbb{Q}^{\times}$ , write  $x = \frac{m}{n} \cdot p^r$ , where  $m, n \in \mathbb{Z} \setminus \{0\}$ ,  $p \nmid mn$ ,  $r \in \mathbb{Z}$ . Then

$$|x| = \frac{|m|}{|n|} \cdot |p|^r = |p|^r.$$

Let  $a := -\log_p |p|$ . Then  $|x| = |x|_p^a$ . ■

**THEOREM 7.12 (Product formula).** For  $p$  a prime or  $\infty$ , let  $|\cdot|_p$  be the normalized absolute value on  $\mathbb{Q}$ . Let  $|\cdot|_0$  be the trivial absolute value. Then

$$\prod_p |a|_p = |a|_0, \quad \forall a \in \mathbb{Q}.$$

### 7.1.1 Places of a number field

Let  $K$  be a number field.

**DEFINITION 7.13.** An equivalence class of absolute values on  $K$  is called a place of  $K$ .

**THEOREM 7.14.** There exists exactly one place of  $K$  for

- any prime ideal  $\mathfrak{p}$ ;
- any real embedding;
- any conjugate pair of complex embeddings.

In each equivalence class, we select a normalized “absolute value”

- any  $\mathfrak{p}$  prime ideal of  $\mathcal{O}_K$ ,

$$|a|_{\mathfrak{p}} := \left( \frac{1}{\mathcal{N}(\mathfrak{p})} \right)^{\text{ord}_{\mathfrak{p}}(a)};$$

- any real embedding,

$$\sigma: K \hookrightarrow \mathbb{R}, \quad |a| := |\sigma(a)|.$$

- any non-real complex embedding

$$\sigma: K \hookrightarrow \mathbb{C}, \quad |a| := |\sigma(a)|^2.$$

(notice that this is not an absolute value, but we ignore it.)

Let  $v$  be a place on  $K$  a number field.

- If  $v$  is from a prime ideal, call  $v$  a finite place.
- If  $v$  is from a (real or non-real) embedding, call  $v$  an infinite (real or complex) place.

**DEFINITION 7.15.** We write  $|\cdot|_v$  for an absolute value in the equivalence class. If  $L \supset K$  and  $w, v$  are places of  $L$  and  $K$  respectively s.t.  $|\cdot|_w|_K$  is equivalent to  $|\cdot|_v$ , then we say  $w$  divides  $v$  or  $w$  lies over  $v$ . Write  $w \mid v$ .

For finite places,  $w \mid v$  means

$$\beta_w \cap \mathcal{O}_K = \mathfrak{p}_v.$$

For infinite place,  $w \mid v$  means

$$\sigma_w: L \hookrightarrow \mathbb{C} \text{ extends the } \sigma_v \text{ or } \overline{\sigma_v}: K \hookrightarrow \mathbb{C}.$$

## 7.2 Note on 20251215

**THEOREM 7.16 (Product formula).** *For any place  $v$ , let  $|\cdot|_v$  be the normalized absolute value. The for any  $\alpha \in K^\times$ ,*

$$\prod_v |\alpha|_v = 1.$$

**Proof.** Product formula for  $\mathbb{Q}$  plus the next result.

**LEMMA 7.17.**  *$L/K$  finite extension:*

- (a) *Any place on  $K$  extends to a finite number of places of  $L$ ;*
- (b) *Any place  $v$  of  $K$  and  $\alpha \in L^\times$ ,*

$$\prod_{w|v} |\alpha|_w = |\mathrm{Nm}_{L/K}(\alpha)|_v.$$

**Proof.** Next part. ■

### 7.2.1 The weak approximation theorem

**LEMMA 7.18.** *If  $|\cdot|_1, \dots, |\cdot|_n$  are distinct inequivalent nontrivial absolute values of  $K$ , then there exists  $a \in K$  s.t.*

$$\begin{cases} |a|_1 > 1, \\ |a|_i < 1, \quad \forall i \neq 1. \end{cases}$$

**Proof.** First let  $n = 2$ . Then  $|\cdot|_1$  and  $|\cdot|_2$  are not equivalent shows that there exist  $b, c \in K$  s.t.

$$\begin{cases} |b|_1 < 1, & |b|_2 \geq 1; \\ |c|_1 \geq 1, & |c|_2 < 1. \end{cases}$$

Take  $a := \frac{c}{b}$ .

By induction, assume this lemma holds for  $n - 1$  absolute values. Then there exists  $b, c \in K$  s.t.

$$\begin{cases} |b|_1 > 1, & |b|_i < 1, \quad i = 2, \dots, n - 1; \\ |c|_1 > 1, & |c|_n < 1. \end{cases}$$

If  $|b|_n \leq 1$ , set

$$a := c \cdot b^r, \quad r \gg 0.$$

If  $|b|_n > 1$ , set

$$a := \frac{c \cdot b^r}{1 + b^r}, \quad r \gg 0.$$

Note

$$\left| \frac{b^r}{1 + b^r} \right| = \begin{cases} 1, & \text{if } |b| > 1; \\ 0, & \text{if } |b| < 1. \end{cases}$$

Easy to check such  $a$  is OK. ■

**LEMMA 7.19.** *Same situation as Lemma 7.18. There exist  $a_r \in K$  for  $r \geq 0$ , s.t.*

$$|a_r|_1 \rightarrow 1 \quad \text{and} \quad |a_r|_i \rightarrow 0, \quad \forall i = 2, \dots, n.$$

**Proof.** Pick  $a$  as in Lemma 7.18. Set

$$a_r := \frac{a^r}{1 + a^r}.$$

■

**THEOREM 7.20.** *Let  $|\cdot|_1, \dots, |\cdot|_n$  be distinct inequivalent nontrivial absolute values of  $K$ , and  $a_1, \dots, a_n \in K$ . For any  $\varepsilon > 0$ , there exists  $a \in K$  s.t.*

$$|a - a_i|_i < \varepsilon, \quad \forall i = 1, \dots, n.$$

**Proof.** By Lemma 7.19, choose  $b_i, i = 1, \dots, n$  s.t.

$$|b_i - 1|_i \approx 0, \quad |b_i|_j \approx 0, \quad \forall j \neq i.$$

Set

$$a = a_1 b_1 + \dots + a_n b_n.$$

■

**REMARK 7.21.** *Let  $K_i := (K, |\cdot|_i)$ .*

*We have the following diagonal embedding:*

$$\tau: K \hookrightarrow \prod_{i=1}^n K_i.$$

*Theorem 7.20  $\implies \tau(K)$  is dense in  $\prod_{i=1}^n K_i$ .*

**COROLLARY 7.22.** *Let  $|\cdot|_1, \dots, |\cdot|_n$  be distinct inequivalent nontrivial absolute values of  $K$ . If*

$$|a|_1^{r_1} \cdots |a|_n^{r_n} = 1, \quad r_i \in \mathbb{R},$$

*for any  $a \in K^\times$ , then  $r_i = 0$  for all  $i$ .*

**Proof.** Assume  $r_i \neq 0$  for all  $i$ . Pick  $a_i \in K$  with  $|a_i|_i^{r_i} > 1$ . Then Theorem 7.20  $\implies$  there exists  $a$  s.t.  $|a - a_i|_i \approx 0$  for all  $i \implies |a_i|_i^{r_i} > 1$  for all  $i \implies \prod_{i=1}^n |a_i|_i^{r_i} > 1$ . ■

## 8 Completions

Let  $K$  be a field with an absolute value  $|\cdot|$ .

**DEFINITION 8.1.** A sequence  $a_n$  of  $K$  is called a Cauchy sequence if  $\forall \varepsilon > 0, \exists N \geq 0$  s.t.

$$|a_n - a_m| < \varepsilon, \quad \forall m, n \geq N.$$

$K$  is called complete if any Cauchy sequence has a limit in  $K$ .

**EXAMPLE 8.2.**

$$a_n = 1 + 2 + \cdots + 2^n = 2^{n+1} - 1$$

is a Cauchy sequence for  $|\cdot|_2$ , and  $\lim_{n \rightarrow \infty} a_n = -1$ .

**THEOREM 8.3.** Let  $K$  be a field with an absolute value  $|\cdot|$ . Then there exists a complete valued field  $(\widehat{K}, |\cdot|)$  and a homomorphism  $K \rightarrow \widehat{K}$  preserving the absolute value that is universal in the following sense:

Any homomorphism  $K \rightarrow L$  from  $K$  into a complete valued field  $(L, |\cdot|)$  preserving the absolute value, extends uniquely to a homomorphism  $\widehat{K} \rightarrow L$ .

**Proof.** Define

$$\widehat{K} := \{\text{Cauchy sequence of } K\} / \sim,$$

where  $(a_n) \sim (b_n)$  if  $|a_n - b_n| \rightarrow 0$ .

Check  $\widehat{K}$  is a field.

For  $a \in \widehat{K}$ , defined by a Cauchy sequence  $a_n \in K$ , define  $|a| := \lim_{n \rightarrow \infty} |a_n|$ .

This is well-defined. Check  $|\cdot|$  is an absolute value on  $\widehat{K}$ , and  $(\widehat{K}, |\cdot|)$  is complete.

Check the map

$$\begin{aligned} K &\longrightarrow \widehat{K} \\ a &\longmapsto (a, a, \dots) \end{aligned}$$

is an isometry.

Let  $(L, |\cdot|)$  be a complete valued field with an isometry  $\phi: K \hookrightarrow L$ . It extends uniquely to  $\widehat{K} \rightarrow L$  by

$$(a_n) \longmapsto \lim_{n \rightarrow \infty} \phi(a_n).$$

■

**REMARK 8.4.**  $K \rightarrow \widehat{K}$  is uniquely determined up to a uniquely isomorphism by the universal property.

**REMARK 8.5.** The image of  $K$  in  $\widehat{K}$  is dense in  $\widehat{K}$ .

Any place  $v$  of  $K$ , write  $K_v$  the completion of  $K$  w.r.t.  $v$ . When  $v$  corresponds to a prime ideal  $\mathfrak{p}$ , we write  $K_{\mathfrak{p}}$  for the completion and  $\widehat{\mathcal{O}}_{\mathfrak{p}}$  for the ring of integers in  $K_{\mathfrak{p}}$ .

For example,  $\mathbb{Q}_p$  is the completion of  $\mathbb{Q}$  w.r.t. the  $p$ -adic absolute value. Write  $\mathbb{Z}_p$  (not  $\widehat{\mathbb{Z}}_p$ ) for the ring of integers in  $\mathbb{Q}_p$ .

### 8.0.1 Completion for discrete valuation field

Let  $|\cdot|$  be a discrete NA absolute value on  $K$ , and  $A$  the valuation ring. Let  $\mathfrak{m}$  be the maximal ideal of  $A$ . Write  $\mathfrak{m} = (\pi)$ , where  $\pi$  is called a local uniformizing parameter. Then

$$|K| = \{c^m \mid m \in \mathbb{Z}\} \cup \{0\},$$

where  $c = |\pi| < 1$ .

Let  $a \in \widehat{K}^\times$  with  $a_n \rightarrow a$ , where  $a_n \in K$ . Then  $|a_n - a| < |a|$  for  $n \gg 0 \implies |a_n| = |a|$  for any  $n \gg 0$ . So  $|\widehat{K}| = |K|$  and  $|\cdot|$  is a discrete absolute value on  $\widehat{K}$ .

Let  $\text{ord}: K^\times \rightarrow \mathbb{Z}$  be a normalized discrete valuation on  $K$ . It extends to a normalized discrete valuation on  $\widehat{K}$ .

Define  $\widehat{A} := \{a \in \widehat{K} \mid |a| \leq 1\}$ , which is the closure of  $A$  in  $\widehat{K}$ , and  $\widehat{\mathfrak{m}} := \{a \in \widehat{K} \mid |a| < 1\}$  is the maximal ideal of  $\widehat{A}$ ,  $\widehat{\pi} = (\pi)$  in  $\widehat{A}$ ,  $\widehat{\pi}$  is the closure of  $\pi$  in  $\widehat{K}$ .

Similarly,  $\widehat{\mathfrak{m}}^n$  is the closure of  $\mathfrak{m}^n$  in  $\widehat{K}$  for any  $n \geq 1$ .

**LEMMA 8.6.** *The map  $A/\mathfrak{m}^n \rightarrow \widehat{A}/\widehat{\mathfrak{m}}^n$  is an isomorphism.*

**Proof.** Easy. ■

**PROPOSITION 8.7.** *Choose a set  $S \ni 0$  of representatives for  $A/\mathfrak{m}$ . The series  $\sum a_i \pi^i$  where every  $a_i \in S$  and  $a_i = 0$  for  $i \ll 0$  converges in  $\widehat{K}$  and each element of  $\widehat{K}$  has a unique representative of the form.*

**Proof.** Easy. ■

**REMARK 8.8.**  $S \ni 0$  is necessary. For example, let  $S^*$  be a set of representatives of  $A/\mathfrak{m} \setminus \{0\}$ . Let  $s_1 \in S^*$ , set  $S := S^* \cup \{s_1 \pi\}$ . Then

$$0 = (s_1 \pi) \cdot \pi^n - (s_1) \cdot \pi^{n+1}, \quad \forall n \in \mathbb{Z}.$$

**EXAMPLE 8.9.** Any element of  $\mathbb{Q}_p$  can be written as

$$\sum a_i p^i, \quad \begin{array}{l} a_i \in \{0, \dots, p-1\}; \\ a_i = 0, \quad \forall i \ll 0. \end{array}$$

**PROPOSITION 8.10.** *We have a natural isomorphism*

$$\widehat{A} \simeq \varprojlim A/\mathfrak{m}^n.$$

**Proof.** Define

$$\widehat{A} \rightarrow \widehat{A}/\widehat{\mathfrak{m}}^n \simeq A/\mathfrak{m}^n.$$

It induces  $\widehat{A} \rightarrow \varprojlim A/\mathfrak{m}^n$ . Define  $\varprojlim A/\mathfrak{m}^n \rightarrow \widehat{A}$  as follows:

$$\overline{a_n} \in A/\mathfrak{m}^n \quad \text{with} \quad \overline{a_{n+1}} \bmod \mathfrak{m}^n = \overline{a_n} \bmod \mathfrak{m}^n.$$

Let  $a_n \in A$  with  $a_n = \overline{a_n} \bmod \mathfrak{m}^n$ . Then  $(a_n)$  is a Cauchy sequence. Define

$$(a_n) \mapsto \lim_{n \rightarrow \infty} a_n \in \widehat{A}.$$

■

### 8.0.2 Newton's lemma

Let  $A$  be a complete discrete valuation ring and  $\pi$  generates its maximal ideal.

**PROPOSITION 8.11.** *Let  $f(x) \in A[x]$  and  $a_0$  be a simple root of  $f(x) \pmod{\pi}$ . Then there exists a unique root  $a$  of  $f(x)$  with  $a \equiv a_0 \pmod{\pi}$ .*

**Proof.** Let

$$U := \{x \in A \mid x \equiv a_0 \pmod{\pi}\} \simeq \pi \cdot A,$$

a complete metric space. Define

$$F: U \longrightarrow U$$

$$x \longmapsto x - \frac{f(x)}{f'(x)}.$$

Since  $a_0$  is a simple root of  $f \pmod{\pi}$ , we have

$$\begin{cases} f(a_0) \equiv 0 \pmod{\pi}; \\ f'(a_0) \not\equiv 0 \pmod{\pi}. \end{cases}$$

Hence, for any  $x \in U$ , we have  $|f'(x)| = 1$  and  $f(x) \equiv 0 \pmod{\pi} \implies F: U \rightarrow U$ .

For  $x_1, x_2 \in U$ ,  $x_2 = x_1 + \Delta$  with  $|\Delta| \leq |\pi|$ . Then

$$f(x_2) = f(x_1) + f'(x_1)\Delta + \varepsilon,$$

where  $\varepsilon \in A \cdot \Delta^2 \implies$

$$|F(x_2) - F(x_1)| = \left| \Delta - \frac{f'(x_1)\Delta + \varepsilon}{f'(x_1)} \right| = \left| \frac{\varepsilon}{f'(x_1)} \right| \leq |\Delta|^2 \leq |\pi| \cdot |\Delta|,$$

$\implies F: U \rightarrow U$  is a contraction map  $\implies F$  has a unique fixed point. ■

**THEOREM 8.12.** *Let  $f(x) \in A[x]$ , and  $a_0 \in A$  satisfying  $|f(a_0)| < |f'(a_0)|^2$ . Then there exists a unique root  $a$  of  $f(x)$  s.t.*

$$|a - a_0| \leq \left| \frac{f(a_0)}{f'(a_0)^2} \right|.$$

**Proof.** Newton's method. ■

## 8.1 Note on 20251217

### 8.1.1 Hensel's lemma

Let  $A$  be a complete discrete valuation ring and  $\mathfrak{m}$  its maximal ideal.

**THEOREM 8.13 (Hensel's lemma).** *Let  $k = A/\mathfrak{m}$  be the residue field of  $K$ . For  $f(x) \in A[x]$ , write  $\bar{f}(x)$  its image in  $k[x]$ .*

*Consider a monic polynomial  $f(x) \in A[x]$ . If  $\bar{f}(x)$  factors as  $\bar{f} = g_0 h_0$  with  $g_0, h_0$  monic and relatively prime in  $k[x]$ , then  $f$  itself factors as  $f = gh$  with  $g$  and  $h$  monic s.t.  $\bar{g} = g_0, \bar{h} = h_0$ .*

*Moreover,  $g$  and  $h$  are uniquely determined and  $(g, h) = A[x]$  (called strictly coprime).*

**LEMMA 8.14.** *If  $f, g \in A[x]$  s.t.  $\bar{f}, \bar{g}$  are relatively prime and  $f$  is monic, then  $(f, g) = A[x]$ .*

*More precisely, there exist  $u, v \in A[x]$  with  $\deg u < \deg g, \deg v < \deg f$  s.t.  $uf + vg = 1$ .*

**Proof.** Set

$$M := A[x] / (f, g).$$

As  $f$  is monic,  $M$  is a finitely generated  $A$ -mod. As  $(\bar{f}, \bar{g}) = k[x]$ , we have

$$(f, g) + \mathfrak{m}A[x] = A[x].$$

$\implies \mathfrak{m}M = M$ . Nakayama's lemma  $\implies M = 0$ . Then there exists  $u, v \in A[x]$  s.t.

$$uf + vg = 1.$$

If  $\deg v \geq \deg f$ , write  $v = fq + r$  with  $\deg r < \deg f \implies$

$$(u + qg)f + rg = 1,$$

where  $\deg(u + qg) < \deg g$ , and  $\deg r < \deg f$ . ■

We next prove the uniqueness.

**LEMMA 8.15.** *Suppose  $f = gh = g'h'$  with  $g, h, g', h'$  monic and  $\bar{g} = \bar{g}', \bar{h} = \bar{h}'$  with  $\bar{g}, \bar{h}$  relatively prime. Then  $g = g', h = h'$ .*

**Proof.** Lemma 8.14 implies  $(g, h') = A[x] \implies$  there exists  $r, s \in A[x]$  s.t.  $gr + h's = 1 \implies$

$$g' = g'gr + g'h's = g'gr + ghs \implies g \mid g'.$$

As  $g, g'$  same degree and monic,  $g = g'$ . ■

Finally, we prove the existence of  $g$  and  $h$ .

**Proof.** There exist monic polynomials  $g_0, h_0 \in A[x]$  s.t.

$$f - g_0 h_0 \in \pi \cdot A[x],$$

where  $\mathfrak{m} = (\pi)$ . Suppose we have constructed monic polynomials  $g_n, h_n$  s.t.

$$f - g_n h_n \equiv 0 \pmod{\pi^{n+1}},$$



and

$$g_n \equiv g_0 \pmod{\pi}, \quad h_n \equiv h_0 \pmod{\pi}.$$

We want to find  $u, v \in A[x]$  with  $\deg u < \deg g_0$  and  $\deg v < \deg h_0$ , s.t.

$$f - (g_n + \pi^{n+1}u)(h_n + \pi^{n+1}v) \equiv 0 \pmod{\pi^{n+2}},$$

i.e.

$$(f - g_n h_n) - \pi^{n+1}(uh_n + g_n v) \equiv 0 \pmod{\pi^{n+2}}.$$

Since  $f - g_n h_n = \pi^{n+1} \cdot r$ , where  $r \in A[x]$ , this is equivalent to

$$r \equiv uh_n + g_n v \pmod{\pi}.$$

Because  $g_0, h_0$  are monic and relatively prime  $(h_n, g_n) = A[x]$ , by Lemma 8.14  $\implies \exists$  such  $u, v$ . ■

**REMARK 8.16.** Theorem 8.13 implies: A factorization of  $f$  into product of relatively prime polynomials in  $k[x]$  lifts to a factorization in  $A[x]$ .

For example, in  $\mathbb{F}_p[x]$ ,  $x^p - x$  splits into  $p$  distinct factors, so it also splits in  $\mathbb{Z}_p[x] \implies \mathbb{Z}_p$  contains the  $(p-1)$ -th roots of 1.

More generally, if  $K$  has a residue field  $k$  with  $q$  elements, then  $K$  contains  $q$  roots of the polynomials  $x^q - x$ . Let  $S$  be the set of these roots, then

$$a \mapsto \bar{a} \quad : \quad S \longrightarrow k$$

is a bijection preserving multiplication. The elements of  $S$  are called the Teichmüller representatives for the elements of the residue field.

### 8.1.2 Extensions of NA absolute values

**THEOREM 8.17.** Let  $K$  be complete w.r.t. a discrete absolute value  $|\cdot|_K$ . Let  $L$  be a finite separable extension of  $K$  of degree  $n$ . Then  $|\cdot|_K$  extends uniquely to a discrete absolute value  $|\cdot|_L$  on  $L$  and  $L$  is complete for the extended absolute value:  $\forall b \in L$ ,

$$|b|_L = |\mathrm{Nm}_{L/K}(b)|_K^{1/n}.$$

**Proof.** Let  $A$  be the valuation ring of  $K$ ,  $B$  the integral closure of  $A$  in  $L$ , and  $\mathfrak{p}$  the maximal ideal of  $A$ . Then  $B$  is a Dedekind domain. Suppose there exist prime ideals  $\beta_1 \neq \beta_2$  in  $B$ . Then there exists  $b \in \beta_1 \setminus \beta_2$ .

Let  $f(x) \in A[x]$  be the minimal polynomial for  $b$ . Then Hensel's lemma implies  $\bar{f}(x) = \bar{g}(x)^l$  power of an irreducible polynomial in  $k[x]$  where  $k = A/\mathfrak{p}$ . Both  $\beta_1 \cap A[b]$  and  $\beta_2 \cap A[b]$  are distinct prime ideals containing  $\mathfrak{p} \implies$

$$\beta_1 \cap A[b] / \mathfrak{p}A[b], \quad \beta_2 \cap A[b] / \mathfrak{p}A[b]$$

are distinct prime ideals of

$$A[b] / \mathfrak{p}A[b] = (A/\mathfrak{p})[x] / (\bar{g}(x)^l) = k[x] / (\bar{g}(x)^l),$$

which only has one prime ideal  $(\bar{g}(x))$ . Contradiction. Hence,  $B$  has only one prime ideal  $\implies B$  is a DVR with a unique prime ideal  $\beta$ .

Therefore,  $|\cdot|_K$  extends to a unique absolute value  $|\cdot|_L$  on  $L$ , which corresponds to  $\beta$ .

Similarly,  $|\cdot|_K$  extends uniquely to an absolute value  $|\cdot|_{L'}$  on a Galois closure  $L'$  of  $L$ . Any  $\sigma \in \text{Gal}(L'/K)$  defines an absolute value on  $L$  by  $b \mapsto |\sigma(b)|_{L'}$ . Uniqueness  $\implies |\sigma(b)|_{L'} = |b|_L = |b|_{L'} \implies$

$$|\text{Nm}(b)|_K = \left| \prod_{\sigma} \sigma(b) \right|_{L'} = |b|_L^n \implies |b|_L = |\text{Nm}_{L/K}(b)|^{1/n}.$$

Finally, we show that  $(L, |\cdot|_L)$  is complete. Let  $e_1, \dots, e_n$  be a basis of  $B$  as  $A$ -mod. Assume  $\mathfrak{p} = \beta^e$ . Consider

$$\begin{aligned} \phi: \bigoplus K e_i &\longrightarrow L \\ (a_i) &\longmapsto \sum a_i e_i \\ \|(a_i)\| &:= \max_i |a_i| \quad |\cdot|_L. \end{aligned}$$

Check

- $\bigoplus K e_i$  is complete;
- $\phi$  is bounded.

Only need to show: if  $\left| \sum a_i e_i \right|_L$  is small, then  $\max_i |a_i|$  is small.

If  $\beta^l \mid b = \sum a_i e_i$ , then  $\mathfrak{p}^{\lfloor l/e \rfloor} \mid b = \sum a_i e_i \implies \mathfrak{p}^{\lfloor l/e \rfloor} \mid a_i$  for any  $i$ .  
 $\implies (L, |\cdot|_L)$  is complete. ■

**COROLLARY 8.18.** *Let  $\Omega$  be a possibly infinite separable algebraic extension of  $K$ , then  $|\cdot|_K$  extends uniquely to an absolute value  $|\cdot|_{\Omega}$  on  $\Omega$ .*

**Proof.**  $\Omega = \bigcup L$ , over all  $L$  subfields of  $\Omega$  with  $L/K$  finite extension. ■

**REMARK 8.19.** *In this corollary,  $|\cdot|_{\Omega}$  is still NA, but it need not be complete and need not be discrete.*

*However, if  $\Omega$  is algebraically closed, then  $\widehat{\Omega}$  is still algebraically closed.*

**EXAMPLE 8.20.** •  $\overline{\mathbb{Q}_p}$  is not discrete:

$$|p^{1/n}| = |p|^{1/n} = p^{-1/n} \rightarrow 1.$$

- $\overline{\mathbb{Q}_p}$  is complete;  $\dim_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$  countable.

Define  $\mathbb{C}_p := \widehat{\overline{\mathbb{Q}_p}}$ .

**COROLLARY 8.21.** *Let  $K, L$  as in Theorem 8.17. Then  $n = ef$ , where  $n = [L : K]$ ,  $e$  is the ramification index, and  $f$  is the degree of residue field extension.*

When  $e = n$ , so  $\mathfrak{p}B = \beta^n$ , we say  $L$  is totally ramified over  $K$ .

When  $f = n$ , we say  $L$  is unramified over  $K$ .

## 8.2 Note on 20251222

### 8.2.1 Newton's polygon

Let  $(K, |\cdot|)$  be complete, discrete, where  $\text{ord}: K^\times \rightarrow \mathbb{Z}$  is the coresponding valuation. It extends to a valuation  $\text{ord}: \overline{K}^\times \rightarrow \mathbb{Q}$ .

**DEFINITION 8.22.** For a polynomial

$$f(x) = x^n + a_1x^{n-1} + \cdots + a_n, \quad a_i \in K,$$

define the Newton polygon of  $f(x)$  to be the lower convex hull of the set to points

$$P_i := (i, \text{ord}(a_i)), \quad i = 0, \dots, n.$$




Figure 1: Newton polygon

**PROPOSITION 8.23.** Assume  $\text{char } K = 0$ . Suppose that the Newton polygon of  $f(x) \in K[x]$  has segments of  $x$ -length  $n_i$  and slope  $s_i$ . Then  $f(x)$  has exactly  $n_i$  roots  $\alpha$  in  $\overline{K}$  with  $\text{ord}(\alpha) = s_i$ . Moreover, the polynomial

$$f_i(x) = \prod_{\text{ord}(\alpha_i)=s_i} (x - \alpha_i)$$

has coefficients in  $K$ .

**Proof.** For the first part, don't need  $f(x)$  has coefficients in  $K$ . It suffices to prove the following statement:

Let  $f(x) = \prod (x - \alpha_j)$ , if exactly  $n_j$  of  $\alpha_j$ 's have order  $s_j$ , then the Newton polygon of  $f(x)$  has a segment of slope  $s_j$  and  $x$ -length  $n_j$ .

Induction on  $n = \deg f$ . If  $n = 1$ , obvious. Assume it holds for  $n$ . Let

$$g(x) = (x - \alpha)f(x) = x^{n+1} + b_1x^n + \cdots + b_{n+1},$$

where  $b_i = a_i - \alpha a_{i-1}$ .

May assume  $\text{ord } \alpha \leq s_1$ . Then for  $i = 1, \dots, n+1$ ,

$$\begin{aligned}
 \text{ord } b_i &= \text{ord}(a_i - a_{i-1}\alpha) \\
 &\geq \min \{ \text{ord } a_i, \text{ord } a_{i-1} + \text{ord } \alpha \} \\
 &\geq \min \{ N(i), N(i-1) + \text{ord } \alpha \} \\
 &\geq N(i-1) + \text{ord } \alpha.
 \end{aligned} \tag{8.11}$$




Figure 2: Newton polygons for  $f$  and  $g$

For those vertex  $i-1$ , the “=” in (8.11) holds, i.e. the Newton polygon for  $g$  is as Figure 2.

Now prove the second statement.

Let  $\alpha$  be any root of  $f(x)$  with  $\text{ord } \alpha = s_i$ . Let  $m_\alpha(x)$  be the minimal polynomial of  $\alpha$ . Then any root  $\alpha'$  of  $m_\alpha(x)$  has  $\text{ord } \alpha' = s_i \implies m_\alpha(x) \mid f(x) \implies f_i(x)$  has coefficients in  $K$ . ■

## 9 Locally compact field

**PROPOSITION 9.1.** *Let  $(K, |\cdot|)$  be complete, discrete. Let  $A$  be the valuation ring of  $K$  and  $\mathfrak{m}$  the maximal ideal of  $A$ . Then  $A$  is compact iff  $A/\mathfrak{m}$  is finite.*

**Proof.** Let  $S$  be the set of representatives for  $A/\mathfrak{m}$ . Then

$$A = \bigsqcup_{x \in S} (x + \mathfrak{m}),$$

where each  $(x + \mathfrak{m})$  is open and nonempty. If  $A$  is compact, then  $\#S < \infty$ .

On the other hand,

$$A = \varprojlim_n A/\mathfrak{m}^n \subseteq \prod_{n \geq 1} A/\mathfrak{m}^n,$$

where the last inclusion is a closed embedding. If  $A/\mathfrak{m}$  is finite, then every  $A/\mathfrak{m}^n$  is finite  $\implies A$  is compact.  $\blacksquare$

**COROLLARY 9.2.** *If  $A/\mathfrak{m}$  is finite, then  $\mathfrak{m}^n$ ,  $1 + \mathfrak{m}^n$  and  $A^\times$  are all compact.*

**DEFINITION 9.3.** A local field is a field  $K$  with a nontrivial absolute value  $|\cdot|$  s.t.  $K$  is locally compact.

### Classification of local fields

- (a) Archimedean case:  $K \simeq \mathbb{C}$  or  $\mathbb{R}$ ;
- (b) NA local field of characteristic 0: finite extension of  $\mathbb{Q}_p$ ;
- (c) NA for char  $p \neq 0$ : isomorphic to the field of formal Laurent series  $k((t))$  over a finite field  $k$ .

#### 9.0.1 Unramified extensions of a field

Let  $(K, |\cdot|)$  be a complete discrete valued field. Assume both  $K$  and the residue field  $k$  are perfect.

Let  $L$  be an algebraic extension of  $K$ . We define

- $B := \{\alpha \in L \mid |\alpha| \leq 1\}$ ;
- $\mathfrak{p} := \{\alpha \in L \mid |\alpha| < 1\}$ ;
- $l := B/\mathfrak{p}$  the residue field of  $L$ .

**PROPOSITION 9.4.** *There is a one-to-one correspondence:*

$$\begin{aligned} \{K' \subset L, \text{ finite and unramified over } K\} &\longleftrightarrow \{k' \subset l, \text{ finite over } k\} \\ K' &\longleftrightarrow k'. \end{aligned}$$

Moreover,

- (a) if  $K' \leftrightarrow k'$  and  $K'' \leftrightarrow k''$ , then  $K' \subset K''$  iff  $k' \subset k''$ ;

(b) if  $K' \leftrightarrow k'$ , then  $K'$  Galois over  $k$  iff  $k'$  is Galois over  $k$ , in which case there is a canonical isomorphism

$$\text{Gal}(K'/K) \simeq \text{Gal}(k'/k).$$

**Proof.** Let  $k'/k$  finite extension. Write  $k' = k[a]$  for  $a \in k' \subset L$ . Set  $f_0(x)$  to be the minimal polynomial of  $a$  over  $k$ . Let  $f(x)$  be any lifting of  $f_0(x)$  in  $A[x]$ . As  $a$  is a simple root of  $f_0(x)$  (since  $K$  and  $k$  are perfect), Newton's lemma shows there exists a **unique**  $\alpha \in L$  s.t.  $f(\alpha) = 0$  with  $\alpha \equiv a \pmod{\mathfrak{p}}$ . Then  $K[\alpha] \subset L$  is a finite unramified extension over  $K$  with residue field  $k' \implies K' \rightarrow k'$  surjective.

Let  $K' \subset L$ ,  $K'/K$  finite unramified with residue field  $k'$ . Then

$$[K' : K] = [k' : k]. \quad (9.12)$$

Newton's method implies there exists  $\alpha_1 \in K' \subseteq L$  s.t.  $f(\alpha_1) = 0$  and  $\alpha_1 \equiv \alpha \pmod{\mathfrak{p}} \implies \alpha_1 = \alpha \implies K' \supset K[\alpha]$ . By (9.12),  $[K' : K] = [k' : k] = [K(\alpha) : K] \implies K' = K[\alpha] \implies$  injective.

(a):  $K' \subset K'' \implies k' \subset k''$ . Assume  $k' \subset k''$ . Then there exists a unique  $K''' \subset K''$  with  $k''' = k'$ . Since both  $K'''$  and  $K' \subset L$ , the uniqueness implies  $K''' = K' \implies K' \subset K'' \implies$  (a) holds.

(b): Assume  $K'/K$  Galois. Since  $\text{Gal}(K'/K)$  preserves the valuation ring  $A'$  in  $K'$  and the maximal ideal, we get  $\text{Gal}(K'/K) \rightarrow \text{Aut}(k'/k)$ . Write  $k' = k[a]$  with  $g(x) \in A[x]$  s.t.  $\bar{g}(x) \in k[x]$  is the minimal polynomial of  $a \implies g$  is irreducible.

Let  $\alpha \in A'$  be the unique root of  $g(x)$  s.t.  $\bar{\alpha} = a$ . Then  $K'/K$  Galois  $\implies g(x)$  splits in  $K' \implies \bar{g}(x)$  splits in  $k' \implies k'$  Galois over  $k$ . Let  $f = [k' : k] = [K' : K]$ , and let  $\alpha_1, \dots, \alpha_f$  be the roots of  $g(x)$ . Then

$$\{\alpha_1, \dots, \alpha_f\} = \{\sigma(\alpha) \mid \sigma \in \text{Gal}(K'/K)\}.$$

As  $\bar{g}(x)$  is separable,  $\alpha_i \pmod{\mathfrak{p}}$  are distinct  $\implies$  the image of  $\text{Gal}(K'/K)$  in  $\text{Gal}(k'/k)$  has order  $f$ . So  $\text{Gal}(K'/K) \rightarrow \text{Gal}(k'/k)$  is an isomorphism.

Conversely, if  $k'/k$  is Galois, write  $k' = k[a]$  and  $\alpha \in A'$  lifts  $a$ . The formal proof shows  $K' = K[\alpha]$ . Hensel's lemma  $\implies A'$  contains the conjugates of  $a \implies K'$  is Galois over  $K$ . ■

**COROLLARY 9.5.** *There exists an unramified extension  $K_0$  of  $K$  contained in  $L$  that contains all unramified extension of  $K$  in  $L$ .*

*When  $K$  is finite, it is obtained from  $K$  by adjoining all roots of 1 of order prime to  $\text{char } k$ .*

**COROLLARY 9.6.** *The residue field of  $\bar{K}$  is  $\bar{k}$ . There exists a subfield  $K^{\text{un}}$  of  $\bar{K}$  s.t. a subfield  $L$  of  $\bar{K}$  finite over  $K$  is unramified iff  $L \subset K^{\text{un}}$ .*

## 9.1 Note on 20251224

### 9.1.1 Totally ramified extension of $K$

Let  $(K, |\cdot|)$  be a complete discrete valued field, and  $\pi$  the local uniformizing parameter of  $K$ .

**DEFINITION 9.7.** A polynomial  $f(x) \in K[x]$  is called Eisenstein if it is Eisenstein for  $(\pi)$ , i.e.

$$f(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n$$

with

$$\text{ord } a_0 = 0, \quad \text{ord } a_i > 0 \text{ for } a_i > 0, \quad \text{ord } a_n = 1,$$

$\iff |a_0| = 1, |a_i| < 1, |a_n| = |\pi| \iff$  the Newton polygon is as follows:

**PROPOSITION 9.8.**  $L/K$  finite extension of  $K$ . Then  $L/K$  is totally ramified iff  $L = K[\alpha]$  with  $\alpha$  a root of an Eisenstein polynomial.

**Proof.**  $\Leftarrow$ :  $L = K[\alpha]$  with  $\alpha$  a root of an Eisenstein polynomial  $f(x)$  of degree  $n$ . Extend  $\text{ord}$  from  $K$  to  $L$ . Then  $\text{ord}(\alpha) = \frac{1}{n} \implies$  the ramification index of  $L/K \geq n$ .  $[L : K] = n \implies$  totally ramified.

$\Rightarrow$ :  $L$  totally ramified over  $K$ . Let  $\alpha$  be a generator of the maximal ideal of  $\mathcal{O}_L$ . Then  $\text{ord } \alpha = \frac{1}{n}$ . The elements  $1, \alpha, \dots, \alpha^{n-1}$  represent different cosets of  $\text{ord}(K^\times)$  in  $\text{ord}(L^\times)$ , i.e.

$$\text{ord}(L^\times) / \text{ord}(K^\times) = \frac{1}{n} \mathbb{Z} / \mathbb{Z}.$$

$\implies 1, \dots, \alpha^{n-1}$  are  $K$ -linearly independent  $\implies$  they are  $K$ -basis of  $L \implies L = K[\alpha]$ , we have a relation

$$\alpha^n + a_1\alpha^{n-1} + \cdots + a_n = 0, \quad a_i \in K.$$

There exist at least two terms having minimal order. Note that

$$1 = \text{ord}(\alpha^n) \equiv \text{ord}(a_n) \text{ in } \text{ord } L / \text{ord } K = \frac{1}{n} \mathbb{Z} / \mathbb{Z},$$

and  $\text{ord}(a_1\alpha^{n-1}), \dots, \text{ord}(a_n)$  differ from each other in  $\frac{1}{n} \mathbb{Z} / \mathbb{Z}$ . This happens only if  $1 = \text{ord}(\alpha^n) = \text{ord}(a_n)$ , and  $\text{ord}(a_i\alpha^i) \geq 1$  for all  $i = 1, \dots, n-1$ . That is, the polynomial  $x^n + a_1x^{n-1} + \cdots + a_n$  is Eisenstein. ■

**REMARK 9.9.** Let  $L/K$  be a finite totally ramified extension, and  $A, B$  the valuation rings for  $K, L$ . Let  $\pi, \Pi$  be prime elements in  $A, B$ . Then  $B = A[\Pi]$ .

**Proof.**  $B, A$  have the same residue field  $\implies B/(\Pi) = A/(\pi) \implies$

$$B = A + \Pi B = A + \Pi(A + \Pi B) = \cdots = A[\Pi] + \Pi^m B, \quad \forall m \geq 1.$$

The discriminant of  $1, \Pi, \dots, \Pi^{n-1}$  equals to a unit times  $\pi^l$  for some  $l \geq 0 \implies \Pi^c B \subset A[\Pi] \subset B$ . Pick  $m = c$ ,  $A[\pi] \supset A[\Pi] + \Pi^c B = B$ . ■

### 9.1.2 Ramification groups

Let  $L/K$  be a finite Galois extension. Assume the residue field  $k$  of  $K$  is perfect. Then  $\mathcal{G} := \text{Gal}(L/K)$  preserves the absolute value on  $L \implies$  it preserves

$$\begin{aligned} B &= \{\alpha \in L \mid |\alpha| \leq 1\} \\ \mathfrak{p} &= \{\alpha \in L \mid |\alpha| < 1\}. \end{aligned}$$

Write  $\mathfrak{p} = (\Pi)$ . We define a sequence of subgroups

$$\mathcal{G} \supset \mathcal{G}_0 \supset \mathcal{G}_1 \supset \dots$$

by the condition

$$\sigma \in \mathcal{G}_i \iff |\sigma(\alpha) - \alpha| < |\Pi|^i, \quad \forall \alpha \in B.$$

**DEFINITION 9.10.** We call

- $\mathcal{G}_0$  the inertia group;
- $\mathcal{G}_1$  the ramification group;
- $\mathcal{G}_i, i > 1$  the higher ramification groups of  $L$  over  $K$ .

**LEMMA 9.11.** The  $\mathcal{G}_i$  are normal subgroups of  $\mathcal{G}$  and  $\mathcal{G}_i = \{1\}$  for  $i \gg 0$ .

**Proof.** Easy. ■

**THEOREM 9.12.** Let  $L/K$  be a Galois extension. Assume the residue field extension  $l/k$  is separable.

(a) The fixed field of  $\mathcal{G}_0$  is the largest unramified extension  $K_0$  of  $K$  in  $L$  and

$$\mathcal{G}/\mathcal{G}_0 = \text{Gal}(K_0/K) = \text{Gal}(l/k).$$

(b) For any  $i \geq 1$ , we have

$$\mathcal{G}_i = \{\sigma \in \mathcal{G}_0 \mid |\sigma\Pi - \Pi| < |\Pi|^i\}.$$

**Proof.** (a): Let  $K_0$  be the largest unramified extension in  $L$ . Then  $\sigma(K_0)$  is also unramified  $\implies \sigma(K_0) \subset K_0 \implies K_0/K$  Galois.

$$\text{Gal}(K_0/K) \longrightarrow \text{Gal}(l/k)$$

is an isomorphism, and  $\mathcal{G}_0 = \ker(\mathcal{G} \longrightarrow \text{Gal}(l/k)) \implies K_0 = L^{\mathcal{G}_0}$ .

(b): Let  $A_0$  be the valuation ring of  $K_0$ . Then  $B = A[\Pi]$ . Since  $\mathcal{G}_0$  leaves  $A_0$  fixed,  $\sigma \in \mathcal{G}_i$  iff  $|\sigma(\Pi) - \Pi| < |\Pi|^i$ . ■

**COROLLARY 9.13.** We have an exhaustive filtration  $\mathcal{G} \supset \mathcal{G}_0 \supset \dots$  s.t.

- $\mathcal{G}/\mathcal{G}_0 = \text{Gal}(l/k)$ ;
- $\mathcal{G}_0/\mathcal{G}_1 \hookrightarrow l^\times$ ;
- $\mathcal{G}_i/\mathcal{G}_{i+1} \hookrightarrow l, i \geq 1$ .



Therefore, if  $k$  is finite,  $\text{Gal}(L/K)$  is solvable.

**Proof.** Let  $\sigma \in \mathcal{G}_0$ , then  $\sigma(\Pi)$  is a prime element  $\implies \sigma(\Pi) = u\Pi$  where  $u \in B^\times$ . The map

$$\sigma \mapsto u \pmod{\Pi}$$

is a homomorphism  $\mathcal{G}_0 \longrightarrow l^\times$  with kernel  $\mathcal{G}_1$ .

Let  $\sigma \in \mathcal{G}_i$ ,  $i \geq 1$ , then  $|\sigma(\Pi) - \Pi| \leq |\Pi|^{i+1} \implies \sigma(\Pi) = \Pi + a\Pi^{i+1}$  for some  $a \in B$ . The map

$$\sigma \mapsto a \pmod{\mathfrak{p}}$$

is a homomorphism  $\mathcal{G}_i \longrightarrow l$  with kernel  $\mathcal{G}_{i+1}$ . ■

**DEFINITION 9.14.** An extension  $L/K$  is said to be widely ramified if  $p \mid e$  where  $p = \text{char } k$ . Otherwise, it is said to be tamely ramified.

Hence, for a Galois extension  $L/K$ ,

$$L/K \text{ unramified} \iff \mathcal{G}_0 = \{1\},$$

and

$$L/K \text{ tamely ramified} \iff \mathcal{G}_1 = \{1\}.$$

### 9.1.3 Krasner's lemma

Let  $(K, |\cdot|)$  be complete discrete. Extend  $|\cdot|$  to an absolute value on  $\overline{K}$ .

**PROPOSITION 9.15 (Krasner's lemma).** Let  $\alpha, \beta \in \overline{K}$  and assume  $\alpha$  is separable over  $K[\beta]$ . If  $\alpha$  is closer to  $\beta$  than to any other conjugate of  $\alpha$  (over  $K$ ), then  $K[\alpha] \subset K[\beta]$ .

**Proof.** See Milne p.132. ■

Now assume  $\text{char } K = 0$ . For  $h(x) = \sum c_i x^i$ , define  $\|h\| := \max\{|c_i|\}$ . If  $h(x)$  monic,

$$h(x) = x^n + \sum_{i=0}^{n-1} c_i x^i.$$

Let  $\alpha$  be any root of  $h(x)$   $|\alpha| \leq \|h\|$ .

Fix a monic irreducible polynomial  $f(x)$  in  $K[x]$ . Let  $g(x)$  be another irreducible polynomial and suppose  $\|f - g\|$  is small. For any root  $\beta$  of  $g(x)$ ,  $|f(\beta) - g(\beta)|$  is small. Write  $f(\beta) = \prod (\beta - \alpha_i)$  where  $\alpha_i$  are roots of  $f \implies$  may assume

$$|\beta - \alpha_1| \leq |f(\beta)|^{1/n}.$$

For  $\|f - g\|$  small enough,

$$|f(\beta)| < \left( \min_{i \neq 1} |\alpha_i - \alpha_1| \right)^n$$

$\implies |\beta - \alpha_1| < |\alpha_1 - \alpha_i|$  for  $i \neq 1$  (called  $\beta$  belongs to  $\alpha_1$ ). Now Krasner's lemma implies  $K[\alpha_1] \subset K[\beta]$ .

If  $\deg f = \deg g$ , then  $K[\alpha_1] = K[\beta]$ .

**PROPOSITION 9.16.** Let  $f(x)$  be a monic irreducible polynomial of  $K[x]$ . Then any monic polynomial  $g(x) \in K[x]$  sufficiently close to  $f(x)$  is also irreducible, and each root  $\beta$  of  $g(x)$  belongs to some root  $\alpha$  of  $f(x)$ . We have  $K[\alpha] = K[\beta]$ .