

NAME : **Mrs. MEGHANA J G** REG # : 25072546 / 52754

AGE/SEX : 22 Yrs / Female DATE OF REPORT : 28-07-2025

REFERRED BY: REFFERRED CENTER: KANGROW HEALTH

TEST PARAMETER RESULT REFERENCE RANGE

BIOCHEMISTRY

RANDOM BLOOD SUGAR (RBS) RANDOM BLOOD SUGAR 73 mg/dl 70 - 140 mg/dl GOD-POD 70 - 140 mg/dl

NOTE:

Criteria for the diagnosis of diabetes (ADA)

FBS ≥126 mg/dL. Fasting is defined as no caloric intake for at least 8 h OR

2-h PPBS ≥200 mg/dL during an OGTT performed using a glucose load containing the equivalent of 75 g anhydrous glucose dissolved in water **OR** HBA1C ≥6.5% **OR**

RBS ≥200 mg/dI In a patient with classic symptoms of hyperglycemia or hyperglycemic crisis.

Categories of increased risk for diabetes (prediabetes) (ADA)

FBS 100 mg/dL -125 mg/dL **OR** 2-h PPBS 140 mg/dL -199 mg/dL **OR** HBA1C 5.7-6.4%

SERUM CREATININE
SERUM CREATININE

SERUM CREATININE	0.8 mg/dL	0.6 - 1.2 mg/dL
Enzymatic	o	ŭ

SERUM CALCIUM

LIVER FUNCTION TEST(LFT)		
TOTAL BILIRUBIN DIAZO	0.8 mg/dl	0.1 - 1.2 mg/dl
DIRECT BILIRUBIN DIAZO	0.2 mg/dl	0 - 0.4 mg/dl
INDIRECT BILIRUBIN Calculated	0.60 mg/dl	0.2 - 1.2 mg/dl
S G O T	28.4 U/L	up to 40 U/L
S G P T	32.3 U/L	up to 40 U/L
ALKALINE PHOSPHATASE	81.0 U/L	42 - 98 U/L
TOTAL PROTEIN BIURET	6.8 g/dl	6 - 8 g/dl
S.ALBUMIN BCG	4.1 g/dl	3 - 4.5 g/dl
S.GLOBULIN Calculated	2.7 g/dl	1.5 - 3 g/dl
A/G RATIO Calculated	1.5	1.3 - 1.4



NAME : **Mrs. MEGHANA J G** REG # : 25072546 / 52754

AGE/SEX : 22 Yrs / Female DATE OF REPORT : 28-07-2025

REFERRED BY: REFFERRED CENTER: KANGROW HEALTH

TEST PARAMETER RESULT REFERENCE RANGE

CLINICAL PATHOLOGY

URINE ROUTINE	
ALBUMIN	NIL
SUGAR	NIL
MICROSCOPY	
PUS CELLS	5 - 6 /hpf
EPITHELIAL CELLS	3 - 4 /hpf
R.B.C	NIL
CAST	ABSENT
CRYST	ABSENT
MISC	ABSENT



NAME : **Mrs. MEGHANA J G** REG # : 25072546 / 52754

AGE/SEX : 22 Yrs / Female DATE OF REPORT : 28-07-2025

REFERRED BY: REFFERRED CENTER: KANGROW HEALTH

TEST PARAMETER RESULT REFERENCE RANGE

HAEMATOLOGY

COMPLETE BLOOD COUNT (CBC) ELECTRIC IMPEDANCE METHOD		
HAEMOGLOBIN	13.9 gm/dl	12 - 15 gm/dl
P.C.V (PACKED CELL VOLUME) /	41.7 %	37 - 47 %
HAEMATOCRIT		
R.B.C COUNT	4.63 mill/cumm	4.5 - 5.5 mill/cumm
M.C.V (Mean Corpuscular Volume)	80.0 fl	80 - 98 fl
M.C.H (Mean Corpuscular Haemoglobin)	29.1 pg	26 - 33 pg
M.C.H.C (Mean Corpuscular Haemoglobin	38.0 %	32 - 38 %
concentration)		
TOTAL W.B.C	8200 cells/cumm	4000 - 11000 cells/cumm
NEUTROPHILS	60 %	40 - 70 %
LYMPHOCYTES	35 %	20 - 45 %
EOSINOPHILS	02 %	0 - 6 %
MONOCYTES	03 %	0 - 10 %
BASOPHIL	00 %	0 - 1 %
PLATELET COUNT	1.95 Lakhs/cumm	1.5 - 4.5 Lakhs/cumm



NAME : **Mrs. MEGHANA J G** REG # : 25072546 / 52754

AGE/SEX : 22 Yrs / Female DATE OF REPORT : 28-07-2025

REFERRED BY: REFFERRED CENTER: KANGROW HEALTH

TEST PARAMETER RESULT REFERENCE RANGE

IMMUNO ASSAY

THYROID PROFILE TEST		
TOTAL TRIIODOTHYRONINE (T3)	1.09 ng/mL	0.69 - 2.15 ng/mL
TOTAL THYROXINE (T4)	11.8 μg/dL	5.2 - 12.7 μg/dL
THYROID STIMULATING HORMONE (TSH)	1.16 µIU/ml	0.4 - 4.5 μlU/ml

Age specific reference ranges

T3 T4		TSH			
Age	Reference Intervals (ng/mL)	Age	Reference Intervals (μg/dL)	Age	Reference Intervals (µIU/mL)
1 - 3 Days	1.00 – 7.40	1- 3 Days	11.8 - 22.6	0 - 4 Days	1.0 - 39.0
1 - 11 Months	1.05 – 2.45	1- 2 Weeks	9.9 - 16.6	2 wks -5 m	1.7 – 9.1
1 - 5 Years	1.05 – 2.69	1 - 4 Months	7.2 - 14.4	6 m – 20 Yrs	0.7 – 6.4
6 – 10 Year	0.94 – 2.41	4 m - 1 Yr	7.8 - 16.5	> 55 years	0.5 – 8.9
11 - 15 Years	0.82 – 2.13	1 - 5 Yrs	7.3 - 15.0		
15 – 20 years	0.80 – 2.10	5 -10 Yrs	6.4 - 13.3		
PREGNANCY				PREGNANCY	
1st Trimester	0.81 -1.90			1st Trimester	0.1 - 2.5
2nd& 3rd	1.00 - 2.60			2nd Trimester	0.2 - 3.0
Trimester				3rd Trimester	0.3 – 3.0

LIMITATIONS: T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin, so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, steroids may falsely affect the T3 and T4 levels. Normal levels of T4 can also be seen in Hyperthyroid patients with T3 Thyrotoxicosis, hypoproteinemia or ingestion of certain drugs. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy. Autoimmune disorders may produce spurious results. Various drugs can interfere with the test result. TSH has a diurnal rhythm so values may vary if sample collection is done at different times of the day.

Dispatched by: Laboratory **** End of Report **** Printed on 28-07-2025 at 06:10 PM

Dr. Chalana

MBBS, DCP

Consultant Pathologist

^{*} For any clarifications or escalations - Please Contact: +91 9108322333