



Class has been archived by your teacher. You can't add or edit anything.

ICE 7.2



Aman Gupta • Feb 21, 2025 (Edited Feb 21, 2025)

6/8

1. (2 marks) This is about public-key cryptography. Let $p = 17$, $q = 11$. We wish to choose an encryption key, e . Which of those below are a valid key?

- 5
- 16
- 30
- All of the above
- **None of the above (correct)**

2. (2 marks) Continuing with question 1 above, let encryption key, $e = 11$. Then which of those below is a decryption key?

- 3
- 67
- **131 (correct)**
- All of the above
- None of the above

3. (2 marks) Continuing with question 1 above, let the encryption key, $e = 17$, what is ciphertext if the plain text is 3?

- **20 (correct)**
- 16
- 69
- None of the above

4. (2 marks) This question is about launching an attack on public-key cryptosystem, where it is known that the public key = $\{17, 209\}$. Can an intruder compute the private key? If so, what is it?

- No, one cannot determine the private key
- **The private key is $\{53, 209\}$ (correct)**
- The private key is $\{53, 180\}$
- The private key is $\{57, 209\}$
- None of the above



ICE 7.2

Google Forms



Class has been archived by your teacher. You can't add or edit anything.

Your work

Graded

ICE 7.2

Google Forms



 Private comments

[Add comment to Aman Gupta](#)

 Class comments

 [Add comment](#)

