

PP 33 : Change of variables in double integrals, Polar coordinates

1. Consider the transformation $T : [0, 2\pi] \times [0, 1] \rightarrow \mathbb{R}^2$ given by $T(u, v) = (2v \cos u, v \sin u)$.

(a) For a fixed $v_0 \in [0, 1]$, describe the set $\{T(u, v_0) : u \in [0, 2\pi]\}$.

(b) Describe the set $\{T(u, v) : (u, v) \in [0, 2\pi] \times [0, 1]\}$.

2. Let R be the region in \mathbb{R}^2 bounded by the straight lines $y = x, y = 3x$ and $x + y = 4$. Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(u, v) = (u - v, u + v)$. Find the set $S \subset \mathbb{R}^2$ satisfying $T(S) = R$. (x, y) ← *

3. Let R be the region in \mathbb{R}^2 bounded by the curve defined in the polar co-ordinates $r = 1 - \cos \theta, 0 \leq \theta \leq \pi$ and the x-axis. Consider the transformation $T : [0, \pi] \times [0, 1] \rightarrow \mathbb{R}^2$ defined by $T(r, \theta) = (r \cos \theta, r \sin \theta)$. Let S be the subset of $[0, \pi] \times [0, 1]$ satisfying $T(S) = R$. Sketch the regions S and R .

4. Using the change of variables $u = x + y$ and $v = x - y$, show that $\int_0^1 \int_0^x (x - y) dy dx =$

$$\int_0^1 \int_v^{2-v} \frac{v}{2} du dv.$$

5. Let R be the region bounded by $x = 0, x = 1, y = x$ and $y = x + 1$. Show that $\iint_R \frac{dx dy}{\sqrt{xy - x^2}} =$

$$\left(\int_0^1 \frac{du}{\sqrt{u}} \right) \left(\int_0^1 \frac{dv}{\sqrt{v}} \right).$$

6. Show that $\int_0^1 \int_0^{1-x} e^{\frac{x-y}{x+y}} dx dy = \frac{1}{2} \int_0^1 \int_{-v}^v e^{\frac{u}{v}} du dv = \frac{1}{2} \sinh(1)$. ?

7. Find the region R in \mathbb{R}^2 satisfying $\int_{\frac{1}{\sqrt{2}}}^1 \int_{\sqrt{1-x^2}}^x xy dy dx + \int_1^{\sqrt{2}} \int_0^x xy dy dx + \int_{\sqrt{2}}^2 \int_0^{\sqrt{4-x^2}} xy dy dx = \iint_R xy dx dy$. Evaluate $\iint_R xy dx dy$. Wrong interpretation

8. Convert $\int_0^1 \int_{x^2}^x dy dx$ in to an iterated integral involving polar coordinates.

9. Evaluate

(a) $\int_0^1 \int_0^{1-y} \sqrt{x+y} (y-2x)^2 dx dy.$

(b) $\int_0^{\frac{1}{\sqrt{2}}} \int_y^{\sqrt{1-y^2}} (x+y) dx dy.$

(c) $\int_1^2 \int_0^y \frac{1}{(x^2+y^2)^{\frac{3}{2}}} dx dy.$

(d) $\int_0^2 \int_0^{\sqrt{2x-x^2}} \sqrt{x^2+y^2} dy dx.$

10. Let $R = \{(x, y) \in \mathbb{R}^2 : 9x^2 + 4y^2 \leq 1\}$. Evaluate $\iint_R \cos(9x^2 + 4y^2) dx dy$.

11. Find the volume of the solid bounded by the surfaces $z = 3(x^2 + y^2)$ and $z = 4 - (x^2 + y^2)$.

* At least one of two integral limits is variable form

12. Find the volume of the solid in the first octant bounded below by the surface $z = \sqrt{x^2 + y^2}$ and above by $x^2 + y^2 + z^2 = 8$ as well as the planes $y = 0$ and $y = x$.

Practice Problems 33: Hints/Solutions

1. (a) If $x = 2v_0 \cos u$ and $y = v_0 \sin u$ then $\frac{x^2}{4} + \frac{y^2}{1} = v_0^2$. The set $\{T(u, v_0) : u \in [0, 2\pi]\}$ is an ellipse.
 (b) The set is the region enclosed by the ellipse $\frac{x^2}{4} + \frac{y^2}{1} = 1$.
2. If $x = u - v$ and $y = u + v$ then $y = x \Rightarrow v = 0$, $y = 3x \Rightarrow v = \frac{u}{2}$ and $x + y = 4 \Rightarrow u = 2$. The region S is bounded by the lines $v = 0$, $v = \frac{u}{2}$ and $u = 2$ in the uv -plane. See Figure 1.
3. See Figure 2.
4. Note that $\int_0^1 \int_0^x (x - y) dy dx = \iint_R (y - x) dx dy$ where R is the region in xy -plane bounded by the lines $y = x$, $x = 1$ and $y = 0$. Since $x = \frac{1}{2}(u + v)$ and $y = \frac{1}{2}(u - v)$, $y = 0 \Rightarrow u = v$, $x = 1 \Rightarrow u + v = 2$ and $x = y \Rightarrow v = 0$. Therefore $\iint_R (y - x) dx dy = \iint_S v \frac{\partial(x, y)}{\partial(u, v)} du dv$ where S is the region in the uv -plane bounded by the lines $u = v$, $v + v = 2$ and $v = 0$.
5. Take $u = x$ and $v = y - x$. Then $y = x \Rightarrow v = 0$ and $y = x + 1 \Rightarrow v = 1$. Therefore $\iint_R \frac{dx dy}{\sqrt{xy - x^2}} = \iint_S \frac{1}{\sqrt{uv}} \frac{\partial(x, y)}{\partial(u, v)} du dv$ where S is the region in the uv -plane bounded by the lines $u = 0$, $u = 1$, $v = 0$ and $v = 1$.
6. Consider $u = x - y$ and $v = x + y$. Then $\int_0^1 \int_0^{1-x} e^{\frac{x-y}{x+y}} dx dy = \iint_S e^{\frac{u}{v}} \frac{\partial(x, y)}{\partial(u, v)} du dv$ where S is the region in the uv -plane bounded by the lines $u = -v$, $u = v$ and $v = 1$.
7. See Figure 3. By polar coordinates, $\iint_D xy dx dy = \int_0^{\frac{\pi}{4}} \int_1^2 r^3 \cos \theta \sin \theta dr d\theta = \frac{15}{4} \int_0^{\frac{\pi}{4}} \sin \theta \cos \theta d\theta$.
8. The integral becomes $\iint_D dx dy$ where D is the region in the first quadrant in \mathbb{R}^2 bounded by the line $y = x$ and the curve $y = x^2$. The equation $y = x^2$ can be converted in polar as $r \sin \theta = r^2 \cos^2 \theta$ which implies $r = \tan \theta \sec \theta$. Therefore $\iint_D dx dy = \int_0^{\frac{\pi}{4}} \int_0^{\sec \theta \tan \theta} r dr d\theta$.
9. (a) Note that $\int_0^1 \int_0^{1-y} \sqrt{x+y}(y-2x)^2 dx dy = \iint_R \sqrt{x+y}(y-2x)^2 dx dy$ where R is the region bounded by the lines $x = 0$, $y = 0$ and $x + y = 1$. Consider $u = x + y$ and $v = y - 2x$. Then $x = 0 \Rightarrow v = u$, $y = 0 \Rightarrow v = -2u$ and $x + y = 1 \Rightarrow u = 1$. Therefore $\iint_R \sqrt{x+y}(y-2x)^2 dx dy = \int_0^1 \int_{-2u}^u \sqrt{uv}^2 \frac{1}{3} dv du$.
 (b) The given integral becomes $\iint_R (x + y) dx dy$ where R is the region bounded by the lines $y = 0$, $y = x$ and the circle $x^2 + y^2 = 1$. By polar coordinates $\iint_R (x + y) dx dy = \int_0^{\frac{\pi}{4}} \int_0^1 (r \cos \theta + r \sin \theta) r dr d\theta$.

(c) See Figure 4. The given integral becomes $\int_0^{\frac{\pi}{4}} \int_{\sec \theta}^{2 \sec \theta} \frac{1}{r^3} r dr d\theta$.

(d) See Figure 5. The given integral becomes $\iint_R \sqrt{x^2 + y^2} dx dy$ where R is the region in the first quadrant bounded by the circle $(x - 1)^2 + y^2 = 1$ and the x -axis. The points on the circle $y^2 = 2x - x^2$ is represented by $r = 2 \cos \theta$ in polar coordinates.

Therefore the integral is given by $\int_0^{\frac{\pi}{2}} \int_0^{2 \cos \theta} r r dr d\theta$.

10. Take $x = \frac{r}{3} \cos \theta$ and $y = \frac{r}{2} \sin \theta$. Then $\frac{\partial(x,y)}{\partial(r,\theta)} = \frac{r}{6}$. Therefore $\iint_R \cos(9x^2 + 4y^2) dx dy =$

$$\int_0^{2\pi} \int_0^1 \cos(r^2) \frac{r}{6} dr d\theta = \int_0^{2\pi} \int_0^1 \cos u \frac{du}{12} d\theta.$$

11. The intersection of the surfaces is the set $\{(x, y, 3) : x^2 + y^2 = 1\}$. Therefore the volume is given by $\iint_R (4 - x^2 - y^2 - 3(x^2 + y^2)) dx dy$ where R is the region in \mathbb{R}^2 enclosed by the

circle $x^2 + y^2 = 1$. By polar coordinate the integral becomes $\int_0^{2\pi} \int_0^1 (4 - 4r^2) r dr d\theta$.

12. The given solid lies above the region R where R is in the first quadrant in \mathbb{R}^2 bounded by the circle $x^2 + y^2 = 4$ and the lines $y = x$ and $y = 0$. Therefore the required volume is

$$\text{given by } \iint_R (\sqrt{8 - x^2 - y^2} - \sqrt{x^2 + y^2}) dx dy = \int_0^{\frac{\pi}{4}} \int_0^2 (\sqrt{8 - r^2} - r) r dr d\theta.$$