

# MA 322: Scientific Computing

## Lecture - 5



# Linear Curve Fitting

# Linear Curve Fitting

- Let  $f$  be real valued function.

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) =$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) =$$



# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 =$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \quad \&$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \quad \& \quad \text{then} \quad b =$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \quad \& \quad \text{then} \quad b = f(x_0) - \frac{(f(x_1) - f(x_0))}{(x_1 - x_0)}x_0.$$

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \quad \& \quad \text{then} \quad b = f(x_0) - \frac{(f(x_1) - f(x_0))}{(x_1 - x_0)}x_0.$$

- If we estimate  $f(x)$  through  $P_1(x)$ , then it is known as linear curve fitting.

# Linear Curve Fitting

- Let  $f$  be real valued function.
- Suppose data  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  are given..
- Then, we can find a polynomial  $P_1(x) = a_0x + a_1$  of degree one such that

$$P_1(x_0) = f(x_0) \quad \text{and} \quad P_1(x_1) = f(x_1).$$

Because, from the conditions, we have

$$f(x_0) = P_1(x_0) = a_0x_0 + b \quad \& \quad f(x_1) = P_1(x_1) = a_0x_1 + b.$$

Subtracting above relations, we obtain

$$a_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \quad \& \quad \text{then} \quad b = f(x_0) - \frac{(f(x_1) - f(x_0))}{(x_1 - x_0)}x_0.$$


- If we estimate  $f(x)$  through  $P_1(x)$ , then it is known as linear curve fitting. Roughly, in linear curve fitting, we find a linear polynomial which intersect given curve at at least two points.

# Linear Curve Fitting Contd..



# Linear Curve Fitting Contd..

Geometrically, linear curve fitting can be described by following figure



# Quadratic Curve Fitting

# Quadratic Curve Fitting

- Similarly, in quadratic curve fitting,

# Quadratic Curve Fitting

- Similarly, in quadratic curve fitting, we find a polynomial of degree two


# Quadratic Curve Fitting

- Similarly, in quadratic curve fitting, we find a polynomial of degree two which intersect given curve at at least three points.

# Quadratic Curve Fitting


- Similarly, in quadratic curve fitting, we find a polynomial of degree two which intersect given curve at at least three points.

Geometrically, quadratic curve fitting can be described by following figure



# Cubic Curve Fitting

# Cubic Curve Fitting





# Motivation to Finite Differences

# Motivation to Finite Differences

- What is Finite Difference (FD)?

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$

# Motivation to Finite Differences

- What is Finite Difference (FD)?
- Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference,

# Motivation to Finite Differences

- What is Finite Difference (FD)?
- Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations

# Motivation to Finite Differences

- What is Finite Difference (FD)?
- Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.



# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting
  - From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting
  - From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial  $P_n$  of degree  $n$

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting

- From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial  $P_n$  of degree  $n$  such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting

- From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial  $P_n$  of degree  $n$  such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

The value  $P_n(x)$ ,

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting

- From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial  $P_n$  of degree  $n$  such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

The value  $P_n(x)$ , for  $x \neq x_i$ ,

# Motivation to Finite Differences

- What is Finite Difference (FD)?
  - Finite difference is a mathematical tool used to study the differences that take place in the value of a function  $y = f(x)$  from a given set of values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1}.$$

More precisely, in finite difference, a function is studied with the help of given set of observations even the function is not known explicitly.

- General Result on Polynomial Fitting

- From a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

we find a polynomial  $P_n$  of degree  $n$  such that

$$P_n(x_i) = f(x_i), \quad 0 \leq i \leq n.$$

The value  $P_n(x)$ , for  $x \neq x_i$ , is called estimate of  $f(x)$ .

# Error in Polynomial Fitting to a Curve



# Error in Polynomial Fitting to a Curve


- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal.

# Error in Polynomial Fitting to a Curve

- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal. Geometrically,


# Error in Polynomial Fitting to a Curve

- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal. Geometrically, a fitted curve and function  $f$  can be described by following figure



# Error in Polynomial Fitting to a Curve

- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal. Geometrically, a fitted curve and function  $f$  can be described by following figure




- Although, for non-polynomial function  $f$ , we have

$$P_n(x) \neq f(x) \text{ for } x \neq x_i, \quad 0 \leq i \leq n.$$

# Error in Polynomial Fitting to a Curve

- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal. Geometrically, a fitted curve and function  $f$  can be described by following figure




- Although, for non-polynomial function  $f$ , we have

$$P_n(x) \neq f(x) \text{ for } x \neq x_i, \quad 0 \leq i \leq n.$$

But, the graphs suggest that  $f(x)$  and  $P_n(x)$  are very closed to each other.

# Error in Polynomial Fitting to a Curve

- In general, the estimate  $P_n(x)$  and  $f(x)$  are not equal. Geometrically, a fitted curve and function  $f$  can be described by following figure



- Although, for non-polynomial function  $f$ , we have

$$P_n(x) \neq f(x) \text{ for } x \neq x_i, \quad 0 \leq i \leq n.$$

But, the graphs suggest that  $f(x)$  and  $P_n(x)$  are very closed to each other. Interpolation is a process of constructing such polynomial  $P_n(x)$ .

# Exact Polynomial Fitting to a Curve

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$



# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

- The answer is yes,

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

- The answer is yes, when the function  $f$  is a polynomial of degree  $n$ .

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

- The answer is yes, when the function  $f$  is a polynomial of degree  $n$ .

Geometrically,

# Exact Polynomial Fitting to a Curve

- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

- The answer is yes, when the function  $f$  is a polynomial of degree  $n$ .

Geometrically, for 10 data points,

# Exact Polynomial Fitting to a Curve


- For a given set of  $n + 1$  values

$$\{(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))\}, \quad x_i > x_{i-1},$$

can we find a polynomial  $P_n$  of degree  $n$  such that  $P_n(x) = f(x)$ .

- The answer is yes, when the function  $f$  is a polynomial of degree  $n$ .

Geometrically, for 10 data points, it can be described by following figure



# Exact Polynomial Fitting to a Curve: An Example



# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given,

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3,

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Due to fundamental theorem, we obtain

$$\Delta^4 P_3(x) = 0$$



# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Due to fundamental theorem, we obtain

$$\Delta^4 P_3(x) = 0 \text{ equivalently } (E - I)^4 f(x) = 0.$$

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Due to fundamental theorem, we obtain

$$\Delta^4 P_3(x) = 0 \text{ equivalently } (E - I)^4 f(x) = 0.$$

Then, expanding  $(E - I)^4$  by binomial theorem, we obtain

$$(E^4 - 4E^3 + 6E^2 - 4E + I)P_3(x) = 0,$$

# Exact Polynomial Fitting to a Curve: An Example

- Consider a cubic polynomial function  $f$  given by

$$f(x) = x^3 + 2x^2 + 3x + 5.$$

Let us generate four data points

$$\{(x, f(x)) : (1, 11), (2, 27), (3, 59), (5, 195)\},$$

and using this data set we wish to estimate  $f(4)$  by curve fitting.

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Due to fundamental theorem, we obtain

$$\Delta^4 P_3(x) = 0 \text{ equivalently } (E - I)^4 f(x) = 0.$$

Then, expanding  $(E - I)^4$  by binomial theorem, we obtain

$$(E^4 - 4E^3 + 6E^2 - 4E + I)P_3(x) = 0,$$

which gives

$$P_3(x + 4h) - 4P_3(x + 3h) + 6P_3(x + 2h) - 4P_3(x + h) + P_3(x) = 0.$$

# Exact Polynomial Fitting to a Curve: An Example Contd..

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x + 4h) - 4P_3(x + 3h) + 6P_3(x + 2h) - 4P_3(x + h) + P_3(x) = 0,$$

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x + 4h) - 4P_3(x + 3h) + 6P_3(x + 2h) - 4P_3(x + h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x+4h) - 4P_3(x+3h) + 6P_3(x+2h) - 4P_3(x+h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x+4h) - 4P_3(x+3h) + 6P_3(x+2h) - 4P_3(x+h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$



# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x + 4h) - 4P_3(x + 3h) + 6P_3(x + 2h) - 4P_3(x + h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$

which gives  $P_3(4) = 113$ .

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x+4h) - 4P_3(x+3h) + 6P_3(x+2h) - 4P_3(x+h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$

which gives  $P_3(4) = 113$ . It is interesting to see that  $f(4) = 113$ .

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x + 4h) - 4P_3(x + 3h) + 6P_3(x + 2h) - 4P_3(x + h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$

which gives  $P_3(4) = 113$ . It is interesting to see that  $f(4) = 113$ .  
Thus  $P_3(4) = f(4)$ .

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x+4h) - 4P_3(x+3h) + 6P_3(x+2h) - 4P_3(x+h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$

which gives  $P_3(4) = 113$ . It is interesting to see that  $f(4) = 113$ .

Thus  $P_3(4) = f(4)$ . Why?

# Exact Polynomial Fitting to a Curve: An Example Contd..

- In the equation

$$P_3(x+4h) - 4P_3(x+3h) + 6P_3(x+2h) - 4P_3(x+h) + P_3(x) = 0,$$

we set  $x = 1$  and  $h = 1$  to have

$$P_3(5) - 4P_3(4) + 6P_3(3) - 4P_3(2) + P_3(1) = 0.$$

Since  $P_3$  agrees with  $f$  at given data points, we have

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

Using given data, we obtain

$$195 - 4P_3(4) + 6 \times 59 - 4 \times 27 + 11 = 0,$$

which gives  $P_3(4) = 113$ . It is interesting to see that  $f(4) = 113$ .

Thus  $P_3(4) = f(4)$ . Why?

- This is due to the fact that the data set is associated with a polynomial of degree three.

# Examples Contd..

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ .

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$



## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given,

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3,

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

and hence,  $P_3(4) = 16.5$ ,

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

and hence,  $P_3(4) = 16.5$ , which is very closed to the actual  $f(4) = 2^4 = 16$ ,

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

and hence,  $P_3(4) = 16.5$ , which is very closed to the actual  $f(4) = 2^4 = 16$ , but not equal.



## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

and hence,  $P_3(4) = 16.5$ , which is very closed to the actual  $f(4) = 2^4 = 16$ , but not equal. Why?

## Examples Contd..

- Consider a non-polynomial function  $f$  given by  $f(x) = 2^x$ . Let us generate four data points

$$\{(x, f(x)) : (1, 2), (2, 4), (3, 8), (5, 32)\},$$

and using this data set we wish to estimate  $f(4)$ .

- Since, four data points are given, we can find a polynomial  $P_3$  of degree 3, which agrees with  $f$  at given points.
- Then, arguing as in previous problem, we obtain

$$f(5) - 4P_3(4) + 6f(3) - 4f(2) + f(1) = 0.$$

and hence,  $P_3(4) = 16.5$ , which is very closed to the actual  $f(4) = 2^4 = 16$ , but not equal. Why?

- This is due to the fact that the data set is associated with a non-polynomial function.

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

Let  $\mathcal{P}_n$  denote the vector space of polynomials of degree at most  $n$ . Let  $\phi_0(x), \dots, \phi_n(x)$  be a basis of  $\mathcal{P}_n$ .

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

Let  $\mathcal{P}_n$  denote the vector space of polynomials of degree at most  $n$ . Let  $\phi_0(x), \dots, \phi_n(x)$  be a basis of  $\mathcal{P}_n$ . Let  $p(x) = a_0\phi_0(x) + \dots + a_n\phi_n(x)$ . Then  $p(x_j) = f_j$  for  $j = 0 : n$  yield

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n]$ ,

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

Let  $\mathcal{P}_n$  denote the vector space of polynomials of degree at most  $n$ . Let  $\phi_0(x), \dots, \phi_n(x)$  be a basis of  $\mathcal{P}_n$ . Let  $p(x) = a_0\phi_0(x) + \dots + a_n\phi_n(x)$ . Then  $p(x_j) = f_j$  for  $j = 0 : n$  yield

$$\begin{bmatrix} \phi_0(x_0) & \phi_1(x_0) & \cdots & \phi_n(x_0) \\ \phi_0(x_1) & \phi_1(x_1) & \cdots & \phi_n(x_1) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_0(x_n) & \phi_1(x_n) & \cdots & \phi_n(x_n) \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_n \end{bmatrix}.$$

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

Let  $\mathcal{P}_n$  denote the vector space of polynomials of degree at most  $n$ . Let  $\phi_0(x), \dots, \phi_n(x)$  be a basis of  $\mathcal{P}_n$ . Let  $p(x) = a_0\phi_0(x) + \dots + a_n\phi_n(x)$ . Then  $p(x_j) = f_j$  for  $j = 0 : n$  yield

$$\begin{bmatrix} \phi_0(x_0) & \phi_1(x_0) & \cdots & \phi_n(x_0) \\ \phi_0(x_1) & \phi_1(x_1) & \cdots & \phi_n(x_1) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_0(x_n) & \phi_1(x_n) & \cdots & \phi_n(x_n) \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_n \end{bmatrix}.$$

Since the coefficient matrix is nonsingular (Check), the linear system has a unique solution.

# Polynomial interpolation

**Problem:** Given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n]$ ,

construct a polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

Let  $\mathcal{P}_n$  denote the vector space of polynomials of degree at most  $n$ . Let  $\phi_0(x), \dots, \phi_n(x)$  be a basis of  $\mathcal{P}_n$ . Let  $p(x) = a_0\phi_0(x) + \dots + a_n\phi_n(x)$ . Then  $p(x_j) = f_j$  for  $j = 0 : n$  yield

$$\begin{bmatrix} \phi_0(x_0) & \phi_1(x_0) & \cdots & \phi_n(x_0) \\ \phi_0(x_1) & \phi_1(x_1) & \cdots & \phi_n(x_1) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_0(x_n) & \phi_1(x_n) & \cdots & \phi_n(x_n) \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_n \end{bmatrix}.$$

Since the coefficient matrix is nonsingular (Check), the linear system has a unique solution. Thus the polynomial interpolation problem has a unique solution.



# Error in Polynomial interpolation

**Interpolation:** For a given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

we find a unique polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ .

# Error in Polynomial interpolation

**Interpolation:** For a given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

distinct nodes:  $[x_0, x_1, \dots, x_n]$

and values:  $[f_0, f_1, \dots, f_n],$

we find a unique polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ . Further, difference between  $f$  and  $p$  is illustrated in the following result:

# Error in Polynomial interpolation

**Interpolation:** For a given a data set  $(x_0, f_0), \dots, (x_n, f_n)$  consisting of

**distinct nodes:**  $[x_0, x_1, \dots, x_n]$

**and values:**  $[f_0, f_1, \dots, f_n],$

we find a unique polynomial  $p(x)$  of degree at most  $n$  such that  $p(x_j) = f_j$  for  $j = 0 : n$ . Further, difference between  $f$  and  $p$  is illustrated in the following result:

Theorem:

Suppose  $x_0, x_1, \dots, x_n$  are distinct real numbers in the interval  $[a, b]$  and  $f \in C^{n+1}[a, b]$ . Then, for each  $x \in [a, b]$ , a number  $\xi_x$  between  $x_0, x_1, \dots, x_n$  and hence in  $(a, b)$  exists such that

$$f(x) - p(x) = \frac{f^{(n+1)}(\xi_x)}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n), \quad (1)$$

where  $p(x)$  is an interpolating polynomial of  $f$  with degree  $n$ .