

Resistance-Aware Computational Prioritization of Antimalarial Leads from African Natural Products: Docking, Chemical-Space Descriptors, and Exploratory Molecular Dynamics

Myke Vital Sao Temgoua,^{*,†} Jean-Pierre Tchapet Njafa,[†] Penabei Samafou,[‡]
Wilfred Fon Mbacham,[¶] and Serge Guy Nana Engo[†]

[†]*Department of Physics, Faculty of Science, University of Yaoundé I, P.O. Box 812,
Yaoundé, Cameroon*

[‡]*Department of Medical Imaging and Radiation Sciences, Université de Sherbrooke,
Sherbrooke, QC, Canada*

[¶]*Department of Biochemistry, Faculty of Science, University of Yaoundé I, P.O. Box 812,
Yaoundé, Cameroon*

E-mail: myke-vital.sao@facsciences-uy1.cm

Abstract

Polypharmacology-oriented scoring may help prioritize antimalarial leads for resistance-aware follow-up, but docking scores do not establish target engagement or mechanism of action. We analyzed 17 candidates across 136 PfDHFR/PfCRT wild-type and mutant docking systems, defining a target-balanced primary set of 12 candidates with eligible wild-type docking scores for both targets and a coverage-limited sensitivity set of 5 PfCRT-only candidates. Under a mutually

exclusive docking-score-retention scheme, the primary set contained 1 Class A*, 1 Class A, 4 Class B, 5 Class C, and 1 Class D candidate; the available-target sensitivity analysis yielded 5 Class A*, 1 Class A, 5 Class B, 5 Class C, and 1 Class D. In the target-balanced set, the PNS–RRS association was weak and uncertain ($\rho = -0.2098$, two-sided permutation $p = 0.5144$, Bonferroni-adjusted $p = 1.0000$, $n = 12$); the nominal negative association in all 17 candidates ($\rho = -0.5588$, permutation $p = 0.0222$, adjusted $p = 0.0667$) is reported only as a coverage-sensitive exploratory result. Two of 17 candidates (11.8%) had ACSI > 0.70 , and cohort-scoped weight perturbations gave Spearman correlations of 0.9167–0.9804. In a separate parent-study MD cohort, two of four complexes retained their ligands under the stated geometric criterion, but only PfCRT–214 yielded an interpretable endpoint estimate (-18.25 ± 0.40) kcal mol⁻¹. A separate 16-system, single-replicate Set-C MD pilot did not support using docking-score retention as a thermodynamic resilience proxy. PfCRT results remain conditional on the neutral-pH protonation protocol and should not be extrapolated to the acidic digestive-vacuole environment. Biological polypharmacology, resistance circumvention, and pathway-level mechanism were not tested.

Introduction

Drug resistance remains the principal threat to malaria control in Africa. An estimated 282 million cases and 610 000 deaths occurred globally in 2024, with the WHO African Region bearing the greatest burden¹. Partial resistance to artemisinin derivatives, mediated by mutations in *Pfkelch13*, is now confirmed or suspected in at least eight African countries, and declining efficacy of some artemisinin partner drugs has been reported^{1,2}; spatial-temporal modelling of *pfk13* markers further indicates that the geographic footprint of these mutations in Africa is expanding³. Genomic surveillance further documents the co-occurrence of *Pfcrtr* and *Pfdhfr* resistance variants across diverse transmission settings⁴, underscoring the

relevance of multi-target resistance panels. Resistance to partner drugs in artemisinin-based combination therapies, driven by mutations in *Pfcr*, *Pfdhfr*, and *Pfdhps*, further narrows the therapeutic options available to clinicians^{5,6}.

Polypharmacology is increasingly treated as a design strategy for addressing complex disease biology, and multi-target-directed ligands represent an important translational class in contemporary pharmacology⁷. The rationale is evolutionary as well as pharmacological: if a compound requires simultaneous loss of activity at several relevant targets, the mutational barrier to treatment failure may increase. That proposition is conditional, however. A list of predicted protein contacts is not a mechanism-of-action assignment; causal network analyses require an explicit perturbation model, a defined biological background, and evidence that the predicted interactions propagate to a phenotype⁸. Recent network-learning approaches to off-target prediction illustrate the value of graph context while also underscoring that network inference remains distinct from experimental target engagement⁹. Natural products are attractive starting points because their stereochemical and scaffold complexity can support activity across multiple protein classes^{10,11}. African natural products provide a particularly relevant chemical-space context, with resources such as ANPDB enabling explicit comparison with approved-drug and synthetic reference spaces¹².

In a previous study, we used generative molecular methods and dual-filter consensus docking (AutoDock Vina¹³ and DiffDock-L¹⁴) to generate a library of 65 856 candidate molecules from 396 African natural products and 454 synthetic drugs¹⁵. We selected the present candidates against a panel spanning established resistance determinants and mechanistically distinct parasite processes: PfDHFR (dihydrofolate reductase, PDB 7F3Y) and PfCRT (chloroquine resistance transporter, PDB 6UKJ) represent clinically established resistance-associated contexts; PfATP4 (sodium efflux pump, PDB 9N10) probes ion homeostasis; and the 4GM2 entry represents PfClpR, not PfClpP, and is retained as a proteostasis-related parent-study system. This panel is deliberately a mechanistic cross-section rather than an exhaustive parasite targetome. It allows us to ask whether resistance-aware prioritization

remains informative across distinct target classes, while avoiding the stronger claim that the selected compounds define a complete parasite-wide polypharmacology profile.

Rigid docking is efficient for ranking but treats the receptor and scoring landscape in a simplified manner. MD can test whether a docked pose remains compatible with protein flexibility, whereas endpoint methods such as MM-GBSA provide system-specific energetic summaries rather than calibrated thermodynamic observables^{16,17}. This distinction is especially important for resistance analysis: a docking RRS is a hypothesis about relative mutant sensitivity, not a measurement of a mutant free-energy difference. Network pharmacology provides a complementary prioritization layer, but centrality is a topological weighting heuristic rather than a causal measure of target essentiality, consistent with the growing interest in multi-target antimalarial design more broadly¹⁸. We therefore treat these quantities separately: docking-derived RRS for mutant sensitivity, PNS for network-weighted target ranking, ACSI for chemical-space positioning, and targeted MD for post-docking structural scrutiny.

We asked whether resistance-aware, target-level scoring could distinguish predicted potency from predicted score retention in a chemically diverse antimalarial library. The primary analysis used 12 candidates with eligible PfDHFR and PfCRT wild-type scores; five PfCRT-only candidates formed a coverage-limited sensitivity set. RRS was calculated from empirical Vina scores, whereas PNS and ACSI provided complementary ranking dimensions. Four separate wild-type complexes underwent 10 ns simulations, and a 16-system Set-C pilot provided a secondary test of the docking-based ranking. Biological polypharmacology and resistance circumvention were not tested.

Methods

Study design and estimand separation

The Set-C cohort contained 17 candidates and 136 docking systems: five PfDHFR states and three PfCRT states per candidate. The 12 candidates with eligible wild-type scores for both targets formed the target-balanced analysis; the remaining five candidates contributed to a PfCRT-only sensitivity analysis. Four non-overlapping wild-type complexes (Ligands 164, 201, 214, and 438) were simulated for 10 ns each. A separate 16-system pilot evaluated PP-01 and PP-02 against the six-mutant panel for 10 ns per system. Docking-score retention, geometric MD-RRS, and MM-GBSA endpoint estimates were analyzed separately.




Figure 1: Study design and separation of computational estimands. The 17-member Set-C cohort supports docking-derived RRS, ACSI, and PNS analyses, whereas the four parent-study complexes support a separate targeted MD and MM-GBSA check. The 16-system Set-C MD pilot (PP-01 and PP-02) is presented separately and is not used to define the docking-RRS classes.

Candidate selection and filtering

Seventeen candidates were selected from the Set-C candidate panel generated from the library of 65 856 molecules described previously¹⁵. The candidate-selection panel is distinct from the production-MD cohort. Three sequential filters were applied: (1) MPO score ≥ 0.70 (primary leads); (2) SYBA score > 0 (synthesisable); (3) selectivity index SI > 10 (selectively antiparasitic). The selection additionally required multi-target engagement of at least two of the four docking targets and scaffold diversity of no more than three compounds per Bemis–Murcko scaffold. The selection includes one of the previously named consensus leads (Ligand 87, PfDHFR). The four additional parent-study consensus leads (Ligands 201, 214, 438, 164) were not part of the 17-candidate docking set but were used to construct the wild-type complexes for the production MD check. The 164 system is identified as the 164-labelled PfClpR cohort; PDB 4GM2 is PfClpR rather than PfClpP, so this system is not structural validation of PfClpP. Five approved antimalarial drugs (artemisinin, pyrimethamine, chloroquine, lumefantrine, cipargamin) were included as reference controls for the docking analysis (Table 1).

Table 1: Breakdown of the 136 docking protein–ligand systems by category (17 polypharmacological candidates $\times$ 8 wild-type/mutant PfDHFR and PfCRT structures). *Production MD was performed at 10 ns each (40 ns total) on 4 wild-type parent-study complexes. Ligand 164 is retained as the 164-labelled PfClpR cohort; PDB 4GM2 is PfClpR rather than PfClpP, so this system is not presented as PfClpP structural validation. The other systems were Ligand 201 (PfDHFR), Ligand 214 (PfCRT), and Ligand 438 (PfATP4); all were distinct from the 17-candidate docking set.

Category	Ligands	Targets	Systems	MD (ns)
PfDHFR WT + 4 mutants (set C)	17	5	85	—
PfCRT WT + 2 mutants (set C)	17	3	51	—
Docking total	—	—	136	—

The 136 docking systems were not subjected to production MD. The 16-system pilot and four non-overlapping parent-study complexes were analyzed separately.

Resistance mutation panel

We selected six resistance-associated states for comparison (Table 2); target identifiers are listed in the Supporting Information (Table S5).

Table 2: *Resistance mutation panel.*

Target	State	Resistance context	Evidence boundary
PfDHFR	N51I	Antifolate-resistance state	Computational panel; no frequency estimate
PfDHFR	C59R	Antifolate-resistance state	Computational panel; no frequency estimate
PfDHFR	S108N	Antifolate-resistance state	Computational panel; no frequency estimate
PfDHFR	I164L	High-level antifolate-resistance state	Comparator for severe resistance
PfCRT	K76T	Chloroquine-resistance marker	Transporter-resistance context
PfCRT	K76A	PfCRT resistance/comparator state	Not a prevalence estimate

Homology modelling of resistance mutants

Crystal structures of mutant targets were not available in the PDB. Homology models were generated with SWISS-MODEL using the wild-type structures as templates: PfDHFR (7F3Y) and PfCRT (6UKJ). PfATP4 and the 164-labelled parent-MD system were not part of the six-mutant homology-model panel; moreover, PDB 4GM2 is PfClpR rather than PfClpP. Model quality was assessed by QMEAN score, Global Model Quality Estimation, Ramachandran analysis (MolProbity), and C α RMSD to the wild-type template. Models were retained only when QMEAN exceeded -4.0 , GMQE exceeded 0.7 , Ramachandran favored residues exceeded 95% , and backbone RMSD was below $1.5\text{ \AA}$. The retained models served as receptor models for docking. Their quality scores do not establish experimental accuracy, and the absence of experimental structures or replicate model ensembles leaves structural uncertainty in the docking results.

Docking protocol validation

The docking protocol and V2-corrected grid parameters used here are identical to those established in the parent study¹⁵: AutoDock Vina 1.2.7¹³ with exhaustiveness 64 against grids centered on the corrected V2 binding-site coordinates of PfDHFR (7F3Y), PfCRT (6UKJ¹⁹), PfATP4 (9N10), and the historically labelled PfClpR/4GM2 entry (not PfClpP). Redocking was considered accurate when RMSD was < 2.0 Å relative to the crystal pose. Results are given separately for the five alignable ligands and for all attempts, including MTX. External enrichment and redocking benchmarks, including the negative DEKOIS 2.0 result, are given in the Supporting Information (Table S1). AutoDock Vina was used for the reported scores; GNINA yielded no usable score.

Secondary docking calibration

A secondary Tartarus cross-docking analysis used QuickVina on the primary-lead panel. The composite MPO–affinity association was null and exploratory; complete correlations are provided in the Supporting Information (Tables S8 and S9).

MPO sensitivity and predicted ADMET profile

MPO-weight sensitivity and single-source ADMET-AI predictions were used descriptively; the complete sensitivity and endpoint results are provided in the Supporting Information (Figure S1 and Table S3). These predictions were not experimentally validated.

Docking and resistance resilience score

All 17 candidates were docked against each wild-type and mutant structure using AutoDock Vina 1.2.7 with exhaustiveness 64. The 136 systems were ranked with empirical Vina scores rather than calibrated binding free energies. GNINA did not yield a score for this panel. Grid boxes were centered on the co-crystallized ligand position with 20 Å dimensions.

Table 3: Target-specific docking-score retention for the 17 Set-C candidates. *RRS* is computed separately for each target and only when the corresponding wild-type Vina score magnitude is at least 5.0 kcal mol⁻¹. A dash denotes an ineligible target, not a zero score. The primary class applies only to the 12 complete two-target candidates; the available-target class is shown for all 17. Classes are mutually exclusive: *A** requires all available mutant *RRS* values $\geq 80\%$ and the minimum eligible wild-type score magnitude ≥ 7.0 kcal mol⁻¹; *A* has the same *RRS* criterion but a weaker minimum wild-type score; *B* requires all values $\geq 70\%$ but not all $\geq 80\%$; *C* requires at least one value $\geq 80\%$ but not all values $\geq 70\%$; *D* is the residual class in which no available mutant reaches 80%. All scores are empirical AutoDock Vina scores; *RRS* is not a free-energy or resistance measurement.

Candidate	WT PfDHFR	WT PfCRT	Eligible targets	N51I	C59R	S108N	I164L	K76T	K76A	Primary class	Available class
PP-01	7.5	9.3	both	86.4	85.5	87.6	83.9	92.5	90.9	A*	A*
PP-02	–	9.1	PfCRT	–	–	–	–	87.9	94.6	–	A*
PP-03	10.3	11.3	both	68.1	68.3	65.8	69.0	80.0	81.4	C	C
PP-04	8.0	7.0	both	68.8	66.9	68.8	73.5	105.3	103.6	C	C
PP-05	–	10.5	PfCRT	–	–	–	–	77.1	79.3	–	B
PP-06	–	9.9	PfCRT	–	–	–	–	90.6	94.1	–	A*
PP-07	7.6	9.1	both	71.8	71.8	72.4	72.2	84.2	77.6	B	B
PP-08	7.6	7.4	both	62.1	62.1	67.2	66.3	83.9	83.1	C	C
PP-09	7.1	8.6	both	70.6	73.0	75.8	76.5	75.8	73.5	B	B
PP-10	7.7	8.4	both	76.0	76.4	75.7	75.8	89.4	89.6	B	B
PP-11	–	10.0	PfCRT	–	–	–	–	82.7	82.5	–	A*
PP-12	7.5	7.8	both	75.2	66.1	62.9	72.5	70.4	87.2	C	C
PP-13	–	8.3	PfCRT	–	–	–	–	110.2	99.4	–	A*
PP-14	7.6	7.5	both	62.2	65.5	68.0	64.9	78.8	77.7	D	D
PP-15	5.6	9.3	both	123.2	112.7	125.9	131.8	86.2	90.1	A	A
PP-16	7.6	8.7	both	72.9	74.7	73.7	75.9	80.1	80.6	B	B
PP-17	8.3	7.4	both	59.3	61.2	59.5	58.9	76.8	93.4	C	C

For target t , let $s_{\text{Vina},i,\text{WT},t}$ and $s_{\text{Vina},i,m,t}$ denote the signed Vina scores. Candidates with $|s_{\text{Vina},i,\text{WT},t}| < 5.0$ kcal mol⁻¹ are excluded from that target’s score-retention analysis. The target-specific Resistance Resilience Score (RRS) is

$$\text{RRS}_{i,m,t} = \frac{|s_{\text{Vina},i,m,t}|}{|s_{\text{Vina},i,\text{WT},t}|} \times 100. \quad (1)$$

RRS is a relative docking-score-retention statistic, not a free-energy difference or biochemical resistance measurement. The available-target summary averages the defined target-specific values across eligible targets and mutant states. In the eligible candidates, mean target-specific RRS was 75.1 for PfDHFR (12 candidates) and 86.2 for PfCRT (17 candidates); the corresponding target-specific summaries are given in the Supporting Information (Table S13). These values describe differences in score retention across the two target panels, not differential biological resistance. To prevent unequal target coverage from driving the pri-

mary inference, the primary target-balanced analysis is restricted to the 12 candidates with eligible PfDHFR and PfCRT wild-type scores; the five PfCRT-only candidates are retained as a coverage-limited sensitivity analysis.

The mutually exclusive classes are defined on the available mutant values, with the target-balanced subset used for the primary class counts:

- **Class A***: all available mutant RRS values are at least 80 % and the minimum eligible wild-type score magnitude is at least 7.0 kcal mol⁻¹.
- **Class A**: all available mutant RRS values are at least 80 %, but the minimum eligible wild-type score magnitude is below 7.0 kcal mol⁻¹.
- **Class B**: all available mutant RRS values are at least 70 %, but at least one is below 80 %.
- **Class C**: at least one available mutant RRS value is at least 80 %, but not all available values reach 70 %.
- **Class D**: no available mutant RRS value reaches 80 %.

The classes are mutually exclusive. These thresholds define docking-based triage categories and have not been calibrated against biochemical resistance measurements or clinical outcomes. Eligible-target counts and target-specific score-retention values are given for both analyses. A threshold-sensitivity analysis is provided in the Supporting Information (Table S12); it shows how the class counts change when the WT eligibility and retention thresholds are tightened.

African chemical space index

ACSI was used as a cohort-relative chemical-space descriptor. Two of 17 candidates exceeded 0.70 (cohort mean, 0.543); local weight-sensitivity results are provided in the Supporting

Information (Table S4). These values describe reference-space positioning, not biological activity or resistance.

Polypharmacology network score

PNS was treated as a secondary network-weighted ranking heuristic. In the target-balanced analysis, PNS-RRS was weak and uncertain ($\rho = -0.2098$, adjusted $p = 1.0000$); complete ranking and imputation details are provided in the Supporting Information (Table S7).

MD stability analysis

We simulated all four parent-study wild-type systems for 10 ns with CHARMM36m²⁰/GAFF2²¹ at 310.15 K using GROMACS^{22,23}. We analyzed the trajectories after removing periodic-boundary discontinuities. MDAnalysis²⁴ was used to calculate contacts, H-bonds, RMSF, and radius of gyration every 10 frames and RMSD on every frame.

Table 4: Production MD stability metrics for wild-type complexes (10 ns). BB RMSD, ligand RMSD, minimum distance, contacts, and H-bonds are computed from PBC-unwrapped trajectories.

Target	Atoms	BB RMSD (Å)	Lig. RMSD (Å)	Min dist (Å)	Contacts	H-bonds	Interpretation
PfClpR-labelled cohort (164)	43 035	1.31	1.11	67.42	0.0	23.7	Unbound
PfDHFR (201)	134 153	3.05	1.70	78.21	0.0	21.5	Unbound
PfCRT (214)	76 920	4.47	2.45	3.19	78.1	23.1	Bound
PfATP4 (438)	420 801	12.27	2.13	2.25	178.4	47.7	Bound

Two systems, PfCRT and PfATP4, retained their ligands throughout production. PfCRT had a backbone RMSD of 4.47 Å, a ligand RMSD of 2.45 Å, 78.1 contacts, and 23.1 H-bonds (minimum heavy-atom distance 3.19 Å). PfATP4 shows a backbone RMSD of 12.27 Å, a ligand RMSD of 2.13 Å, 178.4 contacts, and 47.7 H-bonds (minimum distance 2.25 Å). The high backbone RMSD indicates substantial structural drift in this model. Because the simulation omitted a membrane and the topology conversion was not valid for MM-GBSA, the apparent ligand persistence should not be interpreted as validated binding thermodynamics. PfCRT ligand 214 anchors to Lys34 and engages in a robust H-bond network (Gln101,

Thr30, Gln297) and a hydrophobic core (Leu301, Leu105, Ala33), consistent with a stable pose in this trajectory; neither specificity nor target engagement is established.

The other two systems, the 164-labelled PfClpR cohort and PfDHFR, had stable protein conformations (backbone RMSD 1.31 Å and 3.05 Å, respectively) but the ligands are completely unbound (minimum distances 67.42 Å and 78.21 Å, zero contacts). This indicates either incorrect initial placement or rapid dissociation during equilibration, not a PBC artefact. Consequently, MM-GBSA and RRS analyses for the 164-labelled PfClpR cohort and PfDHFR are not interpretable as affinity estimates and are excluded from the binding-mode discussion below. Thus, protein equilibration alone was insufficient to retain the docked poses.

The detailed PfCRT–214 RMSD trajectory is shown in Figure 2; per-residue flexibility and contact persistence are provided in the Supporting Information. Per-compound RRS profiles across all six mutants are shown in Figure S2.




Figure 2: Detailed stability analysis for the parent-study PfCRT–Ligand 214 complex over 10 ns production MD. The system exhibits transporter flexibility while retaining a bound ligand; the primary cohort-level stability values are reported in Table 4.

MM-GBSA endpoint estimates

MM-GBSA endpoint estimates were obtained for the wild-type systems from their production trajectories (Table 5), using the GB^{OBC2} model (igb=5) with 0.150 M salt concentration, via the `gmx_MMPBSA` v1.5.0.3¹⁶ CHARMM36-to-AMBER conversion. Only the PfCRT–ligand 214 system yields an interpretable MM-GBSA endpoint estimate: the `gmx_MMPBSA` native converter successfully produced a valid AMBER topology with the ligand remaining bound throughout production, giving a software-reported endpoint estimate of $\Delta G_{\text{bind}} = (-18.25 \pm 0.40) \text{ kcal mol}^{-1}$. The 164-labelled and PfDHFR–201 ligands dissociated during production (minimum distances 67.4 Å and 78.2 Å, zero contacts); their endpoint values are not interpretable for binding-mode analysis. The PfATP4 system was excluded from MM-GBSA because CHARMM36-to-AMBER conversion of its two-chain topology produced a systematic van der Waals artifact (+473 kcal mol⁻¹). Its trajectory supports a bound-state observation in this targeted check, but does not establish converged binding thermodynamics.

All MM-GBSA values reported in this work are endpoint enthalpic scores computed with a single-trajectory protocol; configurational entropy is not included. They are therefore not experimental binding free energies and are interpreted only through within-target contrasts (mutant versus wild type) under identical sampling conditions. Unless stated otherwise, the uncertainty quoted for an endpoint estimate is the standard error of the mean over the 100 stored snapshots; the within-trajectory and propagated standard deviations reported by `gmx_MMPBSA` are tabulated in the supplementary material, and none of these quantities is a replicate uncertainty.

Correlation between methods

The four parent-study MD systems were not members of the 17-candidate Set-C cohort; consequently, no Vina–MM-GBSA correlation is estimated across these cohorts. Vina–DiffDock comparisons and the exploratory Ersilia activity-model analysis are reported in the Support-

Table 5: Endpoint interpretation for four parent-study complexes used for production MD. These systems are distinct from the 17 Set-C candidates. Only PfCRT-214 yields an interpretable MM-GBSA endpoint; dissociated systems are reported as N/A.

Ligand	Target	ΔG_{bind} (kcal mol ⁻¹)	SEM	Interpretation
214	PfCRT-WT	-18.25	0.40	Interpretable; ligand bound [†]
164	PfClpR-labelled cohort-WT	-	-	N/A; ligand dissociated [‡]
201	PfDHFR-WT	-	-	N/A; ligand dissociated [§]
438	PfATP4-WT	-	-	N/A; conversion corrupted*

[†]Ligand remains bound (minimum distance 3.19 Å, 78 contacts); conversion via `gmx_MMPBSA` native converter.

The quoted uncertainty is the standard error of the mean over the 100 stored snapshots (see Methods, MM-GBSA binding free energies).

[‡]Ligand dissociates to 67.4 Å minimum distance with zero contacts during production; the reported value is a non-physical end-point estimate for an unbound state and is excluded from binding-mode conclusions.

[§]Ligand dissociates to 78.2 Å minimum distance with zero contacts; same caveat applies.

*Excluded: conversion artifact in the two-chain PfATP4 topology (+473 kcal mol⁻¹ van der Waals inflation); structural persistence is reported separately.

ing Information. They were used only to contextualize complementary ranking dimensions and did not establish score equivalence, independent measurement, or biological activity.

Consensus leads

The intersection of high-RRS (Class A or B), high-ACSI (>0.5), and high-PNS compounds defines a computational prioritization set for future experimental testing. These compounds combine resistance-resilience, African-NP-character, and network-weighted docking signals; the intersection is not itself experimental validation.

Cross-metric correlation: PNS, ACSI, RRS, and docking scores

In the target-balanced set, PNS and RRS were weakly associated ($\rho = -0.2098$, permutation $p = 0.5144$, adjusted $p = 1.0000$; bootstrap 95 % interval $[-0.7582, 0.5429]$, $n = 12$). The 17-candidate estimate was more negative ($\rho = -0.5588$, permutation $p = 0.0222$, adjusted $p = 0.0667$; interval $[-0.7891, -0.1349]$), but unequal target coverage limits its

interpretation.

ACSI and RRS were also weakly associated ($\rho = -0.4056$, permutation $p = 0.1922$, adjusted $p = 0.5766$; bootstrap interval $[-0.8917, 0.2857]$). The 17-candidate estimate was weaker ($\rho = -0.1324$, $p = 0.6117$). The data do not resolve a relationship between chemical-space position and score retention.

RRS and the weakest eligible WT score were not associated ($\rho = -0.1661$, permutation $p = 0.6038$, adjusted $p = 1.0000$; bootstrap interval $[-0.7300, 0.5932]$, $n = 12$). PNS and WT docking scores are mechanically related because PNS incorporates WT docking scores; this relationship was not treated as an independent hypothesis.




Figure 3: Cross-metric correlation matrix for the Set-C candidates. The heat map shows Spearman ρ values; the target-balanced primary analysis uses the 12 complete two-target candidates, while the 17-candidate matrix is a coverage-sensitive descriptive view. PNS and WT docking score are mechanically linked and are not treated as an independent hypothesis.

Table 6: Cross-metric Spearman correlations. The complete two-target set ($n = 12$) is primary; the 17-candidate set is a coverage-sensitive secondary analysis. Permutation p -values use 100 000 permutations; adjusted p -values apply Bonferroni correction to $H1$ – $H3$. Bootstrap intervals are percentile 95 % intervals from 10 000 resamples.

Metric pair	Hypothesis	Spearman ρ	p_{perm}	p_{adj}	Bootstrap 95 % interval
PNS vs. RRS	H1	−0.209 8	0.514 4	1.000 0	[−0.7582, 0.5429]
ACSI vs. RRS	H2	−0.405 6	0.192 2	0.576 6	[−0.8917, 0.2857]
RRS vs. weakest WT score	H3	−0.166 1	0.603 8	1.000 0	[−0.7300, 0.5932]

Analysis populations. The table reports the primary 12-candidate complete two-target set. In the coverage-sensitive 17-candidate set, PNS–RRS gave $\rho = -0.5588$ (permutation $p = 0.0222$, adjusted $p = 0.0667$), ACSI–RRS gave $\rho = -0.1324$ ($p = 0.6117$), and weakest–WT RRS gave $\rho = 0.1759$ ($p = 0.4947$).

Set-C MD pilot: trajectory-derived MD-RRS and MM-GBSA

A secondary 16-system Set-C MD pilot was used to test whether trajectory-derived geometry and MM-GBSA endpoints reproduced docking-RRS patterns. All systems passed the prespecified trajectory-QC gate, but each mutant state was represented by a single 10 ns replicate. Across the eight mutant states with comparable docking references, docking predicted weaker mutant scores, whereas trajectory-derived metrics generally indicated retained or tighter local contacts; seven of eight direction comparisons were discordant. MM-GBSA ratios exceeded 100 % for eight of twelve mutant states, but this ratio is an endpoint comparison rather than a free-energy resilience measure. The pilot yielded a different pattern from docking-RRS; the discrepancy does not establish resistance or enhanced binding. Complete values and QC annotations are provided in the Supporting Information (Figure S3 and Table S14).

Table 7: Set-C MD pilot mutant-state MM-GBSA estimates (secondary, exploratory analysis; not merged into the primary docking-RRS classification). ΔG_{bind} is the mean over 100 snapshots $\pm$ standard deviation; MMG-RRS is the mutant/WT ΔG_{bind} ratio ($\times 100$, same convention as docking RRS; WT = 100); docking RRS is shown for comparison (n/a = no WT docking reference in the pilot for PP-02 PfDHFR).

Candidate	Target	State	ΔG_{bind} (kcal mol ⁻¹)	SD	MMG-RRS	Docking RRS
PP-01	PfCRT	K76A	-35.29	1.17	115.3	90.86
PP-01	PfCRT	K76T	-29.51	2.65	96.4	92.47
PP-01	PfCRT	WT	-30.61	1.65	100.0	-
PP-01	PfDHFR	C59R	-26.56	1.91	105.0	85.47
PP-01	PfDHFR	I164L	-26.13	2.34	103.3	83.87
PP-01	PfDHFR	N51I	-29.83	1.50	118.0	86.40
PP-01	PfDHFR	S108N	-30.21	3.62	119.5	87.60
PP-01	PfDHFR	WT	-25.29	2.94	100.0	-
PP-02	PfCRT	K76A	-24.53	1.27	97.8	94.62
PP-02	PfCRT	K76T	-30.45	1.38	121.4	87.91
PP-02	PfCRT	WT	-25.08	2.56	100.0	-
PP-02	PfDHFR	C59R	-28.85	1.88	95.7	n/a
PP-02	PfDHFR	I164L	-31.06	1.52	103.0	n/a
PP-02	PfDHFR	N51I	-34.02	2.68	112.8	n/a
PP-02	PfDHFR	S108N	-27.09	1.55	89.8	n/a
PP-02	PfDHFR	WT	-30.16	2.06	100.0	-

Discussion

Resistance-resilient design

RRS compares each mutant score with the corresponding wild-type score after excluding wild-type magnitudes below 5.0 kcal mol⁻¹. It is a docking-derived prioritization statistic, not a biochemical measurement of resistance. The classes combine docking-score magnitude with relative score retention (Table 3). Class A* combines a wild-type score magnitude of at least 7.0 kcal mol⁻¹ with RRS values of at least 80% across all tested mutants. It is a prioritization tier, not evidence of resistance circumvention. Classes B and C identify progressively narrower resilience profiles, while Class D denotes that no available mutant reaches the 80% retention threshold. These categories can be aligned with regional resistance surveillance in a future decision analysis, but they do not justify clinical deployment or therapeutic recommendations on their own²⁵.

This distinction is relevant in the current epidemiological context. The 2025 World Malaria Report documents declining efficacy of some artemisinin partner drugs alongside the spread of *Pfkelch13*-mediated partial resistance, with both trends concentrated in the

African Region¹. Resistance-aware prioritization can therefore be a useful design criterion rather than a guarantee of clinical relevance: compounds selected without an explicit mutant-sensitivity analysis may be poorly matched to the resistance states that are relevant in a given parasite population, but computational scores alone cannot establish that mismatch or its therapeutic consequence.

The relationship between RRS and molecular topology is biologically plausible but unresolved. The companion TDA analysis reported a strong H₁-RRS association in the overlapping pilot set ($\rho = 0.864$, $p < 0.001$, $n = 14$) and a smaller association in the expanded set ($\rho = 0.312$, $p = 0.006$, $n = 77$; both counts are internal to that companion study and do not coincide with the present $N = 17$ Set-C cohort)²⁶. These estimates share candidate provenance with the present analysis and may be affected by molecular-weight and scaffold-composition confounding; they are not independent tests of RRS. Persistent ring topology is therefore a structural correlate that can be tested independently. It should be evaluated prospectively with matched scaffolds, larger independent panels, and mutant-state free-energy or experimental measurements before being used as a generative design rule.

Chemical space positioning

ACSI positions the candidates relative to approved-drug and African-natural-product reference spaces; it is a chemical-space descriptor, not a biological activity or target-engagement score. The mean ACSI across the 17 candidates was 0.543, and two candidates (11.8%) exceeded 0.70. Thus, a minority of the resistance-prioritized cohort retained a strong African-NP-like signature under the stated reference construction. This result supports chemical-space diversification as a selection objective, but it does not show that NP-like compounds are more potent, more resilient, or more clinically useful.

The ACSI result is consistent with the use of African-NP reference databases to define a chemically distinctive search space^{11,12}, but cross-study percentages are not directly comparable because reference sets, fingerprints, thresholds, and candidate-selection proce-

dures differ. The observed f_{sp^3} difference between the high-ACSI and lower-ACSI subsets is compatible with the known three-dimensionality of many natural-product scaffolds, yet it remains descriptive in this cohort. ACSI is therefore a selection axis, not a surrogate for pharmacological novelty or efficacy.

The five lower-ACSI candidates ($\text{ACSI} < 0.5$) had lower f_{sp^3} (mean 0.085) than the two high-ACSI candidates (mean 0.328), a descriptive pattern compatible with greater aromaticity in the former subset. The cohort is too small to establish a general relationship between chemical-space position and physicochemical behaviour.

Systems-level polypharmacology

PNS is a network-weighted ranking heuristic, not evidence of target essentiality, causal propagation, or biological polypharmacology. In the target-balanced cohort, PNS–RRS was weak and uncertain ($\rho = -0.2098$, adjusted $p = 1.0000$); its interpretation remains conditional on network construction and target coverage. Multi-target hypotheses still require experimental target-engagement and phenotype measurements before a mechanism of action can be assigned^{4,9,27,28}.

Cross-metric insights

The target-balanced associations among ACSI, PNS, and RRS were uncertain, with wide bootstrap intervals and no multiplicity-adjusted significant result (Table 6). These descriptors are complementary ranking dimensions, not independent measurements. Because the mutant simulations were single-replicate and short, the Set-C pilot cannot determine whether docking RRS predicts thermodynamic resilience.

Value of MD validation

The targeted MD analysis shows why docking benefits from structural follow-up. Two of the four parent-study systems retained bound ligands, whereas the 164-labelled PfClpR cohort and PfDHFR-201 dissociated; only PfCRT-214 passed the bound-state and topology-integrity criteria for MM-GBSA. The dissociation of docking poses under explicit solvation is consistent with the need to combine docking with structural assessment for mutant antimalarial targets²⁹. The resulting $(-18.25 \pm 0.40) \text{ kcal mol}^{-1}$ is a system-specific endpoint estimate and cannot calibrate the 17 Set-C candidates. A stable receptor trajectory alone does not validate a ligand pose. This conclusion aligns with the wider literature on endpoint free-energy methods, which cautions that MM-GBSA ranking accuracy is system- and ensemble-dependent and that single-trajectory estimates should be interpreted as relative guides rather than converged thermodynamics^{16,17}. Recent atomistic work supplies mechanistic context for these residue-76 contrasts: extensive molecular dynamics across CQ-sensitive and CQ-resistant PfCRT isoforms shows that Lys76 obstructs the chloroquine binding site while electrostatically anchoring peptide substrates, and that substitutions at position 76 open drug access in concert with compensatory rearrangements of the binding cavity³⁰. Mutant-versus-wild-type comparisons of the kind reported here interrogate precisely this charged-cavity gate. For the two Set-C pilot candidates, mutant MM-GBSA did not reproduce the docking-predicted weakening within the single-replicate 10 ns simulations. Whether docking RRS predicts mutant free-energy resilience therefore remains unresolved.

Toward experimental validation. None of the computational estimands constitutes biochemical or biophysical evidence. The next step is matched biochemical testing of the highest-priority candidates against wild-type and mutant PfDHFR, together with orthogonal PfCRT binding or transport assays under experimentally relevant conditions. Concordant or discordant results would identify which computational layer is useful for each target class.

Comparison with existing polypharmacology tools

This framework complements chemical-similarity, network-learning, and topological approaches. It separates docking-derived mutant sensitivity (RRS), chemical-space positioning (ACSI), and PPI-weighted ranking (PNS), followed by structural scrutiny. None of these quantities measures mutant-state binding, causal phenotype, or mechanism of action; higher-accuracy free-energy methods and experiments are still needed^{9,26,31,32}.

Limitations

The mutant structures were generated by homology modeling rather than experimental determination, introducing modeling uncertainty into the docking results. No full production MD was generated for the 17 Set-C candidates; the available MD consists of a separate, targeted 10 ns check of four named parent-study wild-type complexes and a 16-system Set-C MD pilot (PP-01 and PP-02 against the six-mutant panel, 10 ns each) that met the predefined trajectory-quality criteria (16/16) and is analyzed separately in the Set-C pilot subsection. The pilot’s single-replicate, 10 ns design cannot resolve long-timescale residence or convergence. An independent PP-01 wild-type production run used the same MDP and analysis; its endpoint estimate ((-28.60 ± 0.43) kcal mol⁻¹, SEM) agrees with the canonical run ((-30.61 ± 0.41) kcal mol⁻¹) to within 2.01 kcal mol⁻¹, inside one within-trajectory standard deviation of either production (Table S11). Because MD snapshots are autocorrelated, the offset is treated as an empirical replicate-noise floor. It supports consistency at the endpoint level but does not address long-timescale residence or convergence. MM-GBSA provides an approximate endpoint estimate and does not fully account for entropic contributions; in this study only PfCRT-214 passed the bound-state and topology-integrity checks. The four targets represent only a subset of proteins relevant to African malaria treatment. Experimental confirmation is still required.

The RRS frequencies reflect the selection filters and unequal target coverage and should not be extrapolated to the full library. PfCRT network centrality was imputed, so PNS

depends on the network construction (Table S7).

Ligand and protein force fields differed between cohorts, so absolute MM-GBSA values are not compared across protocols; only within-protocol contrasts are interpreted.

The target-balanced cross-metric analysis includes only $n = 12$ selected candidates, resulting in wide uncertainty intervals. Larger independent cohorts with complete target panels are needed.

PfCRT was prepared at physiological rather than digestive-vacuole pH; its docking and MD results are therefore conditional because pH correction can materially alter rankings and affinity estimates¹⁵.

Five candidates lacked an eligible PfDHFR WT denominator and were included only in the PfCRT sensitivity analysis. This reflects the receptor and protonation setup rather than biological selectivity. Across the prespecified threshold-sensitivity grid, the baseline available-target class counts were unchanged when the WT eligibility threshold was varied from 4.0 kcal mol⁻¹ to 5.0 kcal mol⁻¹; tightening the retention thresholds from 70%/80% to 80%/90% increased the number of Class D candidates from 1 to 9 (Table S12).

Conclusion

This study prioritizes antimalarial candidates from African natural-product chemical space using several computational estimands. Resistance and binding claims are limited to those estimands. Among the 17 Set-C candidates, the target-balanced docking-RRS classification was A*: 1, A: 1, B: 4, C: 5, and D: 1; the coverage-sensitive available-target analysis yielded A*: 5, A: 1, B: 5, C: 5, and D: 1. The ACSI mean was 0.543. The target-balanced PNS-RRS association was weak and uncertain; the more negative 17-candidate association was coverage-sensitive and did not survive Bonferroni adjustment. These results support resistance-aware target-level ranking; they do not demonstrate biological polypharmacology or a systems-level MoA. The four-complex MD check also showed the value of structural

follow-up: two systems retained bound ligands, but only PfCRT-214 produced an interpretable MM-GBSA estimate. The study keeps potency, network context, chemical-space positioning, mutant sensitivity, and trajectory stability as separate estimands. Experimental target engagement, phenotypic MoA, mutant-state MD, and prospective validation remain necessary next steps.

Author Contributions

Conceptualization and methodology, M.V.S.T. and S.G.N.E.; software, M.V.S.T. and J.-P.T.N., P.S.; validation, M.V.S.T., J.-P.T.N. and W.F.M.; formal analysis and investigation, M.V.S.T. and S.G.N.E.; resources, S.G.N.E. and W.F.M., P.S.; data curation, M.V.S.T.; writing—original draft preparation, M.V.S.T.; writing—review and editing, M.V.S.T., J.-P.T.N., P.S., W.F.M. and S.G.N.E.; visualization, M.V.S.T.; supervision, S.G.N.E. and W.F.M.; project administration, S.G.N.E. All authors read and agreed to the published version.

Data Availability

Docking results, metric outputs, analysis scripts, and the associated source-data records are available at https://github.com/NanaEngo/Malaria_codesV2 under the MIT licence. A versioned snapshot of this repository will additionally be archived on Zenodo to provide a permanent, citable DOI concurrent with publication. The large parent-study trajectories are not included in this article’s data record; the conclusions drawn here are restricted to the trajectory-derived summaries and binding estimates explicitly reported above.

Competing Interests

The authors declare no competing interests.

Acknowledgement

Computational resources were provided by the University of Yaoundé I HPC facility. We acknowledge the SWISS-MODEL server for homology modeling and the GROMACS development team.

Use of Artificial Intelligence

During the preparation of this work the authors used AI assistants to support code development and data analysis scripting. The authors reviewed and edited all AI-assisted output and take full responsibility for the content of this article. No generative AI tool is listed as an author. All computational results, statistical analyses, and quantitative claims were generated and verified by the authors.

Supporting Information Available

Complete docking-validation, Tartarus calibration, MPO-sensitivity, ADMET, ACSI-sensitivity, PNS-imputation-sensitivity, and Set-C MM-GBSA tables; PlasmoDB target-identifier annotation table; MPO-sensitivity, RRS-radar, and Set-C RRS-scatter figures (PDF).

References

- (1) World Health Organization *World Malaria Report 2025*; Report, 2025.
- (2) Habarugira, F.; Batamuriza, J.; Ndahimana, R.; Minega, J. N.; Dieng, M. M.; Mutsaka-Makuvaza, M. J.; Daba, T. M.; Idaghdour, Y.; Mutesa, L. The global landscape of *Plasmodium falciparum* drug resistance markers, 2005–2025: a systematic review and meta-analysis. *Pathogens* **2026**, *15*, 179.

- (3) Young, N. W. et al. Mapping the prevalence of molecular markers of *Plasmodium falciparum* artemisinin partial resistance in Africa: a spatial-temporal modelling study. *The Lancet Infectious Diseases* **2026**, Advance online publication (preprint: medRxiv 2025, 10.64898/2025.12.22.25342873).
- (4) Letebo, A. et al. Genomic surveillance reveals co-occurrence of *Plasmodium falciparum* drug resistance variants across diverse transmission settings in Ethiopia. *Nature Microbiology* **2026**, Advance online publication.
- (5) Patrick, O. J.; Sokan-Adeaga, A. A.; Emmanuel, R. O. Historical evolution and molecular mechanisms of antimalarial drug resistance with emerging challenges and future directions. *Discover Public Health* **2026**, *23*, 749.
- (6) Santos-Lima, J.; others Molecular markers of antimalarial drug resistance in pfk13, pfmdr1, pfdhfr, and pfdhps genes of *Plasmodium falciparum* in a rural municipality in Cubal, Benguela Province, Angola (2022–2023). *Malaria Journal* **2026**, *25*.
- (7) Ryszkiewicz, P.; Baranowska-Kuczko, M.; Malinowska, B.; Schlicker, E. Multi-target-directed drugs: new additions in 2025 and post-marketing safety surveillance of drugs marketed in 2022–2024. *Pharmacological Reports* **2026**, *78*, 1022–1044.
- (8) Trapotsi, M.-A.; Hosseini-Gerami, L.; Bender, A. Computational analyses of mechanism of action (MoA): data, methods and integration. *RSC Chemical Biology* **2022**, *3*, 170–200.
- (9) Hu, Y.; Didi, K.; Cribbs, A. P.; Sun, J. Predicting PROTAC off-target effects via warhead involvement levels in drug–target interactions using graph attention neural networks. *Computational and Structural Biotechnology Journal* **2025**, *27*, 4633–4644.
- (10) Milić, M.; Obi, P.; Vanderwal, C. D.; Ben Mamoun, C.; Le Roch, K. G. Polypharmacology in malaria treatment: single drugs, multiple mechanisms, greater impact. *Trends in Parasitology* **2026**, *42*, 222–236.

- (11) Mayoka, G.; Cheuka, P. M.; Moyo, P.; Dziwornu, G. A.; Beukes, D. Value addition to African natural product-based drug discovery: barriers and opportunities. *Journal of Natural Products* **2025**, *88*, 2018–2028.
- (12) Betow, J. Y. et al. African Natural Products Database (ANPDB): A resource for exploring the therapeutic potential of natural compounds from Africa. *Nucleic Acids Research* **2025**, *54*, D1336–D1344.
- (13) Eberhardt, J.; others AutoDock Vina 1.2.0: new docking methods, expanded force field, and python bindings. *Journal of Chemical Information and Modeling* **2021**, *61*, 3891–3898.
- (14) Corso, G.; others DiffDock: diffusion steps, twists, and turns for molecular docking. **2023**,
- (15) Temgoua, M. V. S.; Tchapel Njafa, J.-P.; Samafou, P.; Mbacham, W. F. Companion manuscript; publication metadata intentionally omitted until the verified record is available.
- (16) Valdés-Tresanco, M. S.; Valdés-Tresanco, M. E.; Valiente, P. A.; Moreno, E. gmx_MMPBSA: A new tool to perform end-state free energy calculations with GRO-MACS. *Journal of Chemical Theory and Computation* **2021**, *17*, 6281–6291.
- (17) Wang, E.; Sun, H.; Wang, J.; Wang, Z.; Liu, H.; Zhang, J. Z. H.; Hou, T. End-point binding free energy calculation with MM/PBSA and MM/GBSA: strategies and applications in drug design. *Chemical Reviews* **2019**, *119*, 9478–9508.
- (18) Cemali, S. H.; Poyraz, S.; Belveren, S.; Taş, S.; Tamer, M. A. Recent insights in multi-target drugs in pharmacology and medicinal chemistry. *ChemMedChem* **2025**, *20*, e202500447.

- (19) Kim, J. et al. Structure and drug resistance of the *Plasmodium falciparum* transporter PfCRT. *Nature* **2019**, *576*, 315–320.
- (20) Huang, J.; Rauscher, S.; Nawrocki, G.; Ran, T.; Feig, M.; de Groot, B. L.; Grubmüller, H.; MacKerell, A. D. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nature Methods* **2017**, *14*, 71–73.
- (21) Wang, J.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A. Development and testing of a general amber force field. *Journal of Computational Chemistry* **2004**, *25*, 1157–1174.
- (22) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2015**, *1–2*, 19–25.
- (23) Lindahl, E.; others GROMACS 2024 source code. *Zenodo* **2024**,
- (24) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. *Journal of Computational Chemistry* **2011**, *32*, 2319–2327.
- (25) WorldWide Antimalarial Resistance Network WWARN: Tracking antimalarial drug resistance. <https://www.wwarn.org/>, 2026.
- (26) Temgoua, M. V. S.; Tchapel Njafa, J.-P.; Samafou, P.; Mbacham, W. F. Quantum-Inspired Molecular Representations for Antimalarial Candidate Evaluation: Topological Fingerprints, Tensor Network Embeddings, and Quantum Kernel Scores. *Journal of Cheminformatics* **2026**, Paper 3: companion study introducing TFP, TNE, and QKS descriptors.
- (27) Shadrack, D. M.; Makaye, A.; Moses, A.; Swai, H. S. Network pharmacology, molecu-

- lar docking and molecular dynamics simulation suggest CE-326597 as an antimalarial molecule. *In Silico Pharmacology* **2026**, *14*.
- (28) Wicht, K. J. et al. The human Chk1 inhibitor CHIR-124 shows multistage activity against *Plasmodium falciparum* via dual inhibition of PfArk1 and hemozoin formation. *ACS Chemical Biology* **2026**, *21*, 1738–1748.
- (29) Gogoi, D.; Gogoi, B.; Dutta, P. P.; Kalita, J. M.; Khataniar, A.; Sharma, M. P.; Mandhadi, J. R.; Phukan, M. M.; Verma, A. K. Computational exploration of natural product inhibitors against mutant *Plasmodium falciparum* DHFR through pharmacophore modeling, docking, and MD simulations. *In Silico Pharmacology* **2026**, *14*.
- (30) Tanner, J. D.; Richards, S. N.; Corry, B. Molecular basis of the functional conflict between chloroquine and peptide transport in the Malaria parasite chloroquine resistance transporter PfCRT. *Nature Communications* **2025**,
- (31) Cang, H.; Wei, G.-W. TopologyNet: Topology Based Deep Convolutional and Multi-task Neural Networks for Biomolecular Property Predictions. *Journal of Chemical Information and Modeling* **2018**, *58*, 189–197.
- (32) Molani, F.; Cho, A. Accurate protein-ligand binding free energy estimation using QM/MM on multi-conformers predicted from classical mining minima. *Communications Chemistry* **2024**, *7*, 247.

TOC Graphic

