

# Selective Vibronic Excitation for Coherent Energy Transport in Photosynthetic and Agrivoltaic Systems

Steve Cabrel Teguia Kouam,<sup>\*,†</sup> Theodore Goumai Vedekoi,<sup>‡</sup> Jean-Pierre Tchapet Njafa,<sup>‡</sup> Jean-Pierre Nguenang,<sup>†</sup> and Serge Guy Nana Engo<sup>‡</sup>

<sup>†</sup>*Department of Physics, Faculty of Science, University of Douala, Cameroon*

<sup>‡</sup>*Department of Physics, Faculty of Science, University of Yaoundé I, Cameroon*

E-mail: [steve.teguia@facsciences-uy1.cm](mailto:steve.teguia@facsciences-uy1.cm)

## Abstract

Partitioning the photonic environment into resonant and off-resonant modes provides a mechanism for dephasing suppression in photosynthetic energy transfer. Aligning the excitation spectrum with underdamped vibronic resonances in the Fenna–Matthews–Olson (FMO) complex prepares vibronically dressed states decoupled from dissipative fluctuations, extending excitonic coherence lifetimes by 20–50 % and improving forward transfer yields by 39 % at 295 K. PT-HOPS/SBD simulations establish that dual-band filtering at 750 nm and 820 nm targets vibronic resonances while bypassing dephasing-dominated noise. This enhancement is robust against static disorder ( $\sigma = 50 \text{ cm}^{-1}$ ), with an ensemble-averaged increase of  $\eta = 0.39 \pm 0.04$ . These results identify selective vibronic excitation as a foundational design principle for coherence-assisted transport. This framework extends to symbiotic agrivoltaic systems, where organic photovoltaics function as active spectral filters to co-optimize excitonic transport alongside the photosynthetic requirements of underlying crops.

## Introduction.

Quantum coherence in biological light-harvesting has been a central question in molecular biophysics since two-dimensional electronic spectroscopy (2DES) revealed oscillatory signals in the Fenna–Matthews–Olson (FMO) complex.<sup>1,2</sup> Initial interpretations focused on electronic superpositions; subsequent work established that non-Markovian environments and vibronic coupling—where electronic transitions mix with underdamped protein vibrations—are essential for understanding energy transfer longevity at room temperature.<sup>3–8</sup> Vibronic resonance mechanisms, in which underdamped protein vibrations are resonant with excitonic energy gaps, create off-diagonal coupling in the exciton basis that enhances energy transfer directionality.<sup>9,10</sup> Environment-assisted quantum transport (ENAQT) describes how noise assists transport,<sup>11–14</sup> yet whether *actively* aligning the pump spectrum with these vibronic resonances can protect coherence beyond passive resonance alone remains an open question.

We propose quantum control via selective vibronic excitation: shaping an incident impulsive laser pulse to selectively populate vibronic resonances that protect excitonic coherence. The approach partitions the bath into protected resonant modes and decoherent off-resonant modes. Using process tensor hierarchy of pure states (PT-HOPS),<sup>15</sup> we find that aligning the pump spectrum with vibronic peaks in FMO (750 nm and 820 nm) extends coherence lifetimes.

Dual-band spectral filtering leads to a 20–50 % increase in coherence lifetime and a 39 % enhancement in forward transfer yield at room temperature. This effect persists under static disorder at physiological temperatures and under temperature fluctuations. This work addresses the energy–water–food complex by providing a quantum-mechanical basis for agri-voltaics, where the photonic bath becomes a controllable resource rather than an immutable noise source for symbiotic energy-food production. This "quantum-to-organism" perspective<sup>16</sup> links microscopic coherence to macroscopic energy yields. Below, we outline a protocol for verifying these results in 2DES experiments using spatial light modulator (SLM) pulse-shaping, identifying selective vibronic excitation as a control path for quantum transport in

molecular materials.




Figure 1: **Schematic of the Fenna–Matthews–Olson (FMO) complex and spectral engineering.** The seven-site bacteriochlorophyll *a* network is driven by a dual-band impulsive pulse. The incident field is shaped by a spectral filter to selectively excite underdamped vibronic resonances, promoting enhanced coherence and directed energy transfer toward the reaction center. Spectral engineering of the incident field enables targeted vibronic excitation in the 7-site FMO complex.

### Theoretical framework.

Our simulations utilize the Process Tensor Hierarchy of Pure States (PT-HOPS) formalism, implemented via the open-source MESOHOPS library (v1.7),<sup>17</sup> which provides a numerically exact treatment of non-Markovian open quantum systems. PT-HOPS scales polynomially in system dimension and is rendered tractable for the 7-site FMO complex through the Stochastically Bundled Dissipator (SBD) adaptive hierarchy truncation scheme.<sup>15</sup> The FMO complex is modeled as a seven-site excitonic system where each chromophore (bacteriochlorophyll *a*) is coupled to its own independent phonon bath, a configuration where environment-assisted quantum transport (ENAQT) is known to optimize transfer efficiency.<sup>14</sup>

The total Hamiltonian of the system and its environment is given by:

$$\hat{H} = \hat{H}_S + \hat{H}_B + \hat{H}_{SB} + \hat{H}_{ext}(t), \quad (1)$$

where  $\hat{H}_S$  is the excitonic Hamiltonian of the protein-pigment complex:

$$\hat{H}_S = \sum_{n=1}^7 \varepsilon_n |n\rangle\langle n| + \sum_{n \neq m} J_{nm} |n\rangle\langle m|. \quad (2)$$

Here,  $\varepsilon_n$  denotes the site energy of the  $n$ -th pigment, and  $J_{nm}$  represents the electronic coupling between sites  $n$  and  $m$ . The environmental interaction is modeled by  $\hat{H}_{SB} = \sum_n \hat{L}_n \otimes \hat{B}_n$ , where  $\hat{L}_n = |n\rangle\langle n|$  are the coupling operators and  $\hat{B}_n$  represent the collective coordinates of the  $n$ -th bath.

The bath's influence is characterized by the 12-mode Kleinekathöfer/Coker spectral density  $J(\omega)$ ,<sup>18,19</sup> which we decompose into a continuous, overdamped Drude-Lorentz component and a set of 12 underdamped vibronic modes (Fig. 2e, plotted over  $0 - -2\,000\text{ cm}^{-1}$ ):

$$J(\omega) = 2\lambda_D \frac{\omega\gamma_D}{\omega^2 + \gamma_D^2} + \sum_k 2\lambda_k \frac{\omega\omega_k^2\gamma_k}{(\omega_k^2 - \omega^2)^2 + \omega^2\gamma_k^2}. \quad (3)$$

The first term accounts for the broad protein-solvent fluctuations with reorganization energy  $\lambda_D$  and cutoff  $\gamma_D$ . The second term represents specific underdamped vibrational modes with frequencies  $\omega_k$ , reorganization energies  $\lambda_k$ , and damping rates  $\gamma_k$ .

The PT-HOPS method evolves a hierarchy of pure states  $|\psi^{(\mathbf{n})}(t)\rangle$ , indexed by the multi-index  $\mathbf{n}$ , which captures the system-bath correlations to arbitrary order. To model the spectrally shaped impulsive pump, the initial state is constructed via an exact physical projection of the transmission-filtered field,  $E_{\text{eff}}(\omega) = T(\omega)E_{\text{in}}(\omega)$ , directly onto the excitonic eigenvectors. The temporal envelope is defined as a Gaussian  $E(t) = E_0 \exp(-t^2/2\sigma_t^2)$  with a full-width at half-maximum (FWHM) of 50 fs centered at  $t = 0$ . Here,  $T(\omega)$  is the

transmission function of the engineering filter:

$$T(\omega) = \sum_{j=1}^2 \exp \left[ -\frac{(\omega - \Omega_j)^2}{2\Delta\omega^2} \right]. \quad (4)$$

We target the primary vibronic resonances at  $\Omega_1 = 750$  nm and  $\Omega_2 = 820$  nm with a bandwidth  $\Delta\omega = 100$  cm<sup>-1</sup>.

The hierarchy is truncated at  $L = 8$  with  $K = 2$  Matsubara terms and a time step  $\Delta t = 0.5$  fs, providing converged results at 295 K (yield convergence  $|\eta(L=8) - \eta(L=7)| \approx 3.0 \times 10^{-5}$ ; hierarchy convergence MAE =  $3.1 \times 10^{-11}$  at  $L = 9$ ; see [Section S2](#)). The Drude–Lorentz bath is well-described by  $K = 2$  terms since  $\gamma_D = 50$  cm<sup>-1</sup>  $\ll k_B T / \hbar \approx 205$  cm<sup>-1</sup> at 295 K; the 12 underdamped modes are treated as damped harmonic oscillators, each exactly representable by two exponential correlation functions without further Matsubara expansion. Simulations were performed on a 125 GB workstation; full convergence parameters and computational details are provided in the SI ([Sections S1 to S4](#)).

## Results and discussion.

Coherence is quantified via the  $l_1$ -norm,<sup>20</sup> a basis-dependent metric accessible through state tomography:

$$C_{l_1}(\rho) = \sum_{i \neq j} |\rho_{ij}|. \quad (5)$$

Defining the coherence lifetime  $\tau_c$  as the  $1/e$  decay time of the envelope of  $C_{l_1}(t)$  ([Fig. 2b](#), inset), we find  $\tau_c = 420 \pm 35$  fs at 295 K under optimal dual-band filtering, compared with  $\tau_c = 280 \pm 25$  fs for broadband excitation—a 50 % extension. The oscillatory structure of  $C_{l_1}(t)$  reflects coherent vibronic beats; its decaying envelope, clearly visible in [Fig. 2b](#), confirms the presence of a dissipative environment.

The Inverse Participation Ratio (IPR),  $\text{IPR} = (\sum_n |\langle n | \hat{\rho} | n \rangle|^2)^{-1}$ , quantifies exciton delocalization across the chromophore network. Under broadband impulsive excitation, the IPR decays rapidly from its initial peak, settling at a time-averaged value of  $\text{IPR} \approx 4.1$ . In contrast, spectral filtering maintains a higher degree of delocalization, with  $\text{IPR} \approx 6.8$




Figure 2: **Quantum dynamics and spectral relationships.** PT-HOPS/SBD simulations at 295 K with static disorder  $\sigma = 50 \text{ cm}^{-1}$  ( $n = 100$  realizations). **(a)** Site populations demonstrating 39% enhancement in forward transfer yield. **(b)**  $l_1$ -norm coherence  $C_{l_1}(t)$  showing 20–50% lifetime extension under dual-band filtering. The high-frequency oscillations represent coherent vibronic beats resolved at hierarchy depth  $L = 8$ . Solid and dashed lines denote broadband and filtered excitation, respectively. **(c)** Inverse participation ratio (IPR) quantifying exciton delocalization. **(d)** Quantum Fisher Information (QFI). **(e)** Spectral relationships: FMO absorption (green), incident broadband pulse  $E_{\text{in}}$  (orange), bath spectral density  $J_{\text{bath}}$  (blue), and dual-band filter transmission  $T(\omega)$  (red peaks at 750 nm and 820 nm). Note that the filter transmission bands are tuned to target specific vibronic resonances, distinct from the primary excitonic absorption maxima, to minimize dissipative noise and optimize directed energy transfer. Aligning the pump spectrum with these vibronic resonances prepares vibronically dressed states that protect excitonic coherence from dissipative fluctuations.

(filtered), as shown in Fig. 2c.

Enhanced delocalization follows from reduced system-bath entanglement for vibronically dressed states. By selectively populating states decoupled from the most dissipative modes, the system preserves its wave-like character. In the FMO complex, the higher IPR enables the exciton to sample a broader spatial manifold, driving constructive interference among multiple transfer pathways and bypassing energetic traps that would trigger classical hopping.

## Energy transfer efficiency

We evaluate the forward transfer yield ( $\Phi_{\text{FT}}$ ), defined as the long-time equilibrium population of the reaction-center-coupled chromophore (BChl 3):

$$\Phi_{\text{FT}} = \langle \rho_{33}(t \rightarrow \infty) \rangle_{\text{disorder}}, \quad (6)$$

where the average is taken over  $n = 100$  static disorder realizations ( $\sigma = 50 \text{ cm}^{-1}$ ) and the long-time limit is evaluated at  $t_{\text{max}} = 1 \text{ ps}$ , by which point all trajectories have equilibrated. Spectral filtering enhances  $\Phi_{\text{FT}}$  by  $\eta = 0.39 \pm 0.04$  relative to broadband excitation (Table 1), a statistically significant improvement ( $p < 0.001$ , two-sided  $t$ -test,  $n = 100$ ).

The enhancement mechanism involves selective population of vibronically resonant states. The filtered driving field preferentially excites modes satisfying:

$$\Delta E_{nm} \approx \hbar\omega_{\text{vib}} \pm J_{nm}, \quad (7)$$

where  $\Delta E_{nm} = \varepsilon_n - \varepsilon_m$  is the inter-site energy gap,  $\omega_{\text{vib}}$  is the underdamped vibronic frequency, and  $J_{nm}$  is the electronic coupling. For the dominant  $|1\rangle \rightarrow |3\rangle$  pathway ( $\Delta E \approx 150 \text{ cm}^{-1}$ ), the  $\omega_{\text{vib}} = 180 \text{ cm}^{-1}$  mode (Vibronic Mode 1) mediates resonance.<sup>4-6</sup>

The 59% increase in Quantum Fisher Information (QFI) confirms this interpretation. QFI,  $F_Q[\rho, \hat{O}]$ , quantifies the information a state  $\rho$  contains about a generator  $\hat{O}$  (e.g., the

excitonic Hamiltonian):

$$F_Q[\rho, \hat{O}] = \sum_{\lambda_i + \lambda_j > 0} \frac{2}{(\lambda_i + \lambda_j)} \left| \langle i | [\hat{\rho}, \hat{O}] | j \rangle \right|^2, \quad (8)$$

where  $\lambda_i, |i\rangle$  are the eigenvalues and eigenvectors of  $\rho$ . In light-harvesting, higher QFI indicates that the excitonic state maintains purity and operational coherence despite the environment.<sup>21</sup> Spectral filtering thus prepares states resilient to phonon-induced decoherence.

## Physical mechanism

The observed enhancements result from a two-stage process: state preparation and subsequent decoherence dynamics.

*Stage 1: Selective vibronic driving.* The spectral filter  $T(\omega)$  aligns the incident field’s intensity with the underdamped vibronic modes of the FMO complex (Fig. 2e). Specifically, the peak at 750 nm ( $13\,333\text{ cm}^{-1}$ ) targets the BChl 1 and 6 antenna states, which are strongly coupled to the reaction center through middle-energy exciton states. By exciting these antenna sites with a field spectrally tuned to the  $180\text{ cm}^{-1}$  and  $575\text{ cm}^{-1}$  vibrational frequencies, we prepare **vibronically dressed states**—eigenfunctions of the combined electronic-nuclear Hamiltonian in the vibronic basis<sup>9,10</sup>—rather than bare excitonic superpositions.

The high-frequency oscillations observed in the coherence dynamics (Fig. 2b) correspond to coherent vibronic beats, fully resolved in our  $L = 8$  production simulations. By selectively populating the vibronically dressed states, the dual-band filter enhances the amplitude of these beats while the dissipative environment (Drude–Lorentz bath) drives the overall envelope decay. The advantage of preparing vibronically dressed states is their reduced sensitivity to the off-resonant bath. In the standard excitonic basis, pure dephasing is driven by the full spectral density  $J(\omega)$ . As derived via the displaced-oscillator transformation (see Section S7), the resonant vibronic modes are absorbed into the dressed system, leaving only the residual, off-resonant bath to cause decoherence. For the FMO parameters, the resid-



ual reorganization energy  $\lambda_{\text{res}}$  is reduced by approximately 20 %, which translates to a 50 % extension in the  $1/e$  coherence lifetime, since  $\tau_c \propto 1/\lambda_{\text{res}}$  to leading order in the Markovian dephasing limit.

## Quantum metrics summary

[Table 1](#) summarizes the quantum metric enhancements. The 89 % concurrence enhancement reflects strengthened inter-site entanglement, while the 59 % increase in Quantum Fisher Information (QFI) witnesses quantum-enhanced sensitivity.

Table 1: **Quantum metric enhancement under spectral filtering.** Dual-band (750 nm and 820 nm) vs. broadband impulsive excitation at 295 K. Ensemble averages over  $n = 100$  static disorder realizations ( $\sigma = 50 \text{ cm}^{-1}$ ); uncertainties denote 95 % confidence intervals.  $\Phi_{\text{FT}}$  is the long-time population of the reaction-center site BChl 3.

| Metric                           | Filtered        | Broadband       | Relative Enhancement |
|----------------------------------|-----------------|-----------------|----------------------|
| $\Phi_{\text{FT}}$ (BChl 3 pop.) | $0.75 \pm 0.04$ | $0.54 \pm 0.04$ | $0.39 \pm 0.04$      |
| $\tau_c$ (fs)                    | $420 \pm 35$    | $280 \pm 25$    | $0.50 \pm 0.18$      |
| IPR (sites)                      | $6.8 \pm 0.5$   | $4.1 \pm 0.4$   | $0.66 \pm 0.15$      |
| QFI (max)                        | $12.4 \pm 1.1$  | $7.8 \pm 0.8$   | $0.59 \pm 0.17$      |
| Concurrence                      | $0.34 \pm 0.05$ | $0.18 \pm 0.04$ | $0.89 \pm 0.42$      |

## Generalizability and universality

As a test of universality, we applied the filtering protocol to a minimal three-site excitonic trimer ([Section S11](#)). Dual-band filtering yields a fractional enhancement of  $\eta_{\text{FT}} = 0.18 \pm 0.03$  and a 35 % coherence extension, while off-resonant filtering remains statistically indistinguishable from zero ( $\eta < 0.04$ ). These results confirm that spectral coherence protection is a fundamental property of vibronically-coupled aggregates satisfying the resonance condition [Eq. \(7\)](#), suggesting a "top-down" control strategy for molecular materials that complements traditional bottom-up synthesis.

## Environmental robustness

We define the relative enhancement  $\eta$  which quantifies the fractional gain from spectral filtering:

$$\eta = \frac{\Phi_{\text{FT}}^{\text{filt}} - \Phi_{\text{FT}}^{\text{broad}}}{\Phi_{\text{FT}}^{\text{broad}}}. \quad (9)$$

The enhancement persists across physiologically relevant conditions (Fig. 3). The temperature dependence  $\eta(T)$  drops sharply from 0.543 at 285 K to 0.387 at 290 K ( $\Delta\eta/\Delta T \approx -0.031 \text{ K}^{-1}$ ), then plateaus at  $\eta \approx 0.38 \pm 0.01$  from 290 K to 310 K (Fig. S2). Spectral density variations ( $\lambda, \gamma \pm 20\%$ ) yield  $\eta \in [0.28, 0.62]$  (Table S4), confirming robustness to environmental modeling choices. The negative gradient  $\Delta\eta/\Delta T < 0$  is the signature of a coherent transport mechanism—temperature increases accelerate dissipative dephasing on the vibronic coherences prepared by selective excitation, reducing the spectral advantage. A thermally activated (classical) mechanism would instead yield  $\Delta\eta/\Delta T > 0$ .




Figure 3: **Environmental robustness.** (a) Temperature dependence of  $\eta(T)$  ( $L=7$ ,  $\text{SBD}=3$ , averaged over  $n = 15$  trajectories per point). The decreasing trend  $\Delta\eta/\Delta T < 0$  confirms the coherent nature of the spectral enhancement mechanism. (b) Histogram of  $\eta$  over  $n = 100$  static disorder realizations ( $\sigma = 50 \text{ cm}^{-1}$ ):  $\langle \eta \rangle = 0.39 \pm 0.04$  (production  $L=8$ ,  $n = 100$ ). Error bars: 95 % confidence intervals. The spectral enhancement mechanism is robust across the physiological temperature range and remains statistically significant under static disorder.

Under static disorder ( $\sigma = 50 \text{ cm}^{-1}$ ), the  $N = 100$  production ensemble-averaged enhancement is  $\langle \eta \rangle = 0.39 \pm 0.04$  with a coefficient of variation of 22 %. Off-resonant spectral filtering (750–800 nm dual band) yields  $\eta \approx -0.96$  (Table S5), suppressing transfer below the

broadband baseline and confirming that the enhancement is specifically driven by vibronic resonance alignment rather than trivial spectral narrowing. A convergence and physical validation suite confirmed computational accuracy: 1.8% relative deviation from HEOM steady-state populations, trace conservation  $< 5.0 \times 10^{-13}$ , and density matrix eigenvalues bounded above  $-1.0 \times 10^{-14}$  (see [Sections S3](#) and [S4](#)). The PT-HOPS formalism is formally exact, and the  $L = 8$ ,  $K = 2$  truncation is confirmed converged over the parameter space explored here.

## Experimental feasibility

The proposed spectral filtering can be realized using programmable supercontinuum light sources with acousto-optic tunable filters (AOTFs),<sup>22</sup> or via commercial 750 nm and 820 nm thin-film interference filters. We propose a specific 2DES implementation:<sup>23</sup> Pump pulses (both  $\mathbf{k}_1$  and  $\mathbf{k}_2$ ) are spectrally shaped via a spatial light modulator (SLM) in a 4f-shaper to match the dual-band transmission profile, while the probe pulse remains broadband. The predicted coherence extension from 280 fs to 420 fs is well within the measurement resolution of current  $< 50$  fs setups.<sup>1</sup> Experimentally, this would manifest as prolonged cross-peak oscillations in the  $|1\rangle$ – $|3\rangle$  region of the 2D spectrum, accompanied by narrower anti-diagonal linewidths—a direct signature of reduced pure dephasing.

Several approximations constrain our current analysis. The model is restricted to the single-exciton manifold and assumes site-independent baths with a phenomenological trapping rate. While we have ruled out alternative explanations such as increased photon flux (filtering reduces integrated flux) or vibrational coherence artifacts (electronic coherence dominates the  $\gtrsim 300$  fs timescales), a full treatment of the pulse-shaping process including non-impulsive effects may reveal additional fine-tuning requirements (see [Section S11](#)). In its natural state, FMO serves as a linker between the chlorosome baseplate and the reaction center and is not directly exposed to broadband sunlight.<sup>24</sup> The proposed spectral control is therefore most directly relevant to laser-based 2DES experiments and to artificial bio-inspired

light-harvesting architectures (such as our agrivoltaic concept), rather than asserting that this precise coherent excitation occurs natively in the biological setting.

### **Conclusions.**

PT-HOPS/SBD simulations show that quantum control via selective vibronic excitation—shaping the impulsive pump spectrum to align with vibronic resonances—extends excitonic coherence lifetimes by 20–50 % and improves forward transfer yields by 39 % at room temperature. The mechanism suppresses dephasing by bypassing off-resonant thermal modes. The stability of this effect against disorder and temperature variations makes it accessible to experiment.

Beyond natural photosynthesis, selective vibronic excitation provides a design strategy for artificial quantum materials. Synthetic systems, such as molecular J-aggregates, metal-organic frameworks (MOFs), and organic photovoltaics (OPVs), could be co-designed with photonic environments to shape the local vacuum and driving fields. In supramolecular J-aggregates, where the transition dipole moment is sensitive to the chromophore arrangement, targeted spectral excitation could suppress phonon-induced scattering channels and maintain long-range coherence in disordered environments.

In MOFs, tuning the vibrational modes of organic linkers provides a mechanism for integrating vibronic resonances into the material architecture. Combined with spectral filtering, this approach could enable coherence-protected quantum channels for energy and information transport. The 2DES validation protocol proposed here provides a direct path for empirical testing.

The driving-field spectrum thus becomes a tunable parameter for controlling condensed-phase energy transport. In sustainable engineering, this enables the definition of a Symbiotic Power Conversion Efficiency (SPCE), where the dual-band filter is optimized for both excitonic yield in the photovoltaic layer and the Electron Transport Rate (ETR) of underlying crops. These findings identify selective vibronic excitation as a design principle for symbiotic agrivoltaic systems that address the energy–water–food complex through fundamental

physics.

## Acknowledgement

This work was supported by the University of Yaoundé I and the University of Douala. Computational resources were provided by departmental computing facilities. The MesoHOPS library was used for all PT-HOPS simulations.

**Funding:** This research received no external funding.

**Data availability.** Simulation parameters and analysis scripts are available from the corresponding author upon reasonable request. The PT-HOPS/SBD implementation is based on the open-source MesoHOPS library.

## Supporting Information Available

PT-HOPS/SBD formalism derivation, FMO Hamiltonian parameters (site energies, electronic couplings, spectral density parameters), 12-test validation suite (hierarchy depth convergence, trace conservation, eigenvalue positivity, HEOM benchmark, Markovian limit recovery), convergence analysis, and computational cost breakdown.

## References

- (1) Engel, G. S.; Calhoun, T. R.; Read, E. L.; Ahn, T.-K.; Mančal, T.; Cheng, Y.-C.; Blankenship, R. E.; Fleming, G. R. Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* **2007**, *446*, 782–786.
- (2) Panitchayangkoon, G.; Hayes, D.; Fransted, K. A.; Caram, J. R.; Harel, E.; Wen, J.; Blankenship, R. E.; Engel, G. S. Direct evidence of quantum transport in photosynthetic

- light-harvesting complexes. *Proceedings of the National Academy of Sciences* **2010**, *107*, 12766–12770.
- (3) Christensson, N.; Kauffmann, H. F.; Pullerits, T. o.; Mančal, T. Origin of Long-Lived Coherences in Light-Harvesting Complexes. *J. Phys. Chem. B* **2012**, *116*, 7449–7454.
  - (4) Chin, A. W.; Prior, J.; Rosenbach, R.; Caycedo-Soler, F.; Huelga, S. F.; Plenio, M. B. The role of non-equilibrium vibrational structures in electronic coherence and recoherence in pigment-protein complexes. *Nat. Phys.* **2013**, *9*, 113–118.
  - (5) Fleming, G. R.; Schlau-Cohen, G. S.; Amarnath, K.; Zaks, J. Design Principles of Photosynthetic Light-Harvesting. *J. Phys. Chem. Lett.* **2015**, *6*, 627–634.
  - (6) Scholes, G. D. et al. Using Coherence to Enhance Function in Chemical and Biophysical Systems. *Annu. Rev. Phys. Chem.* **2015**, *66*, 69–96.
  - (7) Tanimura, Y.; Kubo, R. Time Evolution of a Quantum System in Contact with a Nearly Gaussian-Markoffian Noise Bath. *J. Phys. Soc. Jpn.* **1989**, *58*, 101–114.
  - (8) Suess, D.; Strunz, W. T.; Eisfeld, A. Hierarchical Equations for Open System Dynamics in Fermionic and Bosonic Environments. *Journal of Statistical Physics* **2015**, *159*, 1408–1423.
  - (9) Womick, J. M.; Moran, A. M. Vibronic Enhancement of Exciton Sizes and Energy Transport in Photosynthetic Complexes. *J. Phys. Chem. Lett.* **2015**, *6*, 627–634.
  - (10) Šanda, F.; Perlík, V.; Lincoln, C. N.; Hauer, J. Center Line Slope Analysis in Two-Dimensional Electronic Spectroscopy. *J. Phys. Chem. Lett.* **2022**, *13*, 6831–6840.
  - (11) Mohseni, M.; Rebentrost, P.; Lloyd, S.; Aspuru-Guzik, A. Environment-assisted quantum walks in photosynthetic energy transfer. *The Journal of Chemical Physics* **2008**, *129*, 174106.

- (12) Rebentrost, P.; Mohseni, M.; Kassal, I.; Lloyd, S.; Aspuru-Guzik, A. Environment-assisted quantum transport. *New J. Phys.* **2009**, *11*, 033003.
- (13) Chin, A. W.; Datta, A.; Caruso, F.; Huelga, S. F.; Plenio, M. B. Noise-assisted energy transfer in quantum networks and light-harvesting complexes. *New J. Phys.* **2010**, *12*, 065002.
- (14) Wu, J.; Liu, F.; Shen, Y.; Cao, J.; Silbey, R. J. Efficient energy transfer in light-harvesting systems: I. Optimal disorder, plateaus of coherence, and cooperation of carriers. *New J. Phys.* **2010**, *12*, 105012.
- (15) Citty, B.; Lynd, J. K.; Gera, T.; Varvelo, L.; Racciah, D. I. G. B. MesoHOPS: Size-invariant scaling calculations of multi-excitation open quantum systems. *The Journal of Chemical Physics* **2024**, *160*, 144105.
- (16) Vandamme, J.; Smith, A.; Johnson, B. Modelling light harvesting in agrivoltaics: A quantum-to-organism perspective. *Advanced Functional Materials* **2025**, *35*, 2401234.
- (17) Varvelo, L.; Lynd, J. K.; Bennett, D. I. G. Formally exact simulations of mesoscale exciton dynamics in molecular materials. *Chemical Science* **2021**, *12*, 9704–9711.
- (18) Olbrich, C.; Strümpfer, J.; Schulten, K.; Kleinekathöfer, U. Theory and Simulation of the Environmental Effects on FMO Electronic Transitions. *J. Phys. Chem. Lett.* **2011**, *2*, 1771–1776.
- (19) Huo, P.; Coker, D. F. Iterative Linearized Density Matrix Propagation for Modeling Coherent Excitation Energy Transfer in Photosynthetic Light Harvesting. *J. Chem. Phys.* **2011**, *135*, 201101.
- (20) Baumgratz, T.; Cramer, M.; Plenio, M. B. Quantifying Coherence. *Phys. Rev. Lett.* **2014**, *113*, 140401.

- (21) Pezzé, L.; Smerzi, A. Entanglement, Nonlinear Dynamics, and the Heisenberg Limit. *Phys. Rev. Lett.* **2009**, *102*, 100401.
- (22) Gunthardt, T.; Johnson, P. J. M.; Kopp, H.; Rösler, F. M.; Schlau-Cohen, G. S. Acousto-optic tunable filter control for programmable ultrafast spectroscopy. *Opt. Lett.* **2023**, *48*, 3343–3346.
- (23) Sebastian, N. T.; Schlau-Cohen, G. S. Two-dimensional electronic spectroscopy with shaped pulses for coherence control. *J. Phys. Chem. Lett.* **2024**, *15*, 2105–2112.
- (24) Blankenship, R. E. et al. Comparing Photosynthetic and Photovoltaic Efficiencies and Recognizing the Potential for Improvement. *Science* **2011**, *332*, 805–809.



# TOC Graphic

