

Merge and count step.

- \blacksquare Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- \blacksquare Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and count step.


- \blacksquare Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6

Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6

Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6


Merge and count step.

- $\, \bullet \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3

Merge and count step.


- $\, \bullet \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3

Merge and count step.


- $\, \bullet \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3

Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2

Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2


Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Merge and count step.

- $\mbox{\bf .}$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and count step.


- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2 + 2 + 0

Merge and count step.


- $\mbox{\bf .}$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2 + 2 + 0

Merge and count step.


- $\mbox{\bf .}$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2 + 2 + 0 + 0

Merge and count step.

- $\ \ \,$ Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: 6 + 3 + 2 + 2 + 0 + 0 = 13