



# CS201 DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room903, Nanshan iPark A7 Building

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Number Theory and Cryptography

- Division, Primes
- Congruence
- Greatest Common Divisor (GCD)
- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r$$

- Congruence

- Greatest Common Divisor (GCD)

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

- Greatest Common Divisor (GCD)

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

- Greatest Common Divisor (GCD)

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)  
(extended) Euclidean algorithm

- Euler's Theorem / Fermat's Little Theorem





# Number Theory and Cryptography

## ■ Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

## ■ Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

## ■ Greatest Common Divisor (GCD)

Find the GCD of 286 and 503.

$$\gcd(503, 286) \quad 503 = 1 \cdot 286 + 217$$

$$= \gcd(286, 217) \quad 286 = 1 \cdot 217 + 69$$

$$= \gcd(217, 69) \quad 217 = 3 \cdot 69 + 10$$

$$= \gcd(69, 10) \quad 69 = 6 \cdot 10 + 9$$

$$= \gcd(10, 9) \quad 10 = 1 \cdot 9 + 1$$

$$= 1 \quad 9 = 9 \cdot 1$$

$$1 = 10 - 1 \cdot 9$$

$$1 = 7 \cdot 10 - 1 \cdot 69$$

$$1 = 7 \cdot 217 - 22 \cdot 69$$

$$1 = 29 \cdot 217 - 22 \cdot 286$$

$$1 = 29 \cdot 503 - 51 \cdot 286$$

■ E



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \operatorname{div} d \quad r = a \operatorname{mod} d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)

(extended) Euclidean algorithm

find the modular inverse

solve linear congruence  $ax \equiv b \pmod{m}$  ( $\gcd(a, m) = 1$ )

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \text{ div } d \quad r = a \text{ mod } d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)

(extended) Euclidean algorithm

find the modular inverse

solve linear congruence  $ax \equiv b \pmod{m}$  ( $\gcd(a, m) = 1$ )

Chinese Remainder Theorem / back substitution

- Euler's Theorem / Fermat's Little Theorem



# Number Theory and Cryptography

- Division, Primes

$$a = dq + r \quad q = a \text{ div } d \quad r = a \text{ mod } d$$

- Congruence

$$a \equiv b \pmod{m} \text{ if } m \text{ divides } a - b$$

- Greatest Common Divisor (GCD)

(extended) Euclidean algorithm

find the modular inverse

solve linear congruence  $ax \equiv b \pmod{m}$  ( $\gcd(a, m) = 1$ )

Chinese Remainder Theorem / back substitution

- Euler's Theorem / Fermat's Little Theorem

$$x^{\phi(n)} \equiv 1 \pmod{n} \text{ if } \gcd(x, n) = 1$$

$$x^{p-1} \equiv 1 \pmod{p} \text{ if } x \not\equiv 0 \pmod{p}$$



# RSA Variant

**Q** : Consider the RSA system. Let  $(e, d)$  be a key pair for the RSA. Define

$$\lambda(n) = \text{lcm}(p-1, q-1)$$

and compute  $d' = e^{-1} \bmod \lambda(n)$ . Will decryption using  $d'$  instead of  $d$  still work? (prove  $C^{d'} \bmod n = M$ )



# RSA Variant

**Q** : Consider the RSA system. Let  $(e, d)$  be a key pair for the RSA. Define

$$\lambda(n) = \text{lcm}(p-1, q-1)$$

and compute  $d' = e^{-1} \bmod \lambda(n)$ . Will decryption using  $d'$  instead of  $d$  still work? (prove  $C^{d'} \bmod n = M$ )

Case I:  $\gcd(M, n) = 1$

$$\begin{aligned} C^{d'} \bmod n &= M^{ed'} \bmod n = M^{k\lambda(n)+1} \bmod n \\ &= (M^{k\lambda(n)} \bmod n) M \bmod n \\ &= \left( M^{(p-1)(q-1)/\gcd(p-1, q-1)} \bmod n \right)^k M \bmod n \end{aligned}$$

By Fermat's theorem,  $M^{(p-1)(q-1)/\gcd(p-1, q-1)} \bmod p = \left( M^{(q-1)/\gcd(p-1, q-1)} \right)^{p-1} \bmod p = 1$  and  $M^{(p-1)(q-1)/\gcd(p-1, q-1)} \bmod q = 1$ . Then by Chinese Remainder Theorem, we have  $C^{d'} \bmod n = M$ .



# RSA Variant

**Q** : Consider the RSA system. Let  $(e, d)$  be a key pair for the RSA. Define

$$\lambda(n) = \text{lcm}(p-1, q-1)$$

and compute  $d' = e^{-1} \bmod \lambda(n)$ . Will decryption using  $d'$  instead of  $d$  still work? (prove  $C^{d'} \bmod n = M$ )

Case II:  $\gcd(M, n) = p$

$M = tp$  for some integer  $0 < t < q$ . We have  $\gcd(M, q) = 1$  and  $ed' = k\lambda(n) + 1$  for some integer  $k$ . By Fermat's theorem, we have

$$(M^{k\lambda(n)} - 1) \bmod q = (M^{k(p-1)(q-1)/\gcd(p-1, q-1)} - 1) \bmod q = 0.$$

Then

$$\begin{aligned} (M^{ed'} - M) \bmod n &= M(M^{ed'-1} - 1) \bmod n \\ &= tp(M^{k\lambda(n)} - 1) \bmod pq \\ &= 0 \end{aligned}$$



# RSA Variant

**Q** : Consider the RSA system. Let  $(e, d)$  be a key pair for the RSA. Define

$$\lambda(n) = \text{lcm}(p-1, q-1)$$

and compute  $d' = e^{-1} \bmod \lambda(n)$ . Will decryption using  $d'$  instead of  $d$  still work? (prove  $C^{d'} \bmod n = M$ )

Case III:  $\gcd(M, n) = q$

Similar to Case II.


Case IV:  $\gcd(M, n) = pq$

Trivial.







# Mathematical Induction



# Mathematical Induction



# Mathematical Induction

- We start by reviewing proof by smallest counterexample to try and understand what it is really doing.



# Mathematical Induction

- We start by reviewing proof by smallest counterexample to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of proof by counterexample to *direct proof*. This direct proof technique will be **induction**.



# Mathematical Induction

- We start by *reviewing proof by smallest counterexample* to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of *proof by counterexample* to *direct proof*. This direct proof technique will be **induction**.
- We conclude by distinguishing between the *weak principle* of mathematical induction and the *strong principle* of mathematical induction.



# Mathematical Induction

- We start by *reviewing proof by smallest counterexample* to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of *proof by counterexample* to *direct proof*. This direct proof technique will be **induction**.
- We conclude by distinguishing between the *weak principle* of mathematical induction and the *strong principle* of mathematical induction.

The *strong principle* can actually be derived from the *weak principle*.



# Proof by Smallest Counterexample

- The statement  $P(n)$  is true for all  $n = 0, 1, 2, \dots$



# Proof by Smallest Counterexample

- The statement  $P(n)$  is true for all  $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some  $n > 0$  for which  $P(n)$  is false






# Proof by Smallest Counterexample

- The statement  $P(n)$  is true for all  $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., **There is some  $n > 0$  for which  $P(n)$  is false**
- (ii) Let  $m > 0$  be the **smallest** value for which  $P(n)$  is false




# Proof by Smallest Counterexample

- The statement  $P(n)$  is true for all  $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some  $n > 0$  for which  $P(n)$  is false
- (ii) Let  $m > 0$  be the **smallest** value for which  $P(n)$  is false
- (iii) Then use the fact that  $P(m')$  is true for all  $0 \leq m' < m$  to show that  $P(m)$  is true, **contradicting** the choice of  $m$ .




# Proof by Smallest Counterexample

- The statement  $P(n)$  is true for all  $n = 0, 1, 2, \dots$

We prove this by

- Assume that a counterexample exists, i.e., There is some  $n > 0$  for which  $P(n)$  is false
- Let  $m > 0$  be the **smallest** value for which  $P(n)$  is false
- Then use the fact that  $P(m')$  is true for all  $0 \leq m' < m$  to show that  $P(m)$  is true, **contradicting** the choice of  $m$ .



# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$



# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(\*)** is not always true



# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(\*)** is not always true
- ◇ Then there must be a **smallest**  $n \in \mathbb{N}$  s.t. **(\*)** does not hold for  $n$



# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(\*)** is not always true
- ◇ Then there must be a **smallest**  $n \in \mathbb{N}$  s.t. **(\*)** does not hold for  $n$
- ◇ For any nonnegative integer  $i < n$ ,

$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$



# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(\*)** is not always true
- ◇ Then there must be a **smallest**  $n \in \mathbb{N}$  s.t. **(\*)** does not hold for  $n$
- ◇ For any nonnegative integer  $i < n$ ,
$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$
- ◇ Since  $0 = 0 \cdot 1/2$ , **(\*)** holds for  $n = 0$





# Example 1

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(\*)** is not always true
- ◇ Then there must be a **smallest**  $n \in \mathbb{N}$  s.t. **(\*)** does not hold for  $n$
- ◇ For any nonnegative integer  $i < n$ ,

$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$

- ◇ Since  $0 = 0 \cdot 1/2$ , **(\*)** holds for  $n = 0$
- ◇ The smallest counterexample  **$n$  is larger than 0**



# Example 1

- We now have
  - (i) smallest counterexample  $n$  is greater than 0, and
  - (ii)  $(*)$  holds for  $n - 1$



# Example 1

■ We now have

- (i) smallest counterexample  $n$  is greater than 0, and
- (ii)  $(*)$  holds for  $n - 1$

◇ Substituting  $n - 1$  for  $i$  gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$



# Example 1

■ We now have

- (i) smallest counterexample  $n$  is greater than 0, and
- (ii)  $(*)$  holds for  $n - 1$

◇ Substituting  $n - 1$  for  $i$  gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding  $n$  to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$



# Example 1

■ We now have

- (i) smallest counterexample  $n$  is greater than 0, and
- (ii)  $(*)$  holds for  $n - 1$

◇ Substituting  $n - 1$  for  $i$  gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding  $n$  to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$

◇ Thus,  $n$  is not a counterexample. Contradiction!



# Example 1

■ We now have

- (i) smallest counterexample  $n$  is greater than 0, and
- (ii)  $(*)$  holds for  $n - 1$

◇ Substituting  $n - 1$  for  $i$  gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding  $n$  to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$

◇ Thus,  $n$  is not a counterexample. Contradiction!

◇ Therefore,  $(*)$  holds for all positive integers  $n$ .



# Example 1

- What implication did we have to prove?



# Example 1

- What implication did we have to prove?

The **key step** was proving that

$$P(n - 1) \rightarrow P(n)$$

where  $P(n)$  is the statement

$$1 + 2 + \cdots + n = \frac{n(n + 1)}{2}$$





## Example 2

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,  
$$2^{n+1} \geq n^2 + 2.$$



## Example 2

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,  
$$2^{n+1} \geq n^2 + 2.$$

Let  $P(n) = 2^{n+1} \geq n^2 + 2$ . We start by assuming that the statement

$$\forall n \in \mathbb{N} \ P(n)$$

is **false**.



## Example 2

- Use **proof by smallest counterexample** to show that,  $\forall n \in \mathbb{N}$ ,  
$$2^{n+1} \geq n^2 + 2.$$

Let  $P(n) = 2^{n+1} \geq n^2 + 2$ . We start by assuming that the statement

$$\forall n \in \mathbb{N} \ P(n)$$

is **false**.

When a **for all** quantifier is false, **there must be some  $n$  for which it is false**. Let  $n$  be the **smallest nonnegative integer** for which  $2^{n+1} \not\geq n^2 + 2$ .



## Example 2

- Let  $n$  be the smallest nonnegative integer for which  $2^{n+1} \not\geq n^2 + 2$ .

This means that, for all  $i \in \mathbb{N}$  with  $i < n$ ,

$$2^{i+1} \geq i^2 + 2$$



## Example 2

- Let  $n$  be the smallest nonnegative integer for which  $2^{n+1} \not\geq n^2 + 2$ .

This means that, for all  $i \in \mathbb{N}$  with  $i < n$ ,

$$2^{i+1} \geq i^2 + 2$$

Since  $2^{0+1} \geq 0^2 + 2$ , we know that  $n > 0$ . Thus,  $n - 1$  is a nonnegative integer less than  $n$ .



## Example 2

- Let  $n$  be the smallest nonnegative integer for which  $2^{n+1} \not\geq n^2 + 2$ .

This means that, for all  $i \in \mathbb{N}$  with  $i < n$ ,

$$2^{i+1} \geq i^2 + 2$$

Since  $2^{0+1} \geq 0^2 + 2$ , we know that  $n > 0$ . Thus,  $n - 1$  is a nonnegative integer less than  $n$ .

Then setting  $i = n - 1$  gives

$$2^{(n-1)+1} \geq (n-1)^2 + 2.$$

or

$$(*) \quad 2^n \geq n^2 - 2n + 1 + 2 = n^2 - 2n + 3$$



## Example 2

- Let  $n$  be the **smallest nonnegative integer** for which  $2^{n+1} \not\geq n^2 + 2$ .

We are now given  $2^n \geq n^2 - 2n + 3$ . (\*)



## Example 2

- Let  $n$  be the smallest nonnegative integer for which  $2^{n+1} \not\geq n^2 + 2$ .

We are now given  $2^n \geq n^2 - 2n + 3$ .  $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$





## Example 2

- Let  $n$  be the **smallest nonnegative integer** for which  $2^{n+1} \not\geq n^2 + 2$ .

We are now given  $2^n \geq n^2 - 2n + 3$ .  $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into  $n^2 + 2$  plus an additional nonnegative term.



## Example 2

- Let  $n$  be the **smallest nonnegative integer** for which  $2^{n+1} \not\geq n^2 + 2$ .

We are now given  $2^n \geq n^2 - 2n + 3$ .  $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into  $n^2 + 2$  plus an additional nonnegative term.

Thus, we write

$$\begin{aligned} 2^{n+1} &\geq 2n^2 - 4n + 6 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n - 2)^2 \\ &\geq n^2 + 2. \end{aligned}$$



## Example 2

- Let  $n$  be the **smallest nonnegative integer** for which  $2^{n+1} \not\geq n^2 + 2$ .

We are now given  $2^n \geq n^2 - 2n + 3$ .  $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into  $n^2 + 2$  plus an additional nonnegative term.

Thus, we write

$$\begin{aligned} 2^{n+1} &\geq 2n^2 - 4n + 6 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n - 2)^2 \\ &\geq n^2 + 2. \end{aligned}$$

**contradiction!**



# Example 2

- Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

(a)  $P(0)$  is true

(b) if  $n > 0$ , then  $P(n - 1) \rightarrow P(n)$



## Example 2

- Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)



# Example 2

- Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)
- ◇ Let  $n$  be the smallest counterexample



# Example 2

- Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)
- ◇ Let  $n$  be the smallest counterexample
- ◇ Then, from (a)  $n > 0$ , so  $P(n-1)$  is true



# Example 2

- Let  $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)
- ◇ Let  $n$  be the smallest counterexample
- ◇ Then, from (a)  $n > 0$ , so  $P(n-1)$  is true
- ◇ Therefore, from (b), using direct inference,  $P(n)$  is true





# Example 2

- Let  $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)
- ◇ Let  $n$  be the smallest counterexample
- ◇ Then, from (a)  $n > 0$ , so  $P(n-1)$  is true
- ◇ Therefore, from (b), using direct inference,  $P(n)$  is true
- ◇ This contradicts (\*).



# Example 2

- Let  $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a)  $P(0)$  is true
- (b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some  $n$  for which  $P(n)$  is false (\*)
- ◇ Let  $n$  be the smallest counterexample
- ◇ Then, from (a)  $n > 0$ , so  $P(n-1)$  is true
- ◇ Therefore, from (b), using direct inference,  $P(n)$  is true
- ◇ This contradicts (\*).
- ◇ Thus,  $P(n)$  is true for all  $n \in \mathbb{N}$ .



# Example 2

- What did we really do?

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

(a)  $P(0)$  is true

(b) if  $n > 0$ , then  $P(n - 1) \rightarrow P(n)$



# Example 2

## ■ What did we really do?

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

(a)  $P(0)$  is true

(b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$

We then used proof by smallest counterexample to derive that  $P(n)$  is true for all  $n \in \mathbb{N}$ .



# Example 2

## ■ What did we really do?

Let  $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a)  $P(0)$  is true

(b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$

We then used **proof by smallest counterexample** to derive that  $P(n)$  is true for all  $n \in \mathbb{N}$ .

This is an *indirect proof*. Is it possible to prove this fact *directly*?



# Example 2

## ■ What did we really do?

Let  $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a)  $P(0)$  is true

(b) if  $n > 0$ , then  $P(n-1) \rightarrow P(n)$

We then used **proof by smallest counterexample** to derive that  $P(n)$  is true for all  $n \in \mathbb{N}$ .

This is an *indirect proof*. Is it possible to prove this fact *directly*?

Since  $P(n-1) \rightarrow P(n)$ , we see that

$P(0)$  implies  $P(1)$ ,  $P(1)$  implies  $P(2)$ , ...



# The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.



# The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, *allowing us to use the smallest counterexample*.

This is actually **equivalent** to the *principle of mathematical induction*.





# The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.

This is actually **equivalent** to the *principle of mathematical induction*.

**Principle.** (*the Weak Principle of Mathematical Induction*)

- (a) If the statement  $P(b)$  is true
- (b) the statement  $P(n - 1) \rightarrow P(n)$  is true for all  $n > b$ ,  
then  $P(n)$  is true for all integers  $n \geq b$



# The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.

This is actually **equivalent** to the *principle of mathematical induction*.

**Principle.** (*the Weak Principle of Mathematical Induction*)

- (a) If the statement  $P(b)$  is true
- (b) the statement  $P(n - 1) \rightarrow P(n)$  is true for all  $n > b$ ,  
then  $P(n)$  is true for all integers  $n \geq b$

(a) – *Basic Step*    *Inductive Hypothesis*

(b) – *Inductive Step*    *Inductive Conclusion*



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for  $n = 0$ ,  $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for  $n = 0$ ,  $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that  $n > 0$  and that  $2^n \geq (n-1)^2 + 2 \quad (*)$



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for  $n = 0$ ,  $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that  $n > 0$  and that  $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$



# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for  $n = 0$ ,  $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that  $n > 0$  and that  $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$

Hence, we've just prove that for  $n > 0$ ,  $P(n-1) \rightarrow P(n)$ .





# Proof by Induction

■  $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let  $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for  $n = 0$ ,  $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that  $n > 0$  and that  $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$

Hence, we've just prove that for  $n > 0$ ,  $P(n-1) \rightarrow P(n)$ .

By mathematical induction,  $\forall n > 0, 2^{n+1} \geq n^2 + 2$ .



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$



# Proof by Induction

$$\blacksquare \forall n \geq 2, 2^{n+1} \geq n^2 + 3$$

$$\text{Let } P(n) - 2^{n+1} \geq n^2 + 3$$

$$(i) \text{ Note that for } n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$$



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for  $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that  $n > 2$  and that  $2^n \geq (n-1)^2 + 3 \quad (*)$



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for  $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that  $n > 2$  and that  $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for  $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that  $n > 2$  and that  $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$

Hence, we've just prove that for  $n > 2, P(n-1) \rightarrow P(n)$ .



# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for  $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that  $n > 2$  and that  $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$

Hence, we've just prove that for  $n > 2, P(n-1) \rightarrow P(n)$ .

By mathematical induction,  $\forall n > 2, 2^{n+1} \geq n^2 + 3$ .





# Proof by Induction

■  $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let  $P(n) - 2^{n+1} \geq n^2 + 3$

Base Step

(i) Note that for  $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that  $n > 2$  and that  $2^n \geq (n-1)^2 + 3$  (\*)  
Inductive Hypothesis

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$

Inductive Step

Hence, we've just prove that for  $n > 2, P(n-1) \rightarrow P(n)$ .

By mathematical induction,  $\forall n > 2, 2^{n+1} \geq n^2 + 3$ .

Inductive Conclusion



# Another Form of Induction

- We may have another form of *direct proof* as follows.



# Another Form of Induction

- We may have another form of *direct proof* as follows.
  - ◇ First suppose that we have proof of  $P(0)$



# Another Form of Induction

- We may have another form of *direct proof* as follows.
  - ◇ First suppose that we have proof of  $P(0)$
  - ◇ Next suppose that we have a proof that,  $\forall k > 0$ ,  
$$P(0) \wedge P(1) \wedge P(2) \wedge \cdots \wedge P(k-1) \rightarrow P(k)$$



# Another Form of Induction

- We may have another form of *direct proof* as follows.

- ◇ First suppose that we have proof of  $P(0)$

- ◇ Next suppose that we have a proof that,  $\forall k > 0$ ,

$$P(0) \wedge P(1) \wedge P(2) \wedge \cdots \wedge P(k-1) \rightarrow P(k)$$

- ◇ Then,  $P(0)$  implies  $P(1)$

$P(0) \wedge P(1)$  implies  $P(2)$

$P(0) \wedge P(1) \wedge P(2)$  implies  $P(3) \dots$



# Another Form of Induction

- We may have another form of *direct proof* as follows.

- ◇ First suppose that we have proof of  $P(0)$

- ◇ Next suppose that we have a proof that,  $\forall k > 0$ ,

$$P(0) \wedge P(1) \wedge P(2) \wedge \cdots \wedge P(k-1) \rightarrow P(k)$$

- ◇ Then,  $P(0)$  implies  $P(1)$

$P(0) \wedge P(1)$  implies  $P(2)$

$P(0) \wedge P(1) \wedge P(2)$  implies  $P(3) \dots$

- ◇ Iterating gives us a proof of  $P(n)$  for all  $n$



# Strong Induction

- **Principle** (*The Strong Principle of Mathematical Induction*)
  - (a) If the statement  $P(b)$  is true
  - (b) for all  $n > b$ , the statement  $P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1) \rightarrow P(n)$  is true.then  $P(n)$  is true for all integers  $n \geq b$ .



# Example

- Prove that every positive integer is a power of a prime or the product of powers of primes.



# Example

- Prove that every positive integer is a power of a prime or the product of powers of primes.
  - ◇ **Base Step**: 1 is a power of a prime number,  $1 = 2^0$

# Example

- Prove that every positive integer is a power of a prime or the product of powers of primes.
  - ◇ **Base Step**: 1 is a power of a prime number,  $1 = 2^0$
  - ◇ **Inductive Hypothesis**: Suppose that every number less than  $n$  is a power of a prime or a product of powers of primes.

# Example

- Prove that every positive integer is a power of a prime or the product of powers of primes.
  - ◇ **Base Step**: 1 is a power of a prime number,  $1 = 2^0$
  - ◇ **Inductive Hypothesis**: Suppose that every number less than  $n$  is a power of a prime or a product of powers of primes.
  - ◇ Then, if  $n$  is not a prime power, it is a product of two smaller numbers, each of which is, by the **inductive hypothesis**, a power of a prime or a product of powers of primes.

# Example

- Prove that every positive integer is a power of a prime or the product of powers of primes.
  - ◇ **Base Step**: 1 is a power of a prime number,  $1 = 2^0$
  - ◇ **Inductive Hypothesis**: Suppose that every number less than  $n$  is a power of a prime or a product of powers of primes.
  - ◇ Then, if  $n$  is not a prime power, it is a product of two smaller numbers, each of which is, by the **inductive hypothesis**, a power of a prime or a product of powers of primes.
  - ◇ Thus, by the **strong principle of mathematical induction**, every positive integer is a power of a prime or a product of powers of primes.

# Mathematical Induction

- In practice, we **do not** usually explicitly distinguish between the weak and strong forms.



# Mathematical Induction

- In practice, we **do not** usually explicitly distinguish between the weak and strong forms.
- In reality, they are **equivalent** to each other in that the weak form is a special case of the strong form, and the strong form can be derived from the weak form.



# Summary

- A *typical* proof by mathematical induction, showing that a statement  $P(n)$  is true for all integers  $n \geq b$  consists of three steps:



# Summary

- A *typical* proof by mathematical induction, showing that a statement  $P(n)$  is true for all integers  $n \geq b$  consists of three steps:
  1. We show that  $P(b)$  is true. – Base Step





# Summary

- A *typical* proof by mathematical induction, showing that a statement  $P(n)$  is true for all integers  $n \geq b$  consists of three steps:

1. We show that  $P(b)$  is true. – Base Step

2. We then,  $\forall n > b$ , show either

$$(*) \quad P(n-1) \rightarrow P(n)$$

or

$$(**) \quad P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1) \rightarrow P(n)$$



# Summary

- A *typical* proof by mathematical induction, showing that a statement  $P(n)$  is true for all integers  $n \geq b$  consists of three steps:

1. We show that  $P(b)$  is true. – Base Step

2. We then,  $\forall n > b$ , show either

$$(*) \quad P(n-1) \rightarrow P(n)$$

or

$$(**) \quad P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1) \rightarrow P(n)$$

We need to make the **inductive hypothesis** of either  $P(n-1)$  or  $P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1)$ . We then use  $(*)$  or  $(**)$  to derive  $P(n)$ .



# Summary

- A *typical* proof by mathematical induction, showing that a statement  $P(n)$  is true for all integers  $n \geq b$  consists of three steps:

1. We show that  $P(b)$  is true. – Base Step

2. We then,  $\forall n > b$ , show either

$$(*) \quad P(n-1) \rightarrow P(n)$$

or

$$(**) \quad P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1) \rightarrow P(n)$$

We need to make the **inductive hypothesis** of either  $P(n-1)$  or  $P(b) \wedge P(b+1) \wedge \cdots \wedge P(n-1)$ . We then use  $(*)$  or  $(**)$  to derive  $P(n)$ .

3. We conclude on the basis of the principle of **mathematical induction** that  $P(n)$  is true for all  $n \geq b$ .



# Recursion

- Recursive computer programs or algorithms often lead to inductive analysis.



# Recursion

- Recursive computer programs or algorithms often lead to *inductive analysis*.
- A classical example of *recursion* is the **Towers of Hanoi** Problem.



# Towers of Hanoi



# Towers of Hanoi



- 3 pegs;  $n$  disks of different sizes
- A *legal move* takes a disk from one peg and moves it onto another peg so that **it is not on top of a smaller disk**
- **Problem:** Find a (efficient) way to move all of the disks from one peg to another



# Towers of Hanoi






# Towers of Hanoi




legal move




# Towers of Hanoi



legal move




legal move




# Towers of Hanoi




legal move



legal move




not legal




# Towers of Hanoi




legal move



legal move



not legal



legal move



# Towers of Hanoi

- **Problem:** Start with  $n$  disks on leftmost peg




# Towers of Hanoi

- **Problem:** Start with  $n$  disks on leftmost peg using only legal moves



# Towers of Hanoi

- **Problem:** Start with  $n$  disks on leftmost peg using only legal moves  
move all disks to rightmost peg.



# Towers of Hanoi

- **Problem:** Start with  $n$  disks on leftmost peg  
using only legal moves  
move all disks to rightmost peg.



Given  $i, j \in \{1, 2, 3\}$ , let  
 $\overline{\{i, j\}} = \{1, 2, 3\} - \{i\} - \{j\}$ ,  
i.e.,  $\overline{\{1, 2\}} = \{3\}$ ,  $\overline{\{1, 3\}} = \{2\}$ ,  
 $\overline{\{2, 3\}} = \{1\}$ .





# Towers of Hanoi

- General solution



# Towers of Hanoi

- General solution

Recursion Base:

If  $n = 1$ , moving one disk from  $i$  to  $j$  is easy. Just move it.



# Towers of Hanoi

- General solution

## Recursion Base:

If  $n = 1$ , moving one disk from  $i$  to  $j$  is easy. Just move it.



# Towers of Hanoi



To move  $n > 1$  disks from  $i$  to  $j$



# Towers of Hanoi



1)



To move  $n > 1$  disks from  $i$  to  $j$

move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$


# Towers of Hanoi



1)



2)



To move  $n > 1$  disks from  $i$  to  $j$

move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$

move **largest** disk from  $i$  to  $j$

# Towers of Hanoi



1)



2)



3)



To move  $n > 1$  disks from  $i$  to  $j$

move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$

move **largest** disk from  $i$  to  $j$

move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

```
3 public class Hanoi
4 {
5
6     public void move(int n, char a, char b, char c)
7     {
8         if (n == 1)
9             System.out.println("plate " + n + " from " + a + " to " + c);
10        else
11        {
12            move(n-1,a,c,b);
13            System.out.println("plate " + n + " from " + a + " to " + c);
14            move(n-1,b,a,c);
15        }
16    }
17 }
18
```





# Towers of Hanoi

To move  $n$  disks from  $i$  to  $j$

i) move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$

ii) move largest disk from  $i$  to  $j$

iii) move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

- To prove **Correctness** of solution, we are implicitly using **induction**

To move  $n$  disks from  $i$  to  $j$

- move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$
- move largest disk from  $i$  to  $j$
- move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

- To prove **Correctness** of solution, we are implicitly using **induction**
- $p(n)$  is statement that algorithm is correct for  $n$

To move  $n$  disks from  $i$  to  $j$

- i) move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$
- ii) move largest disk from  $i$  to  $j$
- iii) move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

- To prove **Correctness** of solution, we are implicitly using **induction**
- $p(n)$  is statement that algorithm is correct for  $n$
- $p(1)$  is statement that algorithm works for  $n = 1$  disks, which is obviously true

To move  $n$  disks from  $i$  to  $j$

- move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$
- move largest disk from  $i$  to  $j$
- move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

- To prove **Correctness** of solution, we are implicitly using **induction**

- $p(n)$  is statement that algorithm is correct for  $n$

- $p(1)$  is statement that algorithm works for  $n = 1$  disks, which is obviously true

- $p(n - 1) \rightarrow p(n)$  is **recursion** statement that

if our algorithm works for  $n - 1$  disks, then we can build a correct solution for  $n$  disks

To move  $n$  disks from  $i$  to  $j$

i) move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$

ii) move largest disk from  $i$  to  $j$

iii) move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

## ■ Running time

$M(n)$  is number of disk moves needed for  $n$  disks

To move  $n$  disks from  $i$  to  $j$

i) move top  $n - 1$  disks from  $i$  to  $\overline{\{i, j\}}$

ii) move largest disk from  $i$  to  $j$

iii) move top  $n - 1$  disks from  $\overline{\{i, j\}}$  to  $j$



# Towers of Hanoi

## ■ Running time

$M(n)$  is number of disk moves needed for  $n$  disks

$$M(1) = 1$$

$$\text{if } n > 1, \text{ then } M(n) = 2M(n-1) + 1$$

To move  $n$  disks from  $i$  to  $j$

i) move top  $n-1$  disks from  $i$  to  $\overline{\{i,j\}}$

ii) move largest disk from  $i$  to  $j$

iii) move top  $n-1$  disks from  $\overline{\{i,j\}}$  to  $j$



# Towers of Hanoi

- We saw that  $M(1) = 1$  and that
- $M(n) = 2M(n - 1) + 1$  for  $n > 1$





# Towers of Hanoi

- We saw that  $M(1) = 1$  and that
- $M(n) = 2M(n - 1) + 1$  for  $n > 1$
- Iterating the recurrence gives
$$M(1) = 1, M(2) = 3, M(3) = 7,$$
$$M(4) = 15, M(5) = 31, \dots$$



# Towers of Hanoi

- We saw that  $M(1) = 1$  and that
- $M(n) = 2M(n - 1) + 1$  for  $n > 1$
- Iterating the recurrence gives
$$M(1) = 1, M(2) = 3, M(3) = 7,$$
$$M(4) = 15, M(5) = 31, \dots$$
- We *guess* that  $M(n) = 2^n - 1$



# Towers of Hanoi

- We saw that  $M(1) = 1$  and that
- $M(n) = 2M(n - 1) + 1$  for  $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$

- We *guess* that  $M(n) = 2^n - 1$   
We'll prove this by induction



# Towers of Hanoi

- We saw that  $M(1) = 1$  and that
- $M(n) = 2M(n - 1) + 1$  for  $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$

- We *guess* that  $M(n) = 2^n - 1$

We'll prove this *by induction*

Later, we'll also see how to solve *without guessing*



# Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that  $M(n) = 2^n - 1$ .



# Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that  $M(n) = 2^n - 1$ .

**Proof.** (by induction)

The base case  $n = 1$  is true, since  $2^1 - 1 = 1$ .

For the inductive step, assume that  $M(n-1) = 2^{n-1} - 1$  for  $n > 1$ .



# Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that  $M(n) = 2^n - 1$ .

**Proof.** (by induction)

The base case  $n = 1$  is true, since  $2^1 - 1 = 1$ .

For the inductive step, assume that  $M(n-1) = 2^{n-1} - 1$  for  $n > 1$ .

Then  $M(n) = 2M(n-1) + 1 = 2(2^{n-1} - 1) + 1 = 2^n - 1$



# Towers of Hanoi

- Note that we used induction twice.





# Towers of Hanoi

- Note that we used **induction twice**.
- The first time was to **derive** correctness of algorithm and the recurrence

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$



# Towers of Hanoi

- Note that we used **induction twice**.
- The first time was to **derive** correctness of algorithm and the recurrence

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

- The second time was to **derive** the **closed form solution**  $M(n) = 2^n - 1$  of the recurrence.



# Next Lecture

## ■ recurrence ...

