


SUBSTANCE ABUSE ANALYSIS


● UsualCare ● Intervention



UsualCare
336
Intervention
143


479
Patients

● Male ● Female




Male
300
Female
179

Patients by Race Ethnicity




Patients By Age Group




Patients Admissions


Program — Intervention — UsualCare



Substances Used Most



Mental Health Diagnosis



HOME

DLA&INSIGHTS



ENTERPRISE DNA

DAILY LIVING ACTIVITIES

Intervention

UsualCare


Race Ethnicity	Patients
Non Hispanic White	134
Native American	119
Non Hispanic Black	22
Hispanic	19

3.88
Average DLA1


4.05
Average DLA2

Age Bucket	Patients
46 - 55	134
36 - 45	126
56 - 65	93
26 - 35	79

Daily Living Activities By Age Group



Daily Living Activities 1 & 2



More Than 50%

- patients are between the age **36 - 55**.
- prefer **Usual Care**.
- are **Male**.

- **Alcohol & Opioid** are the **most consumed** substance.
- Patients are mostly diagnosed for **Depression & Anxiety**.
- **Native American & Non-Hispanic White** prefer Usual Care over Intervention.
- Patients who prefer **Intervention** program has improvement rates when comes to the follow up session (DLA2).
- There is a **strong, positive, linear**, relationship between DLA1 & DLA2.
- The age group between 35 and 55 has **slightly higher rates** of attending the follow up sessions.

HOME

DLA&INSIGHTS



ENTERPRISE DNA