


Dealing with Uncertainty

CMPSCI 383
October 25, 2011

Midterm grades



Average: 71

Outline

- Uncertainty
- Probability
- Syntax and Semantics
- Inference
- Independence and Bayes' Rule

Uncertainty

Let action $A_t = \text{leave for airport } t \text{ minutes before flight}$

Will A_t get me there on time?

- Problems
 - Partial observability (road state, other drivers' plans, etc.)
 - Noisy sensors (traffic reports)
 - Uncertainty in action outcomes (flat tire, etc.)
 - Immense complexity of modeling and predicting traffic
- A purely logical approach either...
 - Risks falsehood: “ A_{25} will get me there on time”, or
 - Leads to conclusions that are too weak for decision making
 - “ A_{25} will get me there on time if there's no accident on the bridge and it doesn't rain and my tires remain intact etc etc.” (qualification problem)
 - “ A_{1440} might reasonably be said to get me there on time but I'd have to stay overnight in the airport ...”

Options for handling uncertainty

- Belief states and contingency plans
- Default or nonmonotonic logic
 - Assume my car does not have a flat tire
 - Assume A_{25} works unless contradicted by evidence
 - However...
 - What assumptions are reasonable?
 - How to handle contradiction?
- Rules with fudge factors
 - $A_{25} \mapsto_{0.3}$ get there on time
 - $Sprinkler \mapsto_{0.99} WetGrass$
 - $WetGrass \mapsto_{0.7} Rain$
 - However...
 - Problems with combination, e.g., *Sprinkler causes Rain??*
- Probability
 - Model agent's degree of belief
 - Given the available evidence, A_{25} will get me there on time with probability 0.04

Probability

Probabilistic assertions **summarize** effects of

- **Laziness** — failure to enumerate exceptions, qualifications, etc.
- **Ignorance** — lack of relevant facts, initial conditions, etc.
- **Fundamental stochastic nature** of phenomena

Subjective probability

- Probabilities relate propositions to agent's own state of knowledge
e.g., $P(A_{25} \mid \text{no reported accidents}) = 0.06$
- These are **not** assertions about the world, they are assertions about belief
- Probabilities of propositions change with new evidence:
e.g., $P(A_{25} \mid \text{no reported accidents, 5 a.m.}) = 0.15$

Making decisions under uncertainty

Suppose I believe the following:

$P(A_{25} \text{ gets me there on time} \mid \dots)$	$= 0.04$
$P(A_{90} \text{ gets me there on time} \mid \dots)$	$= 0.70$
$P(A_{120} \text{ gets me there on time} \mid \dots)$	$= 0.95$
$P(A_{1440} \text{ gets me there on time} \mid \dots)$	$= 0.9999$

- Which action to choose?

Depends on my **preferences** for missing flight vs. time spent waiting, etc.

- **Utility theory** is used to represent and infer preferences
- **Decision theory** = probability theory + utility theory
- MEU principle: maximum expected utility

Basics of Probability Theory

Begin with a set Ω —the sample space

e.g., 6 possible rolls of a die.

$\omega \in \Omega$ is a sample point/possible world/atomic event

A probability space or probability model is a sample space with an assignment $P(\omega)$ for every $\omega \in \Omega$ s.t.

$$0 \leq P(\omega) \leq 1$$

$$\sum_{\omega} P(\omega) = 1$$

e.g., $P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = 1/6$.

An event A is any subset of Ω

$$P(A) = \sum_{\{\omega \in A\}} P(\omega)$$

E.g., $P(\text{die roll} < 4) = P(1) + P(2) + P(3) = 1/6 + 1/6 + 1/6 = 1/2$

Sample point/Atomic Event/Possible world

- A *complete* specification of the state of the world about which the agent is uncertain

E.g., if the world consists of only two Boolean variables *Cavity* and *Toothache*, then there are 4 distinct atomic events:

$Cavity = false \wedge Toothache = false$

$Cavity = false \wedge Toothache = true$

$Cavity = true \wedge Toothache = false$

$Cavity = true \wedge Toothache = true$

- Sample points are mutually exclusive and collectively exhaustive

Random Variables

A random variable is a function from sample points to some range, e.g., the reals or Booleans

e.g., $Odd(1) = true$.

P induces a probability distribution for any r.v. X :

$$P(X = x_i) = \sum_{\{\omega: X(\omega) = x_i\}} P(\omega)$$

e.g., $P(Odd = true) = P(1) + P(3) + P(5) = 1/6 + 1/6 + 1/6 = 1/2$

Some Syntax

Propositional or Boolean random variables

e.g., *Cavity* (do I have a cavity?)

Cavity = true is a proposition, also written *cavity*

Discrete random variables (finite or infinite)

e.g., *Weather* is one of $\langle \textit{sunny}, \textit{rain}, \textit{cloudy}, \textit{snow} \rangle$

Weather = rain is a proposition

Values must be exhaustive and mutually exclusive

Continuous random variables (bounded or unbounded)

e.g., *Temp* = 21.6; also allow, e.g., *Temp* < 22.0.

Arbitrary Boolean combinations of basic propositions

Propositions and Probabilities

Think of a proposition as the event (set of sample points) where the proposition is true

Given Boolean random variables A and B :

event a = set of sample points where $A(\omega) = \text{true}$

event $\neg a$ = set of sample points where $A(\omega) = \text{false}$

event $a \wedge b$ = points where $A(\omega) = \text{true}$ and $B(\omega) = \text{true}$

Often in AI applications, the sample points are **defined** by the values of a set of random variables, i.e., the sample space is the Cartesian product of the ranges of the variables

With Boolean variables, sample point = propositional logic model

e.g., $A = \text{true}$, $B = \text{false}$, or $a \wedge \neg b$.

Proposition = disjunction of atomic events in which it is true


e.g., $(a \vee b) \equiv (\neg a \wedge b) \vee (a \wedge \neg b) \vee (a \wedge b)$

$\Rightarrow P(a \vee b) = P(\neg a \wedge b) + P(a \wedge \neg b) + P(a \wedge b)$

Axioms of probability

- For any propositions A, B
 - $0 \leq P(A) \leq 1$
 - $P(\text{true}) = 1$ and $P(\text{false}) = 0$
 - $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$

True



Prior probability

- **Prior or unconditional probabilities** of propositions
e.g., $P(\text{Cavity} = \text{true}) = 0.1$ and $P(\text{Weather} = \text{sunny}) = 0.72$ correspond to belief prior to arrival of any (new) evidence
- **Probability distribution** gives values for all possible assignments:
 $\mathbf{P}(\text{Weather}) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$ (**normalized**, i.e., sums to 1)
- **Joint probability distribution** for a set of random variables gives the probability of every atomic event on those random variables

$\mathbf{P}(\text{Weather}, \text{Cavity})$ = a 4×2 matrix of values:

<i>Weather</i> =	sunny	rainy	cloudy	snow
<i>Cavity</i> = true	0.144	0.02	0.016	0.02
<i>Cavity</i> = false	0.576	0.08	0.064	0.08

- *Every probabilistic question about a domain can be answered by its joint distribution*

Conditional probability

- Conditional or posterior probabilities

e.g., $P(\text{cavity} \mid \text{toothache}) = 0.8$

i.e., given that *toothache* is all I know

NOT “if *toothache* then 80% chance of *cavity*”

- Notation for complete conditional distribution:

$\mathbf{P}(\text{Cavity} \mid \text{Toothache}) = 2\text{-element vector of } 2\text{-element vectors}$

- If we know more, e.g., *cavity* is also given, then we have

$P(\text{cavity} \mid \text{toothache}, \text{cavity}) = 1$

- New evidence may be irrelevant, allowing simplification, e.g.,

$P(\text{cavity} \mid \text{toothache}, \text{sunny}) = P(\text{cavity} \mid \text{toothache}) = 0.8$

Conditional probability

- Definition of conditional probability:
 $P(a \mid b) = P(a \wedge b) / P(b)$ if $P(b) > 0$
- **Product rule** gives an alternative formulation:
 $P(a \wedge b) = P(a \mid b) P(b) = P(b \mid a) P(a)$
- A general version holds for whole distributions, e.g.,
 $\mathbf{P}(\textit{Weather}, \textit{Cavity}) = \mathbf{P}(\textit{Weather} \mid \textit{Cavity}) \mathbf{P}(\textit{Cavity})$
(View as a set of 4×2 equations, **not** matrix mult.)
- **Chain rule** is derived by successive application of product rule:

$$\begin{aligned} \mathbf{P}(X_1, \dots, X_n) &= \mathbf{P}(X_1, \dots, X_{n-1}) \mathbf{P}(X_n \mid X_1, \dots, X_{n-1}) \\ &= \mathbf{P}(X_1, \dots, X_{n-2}) \mathbf{P}(X_{n-1} \mid X_1, \dots, X_{n-2}) \mathbf{P}(X_n \mid X_1, \dots, X_{n-1}) \\ &= \dots \\ &= \prod_{i=1}^n \mathbf{P}(X_i \mid X_1, \dots, X_{i-1}) \end{aligned}$$

Probabilistic Inference

- Unconditional or prior probability: belief in absence of additional information
 - Evidence: additional information
 - Conditional or posterior probability: belief given the evidence in addition to the prior
-
- Probabilistic Inference: computing posterior probabilities for query propositions given prior probabilities and evidence.

Inference by enumeration

- Start with the joint probability distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

- For any proposition φ , sum the atomic events where it is true

Inference by enumeration

- Start with the joint probability distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

- For any proposition φ , sum the atomic events where it is true
- $P(\text{toothache}) = 0.108 + 0.012 + 0.016 + 0.064 = 0.2$

Inference by enumeration

- Start with the joint probability distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

- Can also compute conditional probabilities:

$$\begin{aligned} P(\neg \text{cavity} \mid \text{toothache}) &= \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} \\ &= 0.4 \end{aligned}$$

Normalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

- Denominator can be viewed as a **normalization constant** α

$$\begin{aligned} \mathbf{P}(\text{Cavity} \mid \text{toothache}) &= \alpha \mathbf{P}(\text{Cavity}, \text{toothache}) \\ &= \alpha [\mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch}) + \mathbf{P}(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [<0.108, 0.016> + <0.012, 0.064>] \\ &= \alpha <0.12, 0.08> = <0.6, 0.4> \end{aligned}$$

General idea: compute distribution on query variable by fixing **observed variables** and summing over **unobserved variables**

Inference by enumeration, contd.

Typically, we are interested in
the posterior joint distribution of the **query variables** \mathbf{Y}
given specific values \mathbf{e} for the **evidence (observed) variables** \mathbf{E}

Let the **hidden (unobserved) variables** be $\mathbf{H} = \mathbf{X} - \mathbf{Y} - \mathbf{E}$


Then the required summation of joint entries is done by summing out the hidden variables:

$$\mathbf{P}(\mathbf{Y} \mid \mathbf{E} = \mathbf{e}) = \alpha \mathbf{P}(\mathbf{Y}, \mathbf{E} = \mathbf{e}) = \alpha \sum_{\mathbf{h}} \mathbf{P}(\mathbf{Y}, \mathbf{E} = \mathbf{e}, \mathbf{H} = \mathbf{h})$$

- The terms in the summation are joint entries because \mathbf{Y} , \mathbf{E} and \mathbf{H} together exhaust the set of random variables
- Obvious problems:
 1. Worst-case time complexity $O(d^n)$ where d is the largest arity
 2. Space complexity $O(d^n)$ to store the joint distribution
 3. How to find the numbers for $O(d^n)$ entries?

Independence

- A and B are independent iff
 $P(A|B) = P(A)$ or $P(B|A) = P(B)$ or $P(A, B) = P(A) P(B)$



$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather}) \\ = P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) P(\textit{Weather})$$

- 32 entries reduced to 12; for n independent biased coins, $2^n \rightarrow n$
- Absolute independence powerful but rare
- Dentistry is a large field with hundreds of variables, none of which are independent. What to do?

Conditional independence

- $\mathbf{P}(\textit{Toothache}, \textit{Cavity}, \textit{Catch})$ has $2^3 - 1 = 7$ independent entries
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
(1) $\mathbf{P}(\textit{catch} \mid \textit{toothache}, \textit{cavity}) = \mathbf{P}(\textit{catch} \mid \textit{cavity})$
- The same independence holds if I haven't got a cavity:
(2) $\mathbf{P}(\textit{catch} \mid \textit{toothache}, \neg \textit{cavity}) = \mathbf{P}(\textit{catch} \mid \neg \textit{cavity})$
- *Catch* is **conditionally independent** of *Toothache* given *Cavity*:
 $\mathbf{P}(\textit{Catch} \mid \textit{Toothache}, \textit{Cavity}) = \mathbf{P}(\textit{Catch} \mid \textit{Cavity})$
- Equivalent statements:
 $\mathbf{P}(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) = \mathbf{P}(\textit{Toothache} \mid \textit{Cavity})$
 $\mathbf{P}(\textit{Toothache}, \textit{Catch} \mid \textit{Cavity}) = \mathbf{P}(\textit{Toothache} \mid \textit{Cavity}) \mathbf{P}(\textit{Catch} \mid \textit{Cavity})$

Conditional independence contd.

- Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) \mathbf{P}(\textit{Catch}, \textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) \mathbf{P}(\textit{Catch} \mid \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} \mid \textit{Cavity}) \mathbf{P}(\textit{Catch} \mid \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \end{aligned}$$

i.e., $2 + 2 + 1 = 5$ independent numbers

- In most cases, the use of conditional independence reduces the size of the representation of the joint distribution from exponential in n to linear in n .
- Conditional independence is our most basic and robust form of knowledge about uncertain environments.

Bayes' Rule

- Product rule $P(a \wedge b) = P(a \mid b) P(b) = P(b \mid a) P(a)$

⇒ **Bayes' rule:** $P(a \mid b) = P(b \mid a) P(a) / P(b)$

- or in distribution form

$$\mathbf{P(Y|X) = P(X|Y) P(Y) / P(X) = \alpha P(X|Y) P(Y)}$$

- Useful for assessing **diagnostic** probability from **causal** probability:
 - $P(\text{Cause} \mid \text{Effect}) = P(\text{Effect} \mid \text{Cause}) P(\text{Cause}) / P(\text{Effect})$
 - E.g., let M be meningitis, S be stiff neck:
 $P(m \mid s) = P(s \mid m) P(m) / P(s) = 0.8 \times 0.0001 / 0.1 = 0.0008$
 - Note: posterior probability of meningitis still very small!

Summary

- Probability is a rigorous formalism for uncertain knowledge
- Joint probability distribution specifies probability of every event
- Queries can be answered by summing over events
- For nontrivial domains, we must find a way to reduce the size of the joint distribution
- Independence and conditional independence provide the tools

Next Class

- Bayesian networks
- Sec 4.1 – 4.3