

Internship Report

**Mandelstam-Tamm Quantum Speed Limit  
in a Two Level System**

**SRM University, AP**

**Yamulapalli Nandu Navadeep**  
*IISER Thiruvananthapuram*

Mentor: **Dr. Swetamber P Das**  
*SRM University, AP*

July 23, 2026

## **Abstract**

This study report focused on understanding the Quantum Speed Limit (QSL), which defines the fundamental limit on the time required for a quantum system to evolve from one distinguishable state to another. The project explored the formulation of the Mandelstam-Tamm bound, and its application in a two-level quantum system.

# Contents

|          |                                                                 |           |
|----------|-----------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                             | <b>2</b>  |
| 1.1      | Background . . . . .                                            | 2         |
| 1.2      | Objectives . . . . .                                            | 2         |
| <b>2</b> | <b>Theory</b>                                                   | <b>4</b>  |
| 2.1      | The Energy-Time Uncertainty Principle . . . . .                 | 4         |
| 2.2      | Derivation of the Mandelstam-Tamm Bound . . . . .               | 5         |
| 2.2.1    | The Robertson Uncertainty Relation and Density Matrices . . . . | 5         |
| 2.2.2    | Integrating the Bound for Orthogonal States . . . . .           | 6         |
| 2.3      | Generalized Mandelstam-Tamm Bound . . . . .                     | 6         |
| <b>3</b> | <b>Application to a Two-State System</b>                        | <b>8</b>  |
| 3.1      | System Hamiltonian and Time Evolution . . . . .                 | 8         |
| 3.2      | Energy Variance and the MT Bound . . . . .                      | 9         |
| 3.3      | Generalized Mandelstam-Tamm Bound . . . . .                     | 9         |
| 3.3.1    | Equatorial Pure States and Bound Saturation . . . . .           | 11        |
| 3.4      | Computational Analysis . . . . .                                | 11        |
| <b>4</b> | <b>Results and Conclusion</b>                                   | <b>14</b> |
| 4.1      | Summary . . . . .                                               | 14        |
| 4.2      | Discussion and Physical Meaning . . . . .                       | 14        |
| 4.3      | Future Work . . . . .                                           | 15        |

# Chapter 1

## Introduction

This chapter introduces the background of the study, the motivation behind the project, and the main objectives that were set out to be achieved.

### 1.1 Background

Quantum mechanics places fundamental constraints on the dynamics of physical systems. Determining the minimum time required for a quantum system to evolve into a distinguishable state is a fundamental question that has direct implications for the ultimate speed of quantum computation and the fundamental limits of quantum control.

The energy-time uncertainty relation  $\Delta E \Delta t \geq \hbar/2$  has long been invoked to address this. Rearranging this inequality yields  $\Delta t \geq \frac{\hbar}{2\Delta E}$ , which physically implies that the timescale  $\Delta t$  over which a quantum system undergoes significant evolution is inversely proportional to its energy spread  $\Delta E$ . For example, an atom in an excited state with a very short lifetime ( $\Delta t$ ) will exhibit a broad spread in its measured emission energy (known as the natural linewidth,  $\Delta E$ ). Conversely, a stationary state with a perfectly defined energy ( $\Delta E = 0$ ) has an infinite lifetime ( $\Delta t \rightarrow \infty$ ) and never mathematically evolves. Thus, energy uncertainty acts as the fundamental "fuel" driving quantum evolution.

While practically useful, this relation's interpretation is historically subtle. A rigorous formulation was first provided by Mandelstam and Tamm in 1945, who bypassed these ambiguities by deriving a precise inequality that bounds how fast any dynamical quantity can change based purely on the system's exact energy variance.

### 1.2 Objectives

The primary objectives of this study were:

- To derive the Mandelstam-Tamm quantum speed limit.

- To generalize the MT bound to the case of non-orthogonal initial and final states.
- To apply the MT bound explicitly to a two-level quantum system governed by the Hamiltonian  $H = \frac{\hbar\omega_0}{2}\sigma_z$  (where  $\omega_0$  represents the characteristic transition frequency and  $\sigma_z$  is the Pauli-Z matrix), computing the energy uncertainty, the QSL time, and the time-dependent angle, and visualizing the resulting time evolution computationally.

# Chapter 2

## Theory

This chapter establishes the theoretical foundation for the Quantum Speed Limit, beginning with the historical context of the energy-time uncertainty principle and culminating in the derivation of the Mandelstam-Tamm bound.

### 2.1 The Energy-Time Uncertainty Principle

In standard quantum mechanics, the Heisenberg uncertainty principle places fundamental limits on the precision with which certain pairs of conjugate variables can be simultaneously known (e.g.,  $\Delta x \Delta p \geq \hbar/2$ ) [3].

Historically, a similar relation for energy and time has been widely used:

$$\Delta E \Delta t \geq \frac{\hbar}{2} \quad (2.1)$$

While practically useful, its theoretical interpretation fundamentally differs from the position-momentum uncertainty relation. In this expression,  $\Delta E$  represents the standard statistical uncertainty (standard deviation) of the system's energy ( $\Delta E = \Delta H = \sqrt{\langle H^2 \rangle - \langle H \rangle^2}$ ). However, in standard quantum mechanics, time  $t$  is not an observable represented by a Hermitian operator; it is an external, continuous parameter tracking dynamical evolution [3]. Consequently,  $\Delta t$  does not represent the statistical dispersion of multiple measurements. Instead,  $\Delta t$  represents a characteristic time scale: it is the minimum time required for the expectation value of some dynamical observable to change by an amount equal to its standard deviation.

To formalize this interpretation and apply it to state evolution, Mandelstam and Tamm (1945) circumvented the lack of a time operator by anchoring the concept of “time uncertainty” directly to the survival probability of the system [5]. They derived a lower bound on the time required for an initial quantum state to evolve into a distinguishable, orthogonal state, relying purely on the well-established Robertson uncertainty relation [2].

## 2.2 Derivation of the Mandelstam-Tamm Bound

The core of this project centers on deriving the Mandelstam-Tamm (MT) Quantum Speed Limit from first principles [2].

### 2.2.1 The Robertson Uncertainty Relation and Density Matrices

We begin with the generalized Robertson uncertainty relation. For a quantum system governed by the Hamiltonian  $H$  (the observable corresponding to the total energy of the system) and any arbitrary observable  $A$ , the uncertainties are bounded by their commutator  $[H, A]$ :

$$\Delta H \Delta A \geq \frac{1}{2} |\langle [H, A] \rangle| \quad (2.2)$$

Here, the angle brackets  $\langle . \rangle$  denote the quantum mechanical expectation value: the statistical average of measuring the enclosed operator in the given quantum state. To find a bound on the time evolution of the system, we need to relate the commutator to the time derivative of an observable. The time evolution of the expectation value of any operator  $A$  (that does not explicitly depend on time) is governed by the Heisenberg equation of motion, which is intimately connected to the von Neumann equation:

$$\frac{d\langle A \rangle}{dt} = \frac{i}{\hbar} \langle [H, A] \rangle \implies |\langle [H, A] \rangle| = \hbar \left| \frac{d\langle A \rangle}{dt} \right| \quad (2.3)$$

Substituting this into the uncertainty relation yields:

$$\Delta H \Delta A \geq \frac{\hbar}{2} \left| \frac{d\langle A \rangle}{dt} \right| \quad (2.4)$$

To measure how fast the system leaves its initial state  $|\psi(0)\rangle$ , we strategically choose the observable  $A$  to be the projection operator onto the initial state:  $A = |\psi(0)\rangle\langle\psi(0)|$ . This operator is mathematically equivalent to the density matrix of the initial pure state ( $\rho(0)$ ).

Density matrices are fundamentally involved in this formulation because they naturally encapsulate the statistical state of a quantum system, and their time evolution is governed directly by the von Neumann equation (the quantum analogue of Liouville's theorem in classical mechanics) [1]. A crucial property of the density matrix for a pure state (or any projection operator) is idempotency, meaning  $A^2 = A$ .

Using this property, the variance of  $A$  simplifies significantly:

$$\Delta A = \sqrt{\langle A^2 \rangle - \langle A \rangle^2} = \sqrt{\langle A \rangle - \langle A \rangle^2} \quad (2.5)$$

The expectation value  $\langle A \rangle_t = \langle \psi(t) | A | \psi(t) \rangle = |\langle \psi(0) | \psi(t) \rangle|^2$  represents the exact probability of finding the system still in its initial state at a later time  $t$ .

### 2.2.2 Integrating the Bound for Orthogonal States

Substituting our simplified  $\Delta A$  into the uncertainty relation gives:

$$\Delta H \sqrt{\langle A \rangle - \langle A \rangle^2} \geq \frac{\hbar}{2} \left| \frac{d\langle A \rangle}{dt} \right| \quad (2.6)$$

As the quantum system evolves away from its initial state, the expectation value  $\langle A \rangle = \text{Tr}(A\rho(t))$  (where  $\rho(t)$  is the time-dependent density matrix) decreases from its initial value of 1, meaning  $\frac{d\langle A \rangle}{dt} \leq 0$ . Thus,  $\left| \frac{d\langle A \rangle}{dt} \right| = -\frac{d\langle A \rangle}{dt}$ . Separating variables and integrating from  $t = 0$  (where  $\langle A \rangle_0 = 1$ ) to time  $t$ :

$$\int_0^t \frac{2\Delta H}{\hbar} dt \geq \int_1^{\langle A \rangle_t} \frac{-d\langle A \rangle}{\sqrt{\langle A \rangle(1 - \langle A \rangle)}} \quad (2.7)$$

By substituting  $\langle A \rangle = \sin^2 \theta$ , the integral evaluates exactly to:

$$\frac{2\Delta H t}{\hbar} \geq \pi - 2 \arcsin(\sqrt{\langle A \rangle_t}) \implies \frac{\Delta H t}{\hbar} \geq \frac{\pi}{2} - \arcsin(\sqrt{\langle A \rangle_t}) \quad (2.8)$$

To find the minimum time required for the system to reach a completely *orthogonal* state, which is physically crucial because only perfectly orthogonal states are completely distinguishable from one another upon measurement, representing the time required to complete a distinct quantum operation, we set  $\langle \psi(0) | \psi(\tau) \rangle = 0$ , which means  $\langle A \rangle_\tau = 0$ .

$$\frac{\Delta H \tau_{QSL}}{\hbar} \geq \frac{\pi}{2} - \arcsin(0) = \frac{\pi}{2} \quad (2.9)$$

Rearranging for  $\tau_{QSL}$ , we recover the famous Mandelstam-Tamm Quantum Speed Limit for orthogonal states:

$$\tau_{QSL} \geq \frac{\pi \hbar}{2\Delta H} \quad (2.10)$$

## 2.3 Generalized Mandelstam-Tamm Bound

In many practical scenarios, a quantum system might not evolve to a completely orthogonal state, but rather to a target state separated by some Bures angle  $\Theta$ . The Bures angle is simply a geometric measure of the distinguishability, or the mathematical “distance”, between two quantum states in the Hilbert space. We can explicitly show that the state overlap is equivalent to the trace of the product of the density matrices at  $t = 0$



( $A_0 = |\psi(0)\rangle\langle\psi(0)|$ ) and time  $t$  ( $A_t = |\psi(t)\rangle\langle\psi(t)|$ ). By evaluating the trace, we find:

$$\begin{aligned}
\langle A \rangle_t &= \text{Tr}(A_0 A_t) = \text{Tr}\left(|\psi(0)\rangle\langle\psi(0)|\psi(t)\rangle\langle\psi(t)|\right) \\
&= \langle\psi(t)|\psi(0)\rangle\langle\psi(0)|\psi(t)\rangle \\
&= |\langle\psi(0)|\psi(t)\rangle|^2 \\
&= \cos^2 \Theta
\end{aligned} \tag{2.11}$$

Plugging this non-orthogonal overlap into our integrated bound gives:

$$\begin{aligned}
\frac{\Delta H \Delta t}{\hbar} &\geq \frac{\pi}{2} - \arcsin\left(\sqrt{\cos^2 \Theta}\right) \\
&= \frac{\pi}{2} - \arcsin(\cos \Theta) \\
&= \frac{\pi}{2} - \left(\frac{\pi}{2} - \Theta\right) = \Theta
\end{aligned} \tag{2.12}$$

This provides the generalized Mandelstam-Tamm bound for any arbitrary angle  $\Theta$ :

$$\Delta t \geq \frac{\hbar \Theta}{\Delta H} \tag{2.13}$$

This generalized formulation ensures that the time interval  $\Delta t$  required to traverse an angle  $\Theta$  in the Hilbert space is fundamentally restricted by the energy uncertainty of the system. In other words,  $\Delta t$  represents the absolute minimum duration, the true quantum speed limit, for any physical process to drive the system between two states separated by  $\Theta$ , because no physical mechanism can possibly force the state to evolve faster than this variance dependent timescale.

# Chapter 3

## Application to a Two-State System

This chapter applies the theoretical framework of Mandelstam and Tamm's Quantum Speed Limit discussed in the methodology to a specific two-state quantum system.

### 3.1 System Hamiltonian and Time Evolution

Consider a two-state quantum system governed by the Hamiltonian:

$$H = \frac{\hbar\omega_0}{2}\sigma_z = \frac{\hbar\omega_0}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3.1)$$

where  $\sigma_z$  is the Pauli Z matrix. This time-independent, diagonal Hamiltonian is the standard model for a two-level system, such as a spin-1/2 particle in a constant magnetic field. Its simplicity provides exact analytical solutions for state evolution and energy variance, making it ideal for clearly demonstrating the Mandelstam-Tamm bound.

To compute how the system changes over time, we use the unitary time evolution operator  $U(t)$ . In quantum mechanics, a unitary operator (where  $U^\dagger U = I$ ) is fundamental because it strictly preserves the inner product and the norm of state vectors, ensuring that total probability is perfectly conserved. By applying  $U(t)$ , we can deterministically rotate the initial state through the Hilbert space. This unitary preservation is what mathematically allows us to accurately track the survival probability and Bures angle. For this time-independent Hamiltonian, the evolution operator is given by:

$$U(t) = e^{-iHt/\hbar} = \begin{pmatrix} e^{-i\omega_0 t/2} & 0 \\ 0 & e^{i\omega_0 t/2} \end{pmatrix} \quad (3.2)$$

## 3.2 Energy Variance and the MT Bound

To apply the Mandelstam-Tamm bound, we first compute the energy variance  $\Delta H$  of the initial state. The expectation value of the Hamiltonian is:

$$\langle H \rangle = \langle \psi(0) | H | \psi(0) \rangle = \frac{\hbar\omega_0}{2} \left( \cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2} \right) = \frac{\hbar\omega_0}{2} \cos \theta \quad (3.3)$$

Since  $H^2 = \left(\frac{\hbar\omega_0}{2}\right)^2 I$ , the expectation value of  $H^2$  is simply  $\langle H^2 \rangle = \left(\frac{\hbar\omega_0}{2}\right)^2$ . Taking the square root of the energy variance yields the standard deviation  $\Delta H$ , representing the state's energy uncertainty:

$$\Delta H = \sqrt{\langle H^2 \rangle - \langle H \rangle^2} = \sqrt{\left(\frac{\hbar\omega_0}{2}\right)^2 - \left(\frac{\hbar\omega_0}{2} \cos \theta\right)^2} = \frac{\hbar\omega_0}{2} \sin \theta \quad (3.4)$$

According to the Mandelstam-Tamm bound, the time  $\Delta t$  required for the system to evolve to a state separated by a Bures angle  $\Theta$  from the initial state is bounded by:

$$t \geq \frac{\hbar\Theta}{\Delta H} = \frac{\hbar\Theta}{\frac{\hbar\omega_0}{2} \sin \theta} = \frac{2\Theta}{\omega_0 \sin \theta} \quad (3.5)$$

For the system to reach an orthogonal state, the distance is  $\Theta = \pi/2$ . The minimum time required to reach this orthogonal state is thus:

$$t \geq \frac{2 \cdot (\pi/2)}{\omega_0 \sin \theta} \implies \tau_{QSL} = \frac{\pi}{\omega_0 \sin \theta} \quad (3.6)$$

This indicates that the speed of evolution depends on the initial state parameter  $\theta$ . The fastest evolution to an orthogonal state occurs when  $\theta = \pi/2$  (states on the equator of the Bloch sphere), yielding a minimum time of  $\tau_{QSL} = \pi/\omega_0$ .

The Bloch sphere is a geometrical representation of the pure state space of a two-level quantum mechanical system (a qubit). The north and south poles represent the mutually orthogonal computational basis states  $|0\rangle$  and  $|1\rangle$ , while any point on the equator ( $\theta = \pi/2$ ) represents an equal superposition of these basis states [4].

## 3.3 Generalized Mandelstam-Tamm Bound

Let the initial state of the system be parameterized by an angle  $\theta$  on the Bloch sphere:

$$|\psi(0)\rangle = \cos \frac{\theta}{2} |0\rangle + \sin \frac{\theta}{2} |1\rangle \quad (3.7)$$




Figure 3.1: The Bloch sphere representation of a two-level quantum system state. Figure adapted from [4].

The state at a later time  $t$  is:

$$|\psi(t)\rangle = U(t)|\psi(0)\rangle = \cos \frac{\theta}{2} e^{-i\omega_0 t/2} |0\rangle + \sin \frac{\theta}{2} e^{i\omega_0 t/2} |1\rangle \quad (3.8)$$

The inner product between the initial and evolved state is:

$$\langle\psi(0)|\psi(t)\rangle = \cos^2 \frac{\theta}{2} e^{-i\omega_0 t/2} + \sin^2 \frac{\theta}{2} e^{i\omega_0 t/2} = \cos \left( \frac{\omega_0 t}{2} \right) - i \cos \theta \sin \left( \frac{\omega_0 t}{2} \right) \quad (3.9)$$

The squared magnitude of this overlap gives the fidelity:

$$|\langle\psi(0)|\psi(t)\rangle|^2 = \cos^2 \left( \frac{\omega_0 t}{2} \right) + \cos^2 \theta \sin^2 \left( \frac{\omega_0 t}{2} \right) \quad (3.10)$$

The Bures angle  $\alpha(t)$  between the initial and evolved state is defined as:

$$\alpha(t, \theta) = \arccos (|\langle\psi(0)|\psi(t)\rangle|) = \arccos \sqrt{\cos^2 \left( \frac{\omega_0 t}{2} \right) + \cos^2 \theta \sin^2 \left( \frac{\omega_0 t}{2} \right)} \quad (3.11)$$

Furthermore, adding an arbitrary relative phase  $\phi$  to the initial state:

$$|\psi(0)\rangle = \cos \frac{\theta}{2} |0\rangle + e^{i\phi} \sin \frac{\theta}{2} |1\rangle \quad (3.12)$$

shows that  $e^{i\phi}$  cancels during the inner product calculation. So,  $\alpha(t)$  is independent of  $\phi$ .

### 3.3.1 Equatorial Pure States and Bound Saturation

To determine which initial pure states can ever evolve into a completely orthogonal state ( $\alpha = \pi/2$ ), we must set the argument of the arccosine to zero:

$$\cos^2\left(\frac{\omega_0 t}{2}\right) + \cos^2\theta \sin^2\left(\frac{\omega_0 t}{2}\right) = 0 \quad (3.13)$$

Because this is a sum of two non-negative squares, both terms must equal zero simultaneously. The first term dictates that  $\cos(\omega_0 t/2) = 0$ , meaning the minimum time to reach orthogonality is  $t = \pi/\omega_0$ . At this specific time,  $\sin^2(\omega_0 t/2) = 1$ . Consequently, for the second term to vanish, we must have  $\cos\theta = 0$ , which strictly restricts the initial state to  $\theta = \pi/2$ .

This mathematical condition reveals a profound physical property: only the specific family of pure states lying exactly on the equator of the Bloch sphere can ever reach a state completely orthogonal to themselves. For these equatorial states, the time-dependent Bures angle simplifies perfectly to a linear function:

$$\alpha\left(t, \theta = \frac{\pi}{2}\right) = \frac{\omega_0 t}{2} \quad (3.14)$$

Crucially, we can show that the Mandelstam-Tamm bound saturates (reaches equality) for this specific family of states. Substituting  $\theta = \pi/2$  into our earlier energy variance equation gives the maximum possible uncertainty  $\Delta H = \hbar\omega_0/2$ . The theoretical Quantum Speed Limit time to reach an orthogonal state ( $\Theta = \pi/2$ ) is therefore:

$$\tau_{QSL} = \frac{\pi\hbar}{2\Delta H} = \frac{\pi\hbar}{2(\hbar\omega_0/2)} = \frac{\pi}{\omega_0} \quad (3.15)$$

This strictly matches the dynamical time  $t = \pi/\omega_0$  derived above, proving that equatorial pure states evolve at the absolute maximum physical speed permitted by quantum mechanics.

## 3.4 Computational Analysis

We used Python to plot the Bures angle  $\alpha(t)$  against dimensionless time  $\omega_0 t$  for various initial states ( $\theta$ ) to analyze the generalized MT bound.

Figure 3.2 provides a visualization of the system's evolution for various initial states, defined by different values of the polar angle  $\theta$  on the Bloch sphere, ranging from the poles ( $\theta = 0$ ) to the equator ( $\theta = \pi/2$ ). The horizontal axis represents dimensionless time  $\omega_0 t$ . Using dimensionless time allows the plot to be universally applicable, meaning the fundamental shape of the curves remains the same regardless of the specific physical value of the natural frequency  $\omega_0$ .




Figure 3.2: Generalized Mandelstam-Tamm Bound:  $\alpha(t)$  vs  $\omega_0 t$  for different initial states  $\theta$ . The black dashed line represents the MT bound for orthogonal states ( $\theta = \pi/2$ ), and the shaded gray region is forbidden by the bound.

The plot prominently features a straight diagonal dashed line. This diagonal represents the absolute maximum rate of evolution permitted by the Mandelstam-Tamm bound. Because the generalized bound states  $\alpha \leq \Delta H t / \hbar$ , and the maximum energy variance is  $\Delta H = \hbar \omega_0 / 2$ , this limiting diagonal has a mathematical slope of exactly  $1/2$  (where  $\alpha = \omega_0 t / 2$ ). The shaded gray area above this diagonal is the forbidden region. No physical quantum state can evolve fast enough to cross into this area, as doing so would violate the fundamental energy-time uncertainty principle.

Every curve exhibits a smooth, periodic oscillation. This sinusoidal shape directly reflects the unitary nature of the time evolution operator ( $U(t) = e^{-iHt/\hbar}$ ). Because unitary operators expand mathematically into sine and cosine functions, they strictly conserve the total probability of the system without any decay. Physically, this guarantees that the state vector is continuously and uniformly precessing around the z-axis of the Bloch sphere. Due to this continuous rotation, the state periodically reaches a point of maximum distinguishability at half the oscillation period (when  $\omega_0 t = \pi$ ) before smoothly rotating back to its exact initial configuration to complete the cycle.

Furthermore, the curves clearly show that only the equatorial states ( $\theta = \pi/2$ ) evolve fast enough to touch the diagonal bound and successfully reach a completely orthogonal state ( $\alpha = \pi/2$ ). As the initial state parameter  $\theta$  decreases toward 0, the curves become progressively flatter. This flattening occurs because moving closer to the poles of the Bloch sphere decreases the energy variance  $\Delta H$ . With less energy uncertainty, the system is strictly forced to evolve more slowly, which fundamentally limits the maximum Bures angle it can ever reach. In the extreme case where  $\theta = 0$ , the state is an energy eigenstate

(a stationary state) with zero variance, and the curve remains perfectly flat at  $\alpha = 0$  to indicate that no state evolution occurs.

# Chapter 4

## Results and Conclusion

This chapter summarizes the findings of the study, discusses the physical implications of the derived bound, and outlines potential directions for future study.

### 4.1 Summary

This project explored the fundamental limit of quantum dynamics. We began by reviewing the underlying quantum mechanical formalism, establishing the necessary theoretical foundation in state vectors, operators, and density matrices. With this framework, we derived the Mandelstam-Tamm (MT) Quantum Speed Limit from the Robertson uncertainty relation.

The bound was then generalized to account for the evolution between non-orthogonal initial and final states. Finally, we explicitly applied this generalized bound to a two-level quantum system and verified it computationally. The numerical simulations successfully confirmed that the speed of the system's state evolution is strictly bounded by its energy variance, with the fastest possible evolution occurring for initial states with maximum energy uncertainty (states on the equator of the Bloch sphere).

### 4.2 Discussion and Physical Meaning

The Mandelstam-Tamm bound holds profound implications for the physical reality of dynamic systems. It dictates that quantum states possess a fundamental, inescapable “speed limit” that is entirely governed by their energy variance ( $\Delta H$ ).

In practical terms, this places a hard physical limit on how fast quantum processes can occur. For instance, in the realm of quantum computing, the clock speed of a quantum processor, the minimum time required to reliably flip a qubit from  $|0\rangle$  to  $|1\rangle$  (an orthogonal state), cannot be made arbitrarily short. It is strictly bounded by the energy resources (specifically, the energy spread) available to the system. A faster operation



inherently requires a larger energy variance. This means that designing ultra-fast quantum computers or highly efficient quantum thermodynamic engines will eventually hit a fundamental physical bottleneck dictated by the MT bound, representing the absolute theoretical maximum of quantum control.

### 4.3 Future Work

While this internship focused deeply on the Mandelstam-Tamm formulation, the study of the Quantum Speed Limit is a vast and active field of research. Future work could build upon this foundation in several directions:

- **The Margolus-Levitin Bound:** Exploring the other primary quantum speed limit, the Margolus-Levitin (ML) bound. A fundamental limitation of the MT bound is that for quantum states with a highly dispersed energy spectrum, the variance  $\Delta H$  can become arbitrarily large, which drives the MT minimum time toward zero (falsely suggesting instantaneous evolution). To resolve this, the ML bound provides an alternative limit based on the system's mean energy above the ground state ( $E - E_0$ ). Investigating the crossover regime to determine which bound is tighter for a given state provides a much more complete picture of pure state dynamics.
- **Open Quantum Systems:** The current study assumed a closed quantum system undergoing isolated, unitary time evolution. A natural next step would be to apply the QSL to open quantum systems, where the system interacts with an external environment, leading to decoherence and non-unitary dynamics. Decoherence refers to the loss of a system's quantum superposition as its delicate phase information leaks into the surrounding environment. Consequently, the evolution becomes non-unitary, meaning the system's internal probability is no longer perfectly preserved. Understanding the QSL in this regime is essential because these environmental interactions can fundamentally alter the speed limit.
- **Many-Body Systems:** Extending the application of this bound from a simple two-state system to complex many-body quantum systems. In such systems, particles exhibit entanglement, a phenomenon where their quantum states become intrinsically linked such that one particle cannot be described independently of the others. These quantum correlations are highly important because they can cause the collective system to evolve significantly faster or slower than its individual, isolated components. Understanding how entanglement impacts the Quantum Speed Limit is crucial for accurately predicting the theoretical performance of large-scale, multi-qubit quantum computers.

# Bibliography

- [1] Reinhold A. Bertlmann and Nicolai Friis. Theoretical physics t2: Quantum mechanics. Course of Lectures, Sommersemester 2008, 2008.
- [2] Sebastian Deffner and Steve Campbell. Quantum speed limits: from heisenberg’s uncertainty principle to optimal quantum control. *Journal of Physics A: Mathematical and Theoretical*, 50(45):453001, 2017.
- [3] David J. Griffiths. *Introduction to Quantum Mechanics*. Pearson Prentice Hall, 2nd edition, 2004.
- [4] Farzan Jazaeri, Arnout Beckers, Armin Tajalli, and Jean-Michel Sallese. A review on quantum computing: Qubits, cryogenic electronics and cryogenic mosfet physics. *arXiv preprint arXiv:1908.02656*, 2019.
- [5] Leonid Mandelstam and Igor Tamm. The uncertainty relation between energy and time in non-relativistic quantum mechanics. *J. Phys.(USSR)*, 9:249–254, 1945.