

Policy Gradient Algorithms

Ashwin Rao

ICME, Stanford University

Overview

- 1 Motivation and Intuition
- 2 Definitions and Notation
- 3 Policy Gradient Theorem and Proof
- 4 Policy Gradient Algorithms
- 5 Compatible Function Approximation Theorem and Proof
- 6 Natural Policy Gradient

Why do we care about Policy Gradient (PG)?

- Let us review how we got here
- We started with Markov Decision Processes and Bellman Equations
- Next we studied several variants of DP and RL algorithms
- We noted that the idea of *Generalized Policy Iteration* (GPI) is key
- Policy Improvement step: $\pi(s, a)$ derived from $\operatorname{argmax}_a Q(s, a)$
- How do we do argmax when action space is large or continuous?
- Idea: Do Policy Improvement step with a Gradient Ascent instead

“Policy Improvement with a Gradient Ascent??”

- We want to find the Policy that fetches the “Best Expected Returns”
- Gradient Ascent on “Expected Returns” w.r.t params of Policy func
- So we need a func approx for (stochastic) Policy Func: $\pi(s, a; \theta)$
- In addition to the usual func approx for Action Value Func: $Q(s, a; \mathbf{w})$
- $\pi(s, a; \theta)$ called *Actor* and $Q(s, a; \mathbf{w})$ called *Critic*
- Critic parameters \mathbf{w} are optimized w.r.t $Q(s, a; \mathbf{w})$ loss function min
- Actor parameters θ are optimized w.r.t Expected Returns max
- We need to formally define “Expected Returns”
- But we already see that this idea is appealing for continuous actions
- GPI with Policy Improvement done as **Policy Gradient (Ascent)**

Value Function-based and Policy-based RL

- Value Function-based
 - Learn Value Function (with a function approximation)
 - Policy is implicit - readily derived from Value Function (eg: ϵ -greedy)
- Policy-based
 - Learn Policy (with a function approximation)
 - No need to learn a Value Function
- Actor-Critic
 - Learn Policy (Actor)
 - Learn Value Function (Critic)

Advantages and Disadvantages of Policy Gradient approach

Advantages:

- Finds the best *Stochastic* Policy (Optimal Deterministic Policy, produced by other RL algorithms, can be unsuitable for POMDPs)
- Naturally *explores* due to Stochastic Policy representation
- Effective in high-dimensional or continuous action spaces
- Small changes in $\theta \Rightarrow$ small changes in π , and in state distribution
- This avoids the convergence issues seen in argmax-based algorithms

Disadvantages:

- Typically converge to a local optimum rather than a global optimum
- Policy Evaluation is typically inefficient and has high variance
- Policy Improvement happens in small steps \Rightarrow slow convergence

- Assume episodic with $0 \leq \gamma \leq 1$ or non-episodic with $0 \leq \gamma < 1$
- Usual notation of discrete-time, countable-spaces, stationary MDPs
- We lighten $\mathcal{P}(s, a, s')$ notation to $\mathcal{P}_{s,s'}^a$ and $\mathcal{R}(s, a)$ notation to \mathcal{R}_s^a
- Initial State Probability Distribution denoted as $p_0 : \mathcal{N} \rightarrow [0, 1]$
- Policy Function Approximation $\pi(s, a; \theta) = \mathbb{P}[A_t = a | S_t = s; \theta]$

PG coverage is quite similar for non-discounted non-episodic, by considering average-reward objective (we won't cover it)

“Expected Returns” Objective

Now we formalize the “Expected Returns” Objective $J(\theta)$

$$J(\theta) = \mathbb{E}_{\pi} \left[\sum_{t=0}^{\infty} \gamma^t \cdot R_{t+1} \right]$$

Value Function $V^{\pi}(s)$ and Action Value function $Q^{\pi}(s, a)$ defined as:

$$V^{\pi}(s) = \mathbb{E}_{\pi} \left[\sum_{k=t}^{\infty} \gamma^{k-t} \cdot R_{k+1} \mid S_t = s \right] \text{ for all } t = 0, 1, 2, \dots$$

$$Q^{\pi}(s, a) = \mathbb{E}_{\pi} \left[\sum_{k=t}^{\infty} \gamma^{k-t} \cdot R_{k+1} \mid S_t = s, A_t = a \right] \text{ for all } t = 0, 1, 2, \dots$$

$$\text{Advantage Function } A^{\pi}(s, a) = Q^{\pi}(s, a) - V^{\pi}(s)$$

Also, $p(s \rightarrow s', t, \pi)$ will be a key function for us - it denotes the probability of going from state s to s' in t steps by following policy π

Discounted-Aggregate State-Visitation Measure

$$\begin{aligned} J(\theta) &= \mathbb{E}_{\pi} \left[\sum_{t=0}^{\infty} \gamma^t \cdot R_{t+1} \right] = \sum_{t=0}^{\infty} \gamma^t \cdot \mathbb{E}_{\pi} [R_{t+1}] \\ &= \sum_{t=0}^{\infty} \gamma^t \cdot \sum_{s \in \mathcal{N}} \left(\sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot p(S_0 \rightarrow s, t, \pi) \right) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot \mathcal{R}_s^a \\ &= \sum_{s \in \mathcal{N}} \left(\sum_{S_0 \in \mathcal{N}} \sum_{t=0}^{\infty} \gamma^t \cdot p_0(S_0) \cdot p(S_0 \rightarrow s, t, \pi) \right) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot \mathcal{R}_s^a \end{aligned}$$

Definition

$$J(\theta) = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot \mathcal{R}_s^a$$

where $\rho^{\pi}(s) = \sum_{S_0 \in \mathcal{N}} \sum_{t=0}^{\infty} \gamma^t \cdot p_0(S_0) \cdot p(S_0 \rightarrow s, t, \pi)$ is the key function (for PG) we'll refer to as *Discounted-Aggregate State-Visitation Measure*.

Policy Gradient Theorem (PGT)

Theorem

$$\nabla_{\theta} J(\theta) = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q^{\pi}(s, a)$$

- Note: $\rho^{\pi}(s)$ depends on θ , but there's no $\nabla_{\theta} \rho^{\pi}(s)$ term in $\nabla_{\theta} J(\theta)$
- So we can simply generate sampling traces, and at each time step, calculate $(\nabla_{\theta} \log \pi(s, a; \theta)) \cdot Q^{\pi}(s, a)$ (probabilities implicit in paths)
- Note: $\nabla_{\theta} \log \pi(s, a; \theta)$ is Score function (Gradient of log-likelihood)
- We will estimate $Q^{\pi}(s, a)$ with a function approximation $Q(s, a; \mathbf{w})$
- We will later show how to avoid the estimate bias of $Q(s, a; \mathbf{w})$
- This numerical estimate of $\nabla_{\theta} J(\theta)$ enables **Policy Gradient Ascent**
- Let us look at the score function of some canonical $\pi(s, a; \theta)$

Canonical $\pi(s, a; \theta)$ for finite action spaces

- For finite action spaces, we often use Softmax Policy
- θ is an m -vector $(\theta_1, \dots, \theta_m)$
- Features vector $\phi(s, a) = (\phi_1(s, a), \dots, \phi_m(s, a))$ for all $s \in \mathcal{N}, a \in \mathcal{A}$
- Weight actions using linear combinations of features: $\phi(s, a)^T \cdot \theta$
- Action probabilities proportional to exponentiated weights:

$$\pi(s, a; \theta) = \frac{e^{\phi(s, a)^T \cdot \theta}}{\sum_{b \in \mathcal{A}} e^{\phi(s, b)^T \cdot \theta}} \text{ for all } s \in \mathcal{N}, a \in \mathcal{A}$$

- The score function is:

$$\nabla_{\theta} \log \pi(s, a; \theta) = \phi(s, a) - \sum_{b \in \mathcal{A}} \pi(s, b; \theta) \cdot \phi(s, b) = \phi(s, a) - \mathbb{E}_{\pi}[\phi(s, \cdot)]$$

Canonical $\pi(s, a; \theta)$ for continuous action spaces

- For continuous action spaces, we often use Gaussian Policy
- θ is an m -vector $(\theta_1, \dots, \theta_m)$
- State features vector $\phi(s) = (\phi_1(s), \dots, \phi_m(s))$ for all $s \in \mathcal{N}$
- Gaussian Mean is a linear combination of state features $\phi(s)^T \cdot \theta$
- Variance may be fixed σ^2 , or can also be parameterized
- Policy is Gaussian, $a \sim \mathcal{N}(\phi(s)^T \cdot \theta, \sigma^2)$ for all $s \in \mathcal{N}$
- The score function is:

$$\nabla_{\theta} \log \pi(s, a; \theta) = \frac{(a - \phi(s)^T \cdot \theta) \cdot \phi(s)}{\sigma^2}$$

Proof of Policy Gradient Theorem

We begin the proof by noting that:

$$J(\theta) = \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot V^\pi(S_0) = \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot Q^\pi(S_0, A_0)$$

Calculate $\nabla_\theta J(\theta)$ by parts $\pi(S_0, A_0; \theta)$ and $Q^\pi(S_0, A_0)$

$$\begin{aligned} \nabla_\theta J(\theta) &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_\theta \pi(S_0, A_0; \theta) \cdot Q^\pi(S_0, A_0) \\ &\quad + \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \nabla_\theta Q^\pi(S_0, A_0) \end{aligned}$$

Proof of Policy Gradient Theorem

Now expand $Q^\pi(S_0, A_0)$ as:

$$\mathcal{R}_{S_0}^{A_0} + \sum_{S_1 \in \mathcal{N}} \gamma \cdot \mathcal{P}_{S_0, S_1}^{A_0} \cdot V^\pi(S_1) \text{ (Bellman Policy Equation)}$$

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^\pi(S_0, A_0) + \\ &\quad \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \nabla_{\theta} (\mathcal{R}_{S_0}^{A_0} + \sum_{S_1 \in \mathcal{N}} \gamma \cdot \mathcal{P}_{S_0, S_1}^{A_0} \cdot V^\pi(S_1)) \end{aligned}$$

Note: $\nabla_{\theta} \mathcal{R}_{S_0}^{A_0} = 0$, so remove that term

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^\pi(S_0, A_0) + \\ &\quad \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \nabla_{\theta} \left(\sum_{S_1 \in \mathcal{N}} \gamma \cdot \mathcal{P}_{S_0, S_1}^{A_0} \cdot V^\pi(S_1) \right) \end{aligned}$$

Proof of Policy Gradient Theorem

Now bring the ∇_{θ} inside the $\sum_{S_1 \in \mathcal{N}}$ to apply only on $V^{\pi}(S_1)$

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^{\pi}(S_0, A_0) + \\ &\quad \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \sum_{S_1 \in \mathcal{N}} \gamma \cdot \mathcal{P}_{S_0, S_1}^{A_0} \cdot \nabla_{\theta} V^{\pi}(S_1) \end{aligned}$$

Now bring the outside $\sum_{S_0 \in \mathcal{N}}$ and $\sum_{A_0 \in \mathcal{A}}$ inside the $\sum_{S_1 \in \mathcal{N}}$

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^{\pi}(S_0, A_0) + \\ &\quad \sum_{S_1 \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \gamma \cdot p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \mathcal{P}_{S_0, S_1}^{A_0} \cdot \nabla_{\theta} V^{\pi}(S_1) \end{aligned}$$

Policy Gradient Theorem

Note that $\sum_{A_0 \in \mathcal{A}} \pi(S_0, A_0; \theta) \cdot \mathcal{P}_{S_0, S_1}^{A_0} = p(S_0 \rightarrow S_1, 1, \pi)$

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^{\pi}(S_0, A_0) + \\ &\quad \sum_{S_1 \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \gamma \cdot p_0(S_0) \cdot p(S_0 \rightarrow S_1, 1, \pi) \cdot \nabla_{\theta} V^{\pi}(S_1) \end{aligned}$$

Now expand $V^{\pi}(S_1)$ to $\sum_{A_1 \in \mathcal{A}} \pi(S_1, A_1; \theta) \cdot Q^{\pi}(S_1, A_1)$

$$\begin{aligned} &= \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^{\pi}(S_0, A_0) + \\ &\quad \sum_{S_1 \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \gamma \cdot p_0(S_0) \cdot p(S_0 \rightarrow S_1, 1, \pi) \cdot \nabla_{\theta} \left(\sum_{A_1 \in \mathcal{A}} \pi(S_1, A_1; \theta) \cdot Q^{\pi}(S_1, A_1) \right) \end{aligned}$$

Proof of Policy Gradient Theorem

We are now back to when we started calculating gradient of $\sum_a \pi \cdot Q^\pi$. Follow the same process of splitting $\pi \cdot Q^\pi$, then Bellman-expanding Q^π (to calculate its gradient), and iterate.

$$\nabla_{\theta} J(\theta) = \sum_{S_0 \in \mathcal{N}} p_0(S_0) \cdot \sum_{A_0 \in \mathcal{A}} \nabla_{\theta} \pi(S_0, A_0; \theta) \cdot Q^\pi(S_0, A_0) +$$

$$\sum_{S_1 \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \gamma \cdot p_0(S_0) \cdot p(S_0 \rightarrow S_1, 1, \pi) \cdot \left(\sum_{A_1 \in \mathcal{A}} \nabla_{\theta} \pi(S_1, A_1; \theta) \cdot Q^\pi(S_1, A_1) + \dots \right)$$

This iterative process leads us to:

$$= \sum_{t=0}^{\infty} \sum_{S_t \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \gamma^t \cdot p_0(S_0) \cdot p(S_0 \rightarrow S_t, t, \pi) \cdot \sum_{A_t \in \mathcal{A}} \nabla_{\theta} \pi(S_t, A_t; \theta) \cdot Q^\pi(S_t, A_t)$$

Proof of Policy Gradient Theorem

Bring $\sum_{t=0}^{\infty}$ inside $\sum_{S_t \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}}$ and note that

$$\sum_{A_t \in \mathcal{A}} \nabla_{\theta} \pi(S_t, A_t; \theta) \cdot Q^{\pi}(S_t, A_t) \text{ is independent of } t$$

$$= \sum_{s \in \mathcal{N}} \sum_{S_0 \in \mathcal{N}} \sum_{t=0}^{\infty} \gamma^t \cdot p_0(S_0) \cdot p(S_0 \rightarrow s, t, \pi) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q^{\pi}(s, a)$$

Reminder that $\sum_{S_0 \in \mathcal{N}} \sum_{t=0}^{\infty} \gamma^t \cdot p_0(S_0) \cdot p(S_0 \rightarrow s, t, \pi) \stackrel{\text{def}}{=} \rho^{\pi}(s)$. So,

$$\nabla_{\theta} J(\theta) = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q^{\pi}(s, a)$$

Q.E.D.

Monte-Carlo Policy Gradient (REINFORCE Algorithm)

- Update θ by stochastic gradient ascent using PGT
- Using $G_t = \sum_{k=t}^T \gamma^{k-t} \cdot R_{k+1}$ as an unbiased sample of $Q^\pi(S_t, A_t)$

$$\Delta \theta = \alpha \cdot \gamma^t \cdot \nabla_{\theta} \log \pi(S_t, A_t; \theta) \cdot G_t$$

Algorithm 4.1: REINFORCE(\cdot)

Initialize θ arbitrarily

for each episode $\{S_0, A_0, R_1, S_1, \dots, S_{T-1}, A_{T-1}, R_T, S_T\} \sim \pi(\cdot, \cdot; \theta)$

do $\left\{ \begin{array}{l} \text{for } t \leftarrow 0 \text{ to } T \\ \text{do } \left\{ \begin{array}{l} G \leftarrow \sum_{k=t}^T \gamma^{k-t} \cdot R_{k+1} \\ \theta \leftarrow \theta + \alpha \cdot \gamma^t \cdot \nabla_{\theta} \log \pi(S_t, A_t; \theta) \cdot G \end{array} \right. \end{array} \right.$

Reducing Variance using a Critic

- Monte Carlo Policy Gradient has high variance
- We use a Critic $Q(s, a; \mathbf{w})$ to estimate $Q^\pi(s, a)$
- Actor-Critic algorithms maintain two sets of parameters:
 - Critic updates parameters \mathbf{w} to approximate Q -function for policy π
 - Critic could use any of the algorithms we learnt earlier:
 - Monte Carlo policy evaluation
 - Temporal-Difference Learning
 - $TD(\lambda)$ based on Eligibility Traces
 - Could even use LSTD (if critic function approximation is linear)
 - Actor updates policy parameters θ in direction suggested by Critic
 - This is Approximate Policy Gradient due to *Bias* of Critic

$$\nabla_{\theta} J(\theta) \approx \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q(s, a; \mathbf{w})$$

So what does the algorithm look like?

- Generate a sufficient set of sampling traces
 $S_0, A_0, R_1, S_1, A_1, R_2, S_2 \dots$
- S_0 is sampled from the distribution $p_0(\cdot)$
- A_t is sampled from $\pi(S_t, \cdot; \theta)$
- Receive atomic experience (R_{t+1}, S_{t+1}) from the environment
- At each time step t , update \mathbf{w} proportional to gradient of appropriate (MC or TD-based) loss function of $Q(s, a; \mathbf{w})$
- Sum $\gamma^t \cdot (\nabla_{\theta} \log \pi(S_t, A_t; \theta)) \cdot Q(S_t, A_t; \mathbf{w})$ over t and over paths
- Update θ using this (biased) estimate of $\nabla_{\theta} J(\theta)$
- Iterate with a new set of sampling traces ...

Reducing Variance with a Baseline

- We can reduce variance by subtracting a baseline function $B(s)$ from $Q(s, a; \mathbf{w})$ in the Policy Gradient estimate
- This means at each time step, we replace $\gamma^t \cdot \nabla_{\theta} \log \pi(S_t, A_t; \theta) \cdot Q(S_t, A_t; \mathbf{w})$ with $\gamma^t \cdot \nabla_{\theta} \log \pi(S_t, A_t; \theta) \cdot (Q(S_t, A_t; \mathbf{w}) - B(s))$
- Note that Baseline function $B(s)$ is only a function of s (and not a)
- This ensures that subtracting Baseline $B(s)$ does not add bias

$$\begin{aligned} & \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot B(s) \\ &= \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot B(s) \cdot \nabla_{\theta} \left(\sum_{a \in \mathcal{A}} \pi(s, a; \theta) \right) \\ &= \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot B(s) \cdot \nabla_{\theta} 1 \\ &= 0 \end{aligned}$$

Using State Value function as Baseline

- A good baseline $B(s)$ is state value function $V(s; \mathbf{v})$
- Rewrite Policy Gradient algorithm using advantage function estimate

$$A(s, a; \mathbf{w}, \mathbf{v}) = Q(s, a; \mathbf{w}) - V(s; \mathbf{v})$$

- Now the estimate of $\nabla_{\theta} J(\theta)$ is given by:

$$\sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot A(s, a; \mathbf{w}, \mathbf{v})$$

- At each time step, we update both sets of parameters \mathbf{w} and \mathbf{v}

TD Error as estimate of Advantage Function

- Consider TD error δ^π for the *true* Value Function $V^\pi(s)$

$$\delta^\pi = r + \gamma \cdot V^\pi(s') - V^\pi(s)$$

- δ^π is an unbiased estimate of Advantage function $A^\pi(s, a)$

$$\mathbb{E}_\pi[\delta^\pi | s, a] = \mathbb{E}_\pi[r + \gamma \cdot V^\pi(s') | s, a] - V^\pi(s) = Q^\pi(s, a) - V^\pi(s) = A^\pi(s, a)$$

- So we can write Policy Gradient in terms of $\mathbb{E}_\pi[\delta^\pi | s, a]$

$$\nabla_{\theta} J(\theta) = \sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot \mathbb{E}_\pi[\delta^\pi | s, a]$$

- In practice, we can use func approx for TD error (and sample):

$$\delta(s, r, s'; \mathbf{v}) = r + \gamma \cdot V(s'; \mathbf{v}) - V(s; \mathbf{v})$$

- This approach requires only one set of critic parameters \mathbf{v}

TD Error can be used by both Actor and Critic

Algorithm 4.2: ACTOR-CRITIC-TD-ERROR(\cdot)

Initialize Policy params θ and State VF params \mathbf{v} arbitrarily
for each episode

do {
 Initialize s (first state of episode)
 $P \leftarrow 1$
 while s is not terminal
 {
 $a \sim \pi(s, \cdot; \theta)$
 Take action a , receive r, s' from the environment
 $\delta \leftarrow r + \gamma \cdot V(s'; \mathbf{v}) - V(s; \mathbf{v})$
 do {
 $\mathbf{v} \leftarrow \mathbf{v} + \alpha_{\mathbf{v}} \cdot \delta \cdot \nabla_{\mathbf{v}} V(s; \mathbf{v})$
 $\theta \leftarrow \theta + \alpha_{\theta} \cdot P \cdot \delta \cdot \nabla_{\theta} \log \pi(s, a; \theta)$
 $P \leftarrow \gamma \cdot P$
 $s \leftarrow s'$

Using Eligibility Traces for both Actor and Critic

Algorithm 4.3: ACTOR-CRITIC-ELIGIBILITY-TRACES(\cdot)

Initialize Policy params θ and State VF params \mathbf{v} arbitrarily

for each episode

do {
 Initialize s (first state of episode)
 $\mathbf{z}_\theta, \mathbf{z}_\mathbf{v} \leftarrow 0$ (eligibility traces for θ and \mathbf{v})
 $P \leftarrow 1$
 while s is not terminal
 {
 $a \sim \pi(s, \cdot; \theta)$
 Take action a , observe r, s'
 $\delta \leftarrow r + \gamma \cdot V(s'; \mathbf{v}) - V(s; \mathbf{v})$
 do {
 $\mathbf{z}_\mathbf{v} \leftarrow \gamma \cdot \lambda_\mathbf{v} \cdot \mathbf{z}_\mathbf{v} + \nabla_\mathbf{v} V(s; \mathbf{v})$
 $\mathbf{z}_\theta \leftarrow \gamma \cdot \lambda_\theta \cdot \mathbf{z}_\theta + P \cdot \nabla_\theta \log \pi(s, a; \theta)$
 $\mathbf{v} \leftarrow \mathbf{v} + \alpha_\mathbf{v} \cdot \delta \cdot \mathbf{z}_\mathbf{v}$
 $\theta \leftarrow \theta + \alpha_\theta \cdot \delta \cdot \mathbf{z}_\theta$
 $P \leftarrow \gamma \cdot P, s \leftarrow s'$
 }

Overcoming Bias

- We've learnt a few ways of how to reduce variance
- But we haven't discussed how to overcome bias
- All of the following substitutes for $Q^\pi(s, a)$ in PG have bias:
 - $Q(s, a; \mathbf{w})$
 - $A(s, a; \mathbf{w}, \mathbf{v})$
 - $\delta(s, s', r; \mathbf{v})$
- Turns out there is indeed a way to overcome bias
- It is called the *Compatible Function Approximation Theorem*

Compatible Function Approximation Theorem

Theorem

If the following two conditions are satisfied:

- 1 *Critic gradient is compatible with the Actor score function*

$$\nabla_{\mathbf{w}} Q(s, a; \mathbf{w}) = \nabla_{\theta} \log \pi(s, a; \theta)$$

- 2 *Critic parameters \mathbf{w} minimize the following mean-squared error:*

$$\epsilon = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot (Q^{\pi}(s, a) - Q(s, a; \mathbf{w}))^2$$

Then the Policy Gradient using critic $Q(s, a; \mathbf{w})$ is exact:

$$\nabla_{\theta} J(\theta) = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q(s, a; \mathbf{w})$$

Proof of Compatible Function Approximation Theorem

For \mathbf{w} that minimizes

$$\epsilon = \sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \boldsymbol{\theta}) \cdot (Q^\pi(s, a) - Q(s, a; \mathbf{w}))^2,$$

$$\sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \boldsymbol{\theta}) \cdot (Q^\pi(s, a) - Q(s, a; \mathbf{w})) \cdot \nabla_{\mathbf{w}} Q(s, a; \mathbf{w}) = 0$$

But since $\nabla_{\mathbf{w}} Q(s, a; \mathbf{w}) = \nabla_{\boldsymbol{\theta}} \log \pi(s, a; \boldsymbol{\theta})$, we have:

$$\sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \boldsymbol{\theta}) \cdot (Q^\pi(s, a) - Q(s, a; \mathbf{w})) \cdot \nabla_{\boldsymbol{\theta}} \log \pi(s, a; \boldsymbol{\theta}) = 0$$

$$\begin{aligned} \text{Therefore, } & \sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \boldsymbol{\theta}) \cdot Q^\pi(s, a) \cdot \nabla_{\boldsymbol{\theta}} \log \pi(s, a; \boldsymbol{\theta}) \\ &= \sum_{s \in \mathcal{N}} \rho^\pi(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \boldsymbol{\theta}) \cdot Q(s, a; \mathbf{w}) \cdot \nabla_{\boldsymbol{\theta}} \log \pi(s, a; \boldsymbol{\theta}) \end{aligned}$$

Proof of Compatible Function Approximation Theorem

$$\text{But } \nabla_{\theta} J(\theta) = \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot Q^{\pi}(s, a) \cdot \nabla_{\theta} \log \pi(s, a; \theta)$$

$$\begin{aligned} \text{So, } \nabla_{\theta} J(\theta) &= \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot Q(s, a; \mathbf{w}) \cdot \nabla_{\theta} \log \pi(s, a; \theta) \\ &= \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \nabla_{\theta} \pi(s, a; \theta) \cdot Q(s, a; \mathbf{w}) \end{aligned}$$

Q.E.D.

This means with conditions (1) and (2) of Compatible Function Approximation Theorem, we can use the critic func approx $Q(s, a; \mathbf{w})$ and still have the exact Policy Gradient.

How to enable Compatible Function Approximation

A simple way to enable Compatible Function Approximation

$\frac{\partial Q(s, a; \mathbf{w})}{\partial w_i} = \frac{\partial \log \pi(s, a; \theta)}{\partial \theta_i}, \forall i$ is to set $Q(s, a; \mathbf{w})$ to be linear in its features.

$$Q(s, a; \mathbf{w}) = \sum_{i=1}^m \phi_i(s, a) \cdot w_i = \sum_{i=1}^m \frac{\partial \log \pi(s, a; \theta)}{\partial \theta_i} \cdot w_i$$

We note below that a compatible $Q(s, a; \mathbf{w})$ serves as an approximation of the advantage function.

$$\begin{aligned} \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot Q(s, a; \mathbf{w}) &= \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot \left(\sum_{i=1}^m \frac{\partial \log \pi(s, a; \theta)}{\partial \theta_i} \cdot w_i \right) \\ &= \sum_{a \in \mathcal{A}} \left(\sum_{i=1}^m \frac{\partial \pi(s, a; \theta)}{\partial \theta_i} \cdot w_i \right) = \sum_{i=1}^m \left(\sum_{a \in \mathcal{A}} \frac{\partial \pi(s, a; \theta)}{\partial \theta_i} \right) \cdot w_i \\ &= \sum_{i=1}^m \frac{\partial}{\partial \theta_i} \left(\sum_{a \in \mathcal{A}} \pi(s, a; \theta) \right) \cdot w_i = \sum_{i=1}^m \frac{\partial 1}{\partial \theta_i} \cdot w_i = 0 \end{aligned}$$

Fisher Information Matrix

Denoting $[\frac{\partial \log \pi(s, a; \theta)}{\partial \theta_i}]$, $i = 1, \dots, m$ as the score column vector $\mathbf{SC}(s, a; \theta)$ and assuming compatible linear-approximation critic:

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \sum_{s \in \mathcal{N}} \rho^{\pi}(s) \cdot \sum_{a \in \mathcal{A}} \pi(s, a; \theta) \cdot (\mathbf{SC}(s, a; \theta) \cdot \mathbf{SC}(s, a; \theta)^T \cdot \mathbf{w}) \\ &= \mathbb{E}_{s \sim \rho^{\pi}, a \sim \pi} [\mathbf{SC}(s, a; \theta) \cdot \mathbf{SC}(s, a; \theta)^T] \cdot \mathbf{w} \\ &= \mathbf{FIM}_{\rho^{\pi}, \pi}(\theta) \cdot \mathbf{w}\end{aligned}$$

where $\mathbf{FIM}_{\rho^{\pi}, \pi}(\theta)$ is the Fisher Information Matrix w.r.t. $s \sim \rho^{\pi}, a \sim \pi$.

Natural Policy Gradient

- Recall the idea of Natural Gradient from Numerical Optimization
- Natural gradient $\nabla_{\theta}^{nat} J(\theta)$ is the direction of optimal θ movement
- In terms of the KL-divergence metric (versus plain Euclidean norm)
- Natural gradient yields better convergence (we won't cover proof)

Formally defined as: $\nabla_{\theta} J(\theta) = FIM_{\rho_{\pi}, \pi}(\theta) \cdot \nabla_{\theta}^{nat} J(\theta)$

Therefore, $\nabla_{\theta}^{nat} J(\theta) = \mathbf{w}$

This compact result is great for our algorithm:

- Update Critic params \mathbf{w} with the critic loss gradient (at step t) as:

$$\gamma^t \cdot (R_{t+1} + \gamma \cdot \mathbf{SC}(S_{t+1}, A_{t+1}; \theta)^T \cdot \mathbf{w} - \mathbf{SC}(S_t, A_t; \theta)^T \cdot \mathbf{w}) \cdot \mathbf{SC}(S_t, A_t; \theta)$$

- Update Actor params θ in the direction equal to value of \mathbf{w}