

# Regulator based method to find adiabatic gauge potential for quantum many body systems

Mohit

July 14, 2017

## 1 Goals: what we hope to achieve

Adiabatic gauge potentials are useful for controlling a quantum system when it's driven externally from one configuration to another. These potentials help us in circumventing standard adiabatic limitations which requires infinitesimally small rates [1, 2, 3]. For example, these potentials can be used for arbitrarily fast annealing protocols and implementing fast dissipationless driving.

The goal is to develop a regulator based method to find adiabatic gauge potential for quantum many body systems. If we are successful, then this will be a new method to find these potentials and it will give new insights in quantum control of many body systems.

We will use our method on both quantum integrable and non-integrable systems. For quantum integrable many body systems, exact gauge potential is already known in literature [4, 5]. We hope to derive these results using our new method. For non-integrable systems, exact gauge potentials are very difficult to find. We hope that our method will find an approximate gauge potentials for such systems providing an alternative method to variational approximation scheme recently introduced in [4].

We also hope to use this method to distinguish between quantum integrable and non-integrable systems. Our idea is to use Eigenstate Thermalization Hypothesis (ETH)[6] for this. Since ETH is valid for local operators in non-integrable systems, we expect local approximate gauge potentials to satisfy ETH, and exact gauge potentials (which are non-local) should not satisfy ETH. We can show that using ETH, norm of approximate gauge potential should scale exponentially in system size for non-integrable systems. Whereas for integrable systems (where ETH is not valid), exact gauge potential are supposed to scale like a polynomial in system size. We want to understand this issue into more details using our new method.

## 2 Introduction

### 2.1 Gauge potential

Let's represent a wavefunction in some basis as  $|\psi\rangle = \sum_n \psi_n |n\rangle_0$  where  $|n\rangle_0$  is some fixed, parameter independent basis. Now let's do a unitary basis transformation to  $|m(\lambda)\rangle$  in the parameter  $\lambda$  dependent space using  $U(\lambda)$  by defining  $|m(\lambda)\rangle = \sum_n U_{mn} |n\rangle$ . Hence, now we can express  $|\psi\rangle = \sum_m \tilde{\psi}_m |m(\lambda)\rangle$ , where  $\tilde{\psi}_m = \langle m(\lambda) | \psi \rangle$ .

Quantum gauge potentials  $A_\lambda$  are defined to be generators of continuous unitary transformation. In the lab frame,  $A_\lambda$  is defined as:

$$\boxed{A_\lambda = i\hbar \partial_\lambda} \tag{1}$$

In rotated frame ( $\lambda$ -dependent basis),  $\tilde{A}_\lambda$  is defined as follows:

$$\boxed{\tilde{A}_\lambda = i\hbar U^\dagger \partial_\lambda U} \tag{2}$$

We can show that gauge potentials in these two frames are related by  $A_\lambda = U\tilde{A}_\lambda U^\dagger$ .  
Let's take an example of a shifting transformation  $U$  to understand gauge potentials:

$$U|x'(\lambda)\rangle = |x + \lambda\rangle \quad (3)$$

We know that unitary transformation  $U = \exp(-i\hat{p}\lambda/\hbar)$ . Now,  $\tilde{A}_\lambda = \hat{p}$  and  $A_\lambda = i\hbar\partial_\lambda$ <sup>1</sup>.

Now why do we call it a gauge potential? In [5], they call it gauge potential because there is freedom to choose  $A_\lambda$  like how in EM, we have gauge choice. In [5], they say that “one can show that the gauge potentials for canonical shifts of the momentum appear exactly as the electromagnetic vector potential [see Exercise (III.1)]. Gauge potentials generalize these ideas from electromagnetism to arbitrary parameters”

Here I am listing down some properties:

- They are Hermitian operator.
- $\langle n(\lambda)|A_\lambda|m(\lambda)\rangle = {}_0\langle n|\tilde{A}_\lambda|m\rangle_0$

## 2.2 Adiabatic gauge potential

The gauge potentials become adiabatic gauge potential when unitary transformation generated by  $A_\lambda$  are used to diagonalize Hamiltonian.

Adiabatic gauge potentials are a special subset of these which diagonalize the instantaneous Hamiltonian, attempting to leave its eigenbasis invariant as the parameter is changed. These adiabatic gauge potentials generate non-adiabatic corrections to Hamiltonian in the moving basis ( $\lambda$ -dependent basis).

This is something from Anatoli's lecture notes [5]– “an adiabatic basis is a family of adiabatically connected eigenstates, i.e., eigenstates related to a particular initial basis by adiabatic (infinitesimally slow) evolution of the parameter  $\lambda$ . For example, if two levels cross they will exchange order energetically but the adiabatic connection will be non-singular.”

$H(\lambda)|n(\lambda)\rangle = E_n(\lambda)|n(\lambda)\rangle$ . Let's derive diagonal and off-diagonal elements.

- **n-th diagonal element:**  $A_\lambda^n = \langle n|A_\lambda|n\rangle = i\hbar\langle n|\partial_\lambda|n\rangle$
- **off-diagonal element:** We use the identity  $\langle m|H(\lambda)|n\rangle = 0$ ,  $n \neq m$  and then differentiate with respect to  $\lambda$  to obtain:

$$\boxed{\langle m|A_\lambda|n\rangle = -i\hbar \frac{\langle m|\partial_\lambda H|n\rangle}{E_m - E_n}} \quad (4)$$

where both energies ( $E_m, E_n$ ) and eigenvectors ( $|m\rangle, |n\rangle$ ) depend on  $\lambda$ .

This information can be represented in matrix form [7] as follows:

$$i\hbar\partial_\lambda H = [A_\lambda, H] - i\hbar M_\lambda \quad (5)$$

where

$$M_\lambda = - \sum_n \frac{\partial E_n(\lambda)}{\partial \lambda} |n(\lambda)\rangle \langle n(\lambda)| \quad (6)$$

It's to be noted that for finding  $M_\lambda$ , we need to diagonalize Hamiltonian. We can eliminate  $M_\lambda$  by taking commutator on both sides of equation 5 and obtain:

$$[H, i\hbar\partial_\lambda H - [A_\lambda, H]] = 0 \quad (7)$$

Any  $A_\lambda$  satisfying equation 7 is an exact gauge potential. We note that if  $A_\lambda$  satisfies equation 7, then  $A_\lambda + f(H)$ , where  $f(H)$  is any function that only contains terms involving Hamiltonian  $H$  and other operators that commutes with  $H$ . We note that  $f(H)$  is diagonal in energy basis, i.e.  $f(H)^{nm} = \delta_{n,m}f(H)^{nn}$ .

---

<sup>1</sup>I don't know why there is a minus sign missing as momentum operator is defined as  $\hat{p} = -i\hbar\partial_x$

### 2.2.1 Minimum norm gauge choice

For our future purposes, let's study a gauge choice where we assume diagonal elements of  $A_\lambda$  in energy basis is zero, i. e.  $\langle n|A_\lambda|n\rangle = A_\lambda^{n,n} = 0$ , for  $n = 1, 2 \dots D$ , where  $D$  is the dimension of Hilbert space. Can we always make such a gauge choice?

As we noted above, a family of  $A_\lambda$  satisfies equation 7 – both  $A_\lambda$  and  $A_\lambda + f(H)$  satisfy the equation 7. Using this knowledge, let's suppose we define  $A'_\lambda = A_\lambda + f(H)$ , where in energy basis,  $f(H)$  is diagonal (as we already know),  $A'_\lambda$  is an exact gauge potential with all it's diagonal elements chosen to be zero and  $A_\lambda$  is an exact gauge potential with non-zero diagonal elements. What's the condition on  $f(H)$  so that diagonal elements of  $A'_\lambda$  are zero? The required condition is:

$$f(H)^{nn} = -A_\lambda^{nn}, \quad n = 1, 2, \dots D \quad (8)$$

where  $D$  is the dimension of Hilbert space. Hence, if somebody hands me  $A_\lambda$ , here is the method to obtain  $A'_\lambda$ : I can always cook up a function  $f(H) = \sum_{n=1}^D a_n H^n$  by solving  $D$  number of equations which satisfy equation 8 to find out  $a_n$ . Once, I know  $f(H)$ , I can always subtract it from  $A_\lambda$  to obtain  $A'_\lambda$ . Hereby, I show that this gauge choice can always be made without any loss of generality.

Let's try to understand this gauge choice further by computing the Frobenius norm of  $A_\lambda$ .

$$\|A_\lambda\|^2 = \text{Tr}(A_\lambda^2) = \sum_{n,m} |A_\lambda^{n,m}|^2 = \sum_n |A_\lambda^{n,n}|^2 + \sum_{n \neq m} |A_\lambda^{n,m}|^2 = \sum_n |A_\lambda^{n,n}|^2 + \|A'_\lambda\|^2 \quad (9)$$

where  $A_\lambda^{n,m} = \langle n|A_\lambda|m\rangle$  and  $|m\rangle$  is the energy eigenstate with energy  $E_m$ . Thus, we see that in our gauge choice all the diagonal elements ( $A_\lambda^{n,n}$ ) are zero, and therefore, this choice reduces the norm. Is this the minimum norm of  $A_\lambda$  which satisfies equation 7? The answer is yes as explained below.

Let's compute the norm of  $A_\lambda$  another way by exploiting its' equality to  $A'_\lambda - f(H)$ :

$$\|A_\lambda\|^2 = \text{Tr}((A'_\lambda - f(H))^2) \quad (10)$$

$$= \text{Tr}(A_\lambda'^2) + \text{Tr}(f(H)^2) - 2 \text{Tr}(A'_\lambda f(H)) \quad (11)$$

$$= \text{Tr}(A_\lambda'^2) + \text{Tr}(f(H)^2) \quad (12)$$

$$= \|A'_\lambda\|^2 + \|f(H)\|^2 \quad (13)$$

where we have used the fact that  $f(H)$  is diagonal and  $A'_\lambda$  has no non-zero diagonal elements in energy basis in claiming  $\text{Tr}(A'_\lambda f(H)) = 0$ . We note that since  $f(H)$  is diagonal in energy basis, the only way  $A_\lambda$  will acquire diagonal elements is through  $f(H)$ . Hence, we see that by choosing diagonal elements of an exact adiabatic gauge potential in energy basis to be zero, we are effectively choosing a gauge potential which has no  $f(H)$  term. Hence, this gauge choice is the minimum norm possible.

### 2.2.2 Time evolution in moving frame

Our Hamiltonian would be controlled using a control parameter called  $\lambda$  and our aim is to find time evolution of wave-function is  $\lambda$ -dependent basis called moving frame.

Our Hamiltonian  $H_0(\lambda(t))$  would satisfy the following equation:

$$H_0(\lambda(t))|\psi\rangle = i\partial_t|\psi\rangle \quad (14)$$

Let us go to rotating frame so as to diagonalize our Hamiltonian. Required unitary transformation  $U(\lambda)$  would depend on parameter  $\lambda$ . Wave function in moving frame is  $|\tilde{\psi}\rangle = U^\dagger|\psi\rangle$ . In this basis, Hamiltonian is diagonal:  $\tilde{H}_0 = U^\dagger H_0 U = \sum_n \epsilon(\lambda) |n(\lambda)\rangle \langle n(\lambda)|$ .<sup>2</sup>

How does the wave function evolve in new basis?

$$i\partial_t|\tilde{\psi}\rangle = (\tilde{H}_0(\lambda(t)) - \dot{\lambda}\tilde{A}_\lambda)|\tilde{\psi}\rangle \quad (15)$$

Note that gauge potential should be purely imaginary in a basis in which Hamiltonian is real.

---

<sup>2</sup>Note that expectation value should remain same in both basis, i.e.  $\langle\tilde{\psi}|\tilde{H}_0|\tilde{\psi}\rangle = \langle\psi|H_0|\psi\rangle$

### 3 Regulator based method to find Gauge Potential

Here we would introduce a new method to find Gauge Potential  $A_\lambda$  which includes a regulator  $\mu$ .

Let's start off by writing the off-diagonal elements of exact gauge potential:

$$\langle m|A_\lambda|n\rangle = -i\hbar \frac{\langle m|\partial_\lambda H|n\rangle}{E_m - E_n} \quad (16)$$

For a many-body Hamiltonian, number of states in Hilbert space grows exponentially in system size while energy bandwidth grows linearly with system size (since energy is an extensive quantity). Thus, distance between any two *nearby* eigenvalues is exponentially small in system size. In other words,  $E_m - E_n \sim e^{-S}$ . If there are non-zero off-diagonal elements of  $\partial_\lambda H$ , then  $\langle m|A_\lambda|n\rangle$  is ill-defined. It's called small denominator problem [5].

To resolve this problem, we introduce a regulator/ cutoff  $\mu$  that regularizes our exact gauge potential in large system size  $L$  limit. Once we have taken large  $L$  limit, then we take small  $\mu$  limit. Hence, if this method works, the right way to take limits will be:

$$\langle n|A_\lambda|m\rangle = \lim_{\mu \rightarrow 0} \lim_{L \rightarrow \infty} -i\hbar \frac{\langle n|\partial_\lambda H|m\rangle}{(E_n - E_m)^2 + \mu^2} (E_n - E_m) \quad (17)$$

where we have chosen a gauge choice in which diagonal elements are zero in energy basis, i.e.  $A_\lambda^{nn} = 0$ .

Now let's first recall here the Laplace transform of  $\sin(\omega t)$ :

$$\int_0^\infty dt e^{-st} \sin(\omega t) = \frac{\omega}{\omega^2 + s^2} \quad (18)$$

Thus, we see that with  $s = \mu$ , we can convert our expression of  $A_\lambda$  into an integral. From here on to simplify our expression of integration, we will choose our units such that  $\hbar = 1$ .<sup>3</sup>

$$\langle n|A_\lambda|m\rangle = -i \frac{\langle n|\partial_\lambda H|m\rangle}{(E_n - E_m)^2 + \mu^2} (E_n - E_m) \quad (19)$$

$$= -i \int_0^\infty dt e^{-\mu t} \langle n|\partial_\lambda H|m\rangle \sin((E_n - E_m)t) \quad (20)$$

$$= -\frac{1}{2} \int_0^\infty dt e^{-\mu t} (\langle n|e^{iE_n t} \partial_\lambda H e^{-iE_m t}|m\rangle - \langle n|e^{-iE_n t} \partial_\lambda H e^{iE_m t}|m\rangle) \quad (21)$$

Hence, we can simplify our expression by defining propagator  $U = \exp(-iHt)$ . We note that parameter  $\lambda$  is fixed while we evolve it in the *artificial time*  $t$ .

$$A_\lambda = -\frac{1}{2} \int_0^\infty dt e^{-\mu t} [U^\dagger(t) \partial_\lambda H U(t) - U^\dagger(-t) \partial_\lambda H U(-t)] \quad (22)$$

$$= -\frac{1}{2} \int_0^\infty dt e^{-\mu t} [\partial_\lambda H(t) - \partial_\lambda H(-t)] \quad (23)$$

where  $\partial_\lambda H(t)$  is time-evolved operator  $\partial_\lambda H$  in Heisenberg picture. We note that characteristic time scale in the integration is  $1/\mu$ .

Now using Hadamard (or sometimes called Baker-Campbell-Hausdorff) formula, we find a simplified expression. Detailed calculations are given in appendix B .

$$A_\lambda = -i \int_0^\infty dt e^{-\mu t} \sin([H, \partial_\lambda H]t) = -i\hbar \int_0^\infty dt e^{-\mu t} \sum_{n=0}^\infty \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \quad (24)$$

<sup>3</sup>If we don't choose  $\hbar = 1$ , then for Laplace integral variable, we should use not use  $t$ . Instead we should use  $t/\hbar$  so that we have  $\sin((E_n - E_m)t/\hbar)$  and  $e^{-\mu t/\hbar}$ . Otherwise, we will end up with terms  $\sin((E_n - E_m)t)$  and  $e^{-\mu t}$ , which are dimensionally incorrect.

where  $C^{(n)}$  is  $n$ -commutator of  $H$  and  $\partial_\lambda H$ , i.e.  $C^{(n)} = [H, [H, \text{n times } \dots, [H, \partial_\lambda H]]]$ . We define the first term as  $C^{(1)} = [H, \partial_\lambda H]$ , second term as  $C^{(2)} = [H, [H, \partial_\lambda H]] = [H, C^{(1)}]$  and so on and forth. Properties of  $C^{(n)}$  are noted in appendix A.

Can we further simplify the expression? If we are allowed to change the order of summation and integration <sup>4</sup>, then we can do first Laplace transform of  $t^{2n+1}$  terms and then later the sum. Hence, we get:

$$A_\lambda = -i \int_0^\infty dt e^{-\mu t} \sum_{n=0}^\infty \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \quad (25)$$

$$= -i \sum_{n=0}^\infty (-1)^n C^{(2n+1)} \int_0^\infty dt e^{-\mu t} \frac{t^{2n+1}}{(2n+1)!} \quad (26)$$

$$= -i \sum_{n=0}^\infty (-1)^n \frac{C^{(2n+1)}}{\mu^{2n+2}} \quad (27)$$

Hence, we get another expression where we have integrated before taking the summation:

$$A_\lambda = \frac{-i\hbar}{\mu} \sum_{n=0}^\infty (-1)^n \frac{C^{(2n+1)}}{\mu^{2n+1}} \quad (28)$$

Once we are done with the integration, we can bring back  $\hbar$  factor. We note that  $iC^{(2n+1)}$  is Hermitian, which is consistent with the fact that  $A_\lambda$  is Hermitian.

Now let's think about  $\lim_{L \rightarrow \infty}$  limit which we need to take. I would claim that while doing the infinite summation we have already taken that limit as we have assumed infinite system size. Why is that? In general,  $C^{(n)}$  grows with  $n$  in the sense that it would have operators with larger support over lattice sites as  $n$  increases. At a certain  $n_L$  that is proportional to system length  $L$ , we would find that  $C^{n_L}$  has operators with support on boundary lattice sites. This is where our summation would be truncated for a finite system. Hence, the correct order of limits should be:

$$A_\lambda = \lim_{\mu \rightarrow 0} \lim_{L \rightarrow \infty} \frac{-i\hbar}{\mu} \sum_{n=0}^{n_L} (-1)^n \frac{C^{(2n+1)}}{\mu^{2n+1}} \quad (29)$$

Now one thing which is good is that if we take the wrong order of limit: take  $\lim_{\mu \rightarrow 0}$  before  $\lim_{L \rightarrow \infty}$ , then  $A_\lambda$  diverges. Thus, now divergence is more explicit than the original expression 17.

How does  $\mu$  scale as  $L$ ? It's an important question whose answer we don't know. Allow me to make a heuristic argument: In general, it seems that operators involved in the expression of  $C^{(n)}$  would have support which depend on  $L$ . Let's suppose the support of these operators grow as  $L^\gamma$ , i.e.,  $C^{(n)} \propto L^\gamma$ , where  $\gamma$  is some constant which we don't know. If we assume that  $A_\lambda$  is well-defined in large system size limit for many-body Hamiltonian (both integrable and non-integrable), then  $\mu \propto L^\gamma$ .

Now our task will be use it to find exact/approximate gauge potential for integrable and non-integrable models.

### 3.1 Integrable model

Our goal is to study a integrable model, which is called **Transverse Field Ising model**. It shows quantum phase transition between ferromagnetic and paramagnetic phases. Moreover, it satisfies Ising symmetry  $G = \Pi_i \sigma_i^z$  since  $[H, G] = 0$ , where  $H$  is the Hamiltonian. This model can be

---

<sup>4</sup>If there is some singularity, then the order of summation and integration might be important and we might get two different results.

written in terms of non-interacting spinless fermions  $(c_i, c_i^\dagger)$  using Jordan- Wigner transformation. It's Hamiltonian in spin basis is given by:

$$H = -J \sum_j \sigma_j^x \sigma_{j+1}^x - \lambda \sum_j \sigma_j^z \quad (30)$$

where we have not specified boundary conditions and  $\lambda$  is externally-controlled transverse magnetic field.

This model's exact gauge potential is already known in literature [4, 5] and it's given by:

$$A_\lambda = \sum_l \alpha_l O_l \quad \text{where} \quad \alpha_l = -\frac{1}{4L} \sum_k \frac{\sin(k) \cos(lk)}{(\cos k - h)^2 + \sin^2 k} \quad (31)$$

and where  $O_l$  is given by

$$O_l = 2i \sum_j (c_j^\dagger c_{j+l}^\dagger - \text{h.c.}) = \sum_j (\sigma_j^x \sigma_{j+1}^z \dots \sigma_{j+l-1}^z \sigma_{j+l}^y + \sigma_j^y \sigma_{j+1}^z \dots \sigma_{j+l-1}^z \sigma_{j+l}^x) \quad (32)$$

Let's write a first few terms of  $O_l$  here:

$$\begin{aligned} O_{l=1} &= \sum_j (\sigma_j^x \sigma_{j+1}^y + \sigma_j^y \sigma_{j+1}^x) \\ O_{l=2} &= \sum_j (\sigma_j^x \sigma_{j+1}^z \sigma_{j+2}^y + \sigma_j^y \sigma_{j+1}^z \sigma_{j+2}^x) \end{aligned}$$

It will be good to find either exact or approximate gauge potential using our regulator method. However, before we study this model, we will study much simpler models to learn about this new method.

### 3.1.1 Ising model with local transverse magnetic field

We would take the simplest integrable Hamiltonian with Ising interaction and a local  $x$  magnetic field:

$$H = J \sum_j \sigma_j^z \sigma_{j+1}^z + \lambda \sigma_0^x \quad (33)$$

where boundary conditions are not important. Commutation relation followed by spin operators are:

$$[\sigma_i^a, \sigma_j^b] = 2i\delta_{i,j} \sum_c \epsilon_{abc} \sigma_i^c \quad (34)$$

where  $\epsilon_{abc}$  is the Levi-Civita symbol,  $\delta_{ij}$  is the Kronecker delta.

This model satisfies Ising symmetry  $G = \Pi_i \sigma_i^x$  since  $[H, G] = 0$ .

Let's find out  $A_\lambda$  for this Hamiltonian for which we need to compute different odd-powered commutator  $[H, \partial_\lambda H]$ , where  $\partial_\lambda H = \sigma_0^x$ . Here we begin:

$$C^{(1)} = 2iJ\sigma_0^y(\sigma_{-1}^z + \sigma_1^z) \quad (35)$$

$$C^{(2)} = 8J^2(\sigma_1^z \sigma_0^x \sigma_{-1}^z + \sigma_0^x) - 4J\lambda\sigma_0^z(\sigma_{-1}^z + \sigma_1^z) \quad (36)$$

$$C^{(3)} = (16J^2 + 4\lambda^2)[H, \partial_\lambda H] = \alpha^2 C^{(1)} \quad (37)$$

$$C^{(5)} = [H, [H, C^{(3)}]] = \alpha^2 [H, [H, C^{(1)}]] = \alpha^2 C^{(3)} = \alpha^4 C^{(1)} \quad (38)$$

Hence,  $C^{(2n+1)} = \alpha^{2n} C^{(1)}$ , where  $\alpha^2 = 4(4J^2 + \lambda^2)$ . Now, we would compute  $A_\lambda$  using two methods and compare our results. Using equation (24), we get :

$$\begin{aligned}
A_\lambda &= -i\hbar C^{(1)} \int_0^\infty dt e^{-\mu t} \sum_{n=0}^\infty \frac{(-1)^n t^{2n+1}}{(2n+1)!} \alpha^{2n} \\
&= -i\hbar C^{(1)} \int_0^\infty dt e^{-\mu t} \sum_{n=0}^\infty \frac{(-1)^n \alpha^{2n+1} t^{2n+1}}{\alpha(2n+1)!} \\
&= \frac{-i\hbar C^{(1)}}{\alpha} \int_0^\infty dt e^{-\mu t} \sum_{n=0}^\infty \frac{(-1)^n (\alpha t)^{2n+1}}{(2n+1)!} \\
&= \frac{-i\hbar C^{(1)}}{\alpha} \int_0^\infty dt e^{-\mu t} \sin(\alpha t) \\
&= \frac{-i\hbar C^{(1)}}{\alpha} \frac{\alpha}{\alpha^2 + \mu^2} = \frac{-i\hbar C^{(1)}}{\alpha^2 + \mu^2} = \frac{2\hbar J}{\alpha^2 + \mu^2} \sigma_0^y (\sigma_{-1}^z + \sigma_1^z)
\end{aligned}$$

Using 28, we get:

$$\begin{aligned}
A_\lambda &= \frac{-i\hbar C^{(1)}}{\mu} \sum_{n=0}^\infty (-1)^n \frac{\alpha^{2n}}{\mu^{2n+1}} \\
&= \frac{-i\hbar}{\mu\alpha} C^{(1)} \sum_{n=0}^\infty (-1)^n \left(\frac{\alpha}{\mu}\right)^{2n+1} \\
&= \frac{-i\hbar}{\mu\alpha} C^{(1)} \frac{\mu\alpha}{\mu^2 + \alpha^2} \quad , \text{ if } \alpha^2/\mu^2 < 1 \\
&= C^{(1)} \frac{-i\hbar}{\mu^2 + \alpha^2} = \frac{2\hbar J}{\alpha^2 + \mu^2} \sigma_0^y (\sigma_{-1}^z + \sigma_1^z) \quad , \text{ if } \alpha^2/\mu^2 < 1
\end{aligned}$$

Hence, now we can use analytical continuation to claim that our result is also true when  $\alpha^2/\mu^2 > 1$  since there is no divergence when  $\alpha^2/\mu^2 = 1$ . Hence, both methods give the same answer as they should.

After taking  $\mu \rightarrow 0$  limit, we get an expression for exact gauge potential:

$$\boxed{A_\lambda = \frac{\hbar J}{8J^2 + 2\lambda^2} \sigma_0^y (\sigma_{-1}^z + \sigma_1^z)} \quad (39)$$

This expression is correct because it satisfies equation 7. And it's unique upto any term that commutes with Hamiltonian.

Why  $A_\lambda$  is non-zero in  $\lambda \rightarrow 0$  limit? It need not be zero because additional term in Hamiltonian is  $\lambda A_\lambda$ , which goes to zero in  $\lambda \rightarrow 0$  limit.

I can similarly write an exact expression for additional  $\sum_{j=1}^L h_j \sigma_j^z$  term in the Hamiltonian, although this term breaks Ising symmetry  $G$ .

Now for future purposes, let's study the following Hamiltonian:

$$H = -J \sum_j \sigma_j^x \sigma_{j+1}^x - \lambda \sigma_0^z \quad (40)$$

Apart from change in sign of  $J$  and  $\lambda$ , we have rotated our axes through  $y$  axis such that it interchanges  $z$  and  $x$  axes. In other words, we have done following transformation on spins ignoring the sign changes of couplings  $J$  and  $\lambda$ :

$$\sigma_j^x \rightarrow \sigma_j^z, \sigma_j^z \rightarrow \sigma_j^x, \sigma_j^y \rightarrow \sigma_j^y$$

Hence, we get the following expression for commutator after the required sign changes:

$$\begin{aligned}
C^{(1)} &= -2iJ\sigma_0^y(\sigma_{-1}^x + \sigma_1^x) \\
C^{(2)} &= -8J^2(\sigma_1^x\sigma_0^z\sigma_{-1}^x + \sigma_0^z) + 4J\lambda\sigma_0^x(\sigma_{-1}^x + \sigma_1^x) \\
C^{(3)} &= (16J^2 + 4\lambda^2)[H, \partial_\lambda H] = \alpha^2 C^{(1)} \\
C^{(5)} &= [H, [H, C^{(3)}]] = \alpha^2[H, [H, C^{(1)}]] = \alpha^2 C^{(3)} = \alpha^4 C^{(1)}
\end{aligned}$$

Thus, we conclude that we get an expression for exact gauge potential:

$$A_\lambda = \frac{\hbar J}{8J^2 + 2\lambda^2} \sigma_0^y (\sigma_{-1}^x + \sigma_1^x) \quad (41)$$

### 3.1.2 Ising model with local transverse magnetic fields at two sites

$$H = -J \sum_j \sigma_j^x \sigma_{j+1}^x - \lambda(\sigma_0^z + \sigma_1^z) \quad (42)$$

$$\begin{aligned}
C^{(1)} &= -2iJ\sigma_0^y(\sigma_{-1}^x + \sigma_1^x) - 2iJ\sigma_1^y(\sigma_0^x + \sigma_2^x) \\
C^{(2)} &= -8J^2(\sigma_1^x\sigma_{-1}^x + 1)\sigma_0^z + 4J\lambda\sigma_0^x(\sigma_{-1}^x + \sigma_1^x) \\
&\quad - 8J^2(\sigma_0^x\sigma_2^x + 1)\sigma_1^z + 4J\lambda\sigma_1^x(\sigma_0^x + \sigma_2^x) - 8J\lambda\sigma_1^y\sigma_0^y \\
C^{(3)} &= \alpha^2 C^{(1)} + T_2 + T_3 \\
C^{(5)} &= \alpha^4 C^{(1)} + \beta^2 T_2 + \gamma^2 T_3
\end{aligned}$$

where  $\beta^2 = 20(2J^2 + \lambda^2)$ ,  $\gamma^2 = (32J^2 + 17\lambda^2)$ , and operator terms  $T_2 = -24iJ\lambda^2(\sigma_0^y\sigma_1^x + \sigma_1^y\sigma_0^x)$  and  $T_3 = 32iJ^2\lambda(\sigma_{-1}^x\sigma_0^z\sigma_1^y + \sigma_0^y\sigma_1^z\sigma_2^x)$  are terms involves two body operators and three body operators, respectively.

Hence, using theorem proved in appendix A, I can claim  $C^{(2n+1)} = \alpha^{(2n)}C^{(1)} + \beta^{(2n-2)}T_2 + \gamma^{(2n-2)}T_3$ ,  $\forall n > 0$ .

Now using this information, we find that  $A_\lambda$  is given by:

$$A_\lambda = \frac{-i\hbar C^{(1)}}{\alpha^2 + \mu^2} + \frac{T_2}{\mu^2} \frac{i\hbar}{(\mu^2 + \beta^2)} + \frac{T_3}{\mu^2} \frac{i\hbar}{(\mu^2 + \gamma^2)} \quad (43)$$

Details of the derivation are given in appendix C. We note that the last two terms diverge when we take  $\lim_{\mu \rightarrow 0}$  limit. I am still figuring out how to deal with this.

## A Properties of n-commutators

We define the first term as  $C^{(1)} = [H, \partial_\lambda H]$ . Now,  $C^{(2)} = [H, [H, \partial_\lambda H]] = [H, C^{(1)}]$ .

Hence, we can claim using induction argument:

**Theorem 1.**  $C^{(n)} = [H, C^{(n-1)}], n > 1$

Now we will prove another result, which is useful.

**Theorem 2.** If  $C^{(3)} = \alpha^2 C^{(1)} + T$  and  $C_T^{(2)} = \beta^2 T$ , then  $C^{(2n+1)} = \alpha^{(2n)} C^{(1)} + \beta^{(2n-2)} T$ ,  $\forall n > 0$ , where  $T$  is a term involving some operators and  $C_T^{(2)} = [H, [H, T]]$



*Proof.* We can use principle of mathematical induction to prove the theorem. First, we would check if  $P(0)$  is true. Then our next task is to show if  $P(k)$  holds, then  $P(k+1)$  also holds for some unspecified  $k$ .

Here we start with  $n = 5$ , which is our basis ( $P(0)$ ) of our induction argument.

$$\begin{aligned} C^{(5)} &= [H, C^{(4)}] = [H, [H, C^{(3)}]] \\ &= [H, [H, \alpha^2 C^{(1)} + T]] = \alpha^2 [H, [H, C^{(1)}]] + [H, [H, T]] = \alpha^2 C^{(3)} + C_T^{(2)} = \alpha^4 C^{(1)} + \beta^2 T \end{aligned}$$

Hence,  $C^{(5)} = \alpha^4 C^{(1)} + \beta^2 T$ . Thus, it has been show that  $P(0)$  holds.

Our  $P(k)$  statement is  $C^{(2k-1)} = \alpha^{(2k-2)} C^{(1)} + \beta^{(2k-4)} T$ , which we assume to be true. Using this, we will prove that our  $P(k+1)$  statement is true, i.e.  $C^{(2k+1)} = \alpha^{(2k)} C^{(1)} + \beta^{(2k-2)} T$ .

$$\begin{aligned} C^{(2k+1)} &= [H, [H, C^{(2k-1)}]] \\ &= [H, [H, \alpha^{(2k-2)} C^{(1)} + \beta^{(2k-4)} T]] \\ &= \alpha^{(2k-2)} [H, [H, C^{(1)}]] + \beta^{(2k-4)} [H, [H, T]] \\ &= \alpha^{(2k-2)} C^{(3)} + \beta^{(2k-4)} C_T^{(2)} \\ &= \alpha^{(2k)} C^{(1)} + \beta^{(2k-2)} T \end{aligned}$$

Hence, we proved that  $C^{(2n+1)} = \alpha^{(2n)} C^{(1)} + \beta^{(2n-2)} T$ ,  $\forall n > 0$ .  $\square$

## B Adiabatic gauge potential formula general derivation

$$\langle n | A_\lambda | m \rangle = -i \frac{\langle n | \partial_\lambda H | m \rangle}{(E_n - E_m)^2 + \mu^2} (E_n - E_m) \quad (44)$$

$$= -i \int_0^\infty dt e^{-\mu t} \langle n | \partial_\lambda H | m \rangle \sin((E_n - E_m)t) \quad (45)$$

$$= \frac{-i}{2i} \int_0^\infty dt e^{-\mu t} \langle n | \partial_\lambda H | m \rangle \left( e^{i(E_n - E_m)t} - e^{-i(E_n - E_m)t} \right) \quad (46)$$

$$= -\frac{1}{2} \int_0^\infty dt e^{-\mu t} (\langle n | e^{iE_n t} \partial_\lambda H e^{-iE_m t} | m \rangle - \langle n | e^{-iE_n t} \partial_\lambda H e^{iE_m t} | m \rangle) \quad (47)$$

Hence, we can simplify our expression by defining propagator  $U = \exp(-iHt)$ . We note that parameter  $\lambda$  is fixed while we evolve it in the *artificial time*  $t$ .

$$A_\lambda = -\frac{1}{2} \int_0^\infty dt e^{-\mu t} [U^\dagger(t) \partial_\lambda H U(t) - U^\dagger(-t) \partial_\lambda H U(-t)] \quad (48)$$

$$= -\frac{1}{2} \int_0^\infty dt e^{-\mu t} [\partial_\lambda H(t) - \partial_\lambda H(-t)] \quad (49)$$

where  $\partial_\lambda H(t)$  is time-evolved operator  $\partial_\lambda H$  in Heisenberg picture.

We would be using Hadamard (or sometimes called Baker-Campbell-Hausdorff) formula to simplify  $\partial_\lambda H(t)$ .

$$\partial_\lambda H(t) = U^\dagger(t) \partial_\lambda H U(t) \quad (50)$$

$$= \exp(iHt) \partial_\lambda H \exp(-iHt) \quad (51)$$

$$= \partial_\lambda H + it[H, \partial_\lambda H] + \left(\frac{it}{2!}\right)^2 [H, [H, \partial_\lambda H]] + \left(\frac{it}{3!}\right)^3 [H, [H, [H, \partial_\lambda H]]] + \dots \quad (52)$$

Similarly, for  $\partial_\lambda H(-t)$ , we have:

$$\partial_\lambda H(-t) = \partial_\lambda H - it[H, \partial_\lambda H] + \left(\frac{it}{2!}\right)^2 [H, [H, \partial_\lambda H]] - \left(\frac{it}{3!}\right)^3 [H, [H, [H, \partial_\lambda H]]] + \dots \quad (53)$$

Now we see that  $\partial_\lambda H(t\hbar) - \partial_\lambda H(-t\hbar)$  contains only odd power of time  $t$ :

$$\begin{aligned} \partial_\lambda H(t\hbar) - \partial_\lambda H(-t\hbar) &= 2 \left[ it[H, \partial_\lambda H] + \left(\frac{it}{3!}\right)^3 [H, [H, \partial_\lambda H]] + \left(\frac{it}{5!}\right)^5 [H, [H, [H, [H, \partial_\lambda H]]]] + \dots \right] \\ &= 2 \sum_{n=0}^{\infty} \frac{(it)^{2n+1}}{(2n+1)!} C^{(2n+1)} \end{aligned} \quad (54)$$

$$= 2i \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \quad (55)$$

where  $C^{(n)}$  is  $n$ -commutator of  $H$  and  $\partial_\lambda H$ , i.e.  $C^{(n)} = [H, [H, \dots, [H, \partial_\lambda H]]]$ . We define the first term as  $C^{(1)} = [H, \partial_\lambda H]$ , second term as  $C^{(2)} = [H, [H, \partial_\lambda H]] = [H, C^{(1)}]$  and so on and forth. Properties of  $C^{(n)}$  are noted in appendix A.

We can simplify our expression if we call  $\sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)}$  as  $\sin(C^{(1)}t)$ . Thus, we can write:

$$A_\lambda = -i \int_0^\infty dt e^{-\mu t} \sin([H, \partial_\lambda H]t) = -i\hbar \int_0^\infty dt e^{-\mu t} \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \quad (56)$$

## C Adiabatic gauge potential of Ising model with local transverse magnetic fields at two sites

Using  $C^{(2n+1)} = \alpha^{(2n)} C^{(1)} + \beta^{(2n-2)} T_2 + \gamma^{(2n-2)} T_3$ ,  $\forall n > 0$ , we find  $A_\lambda$  using two methods: equation (24) and (28).

Using equation (24), we get:

$$\begin{aligned} A_\lambda &= -i\hbar \int_0^\infty dt e^{-\mu t} \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \\ &= -i\hbar \int_0^\infty dt e^{-\mu t} \left( t C^{(1)} + \sum_{n=1}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} C^{(2n+1)} \right) \\ &= -i\hbar \int_0^\infty dt e^{-\mu t} \left( t C^{(1)} + \sum_{n=1}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} \left( \alpha^{(2n)} C^{(1)} + \beta^{(2n-2)} T_2 + \gamma^{(2n-2)} T_3 \right) \right) \\ &= -i\hbar C^{(1)} \int_0^\infty dt e^{-\mu t} \left( \sum_{n=0}^{\infty} \frac{(-1)^n \alpha^{2n} t^{2n+1}}{(2n+1)!} \right) - i\hbar \int_0^\infty dt e^{-\mu t} \sum_{n=1}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} \left( \beta^{(2n-2)} T_2 + \gamma^{(2n-2)} T_3 \right) \\ &= \frac{-i\hbar C^{(1)}}{\alpha^2 + \mu^2} - i\hbar \int_0^\infty dt e^{-\mu t} (X_2 + X_3) \end{aligned}$$

where  $X_2 = T_2 \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{(2n-2)} t^{2n+1}}{(2n+1)!}$  and  $X_3 = T_3 \sum_{n=1}^{\infty} (-1)^n \frac{\gamma^{(2n-2)} t^{2n+1}}{(2n+1)!}$ . Let's simplify  $X_2$  below:

$$\begin{aligned}
X_2 &= T_2 \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{(2n-2)} t^{2n+1}}{(2n+1)!} \\
&= \frac{T_2}{\beta^2} \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{2n} t^{2n+1}}{(2n+1)!} \\
&= \frac{T_2}{\beta^2} \left( \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{2n} t^{2n+1}}{(2n+1)!} + t - t \right) \\
&= \frac{T_2}{\beta^2} \left( \sum_{n=0}^{\infty} (-1)^n \frac{\beta^{2n} t^{2n+1}}{(2n+1)!} - t \right) \\
&= \frac{T_2}{\beta^2} \left( \frac{\sin(\beta t)}{\beta} - t \right)
\end{aligned}$$

Similarly,  $X_3 = \frac{T_3}{\gamma^3} (\sin(\gamma t) - \gamma t)$ . Now, let's compute the Laplace transform of  $X_2$ :

$$\begin{aligned}
-i\hbar \int_0^{\infty} dt e^{-\mu t} X_2 &= -i\hbar \frac{T_2}{\beta^3} \int_0^{\infty} dt e^{-\mu t} (\sin(\beta t) - \beta t) \\
&= -i\hbar \frac{T_2}{\beta^3} \left( \frac{\beta}{\beta^2 + \mu^2} - \frac{\beta}{\mu^2} \right) \\
&= -i\hbar \frac{T_2}{\beta^2} \left( \frac{1}{\beta^2 + \mu^2} - \frac{1}{\mu^2} \right) \\
&= \frac{i\hbar}{\beta^2 + \mu^2} \frac{T_2}{\mu^2}
\end{aligned}$$

Hence, after we stitch everything together, we get the final expression of  $A_\lambda$ :

$$\boxed{A_\lambda = \frac{-i\hbar C^{(1)}}{\alpha^2 + \mu^2} + \frac{T_2}{\mu^2} \frac{i\hbar}{(\mu^2 + \beta^2)} + \frac{T_3}{\mu^2} \frac{i\hbar}{(\mu^2 + \gamma^2)}} \quad (57)$$

Using equation (28), we get exactly the same expression as shown below:

$$\begin{aligned}
A_\lambda &= \frac{-i\hbar}{\mu} \sum_{n=0}^{\infty} (-1)^n \frac{C^{(2n+1)}}{\mu^{2n+1}} \\
&= \frac{-i\hbar}{\mu} \frac{C^{(1)}}{\mu^1} + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{C^{(2n+1)}}{\mu^{2n+1}} \\
&= \frac{-i\hbar}{\mu} \frac{C^{(1)}}{\mu^1} + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{\alpha^{(2n)} C^{(1)}}{\mu^{2n+1}} + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{(2n-2)}}{\mu^{2n+1}} T_2 + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{\gamma^{(2n-2)}}{\mu^{2n+1}} T_3 \\
&= \frac{-i\hbar C^{(1)}}{\mu} \sum_{n=0}^{\infty} (-1)^n \frac{\alpha^{2n}}{\mu^{2n+1}} + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{\beta^{(2n-2)}}{\mu^{2n+1}} T_2 + \frac{-i\hbar}{\mu} \sum_{n=1}^{\infty} (-1)^n \frac{\gamma^{(2n-2)}}{\mu^{2n+1}} T_3 \\
&= \frac{-i\hbar C^{(1)}}{\alpha^2 + \mu^2} + \frac{T_2}{\mu^2} \frac{i\hbar}{(\mu^2 + \beta^2)} + \frac{T_3}{\mu^2} \frac{i\hbar}{(\mu^2 + \gamma^2)}
\end{aligned}$$

## D Classical adiabatic gauge potential

Let's start by considering classical systems. For such systems, we specify the system by defining Hamiltonian  $H(\lambda)$  in terms of canonical variables  $q_i(\lambda, t)$  and  $p_j(\lambda, t)$ . where  $\lambda$  is an externally

controlled parameter. These variables satisfy the canonical relations:

$$\{q_i, p_j\} = \delta_{ij} \quad (58)$$

where  $\{\dots\}$  denotes the Poisson bracket.

Canonical transformations are transformations of  $q_i$  and  $p_j$  to new variables  $\bar{q}_i$  and  $\bar{p}_j$  such that it preserves Poisson bracket. Hence,

$$\{\bar{q}_i, \bar{p}_j\} = \delta_{ij} \quad (59)$$

What are gauge potentials? Gauge potential  $A_\lambda$  are the generators of continuous canonical transformations in parameter  $\lambda$  space, which can be defined as :

$$q_j(\lambda + \delta\lambda) = q_j - \frac{\partial A_\lambda}{\partial p_j} \delta\lambda \Rightarrow \frac{\partial q_j}{\partial \lambda} = -\frac{\partial A_\lambda}{\partial p_j} = \{A_\lambda, q_j\} \quad (60)$$

$$p_j(\lambda + \delta\lambda) = p_j + \frac{\partial A_\lambda}{\partial q_j} \delta\lambda \Rightarrow \frac{\partial p_j}{\partial \lambda} = \frac{\partial A_\lambda}{\partial q_j} = \{A_\lambda, p_j\} \quad (61)$$

We can verify that these transformations are canonical upto order  $\delta\lambda^2$  because we can show that:

$$\{q_j(\lambda + \delta\lambda), p_j(\lambda + \delta\lambda)\} = \delta_{ij} + O(\delta\lambda^2) \quad (62)$$

Let's try to understand by taking an example of continuous canonical transformation. We would shift the position coordinate by  $X_i$ . Here our parameter  $\lambda$  is  $X_i$

$$q_i(X_i, t) = q_i(0, t) - X_i \quad (63)$$

$$p_i(X_i, t) = p_i(0, t) \quad (64)$$

Using equation 61, we see that  $\frac{\partial A_{X_i}}{\partial q_j} = 0$  and  $-\frac{\partial A_{X_i}}{\partial p_j} = -\delta_{ij}$ . Hence,  $A_{X_i} = p_j + C_j$ , where  $C_j$  are arbitrary constants of integration. This is the gauge choice we have got in defining these gauge potentials.

## E An example of variational approximation scheme: non-integrable Ising spin chain

Let's consider Ising quantum spin chain with transverse and longitudinal field whose Hamiltonian is given by:

$$H_0 = \sum_{j=1}^{L-1} J(\lambda) \sigma_j^z \sigma_{j+1}^z + \sum_{j=1}^L (Z_j(\lambda) \sigma_j^z + X_j(\lambda) \sigma_j^x) \quad (65)$$

We note that for either  $Z_j = 0$  or  $X_j = 0$ , this model is integrable. Apart from these cases, this model is non-integrable.<sup>5</sup>

Let us consider a Counter-diabatic (CD) protocol for turning on an additional  $x$  magnetic field from  $\lambda_i = 0$  to  $\lambda_f = -10J$  in a periodic chain described by  $H_0 + \lambda \sigma_0^x$ , where  $H_0$  is given by equation 65 with  $J = 1$ ,  $Z_j = 2$  and  $X_j = 0.8$ . Hence, our bare Hamiltonian  $H_b$  (which is a special case of  $H_0$ ) is given by:

$$H_b = \sum_{j=1}^{L-1} \sigma_j^z \sigma_{j+1}^z + \sum_{j=1}^L (2\sigma_j^z + 0.8\sigma_j^x) + \lambda \sigma_0^x \quad (66)$$

where  $\lambda$  is a protocol.

---

<sup>5</sup>In [8], they have mentioned in their paper which parameter are best for this model to be robustly non-integrable. Since our method also depends on exact diagonalization, we should use their results.




Figure 1: Protocol chosen for going from  $\lambda_i = 0$  to  $\lambda_f = -10J$  in time  $\tau$

Initial Hamiltonian is defined by  $\lambda = \lambda_i = 0$  and final Hamiltonian is specified by  $\lambda = \lambda_f = -10J$ . Our problem is to find an approximate gauge potential such that as we tune our  $\lambda$  from 0 to  $-10J$ , we should reach the ground state of our final Hamiltonian with minimal “loss” possible after starting from the ground state of our initial Hamiltonian. If our loss is minimal, then fidelity  $F^2$  of our final state will be *high* and energy of state above ground state  $E - E_0$  would be *small*, where  $F^2 = |\langle \psi(t) | \psi(t)_{GS} \rangle|^2$  and  $E - E_0 = \langle \psi(t) | H | \psi(t) \rangle - \langle \psi_{GS}(t) | H | \psi(t)_{GS} \rangle$

We choose  $\lambda$  protocol (figure 1) that goes from  $\lambda_i = 0$  to  $\lambda_f = -10J$  in time  $\tau$  as:

$$\lambda(t) = \lambda_0 + (\lambda_f - \lambda_0) \sin^2 \left( \frac{\pi}{2} \sin^2 \left( \frac{t\pi}{2\tau} \right) \right), t \in [0, \tau] \quad (67)$$

The *naive* way to drive our system will be take just our bare Hamiltonian  $H_b$  and see the performance by computing  $F^2$  and  $E - E_0$  as we change duration of protocol  $\tau$ . This is shown in blue line of figure 2. We note that increasing  $\tau$  improves our performance no matter how we drive our system because we are going towards adiabatic limit.

For our  $\lambda$  - dependent Hamiltonian  $H_0$ , approximate gauge potential is chosen to be

$$A_\lambda^* = \sum_j \alpha_j \sigma_j^y \quad (68)$$

where  $\alpha_j$  are found using variational approach given in [9]. They find that  $\alpha_j$  for  $H_0$  is given by

$$\alpha_j = \frac{1}{2} \frac{Z_j X'_j - X_j Z'_j}{Z_j^2 + X_j^2 + 2J^2} \quad (69)$$

Now for our  $H_b$ ,  $\alpha_j$  is given by

$$\alpha_j = \delta_{j,0} \frac{1}{6 + (\lambda + 0.8)^2} \quad (70)$$

Hence, our Hamiltonian with gauge potential term (CD term) will be :

$$H_{CD} = H_b + \dot{\lambda} A_\lambda^* \quad (71)$$

$$= H_b + \dot{\lambda} \alpha_0 \sigma_0^y \quad (72)$$

In red line of figure 2, we do find that Hamiltonian with local CD term  $H_{CD}$  does indeed give a better performance by increasing fidelity  $F^2$  and decreasing energy above ground state  $E - E_0$  for short protocol duration  $\tau$ . In Dries’s paper [9], they show similar results in their figure 4, where they have used spin chain of  $L = 15$ .




Figure 2: Fidelity  $F^2$  and final energy above ground state  $E - E_0$  for  $L=12$  spin chains

## F Transverse Field Ising model: calculations in spin basis

We would study another integrable model, which is called Transverse Field Ising model. This model shows quantum phase transition between ferromagnetic and paramagnetic phases. It's Hamiltonian is given by:

$$H = J \sum_j \sigma_j^x \sigma_{j+1}^x + h \sum_j \sigma_j^z + \lambda \sigma_0^z \quad (73)$$

This model satisfies Ising symmetry  $G = \Pi_i \sigma_i^z$  since  $[H, G] = 0$ .

Since this model's exact gauge potential is already known in literature [4, 5], it will be good to find either exact or approximate gauge potential using our regulator method.

Let's find out  $A_\lambda$  for this Hamiltonian for which we need to compute different odd-powered commutator  $[H, \partial_\lambda H]$ , where  $\partial_\lambda H = \sigma_0^z$ . Here we begin:

$$C^{(1)} = -2iJ\sigma_0^y(\sigma_{-1}^x + \sigma_1^x) \quad (74)$$

$$C^{(2)} = 8J^2(\sigma_0^z + \sigma_{-1}^x \sigma_0^z \sigma_1^x) - 4J\lambda(\sigma_{-1}^x + \sigma_1^x)\sigma_0^x - 4hJ((\sigma_{-1}^x + \sigma_1^x)\sigma_0^x - (\sigma_{-1}^y + \sigma_1^y)\sigma_0^y) \quad (75)$$

$$\begin{aligned} C^{(3)} = & -8i(2h^2J\sigma_{-1}^x\sigma_0^y + 2h^2J\sigma_{-1}^y\sigma_0^x + 2h^2J\sigma_0^x\sigma_1^y + 2h^2J\sigma_0^y\sigma_1^x - hJ^2\sigma_{-2}^z\sigma_{-1}^x\sigma_0^y \\ & - 3hJ^2\sigma_{-1}^x\sigma_0^z\sigma_1^y - 3hJ^2\sigma_{-1}^y\sigma_0^z\sigma_1^x - hJ^2\sigma_0^y\sigma_1^z\sigma_2^z \\ & + 2hJ\lambda\sigma_{-1}^x\sigma_0^y + 2hJ\lambda\sigma_{-1}^y\sigma_0^x + 2hJ\lambda\sigma_0^x\sigma_1^y + 2hJ\lambda\sigma_0^y\sigma_1^x + 4J^3\sigma_{-1}^x\sigma_0^y \\ & + 4J^3\sigma_0^y\sigma_1^x + J\lambda^2\sigma_{-1}^x\sigma_0^y + J\lambda^2\sigma_0^y\sigma_1^x) \end{aligned}$$

$$\begin{aligned} C^{(3)} = & -8i(2h^2J(\sigma_{-1}^x + \sigma_1^x)\sigma_0^y + 2h^2J(\sigma_{-1}^y + \sigma_1^y)\sigma_0^x - hJ^2(\sigma_{-2}^z\sigma_{-1}^x + \sigma_1^z\sigma_2^z)\sigma_0^y \\ & - 3hJ^2(\sigma_{-1}^x\sigma_1^y + \sigma_{-1}^y\sigma_1^x)\sigma_0^z + 2hJ\lambda(\sigma_{-1}^x + \sigma_1^x)\sigma_0^y + 2hJ\lambda(\sigma_{-1}^y + \sigma_1^y)\sigma_0^x \\ & + 4J^3(\sigma_{-1}^x + \sigma_1^x)\sigma_0^y + J\lambda^2(\sigma_{-1}^x + \sigma_1^x)\sigma_0^y) \end{aligned}$$

After rearranging terms of  $(\sigma_{-1}^x + \sigma_1^x)\sigma_0^y$ , we get :

$$\begin{aligned} C^{(3)} = & \alpha^2 C^{(1)} - 16ihJ(h + \lambda)(\sigma_{-1}^y + \sigma_1^y)\sigma_0^x + 8ihJ^2(\sigma_{-2}^z\sigma_{-1}^x + \sigma_1^z\sigma_2^z)\sigma_0^y \\ & 24ihJ^2(\sigma_{-1}^x\sigma_1^y + \sigma_{-1}^y\sigma_1^x)\sigma_0^z \end{aligned}$$

where  $\alpha^2 = 4(4J^2 + 2h^2 + \lambda^2 + 2h\lambda) = (4J^2 + h^2 + (h + \lambda)^2)$

## References

- [1] Mustafa Demirplak and Stuart A Rice. Adiabatic population transfer with control fields. *The Journal of Physical Chemistry A*, 107(46):9937–9945, 2003.
- [2] Mustafa Demirplak and Stuart A Rice. Assisted adiabatic passage revisited. *The Journal of Physical Chemistry B*, 109(14):6838–6844, 2005.
- [3] MV Berry. Transitionless quantum driving. *Journal of Physics A: Mathematical and Theoretical*, 42(36):365303, 2009.
- [4] Adolfo del Campo, Marek M Rams, and Wojciech H Zurek. Assisted finite-rate adiabatic passage across a quantum critical point: exact solution for the quantum ising model. *Physical review letters*, 109(11):115703, 2012.
- [5] Michael Kolodrubetz, Pankaj Mehta, and Anatoli Polkovnikov. Geometry and non-adiabatic response in quantum and classical systems. *arXiv preprint arXiv:1602.01062*, 2016.
- [6] Luca D’Alessio, Yariv Kafri, Anatoli Polkovnikov, and Marcos Rigol. From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics. *Advances in Physics*, 65(3):239–362, 2016.
- [7] Christopher Jarzynski. Generating shortcuts to adiabaticity in quantum and classical dynamics. *Physical Review A*, 88(4):040101, 2013.
- [8] Hyungwon Kim and David A Huse. Ballistic spreading of entanglement in a diffusive nonintegrable system. *Physical review letters*, 111(12):127205, 2013.
- [9] Dries Sels and Anatoli Polkovnikov. Minimizing irreversible losses in quantum systems by local counterdiabatic driving. *Proceedings of the National Academy of Sciences*, page 201619826, 2017.