

Quantum control of NV center using counter-diabatic driving

Mohit

January 12, 2018

1 Introduction

The Hamiltonian for the ground state of the NV center can be written as :

$$H_{NV} = \hbar\Delta S_z^2 + \hbar\gamma_e \vec{S} \cdot \vec{B}_{ext} + \hbar\gamma_n \vec{I} \cdot \vec{B}_{ext} + A \vec{S} \cdot \vec{I} \quad (1)$$

where $\Delta = 2\pi \times 2.87$ GHz is zero-field splitting, $\gamma_e = 2.8$ MHz/G is the gyromagnetic ratio of electron in the NV center, $\gamma_n \simeq 1$ kHz/G is the gyromagnetic ratio of nuclear spin, \vec{S} (\vec{I}) is the spin of electron (nucleus) and $A \simeq 3$ MHz is magnitude of spin-spin coupling which leads to hyperfine splitting. The second and third terms are due to Zeeman effect.

Since $\gamma \propto 1/m$ and nucleus is heavier than electron, we have $\gamma_e \gg \gamma_n$. To simplify our model, we will ignore the last two terms resulting in the following Hamiltonian :

$$H_{NV} = \hbar\Delta S_z^2 + \hbar\gamma_e \vec{S} \cdot \vec{B}_{ext} \quad (2)$$

It seems that the above Hamiltonian should be experimentally realizable [1]. Moreover, the ground state of electron in the NV center is a spin triplet with $|0\rangle, |-1\rangle, |1\rangle$ spin sub-levels. They are defined in S_z basis, where \hat{z} direction is along the NV center axis. If there is no external magnetic field, then $|-1\rangle$ and $|1\rangle$ levels are degenerate, and $\hbar\Delta$ is the energy difference between $|0\rangle$ and $|\pm 1\rangle$ energy levels.

2 Static magnetic field

Let's choose magnetic field to be in x-direction¹. Then we have:

$$H_{NV} = \hbar\Delta S_z^2 + \hbar\gamma_e S_x B = \Lambda S_z^2 + \lambda S_x$$

where $\Lambda = \hbar\Delta$ and $\lambda = \hbar\gamma_e B$. Magnetic field is going to be our control parameter in this problem.

2.1 Energy levels as a function of magnetic field

Using spin algebra (appendix B), we obtain Hamiltonian in the S_z basis ($|-1\rangle, |0\rangle, |1\rangle$):

$$H = \begin{pmatrix} \beta & \alpha & 0 \\ \alpha & 0 & \alpha \\ 0 & \alpha & \beta \end{pmatrix} \quad \text{where } \alpha = \lambda/\sqrt{2}, \beta = \Lambda \quad (3)$$

Energy eigenvalues are given by:

$$\lambda_0 = \beta, \quad \lambda_1 = (\beta - \sqrt{\beta^2 + 8\alpha^2})/2, \quad \lambda_2 = (\beta + \sqrt{\beta^2 + 8\alpha^2})/2$$

We should remember that $\alpha \propto B$. Hence, it makes sense that when $\alpha = 0$, there is a two-fold degeneracy and zero field energy gap is given by $\beta = \hbar\Delta$. Now let's have a look at eigenvectors:

$$\nu_0 = (-1, 0, 1), \quad \nu_1 = (1, -(\beta + \sqrt{\beta^2 + 8\alpha^2})/2\alpha, 1), \quad \nu_2 = (1, -(\beta - \sqrt{\beta^2 + 8\alpha^2})/2\alpha, 1)$$

¹If magnetic field is in z-direction, then gauge potential is zero. This is so because increasing B_z does not induce any excitations.




Figure 1: Avoided level crossing as a function of effective magnetic field

2.2 Adiabatic gauge potential

Now let's compute adiabatic gauge potential $A_\lambda = i\hbar\partial_\lambda$. Its' equation of motion is given by:

$$[H, \partial_\lambda H + \frac{i}{\hbar}[A_\lambda, H]] = 0 \quad (4)$$

Eigen-value equation is given by $H(\lambda)|n(\lambda)\rangle = E_n(\lambda)|n(\lambda)\rangle$. Let's derive diagonal and off-diagonal elements:

- **n-th diagonal element:** $A_\lambda^n = \langle n|A_\lambda|n\rangle = i\hbar\langle n|\partial_\lambda|n\rangle$
- **off-diagonal element:** We use the identity $\langle m|H(\lambda)|n\rangle = 0$, $n \neq m$ and then differentiate with respect to λ to obtain:

$$\langle m|A_\lambda|n\rangle = -i\hbar \frac{\langle m|\partial_\lambda H|n\rangle}{E_m - E_n} \quad (5)$$

where both energies (E_m, E_n) and eigenvectors $(|m\rangle, |n\rangle)$ depend on λ .

Here $\partial_\lambda H = S_x$ whose matrix representation is given in appendix B. We find that \tilde{A}_λ in energy basis is given by :

$$\tilde{A}_\lambda = i\hbar\tilde{N} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \text{where } \tilde{N} = \frac{\beta\hbar}{8\alpha^2 + \beta^2} \quad (6)$$

In S_z basis, we have $A_\lambda = U\tilde{A}_\lambda U^T$ where U is the unitary transformation that diagonalizes Hamiltonian, i.e. $H_d = U^T H U$. A_λ is given below:

$$A_\lambda = i\hbar N \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad \text{where } N = \frac{\beta\hbar}{\sqrt{16\alpha^2 + 2\beta^2}}$$

The above matrix of A_λ can be expanded in the basis of $SU(3)$ to expand. This basis is composed of Gell-Mann matrices [2], which are represented as λ_i . For our purpose, most important Gell-Mann matrices are λ_2 and λ_7 whose representation is given below [3, 4]:

$$\lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \frac{1}{\sqrt{2}\hbar^2}(\hbar S_y + S_y S_z + S_z S_y)$$

$$\lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} = \frac{1}{\sqrt{2}\hbar^2}(\hbar S_y - S_y S_z - S_z S_y)$$

Hence, we find that $A_\lambda = \hbar N(\lambda_2 - \lambda_7)$ which can be written as:

$$\boxed{A_\lambda = N_0(S_y S_z + S_z S_y)} \quad (7)$$

where $N_0 = \frac{\beta}{\sqrt{8\alpha^2 + \beta^2}} = \frac{\Delta}{\sqrt{4\gamma_e^2 B^2 + \Delta^2}}$.

3 Periodic magnetic field with noise

Let's choose magnetic field to be $\vec{B} = (B_x(t), 0, 0)$ where $B_x(t) = B_0 \sin(\omega t) + \epsilon(t)$ and $\epsilon(t)$ is an infinitesimal white noise which satisfies $\overline{\epsilon(t)\epsilon(t')} = \kappa\delta(t-t')$ ².

Then we have:

$$\begin{aligned} H_{NV} &= \Lambda S_z^2 + \hbar\gamma_e S_x B_x(t) \\ &= \Lambda S_z^2 + \lambda S_x \end{aligned}$$

where $\Lambda = \hbar\Delta$ and $\lambda = \hbar\gamma_e(B_0 \sin(\omega t) + \epsilon(t))$. For now on, we will work in the unit in which $\hbar\gamma_e = 1$ so that $\lambda = (B_0 \sin(\omega t) + \epsilon(t))$ and $\Lambda = \Delta/\gamma_e$.

Using Fermi Golden rule, we can derive transition rate $\langle \Gamma_n \rangle$ [5] as follows:

$$\Gamma_n(\omega) = \kappa \sum_{m \neq n} |\langle n | G_\lambda | m \rangle|^2 \delta(E_n - E_m - \hbar\omega) \quad (8)$$

where $G_\lambda = \partial_\lambda H + \frac{i}{\hbar}[A_\lambda, H]$ and ω is frequency of the periodic external drive. More details are given in appendix.

A Gell-Mann matrices

$$\begin{aligned} \lambda_1 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & \lambda_2 &= \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & \lambda_3 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ \lambda_4 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} & \lambda_5 &= \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix} & \lambda_6 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \\ \lambda_7 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} & \lambda_8 &= \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \end{aligned}$$

These matrices are traceless and Hermitian.

B Spin Algebra

$$[S_x, S_y] = i\hbar S_z, \quad [S_y, S_z] = i\hbar S_x, \quad [S_z, S_x] = i\hbar S_y \quad (9)$$

$$S^2|s, m\rangle = \hbar^2 s(s+1)|s, m\rangle, \quad S_z|s, m\rangle = \hbar m|s, m\rangle \quad (10)$$

²If the system is in equilibrium, then fluctuation-dissipation theorem dictates $\kappa = T$

$$S_{\pm}|s, m\rangle = \hbar\sqrt{s(s+1) - m(m\pm 1)}|s, m\pm 1\rangle \quad (11)$$

where $S_+ = S_x + iS_y$ and $S_- = S_x - iS_y$. Hence, we get $S_x = (S_+ + S_-)/2$ and $S_y = (S_+ - S_-)/2i$

$$S_+ = \sqrt{2}\hbar \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad S_- = \sqrt{2}\hbar \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad (12)$$

Hence,

$$S_x = \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad S_y = i\frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \quad (13)$$

C Transition rate for periodically driven noisy system

Let's consider a Hamiltonian:

$$\mathcal{H}_{\mathcal{X}} = \mathcal{H}(\lambda) + \dot{\lambda}\mathcal{X} \quad (14)$$

where $\lambda = \lambda_0 + \epsilon(t)$ and $\epsilon(t)$ is an infinitesimal white noise which satisfies $\overline{\epsilon(t)\epsilon(t')} = \kappa\delta(t-t')$ ³. A system under white noise would lead to transitions between all eigenstates. We will see how much there is reduction in transition and dissipation rate with our approximate gauge potential.

We can simplify the above expression:

$$\mathcal{H}_{\mathcal{X}} \approx \mathcal{H}(\lambda_0) + \epsilon\partial_{\lambda}\mathcal{H} + \dot{\epsilon}\mathcal{X} \quad (15)$$

Our expressions will involve G_{λ} which is given as $G_{\lambda}(\mathcal{X}) = \partial_{\lambda}H + \frac{i}{\hbar}[\mathcal{X}, H]$ where $\mathcal{X} = A_{\lambda}$.

Now we would drive the system periodically in addition to the white noise we have in the system. So, in this protocol, time dependence of λ is given as $\lambda(t) = \lambda_0 \sin(\omega t) + \epsilon(t)$. We can use Fermi's golden rule (using results from [6]) to derive transition rate $\langle\Gamma_n\rangle$ [5] as follows:

$$\Gamma_n(\omega) = \kappa \sum_{m \neq n} |\langle n|G_{\lambda}|m\rangle|^2 \delta(E_n - E_m - \hbar\omega) \quad (16)$$

where ω is frequency of the periodic external drive.

References

- [1] Shonali Dhingra and Brian D'Urso. Nitrogen vacancy centers in diamond as angle-squared sensors. *Journal of Physics: Condensed Matter*, 29(18):185501, 2017.
- [2] Murray Gell-Mann. Symmetries of baryons and mesons. *Physical Review*, 125(3):1067, 1962.
- [3] Reinhold A Bertlmann and Philipp Krammer. Bloch vectors for qudits. *Journal of Physics A: Mathematical and Theoretical*, 41(23):235303, 2008.
- [4] Philipp Krammer. *Entanglement beyond two qubits: geometry and entanglement witnesses*. PhD Thesis, 2009.
- [5] Michael Kolodrubetz, Pankaj Mehta, and Anatoli Polkovnikov. Geometry and non-adiabatic response in quantum and classical systems. *arXiv preprint arXiv:1602.01062*, 2016.
- [6] Aashish A Clerk, Michel H Devoret, Steven M Girvin, Florian Marquardt, and Robert J Schoelkopf. Introduction to quantum noise, measurement, and amplification. *Reviews of Modern Physics*, 82(2):1155, 2010.

³If the system is in equilibrium, then fluctuation -dissipation theorem dictates $\kappa = T$