


FIGURE 2.5

Hydration When LiCl dissolves, the ions are hydrated. The attraction between ions and water molecules is strong enough that each ion in solution is surrounded by water molecules.



Dissolving Ionic Compounds in Aqueous Solution

The polarity of water molecules plays an important role in the formation of solutions of ionic compounds in water. The slightly charged parts of water molecules attract the ions in the ionic compounds and surround them to keep them separated from the other ions in the solution. Suppose we drop a few crystals of lithium chloride into a beaker of water. At the crystal surface, water molecules come into contact with Li^+ and Cl^- ions. The positive ends of the water molecules are attracted to Cl^- ions, while the negative ends are attracted to Li^+ ions. The attraction between water molecules and the ions is strong enough to draw the ions away from the crystal surface and into solution, as illustrated in **Figure 2.5**. **This solution process with water as the solvent is referred to as hydration.** The ions are said to be *hydrated*. As hydrated ions diffuse into the solution, other ions are exposed and drawn away from the crystal surface by the solvent. The entire crystal gradually dissolves, and hydrated ions become uniformly distributed in the solution.

When crystallized from aqueous solutions, some ionic substances form crystals that incorporate water molecules. These crystalline compounds, known as *hydrates*, retain specific ratios of water molecules, as shown in figure **Figure 2.6**, and are represented by formulas such as $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$. Heating the crystals of a hydrate can drive off the water of hydration and leave the anhydrous salt. When a crystalline hydrate dissolves in water, the water of hydration returns to the solvent. The behavior of a solution made from a hydrate is no different from the behavior of one made from the anhydrous form. Dissolving either results in a system containing hydrated ions and water.

Nonpolar Solvents

Ionic compounds are generally not soluble in nonpolar solvents such as carbon tetrachloride, CCl_4 , and toluene, $\text{C}_6\text{H}_5\text{CH}_3$. The nonpolar solvent molecules do not attract the ions of the crystal strongly enough to overcome the forces holding the crystal together.

Would you expect lithium chloride to dissolve in toluene? No, LiCl is not soluble in toluene. LiCl and $\text{C}_6\text{H}_5\text{CH}_3$ differ widely in bonding, polarity, and intermolecular forces.

FIGURE 2.6

Hydrate Hydrated copper(II) sulfate has water as part of its crystal structure. Heating releases the water and produces the anhydrous form of the substance, which has the formula CuSO_4 .

