



General Care: Choking (Adult/Child)

1. Position self to the side and slightly behind the choking person.
 - a. For a small child, you may need to kneel behind them rather than stand.
2. Give 5 back blows.
 - a. Use the heel of the hand to strike between the shoulder blades.
3. If no improvement, have the person stand up straight.
4. Move behind the person; bend your knees slightly for balance and support.
5. Give 5 abdominal thrusts.
 - a. Pull inward and upward each time.
6. Continue giving 5 back blows and 5 abdominal thrusts.
 - a. Continue until the person can cough, cry or speak or becomes unresponsive.
7. If the person becomes unresponsive, lower them to a firm, flat surface and begin CPR (starting with compressions) according to your level of training.
 - a. Trained responders: After each set of compressions and before attempting breaths:
 - b. Open the person's mouth.
 - c. Look for an object.
 - d. If seen, remove it using your finger. NEVER do a finger sweep unless you actually see an object.

General Care: Choking Infant



1. Position infant face-down along your forearm using your thigh for support.
 - a. Keep the infant's head lower than their body.
2. Give 5 firm back blows.
 - a. Use the heel of the hand to strike between the shoulder blades.
3. Turn infant face-up with their head lower than their body.
4. Give 5 quick chest thrusts.
 - a. Chests thrusts should be about 1 ½ inches deep.
5. Continue giving 5 back blows and 5 chest thrusts.
 - a. Continue until the infant can cough or cry or becomes unresponsive.
6. If the infant becomes unresponsive, lower them to a firm, flat surface and begin CPR (starting with compressions) according to your level of training.
 - a. Trained responders: After each set of compressions and before attempting breaths:
 - b. Open the infant's mouth.
 - c. Look for an object.
 - d. If seen, remove it using your pinky. NEVER do a pinky sweep unless you actually see the object.