

ADVISORS

A number of experts in the library and medical communities provided invaluable assistance in the formulation of this encyclopedia. Our advisory board performed a myriad of duties, from defining the scope of coverage to reviewing individual entries for accuracy and accessibility. The editor would like to express her appreciation to them.

MEDICAL ADVISORS

Rosalyn Carson-DeWitt, M.D.
Durham, NC

Larry I. Lutwick M.D., F.A.C.P.
Director, Infectious Diseases
VA Medical Center
Brooklyn, NY

Samuel Uretsky, Pharm.D.
Pharmacist
Wantagh, NY

CONTRIBUTORS

Margaret Alic, Ph.D.
Science Writer
Eastsound, WA

Janet Byron Anderson
Linguist/Language Consultant
Rocky River, OH

Lisa Andres, M.S., C.G.C.
*Certified Genetic Counselor and
Medical Writer*
San Jose, CA

Greg Annussek
Medical Writer/Editor
New York, NY

Bill Asenjo, Ph.D.
Science Writer
Iowa City, IA

Sharon A. Aufox, M.S., C.G.C.
Genetic Counselor
Rockford Memorial Hospital
Rockford, IL

Sandra Bain Cushman
*Massage Therapist, Alexander
Technique Practitioner*
Charlottesville, VA

Howard Baker
Medical Writer
North York, Ontario

Laurie Barclay, M.D.
Neurological Consulting Services
Tampa, FL

Jeanine Barone
Nutritionist, Exercise Physiologist
New York, NY

Julia R. Barrett
Science Writer
Madison, WI

Donald G. Barstow, R.N.
Clinical Nurse Specialist
Oklahoma City, OK

Carin Lea Beltz, M.S.
*Genetic Counselor and Program
Director*
The Center for Genetic
Counseling
Indianapolis, IN

Linda K. Bennington, C.N.S.
Science Writer
Virginia Beach, VA

Issac R. Berniker
Medical Writer
Vallejo, CA

Kathleen Berrisford, M.S.V.
Science Writer

Bethanne Black
Medical Writer
Atlanta, GA

Jennifer Bowjanowski, M.S.,
C.G.C.
Genetic Counselor
Children's Hospital Oakland
Oakland, CA

Michelle Q. Bosworth, M.S.,
C.G.C.
Genetic Counselor
Eugene, OR

Barbara Boughton
Health and Medical Writer
El Cerrito, CA

Cheryl Branche, M.D.
Retired General Practitioner
Jackson, MS

Michelle Lee Brandt
Medical Writer
San Francisco, CA

Maury M. Breecher, Ph.D.
Health Communicator/Journalist
Northport, AL

Ruthan Brodsky
Medical Writer
Bloomfield Hills, MI

Tom Brody, Ph.D.
Science Writer
Berkeley, CA

Leonard C. Bruno, Ph.D.
Medical Writer
Chevy Chase, MD

Diane Calbrese
*Medical Sciences and Technology
Writer*
Silver Spring, Maryland

Richard H. Camer
Editor
International Medical News Group
Silver Spring, MD

Rosalyn Carson-DeWitt, M.D.
Medical Writer
Durham, NC

Lata Cherath, Ph.D.
Science Writing Intern
Cancer Research Institute
New York, NY

Linda Chrisman
Massage Therapist and Educator
Oakland, CA

Lisa Christenson, Ph.D.
Science Writer
Hamden, CT

Geoffrey N. Clark, D.V.M.
Editor
Canine Sports Medicine
Update
Newmarket, NH

Rhonda Cloos, R.N.
Medical Writer
Austin, TX

Gloria Cooksey, C.N.E.
Medical Writer
Sacramento, CA

Amy Cooper, M.A., M.S.I.
Medical Writer
Vermillion, SD

David A. Cramer, M.D.
Medical Writer
Chicago, IL

Esther Csapo Rastega, R.N.,
B.S.N.
Medical Writer
Holbrook, MA

Arnold Cua, M.D.
Physician
Brooklyn, NY

Tish Davidson, A.M.
Medical Writer
Fremont, California

Dominic De Bellis, Ph.D.
Medical Writer/Editor
Mahopac, NY

Lori De Milto
Medical Writer
Sicklerville, NJ

Robert S. Dinsmoor
Medical Writer
South Hamilton, MA

Stephanie Dionne, B.S.
Medical Writer
Ann Arbor, MI

Martin W. Dodge, Ph.D.
Technical Writer/Editor
Centinela Hospital and Medical
Center
Inglewood, CA

David Doermann
Medical Writer
Salt Lake City, UT

Stefanie B. N. Dugan, M.S.
Genetic Counselor
Milwaukee, WI

Doug Dupler, M.A.
Science Writer
Boulder, CO

Thomas Scott Eagan
Student Researcher
University of Arizona
Tucson, AZ

Altha Roberts Edgren
Medical Writer
Medical Ink
St. Paul, MN

Karen Ericson, R.N.
Medical Writer
Estes Park, CO

L. Fleming Fallon Jr., M.D.,
Dr.PH
*Associate Professor of Public
Health*
Bowling Green State University
Bowling Green, OH

Faye Fishman, D.O.
Physician
Randolph, NJ

Janis Flores
Medical Writer
Lexikon Communications
Sebastopol, CA

Risa Flynn
Medical Writer
Culver City, CA

Paula Ford-Martin
Medical Writer
Chaplin, MN

Janie F. Franz
Writer
Grand Forks, ND

Sallie Freeman, Ph.D., B.S.N.
Medical Writer
Atlanta, GA

Rebecca J. Frey, Ph.D.
*Research and Administrative
Associate*
East Rock Institute
New Haven, CT

Cynthia L. Frozena, R.N.
Nurse, Medical Writer
Manitowoc, WI

Jason Fryer
Medical Writer
San Antonio, TX

Ron Gasbarro, Pharm.D.
Medical Writer
New Milford, PA

Julie A. Gelderloos
Biomedical Writer
Playa del Rey, CA

Gary Gilles, M.A.
Medical Writer
Wauconda, IL

Harry W. Golden
Medical Writer
Shoreline Medical Writers
Old Lyme, CT

Debra Gordon
Medical Writer
Nazareth, PA

Megan Gourley
Writer
Germantown, MD

Jill Granger, M.S.
Senior Research Associate
University of Michigan
Ann Arbor, MI

Alison Grant
Medical Writer
Averill Park, NY

Elliot Greene, M.A.
*former president, American
Massage Therapy Association*
Massage Therapist
Silver Spring, MD

Peter Gregutt
Writer
Asheville, NC

Laith F. Gulli, M.D.
M.Sc., M.Sc.(MedSci), M.S.A.,
Msc.Psych, MRSNZ
FRSH, FRIPHH, FAIC, FZS
DAPA, DABFC, DABCI
*Consultant Psychotherapist in
Private Practice*
Lathrup Village, MI

Kapil Gupta, M.D.
Medical Writer
Winston-Salem, NC

Maureen Haggerty
Medical Writer
Ambler, PA

Clare Hanrahan
Medical Writer
Asheville, NC

Ann M. Haren
Science Writer
Madison, CT

Judy C. Hawkins, M.S.
Genetic Counselor
The University of Texas Medical
Branch
Galveston, TX

Caroline Helwick
Medical Writer
New Orleans, LA

David Helwig
Medical Writer
London, Ontario

Lisette Hilton
Medical Writer
Boca Raton, FL

Katherine S. Hunt, M.S.
Genetic Counselor
University of New Mexico Health
Sciences Center
Albuquerque, NM

Kevin Hwang, M.D.
Medical Writer
Morristown, NJ

Holly Ann Ishmael, M.S.,
C.G.C.
Genetic Counselor
The Children's Mercy Hospital
Kansas City, MO

Dawn A. Jacob, M.S.
Genetic Counselor
Obstetrix Medical Group of
Texas
Fort Worth, TX

Sally J. Jacobs, Ed.D.
Medical Writer
Los Angeles, CA

Michelle L. Johnson, M.S., J.D.
*Patent Attorney and Medical
Writer*
Portland, OR

Paul A. Johnson, Ed.M.
Medical Writer
San Diego, CA

Cindy L. A. Jones, Ph.D.
Biomedical Writer
Sagescript Communications
Lakewood, CO

David Kaminstein, M.D.
Medical Writer
West Chester, PA

Beth A. Kapes
Medical Writer
Bay Village, OH

Janet M. Kearney
Freelance writer
Orlando, FL

Christine Kuehn Kelly
Medical Writer
Havertown, PA

Bob Kirsch
Medical Writer
Ossining, NY

Joseph Knight, P.A.
Medical Writer
Winton, CA

Melissa Knopper
Medical Writer
Chicago, IL

Karen Krajewski, M.S., C.G.C.
Genetic Counselor
Assistant Professor of Neurology
Wayne State University
Detroit, MI

Jeanne Krob, M.D., F.A.C.S.
Physician, writer
Pittsburgh, PA

Jennifer Lamb
Medical Writer
Spokane, WA

Richard H. Lampert
Senior Medical Editor
W.B. Saunders Co.
Philadelphia, PA

Jeffrey P. Larson, R.P.T.
Physical Therapist
Sabin, MN

Jill Lasker
Medical Writer
Midlothian, VA

Kristy Layman
Music Therapist
East Lansing, MI

Victor Leipzig, Ph.D.
Biological Consultant
Huntington Beach, CA

Lorraine Lica, Ph.D.
Medical Writer
San Diego, CA

John T. Lohr, Ph.D.
*Assistant Director, Biotechnology
Center*
Utah State University
Logan, UT

Larry Lutwick, M.D., F.A.C.P.
Director, Infectious Diseases
VA Medical Center
Brooklyn, NY

Suzanne M. Lutwick
Medical Writer
Brooklyn, NY

Nicole Mallory, M.S.
Medical Student
Wayne State University
Detroit, MI

Warren Maltzman, Ph.D.
*Consultant, Molecular
Pathology*
Demarest, NJ

Adrienne Massel, R.N.
Medical Writer
Beloit, WI

Ruth E. Mawyer, R.N.
Medical Writer
Charlottesville, VA

Richard A. McCartney M.D.
*Fellow, American College of
Surgeons*
*Diplomat American Board of
Surgery*
Richland, WA

Bonny McClain, Ph.D.
Medical Writer
Greensboro, NC

Sally C. McFarlane-Parrott
Medical Writer
Ann Arbor, MI

Mercedes McLaughlin
Medical Writer
Phoenixville, CA

Alison McTavish, M.Sc.
Medical Writer and Editor
Montreal, Quebec

Liz Meszaros
Medical Writer
Lakewood, OH

Betty Mishkin
Medical Writer
Skokie, IL

Barbara J. Mitchell
Medical Writer
Hallstead, PA

Mark A. Mitchell, M.D.
Medical Writer
Seattle, WA

Susan J. Montgomery
Medical Writer
Milwaukee, WI

Louann W. Murray, PhD
Medical Writer
Huntington Beach, CA

Bilal Nasser, M.Sc.
Senior Medical Student
Universidad Iberoamericana
Santo Domingo, Dominican
Republic

Laura Ninger
Medical Writer
Weehawken, NJ

Nancy J. Nordenson
Medical Writer
Minneapolis, MN

Teresa Odle
Medical Writer
Albuquerque, NM

Lisa Papp, R.N.
Medical Writer
Cherry Hill, NJ

Lee Ann Paradise
Medical Writer
San Antonio, TX

Patience Paradox
Medical Writer
Bainbridge Island, WA

Barbara J. Pettersen
Genetic Counselor
Genetic Counseling of Central
Oregon
Bend, OR

Genevieve Pham-Kanter, M.S.
Medical Writer
Chicago, IL

Collette Placek
Medical Writer
Wheaton, IL

J. Ricker Polsdorfer, M.D.
Medical Writer
Phoenix, AZ

Scott Polzin, M.S., C.G.C.
Medical Writer
Buffalo Grove, IL

Elizabeth J. Pulcini, M.S.
Medical Writer
Phoenix, Arizona

Nada Quercia, M.S., C.C.G.C.
Genetic Counselor
Division of Clinical and
Metabolic Genetics
The Hospital for Sick Children
Toronto, ON, Canada

Ann Quigley
Medical Writer
New York, NY

Robert Ramirez, B.S.
Medical Student
University of Medicine &
Dentistry of New Jersey
Stratford, NJ

Kulbir Rangi, D.O.
Medical Doctor and Writer
New York, NY

Esther Csapo Rastegari, Ed.M.,
R.N./B.S.N.
Registered Nurse, Medical Writer
Holbrook, MA

Toni Rizzo
Medical Writer
Salt Lake City, UT

Martha Robbins
Medical Writer
Evanston, IL

Richard Robinson
Medical Writer
Tucson, AZ

Nancy Ross-Flanigan
Science Writer
Belleville, MI

Anna Rovid Spickler, D.V.M., Ph.D.
Medical Writer
Moorehead, KY

Belinda Rowland, Ph.D.
Medical Writer
Voorheesville, NY

Andrea Ruskin, M.D.
Whittingham Cancer Center
Norwalk, CT

Laura Ruth, Ph.D.
*Medical, Science, & Technology
Writer*
Los Angeles, CA

Karen Sandrick
Medical Writer
Chicago, IL

Kausalya Santhanam, Ph.D.
Technical Writer
Branford, CT

Jason S. Schliesser, D.C.
Chiropractor
Holland Chiropractic, Inc.
Holland, OH

Joan Schonbeck
Medical Writer
Nursing
Massachusetts Department of
Mental Health
Marlborough, MA

Laurie Heron Seaver, M.D.
Clinical Geneticist
Greenwood Genetic Center
Greenwood, SC

Catherine Seeley
Medical Writer

Kristen Mahoney Shannon, M.S.,
C.G.C.

Genetic Counselor

Center for Cancer Risk Analysis
Massachusetts General Hospital
Boston, MA

Kim A. Sharp, M.Ln.

Writer

Richmond, TX

Judith Sims, M.S.

Medical Writer

Logan, UT

Joyce S. Siok, R.N.

Medical Writer

South Windsor, CT

Jennifer Sisk

Medical Writer

Havertown, PA

Patricia Skinner

Medical Writer

Amman, Jordan

Genevieve Slomski, Ph.D.

Medical Writer

New Britain, CT

Stephanie Slon

Medical Writer

Portland, OR

Linda Wasmer Smith

Medical Writer

Albuquerque, NM

Java O. Solis, M.S.

Medical Writer

Decatur, GA

Elaine Souder, PhD

Medical Writer

Little Rock, AR

Jane E. Spehar

Medical Writer

Canton, OH

Lorraine Steefel, R.N.

Medical Writer

Morganville, NJ

Kurt Sternlof

Science Writer

New Rochelle, NY

Roger E. Stevenson, M.D.

Director

Greenwood Genetic Center

Greenwood, SC

Dorothy Stonely

Medical Writer

Los Gatos, CA

Liz Swain

Medical Writer

San Diego, CA

Deanna M. Swartout-Corbeil,

R.N.

Medical Writer

Thompsons Station, TN

Keith Tatarelli, J.D.

Medical Writer

Mary Jane Tenerelli, M.S.

Medical Writer

East Northport, NY

Catherine L. Tesla, M.S., C.G.C.

Senior Associate, Faculty

Dept. of Pediatrics, Division of

Medical Genetics

Emory University School of

Medicine

Atlanta, GA

Bethany Thivierge

Biotechnical Writer/Editor

Technicality Resources

Rockland, ME

Mai Tran, Pharm.D.

Medical Writer

Troy, MI

Carol Turkington

Medical Writer

Lancaster, PA

Judith Turner, B.S.

Medical Writer

Sandy, UT

Amy B. Tuteur, M.D.

Medical Advisor

Sharon, MA

Samuel Uretsky, Pharm.D.

Medical Writer

Wantagh, NY

Amy Vance, M.S., C.G.C.

Genetic Counselor

GeneSage, Inc.

San Francisco, CA

Michael Sherwin Walston

Student Researcher

University of Arizona

Tucson, AZ

Ronald Watson, Ph.D.

Science Writer

Tucson, AZ

Ellen S. Weber, M.S.N.

Medical Writer

Fort Wayne, IN

Ken R. Wells

Freelance Writer

Laguna Hills, CA

Jennifer F. Wilson, M.S.

Science Writer

Haddonfield, NJ

Kathleen D. Wright, R.N.

Medical Writer

Delmar, DE

Jennifer Wurges

Medical Writer

Rochester Hills, MI

Mary Zoll, Ph.D.

Science Writer

Newton Center, MA

Jon Zonderman

Medical Writer

Orange, CA

Michael V. Zuck, Ph.D.

Medical Writer

Boulder, CO

A

Abdominal aorta ultrasound see **Abdominal ultrasound**

Abdominal aortic aneurysm see **Aortic aneurysm**

Abdominal hernia see **Hernia**

Abdominal thrust see **Heimlich maneuver**

Abdominal ultrasound

Definition

Ultrasound technology allows doctors to “see” inside a patient without resorting to surgery. A transmitter sends high frequency sound waves into the body, where they bounce off the different tissues and organs to produce a distinctive pattern of echoes. A receiver “hears” the returning echo pattern and forwards it to a computer, which translates the data into an image on a television screen. Because ultrasound can distinguish subtle variations between soft, fluid-filled tissues, it is particularly useful in providing diagnostic images of the abdomen. Ultrasound can also be used in treatment.

Purpose

The potential medical applications of ultrasound were first recognized in the 1940s as an outgrowth of the sonar technology developed to detect submarines during World War II. The first useful medical images were produced in the early 1950s, and, by 1965, ultrasound quality had improved to the point that it came into general medical use. Improvements in the technology, application, and interpretation of ultrasound continue. Its low cost, versatility, safety and speed have brought it into the top drawer of medical imaging techniques.

While **pelvic ultrasound** is widely known and commonly used for fetal monitoring during **pregnancy**, ultrasound is also routinely used for general abdominal imaging. It has great advantage over x-ray imaging technologies in that it does not damage tissues with ionizing radiation. Ultrasound is also generally far better than plain x rays at distinguishing the subtle variations of soft tissue structures, and can be used in any of several modes, depending on the need at hand.

As an imaging tool, abdominal ultrasound generally is warranted for patients afflicted with: chronic or acute abdominal **pain**; abdominal trauma; an obvious or suspected abdominal mass; symptoms of **liver disease**, pancreatic disease, **gallstones**, spleen disease, **kidney disease** and urinary blockage; or symptoms of an abdominal **aortic aneurysm**. Specifically:

- Abdominal pain. Whether acute or chronic, pain can signal a serious problem—from organ malfunction or injury to the presence of malignant growths. Ultrasound scanning can help doctors quickly sort through potential causes when presented with general or ambiguous symptoms. All of the major abdominal organs can be studied for signs of disease that appear as changes in size, shape and internal structure.
- Abdominal trauma. After a serious accident, such as a car crash or a fall, internal bleeding from injured abdominal organs is often the most serious threat to survival. Neither the injuries nor the bleeding are immediately apparent. Ultrasound is very useful as an initial scan when abdominal trauma is suspected, and it can be used to pinpoint the location, cause, and severity of hemorrhaging. In the case of puncture **wounds**, from a bullet for example, ultrasound can locate the foreign object and provide a preliminary survey of the damage. The easy portability and versatility of ultrasound technology has brought it into common emergency room use, and even into limited ambulance service.

- **Abdominal mass.** Abnormal growths—tumors, cysts, abscesses, scar tissue and accessory organs—can be located and tentatively identified with ultrasound. In particular, potentially malignant solid tumors can be distinguished from benign fluid-filled cysts and abscesses. Masses and malformations in any organ or part of the abdomen can be found.
- **Liver disease.** The types and underlying causes of liver disease are numerous, though jaundice tends to be a general symptom. Ultrasound can differentiate between many of the types and causes of liver malfunction, and is particularly good at identifying obstruction of the bile ducts and **cirrhosis**, which is characterized by abnormal fibrous growths and reduced blood flow.
- **Pancreatic disease.** Inflammation and malformation of the pancreas are readily identified by ultrasound, as are pancreatic stones (calculi), which can disrupt proper functioning.
- **Gallstones.** Gallstones cause more hospital admissions than any other digestive malady. These calculi can cause painful inflammation of the gallbladder and also obstruct the bile ducts that carry digestive enzymes from the gallbladder and liver to the intestines. Gallstones are readily identifiable with ultrasound.
- **Spleen disease.** The spleen is particularly prone to injury during abdominal trauma. It may also become painfully inflamed when beset with infection or **cancer**. These conditions also lend themselves well to ultrasonic inspection and diagnosis.
- **Kidney disease.** The kidneys are also prone to traumatic injury and are the organs most likely to form calculi, which can block the flow of urine and cause blood **poisoning** (uremia). A variety of diseases causing distinct changes in kidney morphology can also lead to complete kidney failure. Ultrasound imaging has proven extremely useful in diagnosing kidney disorders.
- **Abdominal aortic aneurysm.** This is a bulging weak spot in the abdominal aorta, which supplies blood directly from the heart to the entire lower body. These aneurysms are relatively common and increase in prevalence with age. A burst aortic aneurysm is imminently life-threatening. However, they can be readily identified and monitored with ultrasound before acute complications result.

Ultrasound technology can also be used for treatment purposes, most frequently as a visual aid during surgical procedures—such as guiding needle placement to drain fluid from a cyst, or to extract tumor cells for biopsy. Increasingly, direct therapeutic applications for ultrasound are being developed.

The direct therapeutic value of ultrasonic waves lies in their mechanical nature. They are shock waves, just like audible sound, and vibrate the materials through which they pass. These vibrations are mild, virtually unnoticeable at the frequencies and intensities used for imaging. Properly focused however, high-intensity ultrasound can be used to heat and physically agitate targeted tissues.

High-intensity ultrasound is used routinely to treat soft tissue injuries, such as **strains**, tears and associated scarring. The heating and agitation are believed to promote rapid healing through increased circulation. Strongly focused, high-intensity, high-frequency ultrasound can also be used to physically destroy certain types of tumors, as well as gallstones and other types of calculi. Developing new treatment applications for ultrasound is an active area of medical research.

Precautions

Properly performed, ultrasound imaging is virtually without risk or side effects. Some patients report feeling a slight **tingling** and/or warmth while being scanned, but most feel nothing at all. Ultrasound waves of appropriate frequency and intensity are not known to cause or aggravate any medical condition, though any woman who thinks she might be pregnant should raise the issue with her doctor before undergoing an abdominal ultrasound.

The value of ultrasound imaging as a medical tool, however, depends greatly on the quality of the equipment used and the skill of the medical personnel operating it. Improperly performed and/or interpreted, ultrasound can be worse than useless if it indicates that a problem exists where there is none, or fails to detect a significant condition. Basic ultrasound equipment is relatively inexpensive to obtain, and any doctor with the equipment can perform the procedure whether qualified or not. Patients should not hesitate to verify the credentials of technicians and doctors performing ultrasounds, as well as the quality of the equipment used and the benefits of the proposed procedure.

In cases where ultrasound is used as a treatment tool, patients should educate themselves about the proposed procedure with the help of their doctors—as is appropriate before any surgical procedure. Also, any abdominal ultrasound procedure, diagnostic or therapeutic, may be hampered by a patient's body type or other factors, such as the presence of excessive bowel gas (which is opaque to ultrasound). In particular, very obese people are often not good candidates for abdominal ultrasound.

KEY TERMS

Accessory organ—A lump of tissue adjacent to an organ that is similar to it, but which serves no important purpose, if functional at all. While not necessarily harmful, such organs can cause problems if they grow too large or become cancerous. In any case, their presence points to an underlying abnormality in the parent organ.

Benign—In medical usage, benign is the opposite of malignant. It describes an abnormal growth that is stable, treatable and generally not life-threatening.

Biopsy—The surgical removal and analysis of a tissue sample for diagnostic purposes. Usually, the term refers to the collection and analysis of tissue from a suspected tumor to establish malignancy.

Calculus—Any type of hard concretion (stone) in the body, but usually found in the gallbladder, pancreas and kidneys. They are formed by the accumulation of excess mineral salts and other organic material such as blood or mucous. Calculi (pl.) can cause problems by lodging in and obstructing the proper flow of fluids, such as bile to the intestines or urine to the bladder.

Cirrhosis—A chronic liver disease characterized by the invasion of connective tissue and the degeneration of proper functioning—jaundice is often an accompanying symptom. Causes of cirrhosis include alcoholism, metabolic diseases, syphilis and congestive heart disease.

Common bile duct—The branching passage through which bile—a necessary digestive enzyme—travels from the liver and gallbladder into the small intestine. Digestive enzymes from the pancreas also enter the intestines through the common bile duct.

Computed tomography scan (CT scan)—A specialized type of x-ray imaging that uses highly focused and relatively low energy radiation to produce detailed two-dimensional images of soft tissue structures, particularly the brain. CT scans are the chief competitor to ultrasound and can yield higher quality images not disrupted by bone or gas. They are, however, more cumbersome, time consuming and expensive to perform, and they use ionizing electromagnetic radiation.

Doppler—The Doppler effect refers to the apparent change in frequency of sound wave echoes returning to a stationary source from a moving target. If the object is moving toward the source, the frequency increases; if the object is moving away, the frequency decreases. The size of this frequency shift can be used to compute the object's speed—be it a car on the road or blood in an artery. The Doppler effect holds true for all types of radiation, not just sound.

Frequency—Sound, whether traveling through air or the human body, produces vibrations—molecules bouncing into each other—as the shock wave travels along. The frequency of a sound is the number of vibrations per second. Within the audible range, frequency means pitch—the higher the frequency, the higher a sound's pitch.

Ionizing radiation—Radiation that can damage living tissue by disrupting and destroying individual cells at the molecular level. All types of nuclear radiation—x rays, gamma rays and beta rays—are potentially ionizing. Sound waves physically vibrate the material through which they pass, but do not ionize it.

Jaundice—A condition that results in a yellow tint to the skin, eyes and body fluids. Bile retention in the liver, gallbladder and pancreas is the immediate cause, but the underlying cause could be as simple as obstruction of the common bile duct by a gallstone or as serious as pancreatic cancer. Ultrasound can distinguish between these conditions.

Malignant—The term literally means growing worse and resisting treatment. It is used as a synonym for cancerous and connotes a harmful condition that generally is life-threatening.

Morphology—Literally, the study of form. In medicine, morphology refers to the size, shape and structure rather than the function of a given organ. As a diagnostic imaging technique, ultrasound facilitates the recognition of abnormal morphologies as symptoms of underlying conditions.

Description

Ultrasound includes all sound waves above the frequency of human hearing—about 20 thousand hertz, or cycles per second. Medical ultrasound generally uses frequencies between one and 10 million hertz

(1-10 MHz). Higher frequency ultrasound waves produce more detailed images, but are also more readily absorbed and so cannot penetrate as deeply into the body. Abdominal ultrasound imaging is generally performed at frequencies between 2-5 MHz.

An ultrasound machine consists of two parts: the transducer and the analyzer. The transducer both produces the sound waves that penetrate the body and receives the reflected echoes. Transducers are built around piezoelectric ceramic chips. (Piezoelectric refers to electricity that is produced when you put pressure on certain crystals such as quartz). These ceramic chips react to electric pulses by producing sound waves (they are transmitting waves) and react to sound waves by producing electric pulses (receiving). Bursts of high frequency electric pulses supplied to the transducer causes it to produce the scanning sound waves. The transducer then receives the returning echoes, translates them back into electric pulses and sends them to the analyzer—a computer that organizes the data into an image on a television screen.

Because sound waves travel through all the body's tissues at nearly the same speed—about 3,400 miles per hour—the microseconds it takes for each echo to be received can be plotted on the screen as a distance into the body. The relative strength of each echo, a function of the specific tissue or organ boundary that produced it, can be plotted as a point of varying brightness. In this way, the echoes are translated into a picture. Tissues surrounded by bone or filled with gas (the stomach, intestines and bowel) cannot be imaged using ultrasound, because the waves are blocked or become randomly scattered.

Four different modes of ultrasound are used in medical imaging:

- **A-mode.** This is the simplest type of ultrasound in which a single transducer scans a line through the body with the echoes plotted on screen as a function of depth. This method is used to measure distances within the body and the size of internal organs. Therapeutic ultrasound aimed at a specific tumor or calculus is also A-mode, to allow for pinpoint accurate focus of the destructive wave energy.
- **B-mode.** In B-mode ultrasound, a linear array of transducers simultaneously scans a plane through the body that can be viewed as a two-dimensional image on screen. Ultrasound probes containing more than 100 transducers in sequence form the basis for these most commonly used scanners, which cost about \$50,000.
- **M-Mode.** The M stands for motion. A rapid sequence of B-mode scans whose images follow each other in sequence on screen enables doctors to see and measure range of motion, as the organ boundaries that produce reflections move relative to the probe. M-mode ultrasound has been put to particular use in studying heart motion.
- **Doppler mode.** **Doppler ultrasonography** includes the capability of accurately measuring velocities of moving material, such as blood in arteries and veins. The principle is the same as that used in radar guns that measure the speed of a car on the highway. Doppler capability is most often combined with B-mode scanning to produce images of blood vessels from which blood flow can be directly measured. This technique is used extensively to investigate valve defects, arteriosclerosis and hypertension, particularly in the heart, but also in the abdominal aorta and the portal vein of the liver. These machines cost about \$250,000.

The actual procedure for a patient undergoing an abdominal ultrasound is relatively simple, regardless of the type of scan or its purpose. **Fasting** for at least eight hours prior to the procedure ensures that the stomach is empty and as small as possible, and that the intestines and bowels are relatively inactive. Fasting also allows the gall bladder to be seen, as it contracts after eating and may not be seen if the stomach is full. In some cases, a full bladder helps to push intestinal folds out of the way so that the gas they contain does not disrupt the image. The patient's abdomen is then greased with a special gel that allows the ultrasound probe to glide easily across the skin while transmitting and receiving ultrasonic pulses.

This procedure is conducted by a doctor with the assistance of a technologist skilled in operating the equipment. The probe is moved around the abdomen to obtain different views of the target areas. The patient will likely be asked to change positions from side to side and to hold their breath as necessary to obtain the desired views. Discomfort during the procedure is minimal.

The many types and uses of ultrasound technology makes it difficult to generalize about the time and costs involved. Relatively simple imaging—scanning a suspicious abdominal mass or a suspected abdominal aortic aneurysm—will take about half an hour to perform and will cost a few hundred dollars or more, depending on the quality of the equipment, the operator and other factors. More involved techniques such as multiple M-mode and Doppler-enhanced scans, or cases where the targets not well defined in advance, generally take more time and are more expensive.

Regardless of the type of scan used and the potential difficulties encountered, ultrasound remains faster and less expensive than computed tomography scans (CT), its primary rival in abdominal imaging. Furthermore, as abdominal ultrasounds are generally undertaken as

“medically necessary” procedures designed to detect the presence of suspected abnormalities, they are covered under most types of major medical insurance. As always, though, the patient would be wise to confirm that their coverage extends to the specific procedure proposed. For nonemergency situations, most underwriters stipulate prior approval as a condition of coverage.

Specific conditions for which ultrasound may be selected as a treatment option—certain types of tumors, lesions, **kidney stones** and other calculi, muscle and ligament injuries, etc.—are described in detail under the appropriate entries in this encyclopedia.

Preparation

A patient undergoing abdominal ultrasound will be advised by their physician about what to expect and how to prepare. As mentioned above, preparations generally include fasting and arriving for the procedure with a full bladder, if necessary. This preparation is particularly useful if the gallbladder, ovaries or veins are to be examined.

Aftercare

In general, no aftercare related to the abdominal ultrasound procedure itself is required.

Risks

Abdominal ultrasound carries with it no recognized risks or side effects, if properly performed using appropriate frequency and intensity ranges. Sensitive tissues, particularly those of the reproductive organs, could possibly sustain damage if violently vibrated by overly intense ultrasound waves. In general though, such damage would only result from improper use of the equipment.

Any woman who thinks she might be pregnant should raise this issue with her doctor before undergoing an abdominal ultrasound, as a fetus in the early stages of development could be injured by ultrasound meant to probe deeply recessed abdominal organs.

Normal results

As a diagnostic imaging technique, a normal abdominal ultrasound is one that indicates the absence of the suspected condition that prompted the scan. For example, symptoms such as a persistent **cough**, labored breathing, and upper abdominal pain suggest the possibility of, among other things, an abdominal aortic aneurysm. An ultrasound scan that indicates the absence of an aneurysm would rule out this life-

threatening condition and point to other, less serious causes.

Abnormal results

Because abdominal ultrasound imaging is generally undertaken to confirm a suspected condition, the results of a scan often will prove abnormal—that is they will confirm the diagnosis, be it kidney stones, cirrhosis of the liver or an aortic aneurysm. At that point, appropriate medical treatment as prescribed by a patient’s doctor is in order. See the relevant disease and disorder entries in this encyclopedia for more information.

Resources

PERIODICALS

Freundlich, Naomi. “Ultrasound: What’s Wrong with this Picture?” *Business Week* September 15, 1997:84-5.

ORGANIZATIONS

American College of Gastroenterology. 4900 B South 31st St., Arlington, VA 22206-1656. (703) 820-7400. <<http://www.acg.gi.org>> .

American Institute of Ultrasound in Medicine. 14750 Sweitzer Lane, Suite 100, Laurel, MD 20707-5906. (800) 638-5352. <<http://www.aium.org>> .

American Society of Radiologic Technologists. 15000 Central Ave., SE, Albuquerque, NM 87123-3917. (505) 298-4500. <<http://www.asrt.org>> .

Kurt Richard Sternlof

Abdominal wall defects

Definition

Abdominal wall defects are birth (congenital) defects that allow the stomach or intestines to protrude.

Description

Many unexpected and fascinating events occur during the development of a fetus inside the womb. The stomach and intestines begin development outside the baby’s abdomen and only later does the abdominal wall enclose them. Occasionally, either the umbilical opening is too large, or it develops improperly, allowing the bowels or stomach to remain outside or squeeze through the abdominal wall.

Causes and symptoms

There are many causes for **birth defects** that still remain unclear. Presently, the cause(s) of abdominal wall defects is unknown, and any symptoms the mother may have to indicate that the defects are present in the fetus are nondescript.

Diagnosis

At birth, the problem is obvious, because the base of the umbilical cord at the navel will bulge or, in worse cases, contain viscera (internal organs). Before birth, an ultrasound examination may detect the problem. It is always necessary in children with one birth defect to look for others, because birth defects are usually multiple.

Treatment

Abdominal wall defects are effectively treated with surgical repair. Unless there are accompanying anomalies, the surgical procedure is not overly complicated. The organs are normal, just misplaced. However, if the defect is large, it may be difficult to fit all the viscera into the small abdominal cavity.

Prognosis

If there are no other defects, the prognosis after surgical repair of this condition is relatively good. However, 10% of those with more severe or additional abnormalities die from it. The organs themselves are fully functional; the difficulty lies in fitting them inside the abdomen. The condition is, in fact, a **hernia** requiring only replacement and strengthening of the passageway through which it occurred. After surgery, increased pressure in the stretched abdomen can compromise the function of the organs inside.

Prevention

Some, but by no means all, birth defects are preventable by early and attentive prenatal care, good **nutrition**, supplemental **vitamins**, diligent avoidance of all unnecessary drugs and chemicals—especially tobacco—and other elements of a healthy lifestyle.

Resources

PERIODICALS

Dunn, J. C., and E. W. Fonkalsrud. "Improved Survival of Infants with Omphalocele." *American Journal of Surgery* 173 (April 1997): 284-7.

J. Ricker Polsdorfer, MD

KEY TERMS

Hernia—Movement of a structure into a place it does not belong.

Umbilical—Referring to the opening in the abdominal wall where the blood vessels from the placenta enter.

Viscera—Any of the body's organs located in the chest or abdomen.

Abnormal heart rhythms see **Arrhythmias**

ABO blood typing see **Blood typing and crossmatching**

ABO incompatibility see **Erythroblastosis fetalis**

Abortion, habitual see **Recurrent miscarriage**

Abortion, partial birth

Definition

Partial birth abortion is a method of late-term (after 20 weeks) abortion that terminates a **pregnancy** and results in the **death** and intact removal of a fetus. This procedure is most commonly referred to as intact dilatation and extraction (D & X). It occurs in a rare percentage of pregnancies.

Purpose

Partial birth abortion, or D&X, is performed to end a pregnancy and results in the death of a fetus, typically in the late second or third trimester. Although D&X is highly controversial, some physicians argue that it has advantages that make it a preferable procedure in some circumstances. One perceived advantage is that the fetus is removed largely intact, allowing for better evaluation and **autopsy** of the fetus in cases of known fetal abnormalities. Intact removal of the fetus also may carry a lower risk of puncturing the uterus or damaging the cervix. Another perceived advantage is that D&X ends the pregnancy without requiring the woman to go through labor, which may be less emotionally traumatic than other methods of late-term abortion. In

addition, D&X may offer a lower cost and shorter procedure time.

Precautions

Women considering D&X should be aware of the highly controversial nature of this procedure. A controversy common to all late-term abortions is whether the fetus is viable, or able to survive outside of the woman's body. A specific area of controversy with D&X is that fetal death does not occur until after most of the fetal body has exited the uterus. Several states have taken legal action to limit or ban D&X and many physicians who perform abortions do not perform D&X. This may restrict the availability of this procedure to women seeking late-term abortions.

In March 2003, the United States Senate passed a bill banning partial birth abortions and implementing fines or maximum two-year jail terms for physicians who perform them. In June 2003, the House approved a ban as well. President George W. Bush signed the legislation into law, but a federal judge declared the law unconstitutional, so that the government had not been able to enforce it. One of the opponents' claims was the legislation did not provide for exceptions for cases in which the procedure was needed to protect the mother's health.

Description

Intact D&X, or partial birth abortion first involves administration of medications to cause the cervix to dilate, usually over the course of several days. Next, the physician rotates the fetus to a footling breech position. The body of the fetus is then drawn out of the uterus feet first, until only the head remains inside the uterus. Then, the physician uses an instrument to puncture the base of the skull, which collapses the fetal head. Typically, the contents of the fetal head are then partially suctioned out, which results in the death of the fetus and reduces the size of the fetal head enough to allow it to pass through the cervix. The dead and otherwise intact fetus is then removed from the woman's body.

Preparation

Medical preparation for D&X involves an outpatient visit to administer medications, such as *laminaria*, to cause the cervix to begin dilating.

In addition, preparation may involve fulfilling local legal requirements, such as a mandatory waiting period, counseling, or an informed consent procedure

KEY TERMS

Cervix—The narrow outer end of the uterus that separates the uterus from the vaginal canal.

Footling breech—A position of the fetus while in the uterus where the feet of the fetus are nearest the cervix and will be the first part of the fetus to exit the uterus, with the head of the fetus being the last part to exit the uterus.

Laminaria—A medical product made from a certain type of seaweed that is physically placed near the cervix to cause it to dilate.

reviewing stages of fetal development, **childbirth**, alternative abortion methods, and adoption.

Aftercare

D&X typically does not require an overnight hospital stay, so a follow up appointment may be scheduled to monitor the woman for any complications.

Risks

With all abortion, the later in pregnancy an abortion is performed, the more complicated the procedure and the greater the risk of injury to the woman. In addition to associated emotional reactions, D&X carries the risk of injury to the woman, including heavy bleeding, **blood clots**, damage to the cervix or uterus, pelvic infection, and anesthesia-related complications. There also is a risk of incomplete abortion, meaning that the fetus is not dead when removed from the woman's body. Possible long-term risks include difficulty becoming pregnant or carrying a future pregnancy to term.

Normal results

The expected outcome of D&X is the termination of a pregnancy with removal of a dead fetus from the woman's body.

Resources

PERIODICALS

"Court Rules Abortion Ban Unconstitutional." *Medicine & Health* (June 7, 2004): 4–6.

"House Approves Partial Birth Abortion Ban." *Medicine and Health* (June 16, 2003): 5.

"Partial-birth Abortion Ban Approved by Senate." *Medical Ethics Advisor* (April 2003): 47.

ORGANIZATIONS

Planned Parenthood Federation of America, Inc., 810
Seventh Ave., New York, NY 10019. (800) 669-0156.
< <http://www.plannedparenthood.org> > .

OTHER

Status of partial-birth abortion laws in the states. Othmer
Institute at Planned Parenthood of NYC. 2000.

Stefanie B. N. Dugan, M.S.
Teresa G. Odle

Abortion, selective

Definition

Selective abortion, also known as selective reduction, refers to choosing to abort a fetus, typically in a multi-fetal **pregnancy**, to decrease the health risks to the mother in carrying and giving birth to more than one or two babies, and also to decrease the risk of complications to the remaining fetus(es). The term selective abortion also refers to choosing to abort a fetus for reasons such as the woman is carrying a fetus which likely will be born with some birth defect or impairment, or because the sex of the fetus is not preferred by the individual.

Purpose

A woman may decide to abort for health reasons, for example, she is at higher risk for complications during pregnancy because of a disorder or disease such as diabetes. A 2004 case reported on an embryo embedded in a **cesarean section** scar. Although rare, it can be life threatening to the mother. In this case, selective abortion was successful at saving the mother and the remaining embryos.

However, selective reduction is recommended often in cases of multi-fetal pregnancy, or the presence of more than one fetus, typically, at least three or more fetuses. In the general population, multi-fetal pregnancy happens in only about 1-2% of pregnant women. But multi-fetal pregnancies occur far more often in women using fertility drugs.

Precautions

Because women or couples who use fertility drugs have made an extra effort to become pregnant, it is possible that the individuals may be unwilling or uncomfortable with the decision to abort a fetus in

KEY TERMS

Multi-fetal pregnancy—A pregnancy of two or more fetuses.

Selective reduction—Typically referred to in cases of multifetal pregnancy, when one or more fetuses are aborted to preserve the viability of the remaining fetuses and decrease health risks to the mother.

cases of multi-fetal pregnancy. Individuals engaging in fertility treatment should be made aware of the risk of multi-fetal pregnancy and consider the prospect of recommended reduction before undergoing fertility treatment.

Description

Selective reduction is usually performed between nine and 12 weeks of pregnancy and is most successful when performed in early pregnancy. It is a simple procedure and can be performed on an outpatient basis. A needle is inserted into the woman's stomach or vagina and potassium chloride is injected into the fetus.

Preparation

Individuals who have chosen selective reduction to safeguard the remaining fetuses should be counseled prior to the procedure. Individuals should receive information regarding the risks of a multi-fetal pregnancy to both the fetuses and the mother compared with the risks after the reduction.

Individuals seeking an abortion for any reason should consider the ethical implications whether it be because the fetus is not the preferred sex or because the fetus would be born with a severe birth defect.

Aftercare

Counseling should continue after the abortion because it is a traumatic event. Individuals may feel guilty about choosing one fetus over another. Mental health professionals should be consulted throughout the process.

Risks

About 75% of women who undergo selective reduction will go into **premature labor**. About 4-5% of women undergoing selective reduction also

miscarry one or more of the remaining fetuses. The risks associated with multi-fetal pregnancy are considered higher.

Normal results

In cases where a multi-fetal pregnancy of three or more fetuses is reduced to two fetuses, the remaining twin fetuses typically develop as they would if they had been conceived as twins.

Resources

PERIODICALS

“Multiple Pregnancy Associated With Infertility Therapy.” *American Society for Reproductive Medicine, A Practice Committee Report* (November 2000): 1-8.

“Selective Reduction Eliminates an Embryo Embedded in a Cesarean Scar.” *Women’s Health Weekly* (April 8, 2004): 117.

ORGANIZATIONS

The Alan Guttmacher Institute. 120 Wall Street, New York, NY 10005. (212) 248-1111. <<http://www.agi-usa.org>>.

The American Society for Reproductive Medicine. 1209 Montgomery Highway, Birmingham, AL 35216-2809. (205) 978-5000. <<http://www.asrm.org>>.

Meghan M. Gourley
Teresa G. Odle

Abortion, spontaneous see **Miscarriage**

Abortion, therapeutic

Definition

Therapeutic abortion is the intentional termination of a **pregnancy** before the fetus can live independently. Abortion has been a legal procedure in the United States since 1973.

Purpose

An abortion may be performed whenever there is some compelling reason to end a pregnancy. Women have abortions because continuing the pregnancy would cause them hardship, endanger their life or health, or because prenatal testing has shown that the fetus will be born with severe abnormalities.

Abortions are safest when performed within the first six to 10 weeks after the last menstrual period. The calculation of this date is referred to as the

gestational age and is used in determining the stage of pregnancy. For example, a woman who is two weeks late having her period is said to be six weeks pregnant, because it is six weeks since she last menstruated.

About 90% of women who have abortions do so before 13 weeks and experience few complications. Abortions performed between 13-24 weeks have a higher rate of complications. Abortions after 24 weeks are extremely rare and are usually limited to situations where the life of the mother is in danger.

Precautions

Most women are able to have abortions at clinics or outpatient facilities if the procedure is performed early in pregnancy. Women who have stable diabetes, controlled epilepsy, mild to moderate high blood pressure, or who are HIV positive can often have abortions as outpatients if precautions are taken. Women with heart disease, previous **endocarditis**, **asthma**, lupus erythematosus, uterine fibroid tumors, blood clotting disorders, poorly controlled epilepsy, or some psychological disorders usually need to be hospitalized in order to receive special monitoring and medications during the procedure.


Description

Very early abortions

Between five and seven weeks, a pregnancy can be ended by a procedure called menstrual extraction. This procedure is also sometimes called menstrual regulation, mini-suction, or preemptive abortion. The contents of the uterus are suctioned out through a thin (3-4 mm) plastic tube that is inserted through the undilated cervix. Suction is applied either by a bulb syringe or a small pump.

Another method is called the “morning after” pill, or **emergency contraception**. Basically, it involves taking high doses of birth control pills within 24 to 48 hours of having unprotected sex. The high doses of hormones causes the uterine lining to change so that it will not support a pregnancy. Thus, if the egg has been fertilized, it is simply expelled from the body.

There are two types of emergency **contraception**. One type is identical to ordinary birth control pills, and uses the hormones estrogen and progestin. This type is available with a prescription under the brand name Preven. But women can even use their regular birth control pills for emergency contraception, after they check with their doctor about the proper dose. About half of women who use birth control pills for



Between 5 and 7 weeks, a pregnancy can be ended by a procedure called menstrual extraction. The contents of the uterus are suctioned out through a thin extraction tube that is inserted through the undilated cervix. (Illustration by Electronic Illustrators Group.)

KEY TERMS

Endocarditis—An infection of the inner membrane lining of the heart.

Fibroid tumors—Fibroid tumors are non-cancerous (benign) growths in the uterus. They occur in 30-40% of women over age 40, and do not need to be removed unless they are causing symptoms that interfere with a woman's normal activities.

Lupus erythematosus—A chronic inflammatory disease in which inappropriate immune system reactions cause abnormalities in the blood vessels and connective tissue.

Prostaglandin—Oxygenated unsaturated cyclic fatty acids responsible for various hormonal reactions such as muscle contraction.

Rh negative—Lacking the Rh factor, genetically determined antigens in red blood cells that produce immune responses. If an Rh negative woman is pregnant with an Rh positive fetus, her body will produce antibodies against the fetus's blood, causing a disease known as Rh disease. Sensitization to the disease occurs when the woman's blood is exposed to the fetus's blood. Rh immune globulin (RhoGAM) is a vaccine that must be given to a woman after an abortion, miscarriage, or prenatal tests in order to prevent sensitization to Rh disease.

emergency contraception get nauseated and 20 percent vomit. This method cuts the risk of pregnancy 75 percent.

The other type of morning-after pill contains only one hormone: progestin, and is available under the brand name Plan B. It is more effective than the first type with a lower risk of **nausea** and vomiting. It reduces the risk of pregnancy 89 percent.

Women should check with their physicians regarding the proper dose of pills to take, as it depends on the brand of birth control pill. Not all birth control pills will work for emergency contraception.

Menstrual extractions are safe, but because the amount of fetal material is so small at this stage of development, it is easy to miss. This results in an incomplete abortion that means the pregnancy continues.

First trimester abortions

The first trimester of pregnancy includes the first 13 weeks after the last menstrual period. In the United States, about 90% of abortions are performed during this period. It is the safest time in which to have an abortion, and the time in which women have the most choice of how the procedure is performed.

MEDICAL ABORTIONS. Medical abortions are brought about by taking medications that end the pregnancy. The advantages of a first trimester medical abortion are:

- The procedure is non-invasive; no surgical instruments are used.
- Anesthesia is not required.
- Drugs are administered either orally or by injection.
- The procedure resembles a natural **miscarriage**.

Disadvantages of a medical abortion are:

- The effectiveness decreases after the seventh week.
- The procedure may require multiple visits to the doctor.
- Bleeding after the abortion lasts longer than after a surgical abortion.
- The woman may see the contents of her womb as it is expelled.

Two different medications can be used to bring about an abortion. Methotrexate (Rheumatrex) works by stopping fetal cells from dividing which causes the fetus to die.

On the first visit to the doctor, the woman receives an injection of methotrexate. On the second visit, about a week later, she is given misoprostol (Cytotec), an oxygenated unsaturated cyclic fatty acid responsible for various hormonal reactions such as muscle contraction (prostaglandin), that stimulates contractions of the uterus. Within two weeks, the woman will expel the contents of her uterus, ending the pregnancy. A follow-up visit to the doctor is necessary to assure that the abortion is complete.

With this procedure, a woman will feel cramping and may feel nauseated from the misoprostol. This combination of drugs is 90-96% effective in ending pregnancy.

Mifepristone (RU-486), which goes by the brand name Mifeprex, works by blocking the action of progesterone, a hormone needed for pregnancy to continue, then stimulates uterine contractions thus ending the pregnancy. It can be taken as much as 49 days after the first day of a woman's last period. On the first visit to the doctor, a woman takes a mifepristone pill. Two days later she returns and, if the miscarriage has not occurred, takes two misoprostol pills, which causes the uterus to contract. Five percent of women won't need to take misoprostol. After an observation period, she returns home.

Within four days, 90% of women have expelled the contents of their uterus and completed the abortion. Within 14 days, 95-97% of women have completed the abortion. A third follow-up visit to the doctor is necessary to confirm through observation or ultrasound that the procedure is complete. In the event that it is not, a surgical abortion is performed.

Studies show that 4.5 to 8 percent of women need surgery or a blood **transfusion** after taking mifepristone, and the pregnancy persists in about 1 percent of women. In this case, surgical abortion is recommended because the fetus may be damaged. Side effects include nausea, vaginal bleeding and heavy cramping. The bleeding is typically heavier than a normal period and may last up to 16 days.

Mifepristone is not recommended for women with ectopic pregnancy, an **IUD**, who have been taking long-term steroidal therapy, have bleeding abnormalities or on blood-thinners such as Coumadin.

Surgical abortions

First trimester surgical abortions are performed using vacuum aspiration. The procedure is also called dilation and evacuation (D & E), suction dilation, vacuum curettage, or suction curettage.

Advantages of a vacuum aspiration abortion are:

- It is usually done as a one-day outpatient procedure.
- The procedure takes only 10-15 minutes.
- Bleeding after the abortion lasts five days or less.
- The woman does not see the products of her womb being removed.

Disadvantages include:

- The procedure is invasive; surgical instruments are used.
- Infection may occur.

During a vacuum aspiration, the woman's cervix is gradually dilated by expanding rods inserted into the cervical opening. Once dilated, a tube attached to a suction pump is inserted through the cervix and the contents of the uterus are suctioned out. The procedure is 97-99% effective. The amount of discomfort a woman feels varies considerably. **Local anesthesia** is often given to numb the cervix, but it does not mask uterine cramping. After a few hours of rest, the woman may return home.

Second trimester abortions

Although it is better to have an abortion during the first trimester, some second trimester abortions may be inevitable. The results of **genetic testing** are often not available until 16 weeks. In addition, women, especially teens, may not have recognized the pregnancy or come to terms with it emotionally soon enough to have a first trimester abortion. Teens make up the largest group having second trimester abortions.

Some second trimester abortions are performed as a D & E. The procedures are similar to those used in the first trimester, but a larger suction tube must be used because more material must be removed. This increases the amount of cervical dilation necessary and increases the risk of the procedure. Many physicians are reluctant to perform a D & E this late in pregnancy, and for some women it is not a medically safe option.

The alternative to a D & E in the second trimester is an abortion by induced labor. Induced labor may require an overnight stay in a hospital. The day before the procedure, the woman visits the doctor for tests, and to either have rods inserted in her cervix to help dilate it or to receive medication that will soften the cervix and speed up labor.

On the day of the abortion, drugs, usually prostaglandins to induce contractions, and a salt water solution, are injected into the uterus. Contractions begin, and within eight to 72 hours the woman delivers the fetus.

Side effects of this procedure include nausea, **vomiting**, and **diarrhea** from the prostaglandins, and **pain** from uterine cramps. Anesthesia of the sort used in **childbirth** can be given to mask the pain. Many women are able to go home a few hours after the procedure.

Very early abortions cost between \$200-\$400. Later abortions cost more. The cost increases about \$100 per week between the thirteenth and sixteenth week. Second trimester abortions are much more costly because they often involve more risk, more services, anesthesia, and sometimes a hospital stay. Insurance carriers and HMOs may or may not cover the procedure. Federal law prohibits federal funds including Medicaid funds, from being used to pay for an elective abortion.

Preparation

The doctor must know accurately the stage of a woman's pregnancy before an abortion is performed. The doctor will ask the woman questions about her menstrual cycle and also do a physical examination to confirm the stage of pregnancy. This may be done at an office visit before the abortion or on the day of the abortion. Some states require a waiting period before an abortion can be performed. Others require parental or court consent for a child under age 18 to receive an abortion.

Despite the fact that almost half of all women in the United States have had at least one abortion by the time they reach age 45, abortion is surrounded by controversy. Women often find themselves in

emotional turmoil when deciding if an abortion is a procedure they wish to undergo. Pre-abortion counseling is important in helping a woman resolve any questions she may have about having the procedure.

Aftercare

Regardless of the method used to perform the abortion, a woman will be observed for a period of time to make sure her blood pressure is stable and that bleeding is controlled. The doctor may prescribe **antibiotics** to reduce the chance of infection. Women who are Rh negative (lacking genetically determined antigens in their red blood cells that produce immune responses) should be given a human Rh immune globulin (RhoGAM) after the procedure unless the father of the fetus is also Rh negative. This prevents blood incompatibility complications in future pregnancies.

Bleeding will continue for about five days in a surgical abortion and longer in a medical abortion. To decrease the risk of infection, a woman should avoid intercourse and not use tampons and douches for two weeks after the abortion.

A follow-up visit is a necessary part of the woman's aftercare. Contraception will be offered to women who wish to avoid future pregnancies, because menstrual periods normally resume within a few weeks.

Risks

Serious complications resulting from abortions performed before 13 weeks are rare. Of the 90% of women who have abortions in this time period, 2.5% have minor complications that can be handled without hospitalization. Less than 0.5% have complications that require a hospital stay. The rate of complications increases as the pregnancy progresses.

Complications from abortions can include:

- uncontrolled bleeding
- infection
- blood clots accumulating in the uterus
- a tear in the cervix or uterus
- missed abortion where the pregnancy continues
- incomplete abortion where some material from the pregnancy remains in the uterus

Women who experience any of the following symptoms of post-abortion complications should call the clinic or doctor who performed the abortion immediately.

- severe pain
- fever over 100.4 °F (38.2 °C)
- heavy bleeding that soaks through more than one sanitary pad per hour
- foul-smelling discharge from the vagina
- continuing symptoms of pregnancy

Normal results

Usually the pregnancy is ended without complication and without altering future fertility.

Resources

BOOKS

Carlson, Karen J., Stephanie A. Eisenstat, and Terra Ziporyn. "Abortion." In *The Harvard Guide to Women's Health*. Cambridge, MA: Harvard University Press, 1996.

Debra Gordon

Abrasions see **Wounds**

Abruptio placentae see **Placental abruption**


Abscess

Definition

An abscess is an enclosed collection of liquefied tissue, known as pus, somewhere in the body. It is the result of the body's defensive reaction to foreign material.

Description

There are two types of abscesses, septic and sterile. Most abscesses are septic, which means that they are the result of an infection. Septic abscesses can occur anywhere in the body. Only a germ and the body's immune response are required. In response to the invading germ, white blood cells gather at the infected site and begin producing chemicals called enzymes that attack the germ by digesting it. These enzymes act like acid, killing the germs and breaking them down into small pieces that can be picked up by the circulation and eliminated from the body. Unfortunately, these chemicals also digest body tissues. In most cases, the germ produces similar chemicals. The result is a thick, yellow liquid—pus—



An amoebic abscess caused by *Entamoeba histolytica*. (Phototake NYC. Reproduced by permission.)

containing digested germs, digested tissue, white blood cells, and enzymes.

An abscess is the last stage of a tissue infection that begins with a process called inflammation. Initially, as the invading germ activates the body's immune system, several events occur:

- Blood flow to the area increases.
- The temperature of the area increases due to the increased blood supply.
- The area swells due to the accumulation of water, blood, and other liquids.
- It turns red.
- It hurts, because of the irritation from the swelling and the chemical activity.

These four signs—heat, swelling, redness, and pain—characterize inflammation.

As the process progresses, the tissue begins to turn to liquid, and an abscess forms. It is the nature of an abscess to spread as the chemical digestion liquefies more and more tissue. Furthermore, the spreading follows the path of least resistance—the tissues most easily digested. A good example is an abscess just beneath the skin. It most easily continues along beneath the skin rather than working its way through the skin where it could drain its toxic contents. The contents of the abscess also leak into the general circulation and produce symptoms just like any other infection. These include chills, **fever**, aching, and general discomfort.

Sterile abscesses are sometimes a milder form of the same process caused not by germs but by non-living irritants such as drugs. If an injected drug like penicillin is not absorbed, it stays where it was

KEY TERMS

Cellulitis—Inflammation of tissue due to infection.

Enzyme—Any of a number of protein chemicals that can change other chemicals.

Fallopian tubes—Part of the internal female anatomy that carries eggs from the ovaries to the uterus.

Flora—Living inhabitants of a region or area.

Pyogenic—Capable of generating pus. *Streptococcus*, *Staphylococcus*, and bowel bacteria are the primary pyogenic organisms.

Sebaceous glands—Tiny structures in the skin that produce oil (sebum). If they become plugged, sebum collects inside and forms a nurturing place for germs to grow.

Septicemia—The spread of an infectious agent throughout the body by means of the blood stream.

Sinus—A tubular channel connecting one body part with another or with the outside.

injected and may cause enough irritation to generate a sterile abscess—sterile because there is no infection involved. Sterile abscesses are quite likely to turn into hard, solid lumps as they scar, rather than remaining pockets of pus.

Causes and symptoms

Many different agents cause abscesses. The most common are the pus-forming (pyogenic) bacteria like *Staphylococcus aureus*, which is nearly always the cause of abscesses under the skin. Abscesses near the large bowel, particularly around the anus, may be caused by any of the numerous bacteria found within the large bowel. Brain abscesses and liver abscesses can be caused by any organism that can travel there through the circulation. Bacteria, amoeba, and certain fungi can travel in this fashion. Abscesses in other parts of the body are caused by organisms that normally inhabit nearby structures or that infect them. Some common causes of specific abscesses are:

- skin abscesses by normal skin flora
- dental and throat abscesses by mouth flora
- lung abscesses by normal airway flora, **pneumonia** germs, or tuberculosis
- abdominal and anal abscesses by normal bowel flora

Specific types of abscesses

Listed below are some of the more common and important abscesses.

- Carbuncles and other **boils**. Skin oil glands (sebaceous glands) on the back or the back of the neck are the ones usually infected. The most common germ involved is *Staphylococcus aureus*. **Acne** is a similar condition of sebaceous glands on the face and back.
- Pilonidal abscess. Many people have as a birth defect a tiny opening in the skin just above the anus. Fecal bacteria can enter this opening, causing an infection and subsequent abscess.
- Retropharyngeal, parapharyngeal, peritonsillar abscess. As a result of throat infections like strep throat and **tonsillitis**, bacteria can invade the deeper tissues of the throat and cause an abscess. These abscesses can compromise swallowing and even breathing.
- Lung abscess. During or after pneumonia, whether it's due to bacteria [common pneumonia], tuberculosis, fungi, parasites, or other germs, abscesses can develop as a complication.
- Liver abscess. Bacteria or amoeba from the intestines can spread through the blood to the liver and cause abscesses.
- Psoas abscess. Deep in the back of the abdomen on either side of the lumbar spine lie the psoas muscles. They flex the hips. An abscess can develop in one of these muscles, usually when it spreads from the appendix, the large bowel, or the fallopian tubes.

Diagnosis

The common findings of inflammation—heat, redness, swelling, and pain—easily identify superficial abscesses. Abscesses in other places may produce only generalized symptoms such as fever and discomfort. If the patient's symptoms and physical examination do not help, a physician may have to resort to a battery of tests to locate the site of an abscess, but usually something in the initial evaluation directs the search. Recent or chronic disease in an organ suggests it may be the site of an abscess. Dysfunction of an organ or system—for instance, seizures or altered bowel function—may provide the clue. **Pain** and tenderness on **physical examination** are common findings. Sometimes a deep abscess will eat a small channel (sinus) to the surface and begin leaking pus. A sterile abscess may cause only a painful lump deep in the buttock where a shot was given.

Treatment

Since skin is very resistant to the spread of infection, it acts as a barrier, often keeping the toxic chemicals of an abscess from escaping the body on their own. Thus, the pus must be drained from the abscess by a physician. The surgeon determines when the abscess is ready for drainage and opens a path to the outside, allowing the pus to escape. Ordinarily, the body handles the remaining infection, sometimes with the help of **antibiotics** or other drugs. The surgeon may leave a drain (a piece of cloth or rubber) in the abscess cavity to prevent it from closing before all the pus has drained out.

Alternative treatment

If an abscess is directly beneath the skin, it will be slowly working its way through the skin as it is more rapidly working its way elsewhere. Since chemicals work faster at higher temperatures, applications of hot compresses to the skin over the abscess will hasten the digestion of the skin and eventually result in its breaking down, releasing the pus spontaneously. This treatment is best reserved for smaller abscesses in relatively less dangerous areas of the body—limbs, trunk, back of the neck. It is also useful for all superficial abscesses in their very early stages. It will “ripen” them.

Contrast **hydrotherapy**, alternating hot and cold compresses, can also help assist the body in resorption of the abscess. There are two homeopathic remedies that work to rebalance the body in relation to abscess formation, *Silica* and *Hepar sulphuris*. In cases of septic abscesses, bentonite clay packs (bentonite clay and a small amount of *Hydrastis* powder) can be used to draw the infection from the area.

Prognosis

Once the abscess is properly drained, the prognosis is excellent for the condition itself. The reason for the abscess (other diseases the patient has) will determine the overall outcome. If, on the other hand, the abscess ruptures into neighboring areas or permits the infectious agent to spill into the bloodstream, serious or fatal consequences are likely. Abscesses in and around the nasal sinuses, face, ears, and scalp may work their way into the brain. Abscesses within an abdominal organ such as the liver may rupture into the abdominal cavity. In either

case, the result is life threatening. Blood **poisoning** is a term commonly used to describe an infection that has spilled into the blood stream and spread throughout the body from a localized origin. Blood poisoning, known to physicians as septicemia, is also life threatening.

Of special note, abscesses in the hand are more serious than they might appear. Due to the intricate structure and the overriding importance of the hand, any hand infection must be treated promptly and competently.

Prevention

Infections that are treated early with heat (if superficial) or antibiotics will often resolve without the formation of an abscess. It is even better to avoid infections altogether by taking prompt care of open injuries, particularly puncture **wounds**. Bites are the most dangerous of all, even more so because they often occur on the hand.

Resources

BOOKS

Fauci, Anthony S., et al., editors. *Harrison's Principles of Internal Medicine*. New York: McGraw-Hill, 1997.

J. Ricker Polsdorfer, MD

Abscess drainage see **Abscess incision and drainage**


Abscess incision & drainage

Definition

An infected skin nodule that contains pus may need to be drained via a cut if it does not respond to **antibiotics**. This allows the pus to escape, and the infection to heal.

Purpose

An **abscess** is a pus-filled sore, usually caused by a bacterial infection. The pus is made up of both live and dead organisms and destroyed tissue from the white blood cells that were carried to the area to fight the infection. Abscesses are often found in the soft tissue under the skin, such as the armpit or the groin.



Although abscesses are often found in the soft tissue under the skin, such as the armpit or the groin, they may develop in any organ, such as the liver. (Illustration by Electronic Illustrators Group.)

However, they may develop in any organ, and are commonly found in the breast and gums. Abscesses are far more serious and call for more specific treatment if they are located in deep organs such as the lung, liver or brain.

Because the lining of the abscess cavity tends to interfere with the amount of the drug that can penetrate the source of infection from the blood, the cavity itself may require draining. Once an abscess has fully formed, it often does not respond to antibiotics. Even if the antibiotic does penetrate into the abscess, it doesn't function as well in that environment.

Precautions

An abscess can usually be diagnosed visually, although an imaging technique such as a computed tomography scan may be used to confirm the extent of the abscess before drainage. Such procedures may also be needed to localize internal abscesses, such as those in the abdominal cavity or brain.

KEY TERMS

White blood cells—Cells that protect the body against infection.

Description

A doctor will cut into the lining of the abscess, allowing the pus to escape either through a drainage tube or by leaving the cavity open to the skin. How big the incision is depends on how quickly the pus is encountered.

Once the abscess is opened, the doctor will clean and irrigate the wound thoroughly with saline. If it is not too large or deep, the doctor may simply pack the abscess wound with gauze for 24–48 hours to absorb the pus and discharge.

If it is a deeper abscess, the doctor may insert a drainage tube after cleaning out the wound. Once the

tube is in place, the surgeon closes the incision with simple stitches, and applies a sterile dressing. Drainage is maintained for several days to help prevent the abscess from reforming.

Preparation

The skin over the abscess will be cleansed by swabbing gently with an antiseptic solution.

Aftercare

Much of the **pain** around the abscess will be gone after the surgery. Healing is usually very fast. After the tube is taken out, antibiotics may be continued for several days. Applying heat and keeping the affected area elevated may help relieve inflammation.

Risks

If there is any scarring, it is likely to become much less noticeable as time goes on, and eventually almost invisible. Occasionally, an abscess within a vital organ (such as the brain) damages enough surrounding tissue that there is some permanent loss of normal function.

Normal results

Most abscesses heal after drainage alone; others require drainage and antibiotic drug treatment.

Resources

BOOKS

Turkington, Carol A., and Jeffrey S. Dover. *Skin Deep*. New York: Facts on File, 1998.

ORGANIZATIONS

National Institute of Arthritis and Musculoskeletal and Skin Diseases. 9000 Rockville Pike, Bldg. 31, Rm 9A04, Bethesda, MD 20892.

Carol A. Turkington

Description

Physical abuse of a child is the infliction of injury by another person. The injuries can include punching, kicking, biting, burning, beating, or pulling the victim's hair. The physical abuse inflicted on a child can result in **bruises**, **burns**, **poisoning**, broken bones, and internal hemorrhages. Physical assault against an adult primarily occurs with women, usually in the form of domestic violence. It is estimated that approximately three million children witness domestic violence every year.

Sexual abuse of a child refers to sexual behavior between an adult and child or between two children, one of whom is dominant or significantly older. The sexual behaviors can include touching breasts, genitals, and buttocks; either dressed or undressed. The behavior also can include exhibitionism, cunnilingus, fellatio, or penetration of the vagina or anus with sexual organs or objects.

Pornographic photography also is used in sexual abuse with children. Reported sex offenders are 97% male. Reports of child pornography have increased since with the popularity of the Internet. Females more often are perpetrators in child-care settings, since children may confuse sexual abuse by a female with normal hygiene care. The 1990s and early 2000s were rocked by reports of sexual abuse of children committed by Catholic priests. Most of the abuse appeared to have occurred during the 1970s and a prominent report released early in 2004 stated that as many as 10,667 children were sexually abused by more than 4,300 priests. Sexual abuse by stepfathers is five times more common than with biological fathers. Sexual abuse of daughters by stepfathers or fathers is the most common form of incest.

Sexual abuse also can take the form of **rape**. The legal definition of rape includes only slight penile penetration in the victim's outer vulva area. Complete erection and ejaculation are not necessary. Rape is the perpetration of an act of sexual intercourse whether:

- will is overcome by force or fear (from threats or by use of drugs).
- mental impairment renders the victim incapable of rational judgment.
- if the victim is below the legal age established for consent.

Substance abuse is an abnormal pattern of substance usage leading to significant distress or impairment. The criteria include one or more of the following occurring within a 12-month period:

Abuse

Definition

Abuse is defined as anything that is harmful, injurious, or offensive. Abuse also includes excessive and wrongful misuse of a substance. There are several major types of abuse: physical and sexual abuse of a child or an adult, **substance abuse**, elderly abuse, and emotional abuse.

KEY TERMS

Encopresis—Abnormalities relating to bowel movements that can occur as a result of stress or fear.

- recurrent substance use resulting in failure to fulfill obligations at home, work, or school.
- using substance in situations that are physically dangerous (i.e., while driving).
- recurrent substance-related legal problems.
- continued usage despite recurrent social and interpersonal problems (i.e., arguments and fights with significant other).

Abuse of the elderly is common and occurs mostly as a result of caretaker burnout, due to the high level of dependency frail, elderly patients usually require. Abuse can be manifested by physical signs, fear, and delaying or not reporting the need for advanced medical care. Elderly patients also may exhibit financial abuse (money or possessions taken away) and abandonment.

Emotional abuse generally continues even after physical assaults have stopped. In most cases it is a personally tailored form of verbal or gesture abuse expressed to illicit a provoked response.

Causes and symptoms

Children who have been abused usually have a variety of symptoms that encompass behavioral, emotional, and psychosomatic problems (body problems caused by emotional or mental disturbance). Children who have been physically abused tend to be more aggressive, angry, hostile, depressed, and have low self-esteem. Additionally, they exhibit fear, **anxiety**, and nightmares. Severe psychological problems may result in suicidal behavior or posttraumatic stress disorder. Physically abused children may complain of physical illness even in the absence of a cause. They also may suffer from eating disorders and **encopresis**, or involuntary defecation caused or psychic origin. Children who are sexually abused may exhibit abnormal sexual behavior in the form of aggressiveness and hyperarousal. Adolescents may display promiscuity, sexual acting out, and—in some situations—homosexual contact.

Physical abuse directed towards adults can ultimately lead to **death**. Approximately 50% of women

murdered in the United States were killed by a former or current male partner. Approximately one-third of emergency room consultations by women were prompted due to domestic violence. Female victims who are married also have a higher rate of internal injuries and unconsciousness than victims of stranger assault (mugging, robbery). Physical abuse or rape also can occur between married persons and persons of the same gender. Perpetrators usually sexually assault their victims to dominate, hurt, and debase them. It is common for physical and sexual violence to occur at the same time. A large percentage of sexually assaulted persons were also physically abused in the form of punching, beating, or threatening the victim with a weapon such as a gun or knife. Usually males who are hurt and humiliated tend to physically assault people whom they are intimately involved with, such as spouses and/or children. Males who assault a female tend to have experienced or witnessed violence during childhood. They also tend to abuse alcohol, to be sexually assaultive, and are at increased risk for assaultive behavior directed against children. Jealous males tend to monitor a women's movements and whereabouts and to isolate other sources of protection and support. They interpret their behavior as betrayal of trust and this causes resentment and explosive anger outbursts during periods of losing control. Males also may use aggression against females in an effort to control and intimidate partners.

Abuse in the elderly usually occurs in the frail, elderly community. The caretaker is usually the perpetrator. Caretaker abuse can be suspected if there is evidence suggesting behavioral changes in the elderly person when the caretaker is present. Additionally, elderly abuse can be possible if there are delays between injuries and treatment, inconsistencies between injury and explanations, lack of hygiene or clothing, and prescriptions not being filled.

Diagnosis

Children who are victims of domestic violence frequently are injured attempting to protect their mother from an abusive partner. Injuries are visible by inspection or self-report. Physical abuse of an adult may also be evident by inspection with visible cuts and/or bruises or self-report.

Sexual abuse of both a child and an adult can be diagnosed with a history from the victim. Victims can be assessed for signs of ejaculatory evidence from the perpetrator. Ejaculatory specimens can be retrieved from the mouth, rectum, and clothing. Tests for sexually transmitted diseases may be performed.

Elderly abuse can be suspected if the elderly patient demonstrates a fear of the caretaker. Additionally, elderly abuse can be suspected if there are signs indicating intentional delay of required medical care or a change in medical status.

Substance abuse can be suspected in a person who continues to indulge in their drug of choice despite recurrent negative consequences. The diagnosis can be made after administration of a comprehensive exam and standardized chemical abuse assessments by a therapist.

Treatment

Children who are victims of physical or sexual abuse typically require psychological support and medical attention. A complaint may be filed with the local family social services agency that will initiate investigations. The authorities usually will follow up the allegation or offense. Children may also be referred for psychological evaluation and/or treatment. The victim also may be placed in foster care pending the investigation outcome. The police also may investigate physical and sexual abuse of an adult. The victim may require immediate medical care and long-term psychological treatment. It is common for children to be adversely affected by domestic violence situations and the local family services agency may be involved.

Substance abusers should elect treatment, either inpatient or outpatient, depending on severity of **addiction**. Long term treatment and/or medications may be utilized to assist in abstinence. The patient should be encouraged to participate in community centered support groups.

Prognosis

The prognosis depends on the diagnosis. Usually victims of physical and sexual abuse require therapy to deal with emotional distress associated with the incident. Perpetrators require further psychological evaluation and treatment. Victims of abuse may have a variety of emotional problems including depression, acts of **suicide**, or anxiety. Children of sexual abuse may enter abusive relationships or have problems with intimacy as adults. The substance abuser may experience relapses, since the cardinal feature of all addictive disorders is a tendency to return to symptoms. Elderly patients may suffer from further medical problems and/or anxiety, and in some cases neglect may precipitate death.

Prevention

Prevention programs are geared to education and awareness. Detection of initial symptoms or characteristic behaviors may assist in some situations. In some cases treatment may be sought before incident. The professional treating the abused persons must develop a clear sense of the relationship dynamics and the chances for continued harm.

Resources

BOOKS

Behrman, Richard E., et al, editors. *Nelson Textbook of Pediatrics*. 16th ed. W. B. Saunders Company, 2000.

PERIODICALS

Plante, Thomas G. "Another Aftershock: What Have We Learned from the John Jay Report?." *America* (March 22, 2004): 10.

ORGANIZATIONS

National Clearinghouse on Child Abuse and Neglect Information. 330 C Street SW, Washington, DC 20447. (800) 392-3366.

OTHER

Elder Abuse Prevention. < <http://www.oaktrees.org/elder> > .

National Institute on Drug Abuse. < <http://www.nida.nih.gov> > .

Laith Farid Gulli, M.D.
Bilal Nasser, M.Sc.
Teresa G. Odle

Acceleration-deceleration cervical injury see **Whiplash**

ACE inhibitors see **Angiotensin-converting enzyme inhibitors**

Acetaminophen

Definition

Acetaminophen is a medicine used to relieve **pain** and reduce **fever**.

Purpose

Acetaminophen is used to relieve many kinds of minor aches and pains—headaches, muscle aches, backaches, toothaches, menstrual cramps, arthritis, and the aches and pains that often accompany colds.

Description

This drug is available without a prescription. Acetaminophen is sold under various brand names, including Tylenol, Panadol, **Aspirin** Free Anacin, and Bayer Select Maximum Strength **Headache** Pain Relief Formula. Many multi-symptom cold, flu, and sinus medicines also contain acetaminophen. The ingredients listing on the container should state if acetaminophen is included in the product.

Studies have shown that acetaminophen relieves pain and reduces fever about as well as aspirin. But differences between these two common drugs exist. Acetaminophen is less likely than aspirin to irritate the stomach. However, unlike aspirin, acetaminophen does not reduce the redness, stiffness, or swelling that accompany arthritis.

Recommended dosage

The usual dosage for adults and children age 12 and over is 325-650 mg every four to six hours as needed. No more than 4 grams (4000 mg) should be taken in 24 hours. Because the drug can potentially harm the liver, people who drink alcohol in large quantities should take considerably less acetaminophen and possibly should avoid the drug completely.

For children ages 6-11 years, the usual dose is 150-300 mg, three to four times a day. A physician should recommend doses for children under age 6 years.

Precautions

In 2004, the U.S. Food and Drug Administration (FDA) launched an advertising campaign aimed at educating consumers about proper use of acetaminophen and other over-the-counter pain killers. Often, acetaminophen is hidden in many cold and flu products and people unexpectedly overdose on the medicine. Some cases have led to **liver transplantation** or **death**. More than the recommended dosage of acetaminophen should not be taken unless told to do so by a physician or dentist.

Patients should not use acetaminophen for more than 10 days to relieve pain (five days for children) or for more than three days to reduce fever, unless directed to do so by a physician. If symptoms do not go away—or if they get worse—a physician should be contacted. Anyone who drinks three or more alcoholic beverages a day should check with a physician before using this drug and should never take more than the recommended dosage. A risk of liver damage exists from combining large amounts of alcohol and

KEY TERMS

Arthritis—Inflammation of the joints. The condition causes pain and swelling.

Fatigue—Physical or mental weariness.

Inflammation—A response to irritation, infection, or injury, resulting in pain, redness, and swelling.

acetaminophen. People who already have kidney or **liver disease** or liver infections should also consult with a physician before using the drug. So should women who are pregnant or breastfeeding.

Many drugs can interact with one another. A physician or pharmacist should be consulted before combining acetaminophen with any other medicine. Two different acetaminophen-containing products should not be used at the same time.

Acetaminophen interferes with the results of some medical tests. Avoiding the drug for a few days before the tests may be necessary.

Side effects

Acetaminophen causes few side effects. The most common one is lightheadedness. Some people may experience trembling and pain in the side or the lower back. Allergic reactions occur in some people, but are rare. Anyone who develops symptoms such as a rash, swelling, or difficulty breathing after taking acetaminophen should stop taking the drug and get immediate medical attention. Other rare side effects include yellow skin or eyes, unusual bleeding or bruising, weakness, **fatigue**, bloody or black stools, bloody or cloudy urine, and a sudden decrease in the amount of urine.

Overdoses of acetaminophen may cause **nausea**, **vomiting**, sweating, and exhaustion. Very large overdoses can cause liver damage. In case of an overdose, immediate medical attention should be sought. In 2004, researchers announced that an injection to counteract the liver injury caused by acetaminophen overdose has been approved by the FDA.

Interactions

Acetaminophen may interact with a variety of other medicines. When this happens, the effects of one or both of the drugs may change or the risk of side effects may be greater. Among the drugs that may interact with acetaminophen are alcohol, nonsteroidal