Table S1. Description and experimental support of the model's Boolean regulatory logic.

\$1.A. Boolean regulatory logic of <u>Growth Signaling</u> nodes.

Target Node	Link	Input	Description & References	
GF_High	←	GF_High	The GFHigh node in our model represents an extracellular environment with saturating levels of growth factors; this input node is self-sustaining in the absence of in silico perturbation.	
GF	←	GF_High	The GF node represents an extracellular environment with low levels of growth factors capable of sustaining survival signaling. Thus, this node is ON in high growth factor as well.	
GF	←	GF	The GF input node is self-sustaining in the absence of in silico perturbation.	
GF	GF or	GF_High		
Growth Factor Si	gnalin	g - Upstream Pl3K	Ccycle	
RTK	F	CAD	Caspase-activated DNAase (CAD) inhibition of receptor tyrosine kinases ensures that apoptotic cells no longer maintain even basal levels of growth signaling.	
RTK	←	GF	The ON state of the RTK node in our model represents basal growth receptor activation by low / medium growth factor availability, encoded by the GF node (required to keep a normal cell alive).	
RTK	←	GF_High	Similarly, high growth factor availability also keeps RTK on.	
RTK	not CAD and (GF_High or GF)			
	(requ	The ON state of the RTK node in our model represents basal growth receptor activation (required to keep a normal cell alive). Thus it requires the absence of CAD and at least low growth levels of growth factors in the extracellular environment [1].		
Grb2	←	RTK	The SH2 domain of Grb2 binds to a phosphotyrosine residue in the activated RTK, where it functions as an adaptor protein [2].	
Grb2	←	GF_High	The ON state of Grb2 node in our model encode the change from basal (or absent) Grb2 activity at weakly stimulated RTKs to the level of recruitment seen in high growth factor environments, capable of mediating Ras activation.	
Grb2	RTK	and GF_High		
		Grb2 is ON when the RTK is activated by high levels of extracellular growth factors (capable of driving proliferation).		
sos	←	Grb2	RTK-bound Grb2 recruits SOS, a guanine nucleotide – exchange protein (GEF) that converts inactive Ras to its active GTP-bound form [2].	
Ras	←	Grb2	RTK-bound Grb2 is required to recruits SOS, the GEF responsible for converting inactive Ras to its GTP-bound active form [2].	
Ras	←	sos	SOS is a GEF that is recruited to activate Ras near ligand-bound, active RTKs [2].	

Target Node	Link	Input	Description & References
Ras	Grb2	and SOS	
	Ras activation requires the GEF activity of SOS and the RTK-linked (active) adaptor protein Grb2 [2].		
RAF	⊢	Casp3	Raf-1 is cleaved and inhibited by Caspase 3 [3].
RAF	←	Ras	Active Ras phosphorylates Raf, enhancing its kinase activity [4].
RAF	not C	asp3 and Ras	
	Raf is active in response to Ras activity in the absence of Caspase 3. As active Raf-1 is continuously dephosphorylated and bound by 14-3-3 which translocates it to the cytoplasm from the plasma membrane (not modeled explicitly), ongoing Ras activity is necessary to keep it ON [4].		
MEK	←	RAF	Raf phosphorylates and activates the MEK kinase [4].
ERK	←	MEK	MEK phosphorylates and activates the ERK kinase [4].
mTORC2	←	PIP3	PtdIns(3,4,5)P3 (PIP3), interacts with the mTORC2 component Sin1 to release its inhibition on the mTOR kinase domain. Thus, PIP3 is necessary for mTORC2 activation [5].
mTORC2	F	S6K	Rictor, a component of the mTORC2 complex, undergoes S6K1-mediated phosphorylation at T1135, dampening mTORC2-dependent phosphorylation of Akt [6]
mTORC2	PIP3 or not S6K		
	Our model assumes that mTORC2 is active in quiescent cells with basal levels of PI3K activity leading to basal PIP3 generation. Alternatively, the absence of high growth factor-stimulated mTORC1 and S6K1 can also increase mTORC2 activity.		
PI3K	←	Ras	Ras binds the catalytic subunit of PI3K and Ras knockdown / over expression decreases /increases the PI3K-dependent generation of PIP3 [7,8].
PI3K	←	RTK	Active RTKs recruit PI3K to the signaling complex they nucleate, where PI3K catalyzes the production of PtdIns(3,4,5)P3 (PIP3) [1].
РІЗК	Ras or RTK		
	In our model, basal PI3K activity can be maintained by active RTKs or active Ras, while high PI3K activity requires both (see below).		
PIP3	←	РІЗК	Active PI3K recruited to the membrane catalyzes the production of membrane-bound PtdIns(3,4,5)P3 (PIP3) from PtdIns(4,5)P2 (PIP2) [1].
PIP3	←	РІЗК_Н	Active PI3K recruited to the membrane catalyzes the production of membrane-bound PtdIns(3,4,5)P3 (PIP3) from PtdIns(4,5)P2 (PIP2) [1].
PIP3	PI3K	_H or PI3K	
	In our	model, PIP3 is ON	as a result of basal or high PI3K activity.

Target Node	Link	Input	Description & References
PDK1	←	РІЗК	The PDK1 kinase is recruited to the plasma membrane by PIP3 at the sites of active PI3K activity [9].
PDK1	←	PIP3	The PDK1 kinase is recruited to the plasma membrane by PIP3 at the sites of active PI3K activity [9].
PDK1	PI3K	and PIP3	
	PDK1	enzyme activation	requires active (at least basal) PI3K and PIP3 [9].
AKT_B	⊢	Casp3	AKT1 is cleaved and inhibited by Caspase 3 [3].
AKT_B	←	PIP3	PIP3 recruits AKT to the plasma membrane and PIP3 binding changes the conformation of AKT such that it becomes accessible for T308 phosphorylation by PDK1 [9].
AKT_B	←	PDK1	Membrane-recruited PDK1 phosphorylates AKT at T308, a critical step in its activation [9].
AKT_B	←	mTORC2	Maximal activation of AKT requires phosphorylation of S473 by mTORC2 [9].
AKT_B	not C	asp3 and PIP3 and	(PDK1 or mTORC2)
	Basal AKT1 activity in our model requires the absence of Caspase 3, the availability of at least basal levels of PIP3, and phosphorylation by PDK1 or mTORC2. In contrast, full mitogen-stimulated AKT1 activation requires phosphorylation by both (see AKT_H) [9].		
Growth Factor Signature	gnalin	g - PI3K cycle	
p110_H	←	FoxO3	FoxO3 is a direct inducer p110 α (PIK3CA), the catalytic subunit of PI3K [10].
p110_H	⊢	NeddL4	p110 α (PIK3CA) is polyubiquitinated by the E3 ligase NEDD4L, leading to its proteasomal degradation. Both free p110 and the regulatory subunit-bound protein is subject to ubiquitination by NEDD4L [11].
p110_H	←	p110_H	Our model assumes that maintaining high p110 levels is easier than driving the re-accumulation of the protein following its rapid destruction.
p110_H	(FoxC	03 and not NeddL4	or [p110_H and (FoxO3 or not NeddL4)]
	In order to capture the cyclic dynamics of p110 protein expression, we make the assumption that high p110 protein levels can be induced by FoxO3 in the absence of the growth factor-activated NeddL4 ubiquitin ligase. Once present, high p110 can be maintained by FoxO3 transcription, or the absence of activated NeddL4.		
PI3K_H	←	p110_H	High levels of PI3K activity in response to strong growth factor stimulation only occur in cells that express high levels of p110 protein [12].
PI3K_H	←	РІЗК	In our model, high PI3K activation is contingent on the ON-state of the basal PI3K node.
РІЗК_Н	←	RTK	High levels of PI3K activation only occur near at growth factor-bound RTKs, which recruit and activate PI3K at the plasma membrane [9].

Target Node	Link	Input	Description & References
PI3K_H	←	Ras	Ras binds the catalytic subunit of PI3K and Ras knockdown / over expression decreases /increases the PI3K-dependent generation of PIP3 [7,8].
РІЗК_Н	p110	_H and PI3K and R	TK and Ras
	Full, peak-level activation of PI3K requires high levels of p110 protein, basal PI3K activation, active RTKs, and active Ras. As the ON-state of Ras in our model represents strong Ras activation in the presence of proliferation-inducing (high) growth factors, PI3K_H activation can only occur in these conditions.		
AKT_H	←	AKT_B	In our model, high AKT activation is contingent on the ON-state of basal AKT (AKT_B).
AKT_H	←	p110_H	Ongoing high p110 availability and PI3K_H activity are required to induce maximal activation of AKT_H [9].
AKT_H	←	РІЗК_Н	Ongoing high p110 availability and PI3K_H activity are required to induce maximal activation of AKT_H [9].
AKT_H	←	PIP3	PIP3 recruits AKT to the plasma membrane and PIP3 binding changes the conformation of AKT such that it becomes accessible for T308 phosphorylation by PDK1 [9].
AKT_H	←	PDK1	Membrane-recruited PDK1 phosphorylates AKT at T308, a critical step in its activation [9].
AKT_H	←	mTORC2	Maximal activation of AKT requires phosphorylation of S473 by mTORC2 [9].
AKT_H	←	Ras	Ras binding to the catalytic subunit of PI3K is required for its full potency in PIP3 generation [7,8]. Active Ras is thus required for inducing peak AKT_H activity.
AKT_H	AKT_	B and p110_H and	PI3K_H and PIP3 and PDK1 and mTORC2 and Ras
	ongoi additi	ing presence of high	high AKT activity in our model requires basal AKT (AKT_B), the n p110 protein levels along with active PI3K_H, and PIP3. In T activation requires phosphorylation by both PDK1 and ve Ras [9].
FoxO3	⊢	AKT_H	AKT mediates the translation of FoxO3 our of the nucleus through direct phosphorylation of three conserved residues. These events create a recognition site for 14-3-3 family proteins, which export and sequester FoxO3 in the cytosol [9].
FoxO3	⊢	AKT_B	AKT mediates the translation of the FoxO3 transcription factor our of the nucleus (see above).
FoxO3	F	ERK	ERK downregulates FOXO3 transactional activity by phosphorylating it at three Serines, inducing its MDM2-mediated ubiquitination and degradation [13].
FoxO3	F	Plk1	Plk1 binds FoxO3, induces its translocation to the cytosol, phosphorylates it and suppresses its activity through most of the the cell cycle, but most significantly during G2 and M [14].
FoxO3	—	Plk1_H	Plk1 binds FoxO3, induces its translocation to the cytosol (see above).

Target Node	Link	Input	Description & References
FoxO3		AKT_B or AKT_H or not (Plk1 and Plk1_	r ERK) or { not [AKT_H and (Plk1 or Plk1_H or AKT_B or ERK)] H and ERK) }
	In order to account for all the influences on FoxO3 activity, we used the following logic. In the absence of basal or high AKT as well as ERK, FoxO3 remains active. In addition, FoxO3 can overcame peak (AKT_H) activation only if no other inhibitor is present and AKT_B is OFF (indicating that AKT levels are falling). Finally, the joint activity of ERK and Plk1 can also block FoxO3.		
PLCgamma	←	RTK	The SH2 domains of PLCγ binds to active RTKs at tyrosine autophosphorylation sites, leading to tyrosine phosphorylation of PLCγ and stimulation its enzymatic activity [15,16].
PLCgamma	←	Grb2	RTK tyrosine autophosphorylation induces PLC _γ binding to the Grb2 adaptor protein and likely aids the translocation of PLC _γ to the plasma membrane [17].
PLCgamma	←	p110_H	Membrane targeting of PLCγ to the sight of growth receptor stimulation requires Pl3K activity and PlP3 generation near growth receptors [18]. Thus, peak PLCγ activity in our model requires high p110 protein expression [19].
PLCgamma	←	PI3K_H	In addition to high p110 protein levels, high PI3K activation is also required to fully activate PLC γ [18,19].
PLCgamma	←	PIP3	Membrane targeting of PLC γ to the sight of growth receptor stimulation is mediated by PIP3 binding of PLC γ [18,19].
PLCgamma	RTK and Grb2 and p110_H and PI3K_H and PIP3		
			requires active an RTK receptor node bound by active Grb2, as r (including high p110 availability and the presence of PIP3).
IP3	←	PLCgamma	Membrane-bound, active PLCγ is responsible for converting phosphatidylinositol(4,5)P2 (PIP2) to the second messenger inositol(1,4,5)P3 (IP3) responsible for triggering a sudden Ca²+ influx from the endoplasmic reticulum, along with DAG (diacylglycerol, another second messenger) [20].
Ca2+	←	IP3	IP3 travels from the cell membrane to the endoplasmic reticulum where it opens IP3-sensitive Ca ²⁺ channels, releasing a sudden Ca ²⁺ efflux from the ER into the cytosol [21].
NeddL4	←	Ca2+	In order to transition to its active form, the E3 ubiquitin ligase NeddL4 binds Ca ²⁺ and inositol 1,4,5-trisphosphate (IP3) [22].
NeddL4	←	IP3	In order to transition to its active form, the E3 ubiquitin ligase NeddL4 binds Ca ²⁺ and inositol 1,4,5-trisphosphate (IP3) [22].
NeddL4	Ca2+ and IP3		
	Activation of NeddL4 requires both Ca2+ and IP3 binding [22].		
Growth Factor Si	gnalin	g - Downstream P	I3K cycle
FoxO1	⊢	AKT_H	AKT mediates the translation of FoxO1 our of the nucleus through direct phosphorylation of three conserved residues. These events create a recognition site for 14-3-3 family proteins, which export and sequester FoxO1 in the cytosol [9].

Target Node	Link	Input	Description & References
FoxO1	F	Plk1	PLK1 interacts with and phosphorylates FOXO1, mainly at the G 2/M phase of the cell cycle. PLK1-mediated phosphorylation leads to the impairment of FOXO1's transcriptional activity in an Akt-independent manner. By immunofluorescence staining and subcellular fractionation, we demonstrate that PLK1-induced FOXO1 phosphorylation causes its nuclear exclusion. [23]
FoxO1	not P	lk1 and not AKT_H	
	FoxO	1 is transcriptionally	active in the absence of peak AKT activation and Plk1 activity.
p21_mRNA	←	FoxO1	p21 ^{Cip1} is a direct transcriptional target of FoxO1 [24].
p21_mRNA	←	FoxO3	p21 ^{Cip1} is a direct transcriptional target of FoxO3 [24].
p21_mRNA	⊢	Мус	Myc is a direct transcriptional repressor of the p21 ^{Cip1} promoter (it is recruited by the DNA-binding Miz-1) [25,26].
p21_mRNA	(FoxC)1 and FoxO3) or [n	not Myc and (FoxO1 or FoxO3)]
	Our model requires both FoxOs to induce p21 ^{Cip1} if Myc is active and one of the two if Myc is OFF. This is based on data showing that both FoxO3 and FoxO1 bind and induce the p21 ^{Cip1} promoter and that loss of Myc repression alone is not sufficient to induce p21 ^{Cip1} [24].		
TSC2	F	AKT_H	TSC2 is phosphorylated by AKT1, inhibiting it by dissociating TSC2 from lysosomal membranes [27], where it stimulates GTP hydrolysis of the small GTPase Rheb, this inactivating it [28].
TSC2	F	AKT_B	TSC2 is phosphorylated by AKT1, inhibiting it by dissociating TSC2 from lysosomal membranes [27], where it stimulates GTP hydrolysis of the small GTPase Rheb, this inactivating it [28].
TSC2	F	ERK	ERK phosphorylates TSC2 directly, causing dissociation of the complex and inhibition of its activity [29]. In addition, the ERK target p90RSK can also inactivate TSC2 [30].
TSC2	not AKT_H or not (AKT_B or ERK)		
	Blocking TSC2 requires ongoing mitogen stimulation through AKT and/or ERK. In our model, TSC2 inhibition requires high (peak) AKT activity, supported by either ERK or basal AKT (assuring that complete loss of AKT activity is not impending) [28].		
PRAS40	F	AKT_H	PRAS40 is an inhibitory component of the mTORC1 complex. It is phosphorylated by AKT, triggering its dissociation from mTORC1 and loss of mTORC1 inhibition [31].
PRAS40	⊢	AKT_B	PRAS40 is phosphorylated by AKT, triggering its dissociation from mTORC1 [31].
PRAS40	F	mTORC1	PRAS40 is a substrate of the mTORC1 kinase; its phosphorylation aids its dissociation from mTORC1 and its sequestration by 14-3-3 proteins [32].
PRAS40	not A	KT_H and (not mT(DRC1 or not AKT_B)
	on its	way down), or ong	eak AKT activity aided by either basal AKT (meaning AKT_H is oing mTORC1 activation. Both Akt and mTORC1 phosphorylate association from mTORC1 [33].

Target Node	Link	Input	Description & References
DAG	←	PLCgamma	Membrane-bound, active PLCγ is responsible for converting phosphatidylinositol(4,5)P2 (PIP2) to the second messenger diacylglycerol (DAG), along with IP3 [20].
Rheb	F	TSC2	TSC2, a key component of the heterotrimeric TSC complex, is a GTPase activating protein (GAP) that induces ATP hydrolysis and deactivation of the small GTPase Rheb [34].
Rheb	←	DAG	The second messenger DAG activates both classical and novel PKCs. One of its targets, PKCη, is responsible for the translocation and accumulation of mTORC1 to perinuclear lysosomes, where the majority of Rheb is anchored. Thus, DAG brings Rheb in proximity with its target, mTORC1 [35].
Rheb	not T	SC2 and DAG	
	activi	ty (to prenuclear lys	ent activation of mTORC1 recruits mTORC1 to the site of Rheb cosomes), while AKT and ERK-mediated TSC2 inhibition mains potent [34-36].
mTORC1	F	Casp3	Raptor, a key component of the mTORC1 complex, is cleaved and inhibited by Caspase 3 [37].
mTORC1	←	Rheb	The Rheb small GTPase binds mTORC1 directly and activates the complex [38].
mTORC1	F	PRAS40	PRAS40 is an inhibitory component of the mTORC1 complex, removed by phosphorylation by AKT or mTORC1 itself [28].
mTORC1	←	E2F1	E2F1 induces mTORC1 activity by inducing mTORC1 translocation to late endosomes. This effect does not require AKT and is not blocked by high levels of TSC2 [39].
mTORC1	←	Cdk1	During mitosis, mTORC1 is activated by the G2/M-specific phosphorylation of Raptor, a component of mTORC1, by CyclinB / Cdk1 complexes, aided by GSK3 [40].
mTORC1	←	CyclinB	mTORC1 is activated by the G2/M-specific phosphorylation of Raptor, a component of mTORC1, by CyclinB / Cdk1 complexes, aided by GSK3 [40].
mTORC1	←	GSK3	mTORC1 is activated by the G2/M-specific phosphorylation of Raptor, a component of mTORC1, by CyclinB / Cdk1 complexes, aided by GSK3 [40].
mTORC1	not Casp3 and [(Rheb and not PRAS40) or E2F1 or (CyclinB and Cdk1 and GSK3)]		
	mTORC1 is activated by mitogenic signals via Rheb in the absence of PRAS40. Independently of this, E2F1 can promote mTORC1 activity. Finally, the mitotic Cyclin B and its Cdk1 kinase can also activate mTORC1, aided by GSK3.		
S6K	-	Casp3	S6K is cleaved and inhibited by Caspase 3 [41].
S6K	←	mTORC1	mTORC1 phosphorylates and activates 40S ribosomal S6 kinases (S6Ks) [42].
S6K	not Casp3 and mTORC1		
	S6K is activated by mTORC1 in the absence of Caspase 3.		

Target Node	Link	Input	Description & References
eIF4E	-	Casp3	eIF4E is cleaved and inhibited by Caspase 3 [43].
eIF4E	←	mTORC1	mTORC1 phosphorylates 4EBP, triggering its dissociation from eIF4F, and thus promoting translation initiation [28].
elF4E	not C	asp3 and mTORC	1
	elF4E	is activated by mT	ORC1-mediated repression in the absence of Caspase 3.
GSK3	F	AKT_H	AKT blocks GSK3 kinase activity via an inhibitory phosphorylation on the amino terminus, which blocks the substrate accessibility of GSK3 [9].
GSK3	F	ERK	ERK binds and phosphorylates GSK3 β at Thr-43, which primes it for subsequent phosphorylation by the ERK target p90RSK at Ser-9, which inactivates GSK3 β [44].
GSK3	-	S6K	GSK3 is a direct phosphorylation target of S6K1, resulting in its inhibition [45].
GSK3	not AKT_H and not (S6K and ERK)		
	GSK3 activity can be completely blocked by peak AKT activation (AKT_H), or by the joint action of S6K and ERK.		

$\textbf{S1.B.} \ \, \textbf{Boolean regulatory logic of} \ \, \underline{\textbf{Restriction Switch}} \ \, \textbf{nodes}.$

Target Node	Link	Input	Description & References	
p21	←	p21_mRNA	p21 protein activity requires the presence of p21 transcription and mRNA.	
p21	-	Casp3	Caspase 3 cleaves and deactivates p21Cip1 [46].	
p21	F	CyclinE	p21 and Cyclin E / Cdk2 form a positive (double-negative) feedback loop in which Cyclin E / Cdk2 activates the SCF/Skp2 complex responsible for the degradation of Cyclin E / Cdk2-bound, phosphorylated p21 [47]. p21, in turn, not only blocks Cyclin E / Cdk2 activity, but it also inhibits Cyclin D1. Thus, p21 interferes with the mitogen signal that turns on Cyclin E in the first place. In quiescent cells with high basal p21 levels, this positive feedback renders cell cycle entry stochastic [48].	
p21	p21_i	mRNA and not Cas	p3 and not CyclinE	
	p21 a	In this model, the p21 node corresponds to nuclear p21 in cells with relatively high basal p21 activity. p21 activity and or localization can be lowered by loss of FoxO mediated transcription (see p21_mRNA node) and via feedback from Cyclin E / Cdk2 [48].		
pRB	-	Casp3	Caspase 3 cleaves RB, generating fragments that do not associate with E2F, rendering RB inactive [49].	
pRB	-	CyclinD1	Cyclin D1 / Cdk4,6 complexes bind and phosphorylate RB, inhibiting its activity [50,51].	
pRB	F	CyclinA	Cyclin A / Cdk2 complexes phosphorylate and deactivate RB [52].	

Target Node	Link	Input	Description & References
pRB	←	p27Kip1	Active p27Kip1 can counteract the inhibitory effects of active CyclinE/Cdk2 complexes [53].
pRB	-	CyclinE	Cyclin E / Cdk2 complexes bind and phosphorylate RB, inhibiting its activity [51,54].
pRB	not C	asp3 and not Cycli	nD1 and not CyclinA and [p27Kip1 or not CyclinE]
			ence of Caspase 3, CyclinD1 and CyclinA. In addition, pRB en active p27Kip1 counteracts the effects of CyclinE [53].
p27Kip1	F	Casp3	Caspase 3 cleaves p27Kip1 [55]; the cleaved fragments can no longer associate with Cdk2 / Cyclin complexes [56].
p27Kip1	F	CyclinD1	Active Cyclin D1 / Cdk4,6 complexes competitively bind to p27Kip1 and progressively inhibit its ability to keep Cyclin E / Cdk2 inactive [53].
p27Kip1	⊢	Cdk1	Cyclin B / Cdk1 complexes phosphorylate p27Kip1 [57] and although they do not promote its degradation, phosphorylated p27Kip1 is exported from the nucleus and looses its ability to inhibit Cdk activity [58].
p27Kip1	F	CyclinB	Cyclin B / Cdk1 complexes phosphorylate p27Kip1 [57] and although they do not promote its degradation, phosphorylated p27Kip1 is exported from the nucleus and looses its ability to inhibit Cdk activity [58].
p27Kip1	F	CyclinE	Active Cyclin E / Cdk2 phosphorylate p27Kip1 [59], which marks it for degradation by the SCFSKP2 complex at the onset of S-phase [60]. (Cyclin- E/Cdk2 complexes can remain active in the presence of p27Kip1 when Cyclin A is also active).
p27Kip1	-	CyclinA	Cyclin A / Cdk2 complexes sequester and inactivate p27Kip1, phosphorylate it, and promote its degradation [57].
p27Kip1	←	FoxO1	FoxO factors are direct inducers of p27Kip1 expression. [61].
p27Kip1	←	FoxO3	FoxO factors are direct inducers of p27Kip1 expression. [61].
p27Kip1			nD1 and not (Cdk1 and CyclinB) and { [not CyclinE and (FoxO3 clinA and (FoxO3 or FoxO1)] or (not CyclinE and not CyclinA) }
	Active p27Kip1 is cleaved by Caspase 3 and inhibited (sequestered) by Cyclin D1/Cdk4,6 [53] or Cyclin B / Cdk1 [58]. In addition, maintenance of p27Kip1 requires one or both FoxO factors when sequestered by Cyclin E / Cdk2 (one FoxO factor) or Cyclin A / Cdk2 (both FoxO factors), but it cannot keep pace with the simultaneous activity of Cyclin E / Cdk2 and Cyclin A / Cdk2 [62].		
Мус	←	ERK	Ser-62 phosphorylation by ERK increases its half life, leading to Myc accumulation [63,64].
Мус	F	GSK3	Thr-58 phosphorylation by GSK-3 promotes Myc degradation [63,65]
Мус	←	elF4E	Increased translational initiation in the presence of activated eIF4E leads to an increase in Myc protein levels [66].

Target Node	Link	Input	Description & References
Мус	←	E2F1	E2F1 binds and activates the c-Myc promoter [67,68] [34, 33, 58, 59].
Мус	F	pRB	E2F1's ability to induce Myc is blocked by active (hypophosphorylated pRB) [69].
Мус	[ERK	and (eIF4E or not 0	GSK3)] or [E2F1 and not pRB and (eIF4E or ERK or not GSK3)]
	either phosi prom	r an increase in tran phorylation by GSK ote Myc accumulat	by stabilization of the protein via ERK phosphorylation, aided by slation initiated by eIF4E or loss of degradation-promoting 3 [63]. Alternatively, increased transcription by E2F1 can also ion in the absence of active pRB [70], provided that the protein is , or the absence of GSK3.
CyclinD1	⊢	CHK1	During replication, checkpoint kinases such as CHK1 (active during normal DNA synthesis) suppress Cyclin D1 [71], which has a very short half- life (~ 24 min) [72].
CyclinD1	F	p21	p21Cip1 is a Cyclin Dependent kinase inhibitor which binds to and blocks the activity of Cdk2, Cdk3, Cdk4 and Cdk6 kinases [73] and thus inhibits CyclinD1/Cdk4,6 [74].
CyclinD1	←	E2F1	The Cyclin D1 promoter is bound by E2F factors including E2F1; E2F1 over-expression can increase Cyclin D1 (though its effects are context-dependent, as E2F1 over-expression can also lead to apoptosis) [75]. Dominant negative E2F1 over-expression results in a 2-3 fold decrease in Cyclin D1 expression and Cyclin D1/Cdk4,6 activity [76].
CyclinD1	-	pRB	E2F1's ability to induce CyclinD1 is blocked by active (hypophosphorylated pRB) [77].
CyclinD1	←	Мус	Extracellular growth signals activate the MAPK pathway, leading to transcriptional activation of Cyclin D1 by Myc [78,79]. Myc overexpression leads to rapid Cyclin D1 induction and subsequent cell cycle entry [80], while its absence halves Cyclin D1 levels [81]. In addition, Myc induces Cdk4, aiding the assembly of active Cyclin D1 / Cdk4,6 complexes [81,82].
CyclinD1	F	GSK3	GSK-3β phosphorylates Cyclin D1 on Thr-286, promoting its ubiquitination and degradation [83].
CyclinD1	←	CyclinD1	In order to take into account both production and stability of Cyclin D1, we assumed that the presence of active CyclinD/Cdk2,4 complexes renders transcriptional maintenance of their levels easier.
CyclinD1	{Myc	and E2F1 } or { E2F	1 and ({not GSK3 and [Myc or E2F1]} or {Myc and CyclinD1} or 1 and CyclinD1})] or [not pRB and E2F1 and ({Myc and not GSK3})] }

Target Node	Link	Input	Description & References	
	Ongoing DNA synthesis keeps the CHK1 kinase active, which inhibits CyclinD1. The precise regulatory logic of Cyclin D1 as a function of transcriptional control by Myc and E2F, combined with the regulation of its protein stability / activity by GSK3/ basal p21 is not known. Here, we assume that in the absence of p21 (once p21 levels drop due to growth factor signals and/or Cdk2 activation), Cyclin D1 can be activated by either Myc or E2F1 in the absence of GSK3. In the presence of GSK3, we assume that Cyclin D1 can be induced by the combined action of both Myc and E2F1 [84], but sustained in an ON state by either. In the presence of basal (normal quiescent) levels of p21, we assume that Cyclin D1 transcription requires E2F1 unencumbered by pRB, as well as any two of the following: Myc, already active Cyclin D1, and no GSK3.			
E2F1	-	CAD	This link from Caspase-activated DNAase (CAD) to E2F1 ensures that apoptotic cells settle into an E2F1-negative attractor regardless of their initial state. The rationale for this is that E2F1 cannot maintain its activity if DNA is fragmented.	
E2F1	-	CyclinA	The phosphorylation of the E2F1-binding DP-1 protein by Cyclin A, which binds directly to E2F1 (as well as E2F-2,3) down-regulates E2F1 transcriptional activity in S phase [85,86].	
E2F1	⊢	pRB	RB binds to E2F/DP1 complexes and switches their DNA binding activity from an activating to a repressive form [77,87].	
E2F1	←	E2F1	E2F1 binds to its own promoter and up regulates transcription (as long as Cyclin D1/E activity blocks RB-E2F1 binding) [88].	
E2F1	←	Мус	Myc is required for growth-factor mediated induction of E2F1 [56, 47]. It binds to and remodels the E2F1 promoter, facilitating E2F1 transcription [89]. In addition, Myc augments protein expression of E2F1 [90]. Single-cell experiments show that Myc is a critical modulator of the amplitude of E2F activation [91].	
E2F1	not (not (CAD or CyclinA or pRB) and (E2F1 or Myc)		
		In the absence of both CyclinA and pRB, E2F1 transcription can be induced by Myc or maintained by active E2F1. CAD deactivates E2F1 as it destroys the cell's DNA.		
CyclinE	←	E2F1	E2F1 is a potent transcriptional activator of Cyclin E [92].	
CyclinE	←	Cdc6	Chromatin association and full activation of Cyclin E / Cdk2 requires Cdc6 [93].	
CyclinE	←	Pre-RC	At the G1/S transition, Cyclin E is loaded onto chromatin by pre-RC complexes (Cdc6 and Cdt1 binding), where it is required for MCM2 loading, origin firing and the start of DNA synthesis [94]. In addition, activation of its partner Cdk2 by Cdc6 is contingent on this localization [93].	
CyclinE	F	pRB	CyclinE transcription by E2F1 is inhibited by hypophosphorylated (active) pRB protein [95].	
CyclinE	F	p27Kip1	p27Kip1 binds to Cyclin E / Cdk2 complexes and prevents their activation [53].	
CyclinE	F	СНК1	Chk1 activation during normal S-phase progression keeps Cdk2 activity in a physiological range by binding to both Cdk2 and Cdc25A, aiding the loss of Cyclin E / Cdk1 activity [96].	

Target Node	Link	Input	Description & References	
CyclinE	-	Casp3	Caspase 3 cleaves and deactivates Cyclin E, which is then rapidly degraded [97].	
CyclinE	E2F1	E2F1 and Cdc6 and Pre-RC and not (pRB or p27Kip1 or CHK1 or Casp3)		
	In our model, the ON state of Cyclin E represents active Cyclin E / Cdk2 complexes. Thus, its full activation requires transcription via E2F1 not blocked by active pRB, binding to Cdc6 and pre-RC complexes, and the absence of its inhibitors p27Kip1, CHK1 and Caspase 3.			

$\textbf{S1.C.} \ \, \textbf{Boolean regulatory logic of } \underline{\textbf{Phase Switch}} \ \, \textbf{nodes}.$

Target Node	Link	Input	Description & References
CyclinA_mRNA	F	CAD	This link from Caspase-activated DNAase (CAD) to Cyclin A mRNA ensures that apoptotic cells settle into a G0-like attractor regardless of their initial state. The rationale for this is that no mRNA synthesis can be maintained if DNA is fragmented (we only use these links from CAD if needed).
CyclinA_mRNA	←	E2F1	Cyclin A is transcriptionally activated by E2F factors [95].
CyclinA_mRNA	Н	pRB	Active RB blocks E2F1's ability to transcribe Cyclin A [98].
CyclinA_mRNA	←	FoxM1	Depletion of FoxM1 results in reduced Cyclin A2 expression (it is not clear whether FoxM1 is a direct transcriptional inducer of Cyclin A) [99-102].
CyclinA_mRNA	not C	AD and [(E2F1 and	d not pRB) or FoxM1]
	In non-apoptotic cells Cyclin A is transcribed by E2F1 in the absence of active RB or by FoxM1.		
Emi1	←	E2F1	Emi1 is a direct transactional target of E2F1 [103,104].
Emi1	F	pRB	Active retinoblastoma protein can block Emi1 transcription mediated by E2F1 [104].
Emi1	F	p21	p21 activation during DNA damage lead to a substantial decrease of Emi1 levels, not observed in p21-null cells [105].
Emi1	F	Plk1	Plk1 phosphorylates Emi1 at mitotic spindle poles, stimulating its β TrCP binding and ubiquitination [106].
Emi1	-	CyclinB	Cyclin B / Cdk1 enhances the ability of Plk1 to mediate Emi1 destruction [106].
Emi1	-	Cdk1	Cyclin B / Cdk1 enhances the ability of Plk1 to mediate Emi1 destruction [106].
Emi1	F	U_Kinetochores	As Plk1-mediated phosphorylation of Emi1 occurs at mitotic spindle poles, our model requires ongoing mitosis for this interaction [106].
Emi1	F	A_Kinetochores	As Plk1-mediated phosphorylation of Emi1 occurs at mitotic spindle poles, our model requires ongoing mitosis for this interaction [106].

Target Node	Link	Input	Description & References	
Emi1		or not pRB or not netochores or A_K	p21) and not [Plk1 and CyclinB and Cdk1 and (inetochores)]	
	transo media	Our model allows the sustained presence of Emi1 protein when it is either actively transcribed by E2F1 or lacks joint inhibition by pRB and p21. Degradation of Emi1 is mediated by Plk1 and CyclinB/Cdk1 complexes. In addition, it requires at least temporary co-localization of Emi1 with Plk1 at mitotic spindle poles [106].		
FoxM1	←	Мус	FoxM1 is a direct transcriptional target of c-Myc [107].	
FoxM1	←	CyclinE	Cyclin E / Cdk2 complexes bind and phosphorylate FoxM1, potently inducing its transcriptional activity, which starts during S-phase [108].	
FoxM1	←	CyclinA	In addition to Cyclin E / Cdk2, Cyclin A / Cdk2 complexes can also keep FoxM1 transcriptionally active by phosphorylating its autoinhibitory N-terminal region [109].	
FoxM1	←	Cdc25A	Active Cdc25A enhances the transcriptional activity of FoxM1, likely via keeping a strong pool of active Cdk2 [110].	
FoxM1	←	Cdc25B	Cdc25B overexpression can increase FoxM1-dependent transcription, likely via keeping aiding Cdk1 activity [111].	
FoxM1	←	CyclinB	FoxM1 binds Plk1, and phosphorylation of two key residues in this domain by Cyclin B / Cdk1 primes it for Plk1 binding [112].	
FoxM1	←	Cdk1	FoxM1 binds Plk1, and phosphorylation of two key residues in this domain by Cyclin B / Cdk1 primes it for Plk1 binding [112].	
FoxM1	←	Plk1	Plk1 binds and phosphorylates FoxM1, which activates the FoxM1-mediated transcription in early mitosis [112].	
FoxM1	(Myc Cdk1	, , ,	yclinA and Cdc25A and Cdc25B) or (Plk1 and CyclinB and	
	phosp Cdk2	In our model, FoxM1 activity requires increased expression by Myc and activating phosphorylation by Cyclin E. Alternatively, FoxM1 activity can be sustained by potent Cdk2 / Cdk1 activity in G2 (supported by Cdc25A or Cdc25B), or a serial phosphorylation by Cyclin B / Cdk1 and Plk1 during mitosis.		
Cdc25A	←	FoxM1	FoxM1 is a direct transcriptional inducer of Cdc25A [110].	
Cdc25A	←	E2F1	E2F1 is a direct transcriptional inducer of Cdc25A [113].	
Cdc25A	⊢	pRB	RB blocks E2F1's ability to drive Cdc25A transcription [113,114].	
Cdc25A	⊢	Cdh1	APC/C ^{Cdh1} degrades Cdc25A at mitotic exit [115].	
Cdc25A	—	GSK3	GSK3β phosphorylates Cdc25A, promoting its proteolysis [116].	
Cdc25A	F	СНК1	CHK1 phosphorylates Cdc25A, promoting its proteolysis and inhibiting its interaction with Cyclin B / Cdk1 [117].	
Cdc25A	←	CyclinE	Cdc25A protein levels are stabilized during S-phase by CyclinE/Cdk2 dependent phosphorylation [118].	

Target Node	Link	Input	Description & References	
Cdc25A	←	CyclinA	Cdc25A protein levels are stabilized during S and G2 by Cdk2-dependent phosphorylation. Cdk2 first partners with Cyclin E [118], then continues to stabilize Cdc25A past the point of Cyclin E expression by partnering with Cyclin A [119].	
Cdc25A	←	Cdk1	During mitosis, Cdc25A is stabilizes by Cyclin B / Cdk1 phosphorylation, which protects it from the proteasome [120].	
Cdc25A	←	CyclinB	During mitosis, Cdc25A is stabilizes by Cyclin B / Cdk1 phosphorylation, which protects it from the proteasome [120].	
Cdc25A			not pRB) or [not Cdh1 and { FoxM1 or (E2F1 and not pRB) }] }) or CyclinE or CyclinA or (CyclinB and Cdk1)]	
	our mactiva APC/ accur requir	As the precise combinatorial regulation of Cdc25A throughput the cell cycle is unknown, our model assumes that accumulation of the Cdc25A protein requires transcriptional activation by both E2F1 in the absence of pRB and FoxM1 to override destruction by APC/C ^{Cdh1} . Alternatively, one of the two transcription factors can drive Cdc25A accumulation in the absence of APC/C ^{Cdh1} . In addition, stabilization of Cdc25A either requires the absence of GSK3 and CHK1 (both of which promote its degradation), or stabilization by Cdk activity.		
CyclinA	←	CyclinA_mRNA	Our model, sustained availability of Cyclin A requires transcription.	
CyclinA	F	pAPC	Cyclin A is degraded by the APC/C ^{Cdc20} in prometaphase (as soon as the APC/C components are phosphorylated by Cdk1) [121,122], before the full activation of the complex at SAC passage [123]. In our model, this stage of mitotic APC/C ^{Cdc20} activation is represented by Cdk1-phosphorylated APC/C (pAPC).	
CyclinA	←	Cdc25A	Cdc25A promotes active Cyclin A / Cdk2 complex formation by removing inhibitory phosphorylation from Cdk2 [124,125].	
CyclinA	F	Cdh1	Cyclin A is degraded by APC/C ^{Cdh1} in the presence of the UbcH10 protein [121,126].	
CyclinA	←	Emi1	Emi1 binding to Cdh1 is required to stabilize Cyclin A levels at the G1/S transition, allowing Cyclin A / Cdk2 to block Cdh1 [127-129].	
CyclinA	-	UbcH10	Cyclin A degradation by APC/C ^{Cdh1} requires UbcH10 [126].	
CyclinA	←	CyclinA	We assume that once activated, Cyclin A / Cdk2,1 complexes can sustain their activity under favorable conditions until Cyclin A is degraded.	
CyclinA	_	CyclinA_mRNA and not pAPC and { [Cdc25A and (not Cdh1 or Emi1)] or [CyclinA and ((not Cdh1 and [Emi1 or not UbcH10]) or {Emi1 and not UbcH10})] }		

Target Node	Link	Input	Description & References		
	by phactive CCdh1 influe of Ub CCdh1 of Cy	Cyclin A activity requires transcription (Cyclin_A mRNA) and the absence of degradation by phosphorylated (mitotic) pAPC. In addition, turning ON inactive Cyclin A requires activation of Cdk2 by Cdc25A [124] and the absence / Emi1-mediated inhibition of APC/CCdh1. Once active, Cyclin A maintains its activity in the absence of overpowering influences driving its degradation. Namely, Cyclin A relies on either Emi1 or the absence of UbcH10 for its ability to keep inactive APC/CCdh1 in check. To overpower active APC/CCdh1, Cyclin A requires both Emi1 and no UbcH10. The precise combinatorial regulation of Cyclin A is not known; the above logic is consistent with Cyclin A activity pattern during cell cycle progression.			
Wee1	-	Casp3	Caspase 3 cleaves and deactivates Wee1 [130].		
Wee1	←	СНК1	During DNA replication Wee1 is activated by the checkpoint kinase Chk1 [131].		
Wee1	←	Replication	To model the sensitivity of Wee1 activation to ongoing DNA synthesis even in the absence of damage, our model turns on Wee1 immediately upon the start of DNA replication and maintains it until both Replication and the checkpoint kinase Chk1 is OFF [131]. In addition, Wee1 activity has been implicated in maintaining normal replication fork procession, linking its activity directly to ongoing replication [132].		
Wee1	F	Cdk1	The somatic Wee1 protein is an order of magnitude more sensitive to Cdk1 activity than Cdc25C. Thus, both Cyclin B / Cdk1 and Cyclin A / Cdk1 induce Wee1 phospho- rylation and deactivation [133,134].		
Wee1	-	CyclinB	Cyclin B / Cdk1 is a strong inducer of Wee1 phosphorylation and deactivation [133,134].		
Wee1	F	CyclinA	Cyclin A / Cdk1 can also induce Wee1 phosphorylation and deactivation [133,134].		
Wee1	F	Plk1	Plk1 phosphorylation at S53 promotes Wee1 degradation [133]. This event is primed by Cdk1 phosphorylation of Wee1 at S123 [133]. As the main partner of Cdk1 in mitosis is Cyclin B, we assume that assistance from Plk1 to block Wee1 is more relevant when paired with Cyclin A / Cdk1 complexes.		
Wee1		not Casp3 and (Replication or CHK1) and not (Cdk1 and CyclinB) and [CHK1 or not (Cdk1 and CyclinA and Plk1)]			
	Wee1 is active in non-apoptotic cells undergoing DNA synthesis or subject to the single-strand damage checkpoint signaled by active CHK1. That said, Wee1 is blocked by Cyclin B / Cdk1 phosphorylation or Cyclin A / Cdk1 / Plk1 in the absence of CHK1.				
UbcH10	-	Cdh1	UbcH10 is degraded by APC/C ^{Cdh1} [84].		
UbcH10	←	UbcH10	Active UbcH10 cannot be autoubiquitinated in the presence of APC/C ^{Cdh1} substrates and thus remains active [84].		
UbcH10	←	Cdc20	The presence of APC/C ^{Cdh1} substrates, including Cdc20, inhibit the autoubiquitination of UbcH10 but not its function, thus preserving APC activity [84].		

Target Node	Link	Input	Description & References	
UbcH10	←	CyclinA	The presence of APC/C ^{Cdh1} substrates, including Cyclin A, inhibit the autoubiquitination of UbcH10 but not its function, thus preserving APC activity [84].	
UbcH10	←	CyclinB	The presence of APC/C ^{Cdh1} substrates, including CyclinB, inhibit the autoubiquitination of UbcH10 but not its function, thus preserving APC activity [84].	
UbcH10	not C	dh1 or [UbcH10 an	nd (Cdc20 or CyclinA or CyclinB)]	
	native	ely, active UbcH10 i	g enzyme (E2) UbcH10 is active in the absence of Cdh1. Alters maintained in the presence of Cdh1 when some of its targets CyclinA OR CyclinB [126].	
CyclinB	←	FoxM1	FoxM1 is a direct transcriptional regulator of Cyclin B1 [102,135].	
CyclinB	←	FoxO3	FoxO3 is a direct transcriptional regulator of Cyclin B; its activation in G2 helps increase/maintain Cyclin B levels [136].	
CyclinB	←	CyclinB	Here we assume that FoxO3 alone can only maintain, but not independently induce Cyclin B1 expression.	
CyclinB	—	Cdh1	Cyclin B is degraded by APC/C ^{Cdh1} [137].	
CyclinB	⊢	pAPC	Cyclin B is degraded by APC/C ^{Cdc20} [137].	
CyclinB	⊢	Cdc20	Cyclin B is degraded by APC/C ^{Cdc20} [137].	
CyclinB	[FoxM1 or (FoxO3 and CyclinB)] and not [Cdh1 or (pAPC and Cdc20)]			
	repre	Cyclin B node is ON when the concentration of Cyclin B proteins is high (does not represent the activity of CyclinB/Cdk1 complexes). This occurs when Cyclin B is transcribed by FoxM1, maintained by FoxO3 transcription, and not undergoing APC-mediated degradation by APC/C ^{Cdc20} or APC/C ^{Cdh1} .		
Cdc25B	←	FoxM1	FoxM1 is an essential inducer or Cdc25B [138].	
Cdc25B	←	4N_DNA	Cdc25B is localized at centrosomes, where it is activated by Aurora A kinase [139]. As Aurora A itself is only recruited to duplicated, centrosomes before their separation [140], Cdc25B activation requires duplicated centrosomes. As our model does not directly account for centrosome dynamics, we account for this by requiring the completion of S-phase (4N_DNA).	
Cdc25B	FoxM1 and 4N_DNA			
	Cdc25B activation requires transcription by FoxM1, centrosomal localization, and activation by Aurora A kinase on replicated centrosomes.			
Plk1	-	Cdh1	The majority of Plk1 is degraded in anaphase by the APC/CCdh1 complex [141]	
Plk1	←	FoxM1	Plk1 is a direct transcriptional target of FoxM1 [112].	

Target Node	Link	Input	Description & References	
Plk1	←	Plk1_H	Our model tracks the accumulauton of high-enough levels of Plk1 to survive APC/C ^{Cdh1} mediated destruction into telophase via the Plk1_H node. Its ON state represents strong prior Plk1 activation. Thus, it sustains the Plk1 node in the absence of FoxM1-mediated transcription until Plk1_H itself is lost as Plk1 levels fall.	
Plk1	←	CyclinB	Plk1 is activated by Cyclin B / Cdk1 phosphorylation [142-144].	
Plk1	←	Cdk1	Plk1 is activated by Cyclin B / Cdk1 phosphorylation [142-144].	
Plk1	←	CyclinA	Plk1 activation at the G2/M boundary, before Cdk1/Cuclin B complexes are activated, requires active Cyclin A / Cdk [145].	
Plk1	F	Wee1	Cyclin A-mediated induction of Plk1 is blocked by Wee1 kinase, which specifically inhibits Cdk2 activity [145].	
Plk1	←	Cdc25A	As we do not include a separate Cdk2 node in our model, strong Cyclin A / Cdk2 activity requires ongoing dephosphorylation of Cdk2 by Cdc25A [125].	
Plk1	not Cdh1 and (FoxM1 or Plk1_H) and [(CyclinB and Cdk1) or (CyclinA and not Wee1 and Cdc25A)]			
	levels Plk1 a	Plk1 activity requires the absence of APC/C ^{Cdh1} , transcription by FoxM1, or high Plk1 levels transcribed earlier by both FoxM1 and FoxO3 (see Plk1_H below) [136]. In addition, Plk1 activation requires phosphorylation by either CyclinB/Cdk1 during mitosis or Cyclin A / Cdk2 (aided by lack of Wee1 and Cdc25A) at the G2/M boundary [145].		
Cdc25C	←	Plk1	Plk1-mediated phosphorylation of Cdc25C before mitotic entry is required as the initiating step of the Cdc25C / Cdk1 feedback loop [145]. In addition, Plk1 induces nuclear transport of CDC25B, where it contributes to the initiation of Cdk1 activity [146]. During mitosis, Plk1 helps maintain strong Cdc25C activation by phosphorylating it on the same site as Cyclin B / Cdk1 [147], as indicated by the profound decrease of Cdc25C activity in Plk1-inhibited mitotic cells [143,148].	
Cdc25C	←	Cdc25B	CDC25B starts the cascade leading to mitotic entry by activating a small centrosomal pool of Cyclin B / Cdk1, leading to their nuclear translocation where they trigger the activation of Cdc25C and eventually the larger nuclear Cyclin B / Cdk1 pool [149-151].	
Cdc25C	F	СНК1	Chk1 phosphorylates Cdc25C, leading to its nuclear exclusion, loss of access to its main target, Cdk1 [152]. In addition, CHK1 blocks the ability of Cdc25B to activate Cdc25C at the centrosomes by phosphorylating it and blocking it's Cdk1 activity [153,154].	
Cdc25C	←	Cdk1	Cyclin B / Cdk1 are potent Cdc25C activators, creating positive feedback that causes switch-like mitotic entry [155].	
Cdc25C	←	CyclinB	Cyclin B / Cdk1 are potent activators of Cdc25C [155].	

Target Node	Link	Input	Description & References	
Cdc25C	←	4N_DNA	The nature and localization of the signals responsible for the onset and maintenance of Cdc25C activity require replicated DNA (4N_DNA) [146,151]. Namely, Cdc25C is initially activated by a small pool of Cyclin B / Cdk1 (below the ON-threshold of Cdk1 in our model) which starts our at the replicated centrosome (see above). Moreover, the pool of mitotic Cdc25C co-localized with active Chk1/Cyclin B is found on condensed chromosomes, again requiring the presence of 4N_DNA [152].	
Cdc25C	4N_D	NA and Plk1 and [(Cdc25B and not CHK1) or (CyclinB and Cdk1)]	
	link). activa Cdc2	In our model, Cdc25C is active in cells with replicated DNA (see 4N_DNA → Cdc25C link). Its activation is initiated by a small, initially cytoplasmic pool of Cyclin B / Cdk1 activated by Cdc25B (not directly represented in our model) and further increased by Cdc25B itself, which translocates to the nucleus with the aid of Plk1. During mitosis, Plk1 potentiates the ability of Cyclin B / Cdk1 to maintain Cdc25C activity.		
Cdk1	←	CyclinB	Full kinase activation of Cdk1 in our model requires it to complex with Cyclin B [156].	
Cdk1	←	Cdc25C	Cdk1 is subject to inhibitory phosphorylation by Wee1 or Myt1, while its dephosphorylation is carried out by activated Cdc25C [155,156].	
Cdk1	←	Cdk1	We assume that the presence of fully activated, nuclear Cdk1 is able to overcome the effect of active Wee1 (as long as CHK1 is off), given that Wee1 is very sensitive to Cdk1-mediated inhibitory phosphorylation [134].	
Cdk1	F	Wee1	Wee1 is a nuclear protein that ensures the completion of DNA replication prior to mitosis by blocking Cdk1 activation [157].	
Cdk1	F	СНК1	In the absence of CHK1 kinase, a small cytosolic (centrosomal) pool of Cyclin B / Cdk1 can be activated by Cdc25B, the nuclear translocation of which can trigger a positive feedback loop that activates the full Cdk1 pool (assuming nuclear Wee1 is also inactive). Thus, CHK1 can maintain the OFF state of inactive Cdk1 [154].	
Cdk1	Cycli	nB and Cdc25C an	d [not CHK1 or (not Wee1 and Cdk1)]	
	Full Cdk1 kinase activation requires its binding partner Cyclin B and the Cdc25C phosphatase, which maintains Cdk1 in an active dephosphorylated state. Cdk1 is inhibited by the checkpoint kinase CHK1, unless it is already full active and Wee1 kinase is inhibited.			
pAPC	←	CyclinB	CyclinB/Cdk1 activation promotes APC/C ^{Cdc20} activity via APC/C subunit phosphorylation [158].	
pAPC	←	Cdk1	CyclinB/Cdk1 activation promotes APC/C ^{Cdc20} activity via APC/C subunit phosphorylation [158].	
рАРС	←	Plk1	In addition to Cyclin B / Cdk1 phosphorylation, full activation of the APC/C ^{Cdc20} complex also requires the kinase activity of Plk1 [159].	

Target Node	Link	Input	Description & References	
рАРС	←	pAPC	Activated APC/C ^{Cdc20} initiates the Metaphase! Anaphase transition by degrading Cyclin B and securing [87, 68]. Once active, APC/C ^{Cdc20} no longer requires sustained CyclinB/Cdk1 phosphorylation.	
рАРС	←	Cdc20	Once catalytically active, APC/C ^{Cdc20} no longer requires ongoing CyclinB/Cdk1 or Plk1 phosphorylation.	
pAPC	(Cycl	inB and Cdk1 and	Plk1) or (CyclinB and Cdk1 and pAPC) or (pAPC and Cdc20)	
	the m	ore potent inducer, C from an OFF state	Plk1 can aid full activation of APC/C, but Cdk1 appears to be our model requires both Cyclin B / Cdk1 and Plk1 to activate e, but only Cdk1 activity to maintain it. In addition, ongoing unctional APC/C ^{Cdc20} complex is no longer required.	
Cdc20	F	Cdh1	APC/C ^{Cdh1} complexes degrade Cdc20, leading to a complete switch from APC/C ^{Cdc20} to APC/C ^{Cdh1} during mitotic exit [160,161].	
Cdc20	F	Cdk1	Cdk1-phosphorylated Cdc20 interacts with Mad2 rather than APC/C, resulting in a block on APC/C ^{Cdc20} activation until completion of spindle assembly [162].	
Cdc20	F	CyclinB	Cyclin B partners with Cdk1 to keep Cdc20 phosphorylated, increasing its interaction with Mad2 rather than APC/C [162].	
Cdc20	F	CyclinA	Cyclin A / Cdk2 complexes phosphorylate Cdc20 and inactivate the APC/C ^{Cdc20} complex during S and G2 [163].	
Cdc20	F	Emi1	Emi1 binds Cdc20 and inhibits the ubiquitin ligase activity of APC/C ^{Cdc20} [128].	
Cdc20	F	Mad2	The physical alignment of chromosomes along the mitotic spindle is monitored by Mad checkpoint proteins [164]. In cells with a normal spindle assembly checkpoint (SAC), even a single unattached, Mad2-bound kinetochore can sequester Cdc20 and thus inhibit APC/C ^{Cdc20} [165].	
Cdc20	←	pAPC	Cdc20 becomes active in early mitosis by binding to APC/C, an event that requires Cyclin B / Cdk1 phosphorylation of several APC/C subunits [160,166].	
Cdc20	-	pAPC and not Emi1 and not Cdh1 and { not Mad2 or [not CyclinA and not (CyclinB and Cdk1)] }		
	In our model, APC/C ^{Cdc20} complex formation is represented by the joint activity of Cdc20 and phosphorylated APC/C (pAPC). Cdc20 is thus ON in the presence of pAPC when both Emi1 and Cdh1 are absent (APC/C ^{Cdh1} is represented by the Cdh1 node, see below). In addition, Cdc20 activity requires either the absence of Mad2 at unattached kinetochores, or the absence of Cdc20 phosphorylation by Cyclin B / Cdk1 or by Cyclin A / Cdk2 complexes to potentiate the interaction between Mad2 and Cdc20. and pAPC is ON (present and phosphorylated) [162].			
Cdh1	-	CyclinB	Cyclin B / Cdk1 complexes phosphorylate Cdh1, blooming its ability to bind APC/C and inhibiting APC/C ^{Cdh1} activity [167].	
Cdh1	F	Cdk1	Cyclin B / Cdk1 complexes phosphorylate Cdh1, blooming its ability to bind APC/C and inhibiting APC/C ^{Cdh1} activity [167].	

Target Node	Link	Input	Description & References
Cdh1	-	CyclinA	Active Cyclin A / Cdk1,2 complexes phosphorylate Cdh1 during S, G2 and early mitosis, impairing its interaction with APC/C until late stages of mitosis when Cdk1/2 activity falls [161].
Cdh1	⊢	Cdc25A	As we do not include a separate Cdk2 node in our model, strong Cyclin A / Cdk2 activity capable of overriding Cdh1 activity even in the presence of Emi1 requires ongoing dephosphorylation of Cdk2 by Cdc25A [125].
Cdh1	-	Emi1	Emi1 blocks APC/C ^{Cdh1} binding to its substrates [129], as well as its ability to add ubiquitin chains to them [168].
Cdh1	not (CyclinB and Cdk1) and not [CyclinA and (Emi1 or Cdc25A)]		
	APC/C ^{Cdh1} activity requires the absence of Cyclin Dependent kinase phosphorylation by Cyclin B / Cdk1, or Cyclin A / Cdk2 aided by further inhibition of Cdh1 by Emi1, or ongoing Cdk2 activation by Cdc25A in the absence of Emi1.		

S1.D. Boolean regulatory logic of <u>Origin of Replication Licensing Switch</u> nodes.

Target Node	Link	Input	Description & References	
ORC	-	E2F1	Expression of the ORC1 gene is regulated by E2F1 [169].	
ORC	←	Pre-RC	Licensed but not yet fired replication complexes (Pre-RCs containing ORC, Cdc6, Cdt1 and inactive MCMs) remain stable at sites of replication origin until fired by the activation of the MCM helicase [170].	
ORC	←	Cdc6	Availability of stable (unphosphorylated) Cdc6 in the Pre-RC is necessary for the maintenance of licensed origins [170].	
ORC	←	Cdt1	Active (unphosphorylated and not geminin-bound) Cdt1 bound to the Pre-RC is necessary for the maintenance of licensed origins [170].	
ORC	E2F1 or (Pre-RC and Cdt1 and Cdc6)			
		ORC proteins can bind at origins of replication when transcribed by E2F1 or as part of a fully assembled and licensed Pre-RC complex (including active Cdc6 and Cdt1).		
Cdc6	-	Casp3	Caspase 3 cleaves and deactivates Cdc6 [171].	
Cdc6	F	CyclinA	Phosphorylation of CDC6 by Cyclin A / Cdk2 during DNA replication leads to its re-localization to the cytoplasm [172].	
Cdc6	F	4N_DNA	In our model, full deactivation of Cdc6 represents the firing of all ORCs as DNA replication is completed. Thus Cyclin A's inhibitory action takes full effect once the cell reaches 4N DNA content [172].	
Cdc6	←	E2F1	Transcription of Cdc6 is directly induced by E2F1 [173]	
Cdc6	←	ORC	ORC recruits Cdc6 to origins of replication [170].	

Target Node	Link	Input	Description & References
Cdc6	F	Plk1	Plk1 binds, phosphorylated and strongly recruits Cdc6 to the spindle pole during metaphase, then to the central spindle in anaphase, leading to its exclusion from chromosomes until telophase, when the majority of Plk1 is degraded by APC/C ^{Cdh1} [174].
Cdc6	←	Pre-RC	Licensed but not yet fired replication complexes (Pre-RCs containing ORC, Cdc6, Cdt1 and inactive MCMs) remain stable and Cdc6-bound until fired by the activation of the MCM helicase [170].
Cdc6	←	Cdc6	Stable (unphosphorylated) Cdc6 in the Pre-RC is necessary for the maintenance of licensed origins [170].
Cdc6	←	Cdt1	Active (unphosphorylated and not geminin-bound) Cdt1 bound to the Pre-RC is necessary for the maintenance of licensed origins [170].
Cdc6		asp3 and not (4N_Ind ORC and Cdc6	DNA and CyclinA) and [(E2F1 and ORC and not Plk1) or (Preand Cdt1)]
	In our model the Cdc6 node represents nuclear, chromatin-bound Cdc6. Thus, the node is only active during the assembly of pre-replication complexes, or their ongoing presence during DNA replication. Cdc6 is ON in the absence of Caspase 3 or CyclinA / Cdk2 phosphorylation of Cdc6 in all origins required for the completion of DNA replication (thus, its inhibition by Cyclin A also requires 4N_DNA). In addition, active Cdc6 requires either transcription by E2F1 and recruitment by origin-bound ORC proteins in the absence of mitotic Plk1 or maintenance of Pre-RCs by the presence of all of its components.		
Cdt1	←	ORC	ORC-bound origin of replication sites are the point of pre- replication complex assembly, where Cdt1 is recruited by ORC- bound Cdc6 [170].
Cdt1	←	Cdc6	ORC-bound Cdc6 recruits Cdt1 to pre-RC complexes [170].
Cdt1	F	CyclinE	Sustained Cdk2 activity during S-phase, responsible for the firing of all origins required to complete DNA synthesis (modeled as simultaneous Cyclin E, Cyclin A and Cdc25A activity), also leads to the phosphorylation and proteasomal degradation of Cdt1 [170].
Cdt1	F	CyclinA	Sustained Cdk2 activity leads to phosphorylation and degradation of Cdt1 (see above) [170].
Cdt1	F	Cdc25A	Sustained Cdk2 activity leads to phosphorylation and degradation of Cdt1 (see above) [170].
Cdt1	←	Pre-RC	Licensed but not yet fired replication complexes (Pre-RC) remain stable until fired during DNA replication [170].
Cdt1	←	Мус	Cdt1 is a direct transcriptional target of the Myc-Max complex [175], ensuring its availability for Pre-RC formation and maintenance.
Cdt1	←	E2F1	Cdt1 is a direct transcriptional target of E2F1 [176], ensuring its availability for Pre-RC formation and maintenance.

Target Node	Link	Input	Description & References	
Cdt1	-	pRB	E2F1-mediated transcription of Cdt1 is blocked by hypophosphorylated (active) pRB [176].	
Cdt1	F	geminin	Geminin binds to Cdt1 at pre-replication complexes, where it blocks Cdt1 binding to DNA, sequestering it away from Pre-RCs [177].	
Cdt1			nd Cdc6 and not (CyclinE and CyclinA and Cdc25A) and { [Pre- or [E2F1 and (Myc or not pRB)] }	
	bound firing Cdc2 stable comp Altern	Replication-origin bound Cdt1 requires the absence of geminin, the presence of origin-bound ORC and Cdc6, and the absence of sustained Cdk2 activity responsible for the firing of all origins during DNA synthesis (modeled as simultaneous Cyclin E, Cyclin A and Cdc25A activity). Bound into a licensed pre-replication complex (Pre-RC), Cdt1 remains stable as long as it is transcribed by E2F1 or Myc (this guarantees that Pre-RC complexes cannot persist indefinitely in the absence of <i>de novo</i> transcription). Alternatively, it can be turned on by E2F1, aided by Myc or the absence of RB, and FoxO3 in cells with 4N_DNA.		
Pre-RC	←	ORC	Pre-RC complexes assemble when ORC, Cdc6 and Cdt1 are all bound to sites of replication origin along the DNA [170].	
Pre-RC	←	Cdc6	Pre-RC complexes assemble when ORC, Cdc6 and Cdt1 are all bound to sites of replication origin along the DNA [170].	
Pre-RC	←	Cdt1	Pre-RC complexes assemble when ORC, Cdc6 and Cdt1 are all bound to sites of replication origin along the DNA, leading to the recruitment of the MCM helicase [170].	
Pre-RC	—	Replication	Pre-RCs fire and fall apart during DNA replication [170].	
Pre-RC	F	4N_DNA	In our model Pre-RC turns OFF when Replication is completed, marked by the time-point when both Replication and 4N_DNA are ON.	
Pre-RC	ORC	and Cdc6 and Cdt	1 and not (Replication and 4N_DNA)	
	Pre-RC complexes assemble when ORC, Cdc6, and Cdt1 are all bound to sites of replication origin along the DNA. The node denoting their licensing turns OFF at the moment of transition from ongoing Replication to 4N_DNA (it is blocked in the one time-point when both of these nodes are ON).			
geminin	←	E2F1	Geminin is a direct transcriptional target of E2F1 [176].	
geminin	-	Cdh1	Geminin is a target of APC/C ^{Cdh1} ubiquitin ligase [178].	
geminin	-	pAPC	Geminin is a target of APC/C ^{Cdc20} at the metaphase/anaphase transition [179].	
geminin	F	Cdc20	Geminin is a target of APC/C ^{Cdc20} at the metaphase/anaphase transition [179].	
geminin	E2F1	and not Cdh1 and	not (pAPC and Cdc20)	
		nin is present when or APC/C ^{Cdc20} .	transcribed by E2F1 and not targeted for degradation by APC/	

 $\textbf{S1.E.} \ \ \textbf{Boolean regulatory logic of } \underline{\textbf{Cell Cycle Process}} \ \ \textbf{nodes}.$

Target Node	Link	Input	Description & References	
Replication	F	CAD	Caspase-activated DNAase (CAD) destroys DNA, preventing ongoing replication.	
Replication	←	Pre-RC	Ongoing DNA replication requires licensed replication origins, which fire throughout DNA synthesis [180].	
Replication	←	E2F1	In addition to E2F1 target genes directly included in our model, E2F1 transcribes an array of critical S-phase genes responsible for carrying out DNA synthesis (e.g, POLA1, POLA2, MCM3, MCM5, MCM6, PCNA, TOP2A, RFC2, TK1) [181,182].	
Replication	←	CyclinE	DNA replication is initiated by fully active CyclinE/Cdk2 [183].	
Replication	←	Cdc25A	Active Cdc25A is required for onset as well as progression through S-phase [184,185].	
Replication	←	CyclinA	Cyclin A / Cdk1 complexes regulate the origin firing program in mammalian cells and are required for the completion of DNA replication [185,186].	
Replication	←	Replication	Once ongoing, DNA synthesis continues in the presence of active Cyclin A / Cdk2, only ending when DNA content is doubled.	
Replication	F	4N_DNA	Complete duplication of a cell's DNA, represented in our model by 4N DNA = ON, marks the end of active Replication.	
Replication		not CAD and Pre-RC and { (E2F1 and CyclinE and Cdc25A) or (Replication and CyclinA and Cdc25A and (E2F1 or not 4N_DNA)]}		
	cell, E2F1 first re susta	The Replication node represents ongoing DNA synthesis. This requires a non-apoptotic cell, licensed pre-replication complexes (Pre-RC). The start of DNA synthesis requires E2F1-meditated transcription of the genes that help execute it, as well as the firing of the first round of replication origins by Cyclin E / Cdk2. Once ongoing, replication is sustained by Cyclin A / Cdk2 and Cdc25A, aided by E2F1 and terminated by completion of a full round of synthesis (4N_DNA).		
ATR	←	Replication	ATR accumulates at replication forks during unperturbed DNA synthesis [187].	
CHK1	←	ATR	ATR kinase activates CHK1 at replication forks, which not only blocks premature mitosis but also regulates the rate of origin firing by keeping Cdc25 protein levels from increasing above their physiological range [187]	
4N_DNA	F	CAD	Caspase-activated DNAase (CAD) destroys DNA, preventing maintenance of a double DNA content.	
4N_DNA	←	Replication	DNA content is doubled by the process of Replication.	
4N_DNA	←	Pre-RC	Replication can only complete DNA synthesis and produce double DNA content if the availability of licensed replication origins is not blocked [180].	
4N_DNA	←	CyclinA	Cyclin A / Cdk1 complexes regulate the origin firing program in mammalian cells and are required for the completion of DNA replication [183,186].	

Target Node	Link	Input	Description & References	
4N_DNA	←	4N_DNA	Once achieved, a cell's 4N DNA content is sustained up to the point of cytokinesis.	
4N_DNA	F	Ect2	At the start of cytokinesis, the Ect2 RhoGEF is recruited to the central spindle [188]. Ect2 aids the accumulation of GTP-bound RhoA [189] and the formation of the contractile ring. In our model, Ect2 recruitment to the central spindle marks the start of cytokinesis and the subsequent resetting of daughter cell DNA content to 2N.	
4N_DNA		not CAD and { [Replication and ({Pre-RC and CyclinA} or $4N_DNA$)] or $(4N_DNA)$ and not Ect2)}		
	of the	4N DNA content in our model is reached via the completion of Replication (via the firing of the last round of replication origins by Cyclin A / Cdk complexes) and maintained in non-apoptotic cells the absence of a contractile ring (marked by Ect2), driving cytokinesis.		
U_Kinetochores	←	4N_DNA	Metaphase requires replicated sister chromatids (4N_DNA), held together by their kinetochores, face in opposing directions and can be attached to opposite poles of the mitotic spindle.	
U_Kinetochores	F	Cdh1	Premature activation of APC/C ^{Cdh1} destroys the incomplete spindle by triggering premature, aberrant anaphase. This occurs due to premature degradation of APC/C targets including Securin (responsible for keeping sister chromatids attached [190]), Cyclin B, CDC20, and Aurora kinase A (AURKA) [191].	
U_Kinetochores	F	A_Kinetochores	In our model, the transition from unattached (U_Kinetochores) to all attached kinetochores (A_Kinetochores) marks the completion of the mitotic spindle and Spindle Assembly Checkpoint (SAC) passage.	
U_Kinetochores	←	CyclinB	The start of mitotic spindle assembly is initiated by active CyclinB/Cdk1 [192].	
U_Kinetochores	←	Cdk1	The start of mitotic spindle assembly is initiated by active CyclinB/Cdk1 [192].	
U_Kinetochores	←	U_Kinetochores	Once metaphase starts, the mitotic spindle remains incomplete as long as some of the kinetochores remain unattached.	
U_Kinetochores	4N_DNA and not Cdh1 and not A_Kinetochores and [(CyclinB and Cdk1) or U_Kinetochores]			
	The U_Kinetochores node in our model is on from the moment the nuclear envelope is dissolved in prometaphase and the mitotic spindle starts to form, until all kinetochores are properly attached. In addition to the presence of unattached kinetochores, U_Kinetochores = ON requires attached sister chromatics, the absence of APC/CCdh1 activity. It is turned on my CyclinB/Cdk1 and remains on until the spindle is complete (or it is destroyed by APC/CCdh1).			
Mad2	←	U_Kinetochores	The Mad2 SAC protein is active and potent in the presence of even a single unattached kinetochore [165].	
Mad2	F	A_Kinetochores	Mad2 is inhibited by SAC passage, marked by the completion of the spindle and proper attachment of all kinetochore [165].	

Target Node	Link	Input	Description & References		
Mad2	U_Ki	netochores and no	t A_Kinetochores		
	as lor keepi	Our model represents the SAC via the Mad2 kinetochore-binding protein. Mad2 is active as long as the cell has at least one unattached kinetochore and it is responsible for keeping Cdc20 sequestered from APC/C. By keeping APC at bay until the spindle is complete, Mad2 is required for the proper timing of anaphase [165].			
A_Kinetochores	←	4N_DNA	Completion of the mitotic spindle requires replicated and attached sister chromatids (4N_DNA).		
A_Kinetochores	⊢	Cdh1	APC/C ^{Cdh1} destroys the spindle by triggering anaphase via the degradation of APC/C targets, including Securin [191].		
A_Kinetochores	F	pAPC	During normal mitosis, the completed spindle is pulled apart in response to APC/C ^{Cdc20} -mediated degradation of Securin, which normally blocks separate from severing the cohesin rings keeping sister chromatids attached [190].		
A_Kinetochores	F	Cdc20	During normal mitosis, the completed spindle is pulled apart in response to APC/C ^{Cdc20} -mediated degradation of Securin, which normally blocks separate from severing the cohesin rings keeping sister chromatids attached [190].		
A_Kinetochores	←	A_Kinetochores	Once assembled, separation of the mitotic spindle requires APC/C activity to promote the destruction of sister chromatid cohesion [190].		
A_Kinetochores	←	U_Kinetochores	The mitotic spindle is assembled gradually, as the number of unattached kinetochores gradually decreased by the formation of microtubule attachments.		
A_Kinetochores	←	Plk1	Plk1 activity at unattached kinetochores is required for promoting their attachment [193]. In its absence, kinetochores remain unattached and cells eventually undergo mitotic catastrophe and apoptosis [194].		
A_Kinetochores	←	CyclinB	Ongoing Cyclin B / Cdk1 at unattached kinetochores is necessary to keep Plk1 active and allow the completion of mitosis [195].		
A_Kinetochores	←	Cdk1	Ongoing Cyclin B / Cdk1 at unattached kinetochores is necessary to keep Plk1 active and allow the completion of mitosis [195].		
A_Kinetochores		4N_DNA and not Cdh1 and not (pAPC and Cdc20) and [A_Kinetochores or (U_Kinetochores and Plk1 and CyclinB and Cdk1)]			
	attac the p to un	The completed spindle, represented by the A_Kinetochores node, requires replicated and attached sister chromatids (4N_DNA) and the absence of APC/C activity. It turns on when the process of spindle assembly (U_Kinetochores) is completed by active Plk1 localized to unattached kinetochores in the presence of ongoing Cyclin B / Cdk1 activity and it remains on until anaphase (APC/C activation).			
Plk1_H	←	Plk1	Active mitotic Plk1 is a prerequisite for the accumulation of the larger active Plk1 pool denoted by Plk1_H.		
Plk1_H	←	FoxM1	Plk1 is a direct transcriptional target of FoxM1; loss of FoxM1 severely reduces Plk1 protein levels [101,112].		

Target Node	Link	Input	Description & References
Plk1_H	←	FoxO3	Plk1 is a direct transcriptional target of FoxO3, but Plk1 appears to be sufficiently induced in the absence of FoxO preteens to aid its G2/M and mitotic functions. In contrast, accumulation if a large enough Plk1 pool to briefly outlast APC/C ^{Cdh1} activation (modeled by the Plk1_H node), requires FoxO activity in G2 [136].
Plk1_H	←	FoxO1	In addition to FoxO3, FoxO1 also binds the Plk1 promoter, potentially aiding its accumulation during G2 [196].
Plk1_H	←	Plk1_H	Once accumulated, we assume that the Plk1_H pool of active Plk1 remains stable in the absence of FoxO-mediated transcription. This is supported by negative feedback regulation of FoxO proteins by Plk1 [14], indicating that ongoing high FoxO activity is likely not required for the maintenance of Plk1_H.
Plk1_H	Plk1	and FoxM1 and (PI	k1_H or FoxO3 or FoxO1)
	The ON state of Plk1_H encodes the short-lived memory of a sufficiently large active Plk1 pool to temporarily survive Plk1 destruction by APC/CCdh1 [141], recruit Ect2 to the central spindle, and thus aid the completion of cytokinesis [188]. Thus, Plk1_H requires ongoing Plk1 activation and transcription by FoxM1, and either induction by FoxO3 or FoxO1, or prior accumulation.		
Ect2	←	4N_DNA	Formation of a spindle midzone, where Ect2 accumulates in preparation of cytokinesis requires recently separated sister chromatids (4N DNA content).
Ect2	←	Plk1_H	Plk1 activity in telophase (Plk1_H) is required for the recruitment of Ect2 to the central spindle [141].
Ect2	←	Cdh1	APC/C ^{Cdh1} -mediated destruction of Aurora kinase is required for the assembly of a robust spindle midzone at anaphase and for the normal timing of cytokinesis [197].
Ect2	F	U_Kinetochores	Formation of a spindle midzone requires the separation of sister chromatids; thus it cannot occur before anaphase.
Ect2	F	A_Kinetochores	Formation of a spindle midzone requires the separation of sister chromatids; thus it cannot occur before anaphase.
Ect2	4N_D	NA and Plk1_H and	d Cdh1 and not U_Kinetochores and not A_Kinetochores
	Ect2 activation at the spindle midzone represents the step of cytokinesis in our model. Thus, Ect2 requires 4N_DNA, high Plk1 activity, as well as Cdh1 for the assembly of a normal spindle midzone. Finally, Ect2 cannot be recruited to the mid zone before anaphase is completed.		

 $\textbf{S1.F.} \ \ \textbf{Boolean regulatory logic of} \ \underline{\textbf{Apoptotic Switch}} \ \ \textbf{nodes}.$

Target Node	Link	Input	Description & References
Trail	←	Trail	The Trail node represents environmental availability of the Trail protein outside the cell; this input node remains on/off if set ON/OFF in the absence of in silico perturbation.
DR4_5	←	Trail	The DR4 and DR5 death receptors, represented by the DR4_5 node, are activated by extracellular Trail [198].
Casp8	←	DR4_5	Trail-bound (active) DR4 and DR5 receptors trigger the assembly of the pro-apoptotic death-inducing signaling complex (DISC), which binds a cluster of pro-Caspase 8 proteins and initiates their cleavage into active Caspase 8 [199].
Casp8	←	Casp3	Caspase 3 indirectly activates Caspase 8 by cleaving Caspase 6 [200], which, in turn, cleaves Caspase 8 [201].
Casp8	DR4_	5 or Casp3	
	Pro-C	Caspase 8 may be o	eleaved independently by DISC (DR4_5) or Caspase 3.
Casp2	←	Casp3	Caspase 2 is a target of Caspase 3, as its inhibition severely limits Caspase 2 cleavage during apoptosis [202,203].
Casp2	←	U_Kinetochores	Although the precise molecular mechanism by which Caspase 2 is activated during prolonged or stalled mitosis is unclear, its activation platform, the PIDDosome, has been localized to unattached kinetochores [204]. Even though a checkpoint protein keeps the PIDDosome unresponsive to DNA damage signals, the loss of protective Cyclin B / Cdk1 phosphorylation only leads to Caspase 2 activation in the presence of a partially assembled mitotic spindle, and requires active SAC.
Casp2	←	Mad2	A functional spindle assembly checkpoint is required for mitotic cell death upon prolonged mitotic arrest [205] or spindle damage [206].
Casp2	F	Cdk1	Cyclin B1/Cdk1 phosphorylate caspase-2 at Ser 340, preventing its activation [207].
Casp2	F	CyclinB	Cyclin B1/Cdk1 phosphorylate caspase-2 at Ser 340, preventing its activation [207].
Casp2	Casp3 or [U_Kinetochores and Mad2 and not (CyclinB and Cdk1)]		
	Pro-caspase 2 is cleaved and activated by Caspase 3, or by failed cytokinesis marked by the presence of unattached kinetochores, an active SAC, and the absence of active Cyclin B / Cdk1 complexes to phosphorylate and inhibit Caspase 2.		
MCL-1	-	Casp3	Caspase 3 cleaves and deactivated MCL-1 [208].
MCL-1	-	Casp2	Caspase 2 activation destabilizes the MCL-1 protein [209].
MCL-1	F	U_Kinetochores	During prolonged mitotic arrest (U_kinetochores), MCL-1 levels drop steadily due to phosphorylation by JNK, p38 and/or CKII and its subsequent degradation by the E3 ubiquitin ligase SCF(FBW7) [210].

Target Node	Link	Input	Description & References
MCL-1	-	Cdk1	Phosphorylation by CyclinB / Cdk1 in cells arrested in mitosis initiates MCL-1 degradation [211].
MCL-1	F	CyclinB	In cells arrested in mitosis, phosphorylation by CyclinB / Cdk1 on T92 initiates MCL-1 degradation [211].
MCL-1	F	E2F1	E2F1 is a direct transcriptional repressor of MCL-1 [212].
MCL-1	←	ERK	ERK phosphorylates McI-1, promoting its interaction with Pin1, which stabilizes it [213,214].
MCL-1	-	GSK3	MCL-1 is phosphorylated by GSK-3, leading to ubiquitinylation and degradation of MCL-1 [215].
MCL-1	←	AKT_B	In order to account for the loss of MCL-1 in the complete absence of growth factors vs. its presence in low growth factor environments, we required basal AKT to modulate the strength of GSK3 inhibition [215].
MCL-1		asp3 and not Casp 1 and CyclinB and	o2 and {not GSK3 or [AKT_B and (ERK or not E2F1)] } and [not U_Kinetochores)]
	Avoic activi repre Cyclin	ling degradation via ty (AKT_B) [215] an ssor E2F1. Finally, o nB / Cdk1 phospho	d destruction of MCL-1 must be absent for MCL-1 to be ON. GSK3 requires the GSK3-weakenjng presence of basal AKT deither ERK-mediated stabilization, or the absence of its during mitotic arrest (U_Kinetochores), MCL-1 is deactivated by rylation, which shields it from the PPA2-mediated degradation-targeting sites [210].
BCLXL	-	Casp3	BCL-xL is cleaved and deactivated by Caspase 3 [216].
BCLXL	-	BAD	Bad can bind BCL-xL and displace it from BAX, thus deactivating it [217].
BCLXL	←	BCL2	BCL2 competes with BCL-xL for BAD binding. Although BCL-xL is a stronger binding partner, we assume that BAD alone cannot fully block BCL-xL in the presence of BCL2 [217].
BCLXL	←	MCL-1	MCL-1 competes with BCL-xL for BAK binding; the presence of MCL-1 can keep part of the BCL-xL pool active [218].
BCLXL	-	U_Kinetochores	Prolonged mitosis is required for the accumulation of BCL-xL phosphorylation, weakening its interaction with Bax [219].
BCLXL	←	Plk1	In addition to other effects of prolonged mitotic arrest on BCL-2 proteins, Plk1 inhibition synergistically enhances the inhibitory phosphorylation of BCL-2 and BCL-xL, as well as downregulation of MCL-1 [220].
BCLXL	F	Cdk1	During normal mitosis, Cyclin B / Cdk1 only transiently phosphorylates part of the BCLXL pool. Prolonged mitosis, however, results in high levels of BCL-xL (and Bcl-2) phosphorylation, priming the system for Caspase 2-mediated apoptosis [221,222].
BCLXL	F	CyclinB	Cyclin B / Cdk1 phosphorylates BCL-xL (and Bcl-2) during mitosis [221,222].

Target Node	Link	Input	Description & References	
BCLXL	not Casp3 and (BCL2 or not BAD) and { not U_Kinetochores or [Plk1 and (not {CyclinB and Cdk1} or {BCL2 and MCL-1})] or [BCL2 and MCL-1 and not (CyclinB and Cdk1)] }			
	While the precise combinatorial logic governing BCLXL activity is not clear from literature. BCLXL activity requires the absence of Caspase 3. In addition, BAD can block BCLXL when BCL2 is also OFF. Finally, mitotic BCLXL can be inhibited by Cdk1 activity if either BCL2, MCL-1, or Plk1 are OFF. In the absence of Plk1, loss of either BCL2 or MCL-1 can result in BCLXL inhibition (even without Cdk1 phosphorylation), as we assume its targets are no longer competitively bound by its family members.			
BCL2	-	Casp3	BCL2 is cleaved and deactivated by Caspase 3 [223].	
BCL2	⊢	BAD	BCL2 competes with BCL-xL for BAD binding. BAD displaces BCL2 from its Although BCL-xL is a stronger binding partner, we assume that BAD alone cannot fully block BCL-xL in the presence of BCL2 [217].	
BCL2	F	ВІМ	BIM binds BCL2 and they mutually inhibit each other's ability to activate further targets [224].	
BCL2	-	вік	BIK also binds BCL2 and they mutually inhibit each other's activity [225].	
BCL2	←	BCLXL	BCL2 competes with BCL-xL for binding most of their apoptotic partners, including BIK, BIM, BID, BAX and BAK.	
BCL2	←	MCL-1	MCL-1 competes with BCL2 for binding most of their apoptotic partners, including BIK, BIM, BID, BAX and BAK.	
BCL2	-	U_Kinetochores	Prolonged mitosis is required for the accumulation of BCL-2 phosphorylation [221,222,226].	
BCL2	←	Plk1	In addition to other effects of prolonged mitotic arrest on BCL-2 proteins, Plk1 inhibition synergistically enhances the inhibitory phosphorylation of BCL-2 and BCL-xL, as well as downregulation of MCL-1 [220].	
BCL2	F	Cdk1	Prolonged mitosis results in high levels of BCL-xL and Bcl-2 phosphorylation, priming the system for Caspase 2-mediated apoptosis [221,222,226].	
BCL2	F	CyclinB	Cyclin B / Cdk1 phosphorylates Bcl-2 (and BCL-xL) during mitosis [221,222].	
BCL2	not (Casp3 or BAD or BIM or BIK) and {not U_Kinetochores or (MCL-1 and BCLXL) or [Plk1 and (BCLXL or MCL-1 or not {Cdk1 and CyclinB})]}			
	While the precise combinatorial logic governing BCL2 activity is not clear from literature, we modeled BCL2 as ON in the absence of Caspase 3, BAD, BIM or BIK. This choice makes BCL2 the most sensitive of the three family members to activation of its three inhibitors. In addition, mitotic BCL2 is blocked by Cdk1 if both BCL-xL and MCL-1 are OFF. In the absence of Plk1, loss of either BCL2 or MCL-1 can result in BCL-2 inhibition (even without Cdk1 phosphorylation), as we assume its targets are no longer competitively bound by its family members.			
BAD	←	Casp3	Caspase 3 cleaves BAD, generating a more potently apoptotic fragment [227].	

Target Node	Link	Input	Description & References		
BAD	←	Casp8	Caspase 8 is also able to cleave BAD, generating a more potently apoptotic fragment [227]. In addition, TRAIL-mediated apoptosis results in BAD cleavage by a Caspase upstream of MOMP, creating a potent apoptotic inducer before full Caspase 3 activation [228].		
BAD	-	AKT_B	Akt phosphorylates BAD at Ser-136, inducing its sequestration away from the mitochondrial membrane where its BCL-2 family targets are located (e.g., BCL-2, BCL-xL) [229]		
BAD	-	AKT_H	Akt phosphorylates BAD at Ser-136, inducing its sequestration away from the mitochondrial membrane where its BCL-2 family targets are located (e.g., BCL-2, BCL-xL) [229]		
BAD	-	ERK	ERK phosphorylates BAD at Ser-112, inducing its sequestration away from the mitochondrial membrane where its BCL-2 family targets are located (e.g., BCL-2, BCL-xL)[230].		
BAD	⊢	S6K	S6K1 phosphorylates BAD at Ser-155, directly blocking its binding to BCL-xL [231].		
BAD		Casp3 or not (AKT_H or AKT_B or ERK or S6K) or {Casp8 and [not (AKT_B and ERK and S6K) and not (AKT_H and {AKT_B or ERK})] }			
	BAD in our model is ON when cleaved by Caspase 3, or in the complete absence of survival signals (AKT, ERK or S6K). Alternatively, BAD can be cleaved and activated by Caspase 8 in the absence of strong survival signaling. We modeled this inhibitory survival signal as either the combined activity of ERK, S6K and (at least) basal AKT, or high AKT in the joint presence of ERK and basal AKT (indicating that AKT_H will not drop by the next time-step).				
BIK	-	BCL2	BCL2 binds BIK; they mutually inhibit each other [225].		
BIK	-	BCLXL	BCLXL also binds BIK; they mutually inhibit each other [232].		
BIK	-	MCL-1	MCL-1 also binds BIK; they mutually inhibit each other [233].		
BIK	not (MCL-1 or BCLXL or	BCL2)		
		s free to activate its 2 family proteins in	target, BAX, only when it is not sequestered by any of the three our model [233].		
ВІМ	←	FoxO3	FoxO3 is a transcriptional activator of BIM [234].		
ВІМ	←	GSK3	GSK3 kinase is likely required for the AP1-dependent expression of BIM [235].		
ВІМ	-	ERK	The MEK/ERK pathway represses BIM protein levels, likely via transcriptional repression [236].		
ВІМ	-	BCL2	BLC2 binds BIM and inhibits its apoptotic activity [224].		
ВІМ	-	BCLXL	BLC-XL binds BIM and inhibits its apoptotic activity [224].		
ВІМ	-	MCL-1	MCL-1 binds BIM and inhibits its apoptotic activity [237].		
ВІМ	FoxO	3 and GSK3 and no	ot (ERK or MCL-1 or BCLXL or BCL2)		

Target Node	Link	Input	Description & References
			vity requires expression driven by FoxO3 and aided by GSK3 ree inhibitory BCL2 family proteins.
BID	←	Casp8	In response to TRAIL (or FAS ligand), the initiator caspase 8 cleaves BID to its active truncated form [238-240].
BID	←	Casp2	Caspase 2 cleaves BID to its active truncated form [241]
BID	F	BCL2	All three anti-apoptotic BCL2 proteins (BCL2, BCL-xL and MCL-1) sequesters BID into stable complexes, preventing them from activating BAX or BAK [242].
BID	F	BCLXL	All three anti-apoptotic BCL2 proteins (BCL2, BCL-xL and MCL-1) sequesters BID into stable complexes, preventing them from activating BAX or BAK [242].
BID	-	MCL-1	All three anti-apoptotic BCL2 proteins (BCL2, BCL-xL and MCL-1) sequesters BID into stable complexes, preventing them from activating BAX or BAK [242].
BID	Casp	8 or { Casp2 and n	ot (BCL2 or BCLXL or MCL-1) }
			nse to Caspase 8 activation. In addition, Caspase 2 can also once all three pro-apoptotic BCL2 family proteins are blocked.
BAK	←	BID	Activated (truncated) BID binds to mitochondrial BAK, resulting in its activation and oligomerization in the mitochondrial membrane, followed by cytochrome c release [243].
BAK	←	BIM	BAK is preferentially activated by BID compared to BIM, but BIM can also promote BAK oligomerization [244]
BAK	←	вік	BIK can aid the activation of both BAK and BAX by triggering BAK oligomerization on the ER membrane and promoting a Ca2+ efflux required for the fragmentation of hyper fused mitochondrial tubules, aiding BAK and BAX activation [245].
BAK	F	BCLXL	BCLXL binds BAK and prevent its oligomerization in the mitochondrial membrane [246-248].
BAK	F	MCL-1	MCL-1 binds BAK and prevent its oligomerization in the mitochondrial membrane [247,248].
BAK	F	BCL2	BCL2 can also bind BAK to prevent its oligomerization, but it does so less potently than the other two BCL-2 family members [247-249].
BAK	{BID and [BIM or BIK or not (BCL2 and BCLXL and MCL-1)]} or { (BIM or BIK) and not (BCLXL or MCL-1)}		
	Given that BAK is preferentially activated by BID compared to BIM [244] and that it is less responsive to sequestration by BCL2 than the other two anti-apoptotic BCL2 family proteins [247,248], BAK in our model turns on when stimulated by BID if one or more BCL2 family proteins are absent, or if BIM or BIK are also present. In contrast, BIM or BIK only activate BAK if BCLXL and MCL-1 are absent (BCL-2 alone cannot block them).		
ВАХ	←	ВІМ	Activated BIM binds to mitochondrial BAX, resulting in its allosteric activation and oligomerization in the mitochondrial membrane, leading to cytochrome c release [244].

Target Node	Link	Input	Description & References	
ВАХ	←	BID	BAX is preferentially activated by BIM compared to BID, but BID can also promote BAK oligomerization [244].	
ВАХ	←	вік	BIK can aid the activation of both BAK and BAX by triggering BAK oligomerization on the ER membrane and promoting a Ca2+ efflux required for the fragmentation of hyper fused mitochondrial tubules, aiding BAK and BAX activation [245].	
BAX	F	BCL2	BCL2 binds BAX and prevent its oligomerization in the mitochondrial membrane [247,248].	
BAX	F	BCLXL	BCL-xL binds BAK and prevent its oligomerization in the mitochondrial membrane [247,248].	
BAX	F	MCL-1	MCL-1 can also bind BAK to prevent its oligomerization, but it does so less potently than the other two BCL-2 family members [247,248,250].	
ВАХ		and [(BID or BIK) or BCL2 or BCLXL) }	or not (BCL2 and BCLXL and MCL-1)] } or { (BID or BIK) and	
	In contrast to BAK, BAX is preferentially activated by BIM compared to BID [244] and it is less responsive to sequestration by MCL-1 than the other two anti-apoptotic BCL2 family proteins [247,248]. BAX in our model turns on when stimulated by BIM if one or more BCL2 family proteins are absent, or if BID or BIK are also present. In contrast, BID or BIK only activate BAK if BCL2 and BCLXL are absent (MCL-1 alone cannot block them).			
Cyto_C	←	BAX	BAX oligomerization at the mitochondrial membrane triggers MOMP, which results in the release of cytochrome C from mitochondria [251,252].	
Cyto_C	←	ВАК	BAK oligomerization at the mitochondrial membrane triggers MOMP, which results in the release of cytochrome C from mitochondria [253].	
Cyto_C	BAX or BAK			
		Cytochrome C release from mitochondria requires the oligomerization of either BAK or BAX [253].		
SMAC	←	BAX	BAX oligomerization at the mitochondrial membrane triggers MOMP, which results in the release of cytochrome C from mitochondria [253,254].	
SMAC	←	ВАК	BAK oligomerization at the mitochondrial membrane triggers MOMP, which results in the release of cytochrome C from mitochondria [253,254].	
SMAC	BAX	or BAK		
	SMAC/Diablo release from mitochondria requires the oligomerization of either BAK or BAX [253,254].			
IAPs	←	AKT_H	cIAP-2 and XIAP are both transcriptionally up-regulated in response to strong PI3K/AKT1 activation [255]	
IAPs	F	SMAC	SMAC / Diablo binds tightly to IAP proteins and blocks their ability to inhibit Caspase 3 [256].	

Target Node	Link	Input	Description & References	
IAPs	not S	MAC or AKT_H		
	Inhibitor of Apoptosis Proteins (IAPs) are active in the absence of SMAC/Diablo inhibition, or following AKT_H mediated up regulation (this protection from SMAC requires peak or oncogenic AKT activity).			
Casp9	←	Casp3	Procaspase 9 is a direct cleavage target of Caspase 3 [203].	
Casp9	⊢	IAPs	XIAP, cIAP1 and cIAP2 inhibitor the the cytochrome c-induced activation of procaspase-9 [257].	
Casp9	←	Cyto_C	Cytochrome c binds to APAF-1 proteins, promoting their assembly into the apoptosome, a platform for procaspase 9 binding and cleavage into its active form [258].	
Casp9	Casp	3 or (not IAPs and	Cyto_C)	
		Procaspase 9 is cleaved into active Caspase 9 by Caspase 3, or by the apoptosome (which relies on Cytochrome C for its assembly) in the absence of IAP proteins.		
Casp3	-	IAPs	IAPs bind tightly to the active site of Caspase 3, keeping its activity in check [257,259].	
Casp3	←	Casp9	Active Caspase 9 cleaves Caspase 3 [260].	
Casp3	←	Casp8	Caspase 8 can cleave Caspase 3 [261], but full Caspase 3 activation also requires MOMP (potentially due to a need for IAP inhibition) [262].	
Casp3	←	Casp3	Once activated, Caspase 3 helps sustain its own activation by cleaving procaspase 8 and 6. Caspase 6, in turn, generates additional active caspase 8 and 9. Together they all sustains a continuing active pool of Caspase 3.	
Casp3	(Casp9 and Casp8) or [Casp3 and (Casp9 or Casp8)] or [not IAPs and (Casp9 or Casp8 or Casp3)]			
	Activation of Caspase 3 requires proteolytic cleavage of procaspase-3 by initiator caspases such as Caspase 9 or Caspase 8. In our model, cooperation of two of the three caspases (Casp9, Casp8, Casp3) is required in the presence of IAPS, which inhibit the proteolytic activity of Caspase 3 by bind tightly to its active site. In the absence of IAPS, either of the three caspases can cleave and activate Caspase 3.			
CAD	←	Casp3	Caspase 3 relives CAD inhibition by cleaving its inhibitor ICAD [263].	
CAD	←	Casp9	In addition of Caspase 3, CAD inhibition can also be relieved by ICAD cleavage by Caspase 7, which is a direct target of Caspase 9 [263].	
CAD	Casp	3 and Casp9		
	Caspase-activated DNase (CAD) is activated when its inhibition is released via the cleavage of ICAD (inhibitor of caspase-activated DNase). While Capsase 3 and 7 (a direct target of Caspase 9) can inhibit ICAD [263], in our model they are both required, as CAD = ON is represents terminal, irreversible apoptotic commitment, which is fully locked in when both Caspase 3 and 9 are on.			

References

- 1. Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. Cell. 2010;141: 1117–1134. doi:10.1016/j.cell.2010.06.011
- 2. Uzman A. Molecular Cell Biology (4th edition) Harvey Lodish, Arnold Berk, S. Lawrence Zipursky, Paul Matsudaira, David Baltimore and James Darnell; Freeman & Co., New York, NY, 2000, 1084 pp., ISBN 0-7167-3136-3. Biochemistry and Molecular Biology Education. 2001;29: 126–128. doi: 10.1016/S1470-8175(01)00023-6
- 3. Widmann C, Gibson S, Johnson GL. Caspase-dependent cleavage of signaling proteins during apoptosis. A turn-off mechanism for anti-apoptotic signals. J Biol Chem. 1998;273: 7141–7147.
- 4. Chang F, Steelman LS, Lee JT, Shelton JG, Navolanic PM, Blalock WL, et al. Signal transduction mediated by the Ras/Raf/MEK/ERK pathway from cytokine receptors to transcription factors: potential targeting for therapeutic intervention. Leukemia. 2003;17: 1263–1293. doi:10.1038/sj.leu. 2402945
- Liu P, Gan W, Chin YR, Ogura K, Guo J, Zhang J, et al. Ptdlns(3,4,5)P3-Dependent Activation of the mTORC2 Kinase Complex. Cancer Discov. 2015;5: 1194–1209. doi: 10.1158/2159-8290.CD-15-0460
- 6. Dibble CC, Asara JM, Manning BD. Characterization of Rictor phosphorylation sites reveals direct regulation of mTOR complex 2 by S6K1. Mol Cell Biol. 2009;29: 5657–5670. doi:10.1128/MCB. 00735-09
- 7. Rodriguez-Viciana P, Warne PH, Dhand R, Vanhaesebroeck B, Gout I, Fry MJ, et al. Phosphatidylinositol-3-OH kinase as a direct target of Ras. Nature. 1994;370: 527–532. doi: 10.1038/370527a0
- 8. Gupta S, Ramjaun AR, Haiko P, Wang Y, Warne PH, Nicke B, et al. Binding of ras to phosphoinositide 3-kinase p110alpha is required for ras-driven tumorigenesis in mice. Cell. 2007;129: 957–968. doi:10.1016/j.cell.2007.03.051
- Manning BD, Toker A. AKT/PKB Signaling: Navigating the Network. Cell. 2017;169: 381–405. doi: 10.1016/j.cell.2017.04.001
- 10. Hui RC-Y, Gomes AR, Constantinidou D, Costa JR, Karadedou CT, Fernández de Mattos S, et al. The forkhead transcription factor FOXO3a increases phosphoinositide-3 kinase/Akt activity in drug-resistant leukemic cells through induction of PIK3CA expression. Mol Cell Biol. 2008;28: 5886–5898. doi:10.1128/MCB.01265-07
- 11. Wang Z, Dang T, Liu T, Chen S, Li L, Huang S, et al. NEDD4L Protein Catalyzes Ubiquitination of PIK3CA Protein and Regulates PI3K-AKT Signaling. Journal of Biological Chemistry. 2016;291: 17467–17477. doi:10.1074/jbc.M116.726083
- 12. Yuan TL, Wulf G, Burga L, Cantley LC. Cell-to-Cell Variability in Pl3K Protein Level Regulates Pl3K-AKT Pathway Activity in Cell Populations. Curr Biol. 2011;21: 173–183. doi:10.1016/j.cub. 2010.12.047
- 13. Yang J-Y, Zong CS, Xia W, Yamaguchi H, Ding Q, Xie X, et al. ERK promotes tumorigenesis by inhibiting FOXO3a via MDM2-mediated degradation. Nat Cell Biol. 2008;10: 138–148. doi: 10.1038/ncb1676
- 14. Bucur O, Stancu AL, Muraru MS, Melet A, Petrescu SM, Khosravi-Far R. PLK1 is a binding partner and a negative regulator of FOXO3 tumor suppressor. Discoveries (Craiova). 2014;2: e16. doi:10.15190/d.2014.8
- 15. Cockcroft S, Thomas GM. Inositol-lipid-specific phospholipase C isoenzymes and their differential regulation by receptors. Biochem J. 1992;288 (Pt 1): 1–14.
- 16. Kim HK, Kim JW, Zilberstein A, Margolis B, Kim JG, Schlessinger J, et al. PDGF stimulation of inositol phospholipid hydrolysis requires PLC-gamma 1 phosphorylation on tyrosine residues 783 and 1254. Cell. 1991;65: 435–441.

- 17. Zhang W, Trible RP, Zhu M, Liu SK, McGlade CJ, Samelson LE. Association of Grb2, Gads, and phospholipase C-gamma 1 with phosphorylated LAT tyrosine residues. Effect of LAT tyrosine mutations on T cell angigen receptor-mediated signaling. J Biol Chem. 2000;275: 23355–23361. doi:10.1074/jbc.M000404200
- 18. Falasca M, Logan SK, Lehto VP, Baccante G, Lemmon MA, Schlessinger J. Activation of phospholipase C gamma by PI 3-kinase-induced PH domain-mediated membrane targeting. EMBO J. 1998;17: 414–422. doi:10.1093/emboj/17.2.414
- 19. Rameh LE, Rhee SG, Spokes K, Kazlauskas A, Cantley LC, Cantley LG. Phosphoinositide 3-kinase regulates phospholipase Cgamma-mediated calcium signaling. J Biol Chem. 1998;273: 23750–23757.
- 20. Gresset A, Sondek J, Harden TK. The phospholipase C isozymes and their regulation. Subcell Biochem. 2012;58: 61–94. doi:10.1007/978-94-007-3012-0 3
- 21. Michell RH, Kirk CJ, Jones LM, Downes CP, Creba JA. The stimulation of inositol lipid metabolism that accompanies calcium mobilization in stimulated cells: defined characteristics and unanswered questions. Philos Trans R Soc Lond, B, Biol Sci. 1981;296: 123–138.
- 22. Escobedo A, Gomes T, Aragón E, Martín-Malpartida P, Ruiz L, Macias MJ. Structural basis of the activation and degradation mechanisms of the E3 ubiquitin ligase Nedd4L. Structure. 2014;22: 1446–1457. doi:10.1016/j.str.2014.08.016
- 23. Yuan C, Wang L, Zhou L, Fu Z. The function of FOXO1 in the late phases of the cell cycle is suppressed by PLK1-mediated phosphorylation. Cell Cycle. 2014;13: 807–819. doi:10.4161/cc. 27727
- 24. Seoane J, Le H-V, Shen L, Anderson SA, Massagué J. Integration of Smad and forkhead pathways in the control of neuroepithelial and glioblastoma cell proliferation. Cell. 2004;117: 211–223.
- 25. Staller P, Peukert K, Kiermaier A, Seoane J, Lukas J, Karsunky H, et al. Repression of p15INK4b expression by Myc through association with Miz-1. Nature Cell Biology. 2001;3: 392–399. doi: 10.1038/35070076
- 26. Seoane J, Le H-V, Massagué J. Myc suppression of the p21(Cip1) Cdk inhibitor influences the outcome of the p53 response to DNA damage. Nature. 2002;419: 729–734. doi:10.1038/nature01119
- 27. Inoki K, Li Y, Zhu T, Wu J, Guan K-L. TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. Nature Cell Biology. 2002;4: 648–657. doi:10.1038/ncb839
- 28. Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. Cell. 2017;169: 361–371. doi:10.1016/j.cell.2017.03.035
- 29. Phosphorylation and Functional Inactivation of TSC2 by Erk: Implications for Tuberous Sclerosisand Cancer Pathogenesis. Cell. 2005;121: 179–193. doi:10.1016/j.cell.2005.02.031
- 30. Roux PP, Ballif BA, Anjum R, Gygi SP, Blenis J. Tumor-promoting phorbol esters and activated Ras inactivate the tuberous sclerosis tumor suppressor complex via p90 ribosomal S6 kinase. Proc Natl Acad Sci U S A. 2004;101: 13489–13494. doi:10.1073/pnas.0405659101
- 31. Haar EV, Lee S-I, Bandhakavi S, Griffin TJ, Kim D-H. Insulin signalling to mTOR mediated by the Akt/PKB substrate PRAS40. Nature Cell Biology. 2007;9: 316–323. doi:10.1038/ncb1547
- 32. Fonseca BD, Smith EM, Lee VH-Y, MacKintosh C, Proud CG. PRAS40 is a target for mammalian target of rapamycin complex 1 and is required for signaling downstream of this complex. J Biol Chem. 2007;282: 24514–24524. doi:10.1074/jbc.M704406200
- 33. Wiza C, Nascimento EBM, Ouwens DM. Role of PRAS40 in Akt and mTOR signaling in health and disease. Am J Physiol-Endoc M. 2012;302: E1453–60. doi:10.1152/ajpendo.00660.2011
- 34. Inoki K, Li Y, Xu T, Guan K-L. Rheb GTPase is a direct target of TSC2 GAP activity and regulates mTOR signaling. Genes & Development. 2003;17: 1829–1834. doi:10.1101/gad.1110003

- 35. Liu M, Clarke CJ, Salama MF, Choi YJ, Obeid LM, Hannun YA. Co-ordinated activation of classical and novel PKC isoforms is required for PMA-induced mTORC1 activation. PLoS ONE. 2017;12: e0184818. doi:10.1371/journal.pone.0184818
- 36. Demetriades C, Plescher M, Teleman AA. Lysosomal recruitment of TSC2 is a universal response to cellular stress. Nat Commun. 2016;7: 10662. doi:10.1038/ncomms10662
- 37. Martin R, Desponds C, Eren RO, Quadroni M, Thome M, Fasel N. Caspase-mediated cleavage of raptor participates in the inactivation of mTORC1 during cell death. Cell Death Discov. 2016;2: 16024. doi:10.1038/cddiscovery.2016.24
- 38. Long X, Lin Y, Ortiz-Vega S, Yonezawa K, Avruch J. Rheb Binds and Regulates the mTOR Kinase. Current Biology. 2005;15: 702–713. doi:10.1016/j.cub.2005.02.053
- 39. Real S, Meo-Evoli N, Espada L, Tauler A. E2F1 regulates cellular growth by mTORC1 signaling. PLoS ONE. 2011;6: e16163. doi:10.1371/journal.pone.0016163
- 40. Ramirez-Valle F, Badura ML, Braunstein S, Narasimhan M, Schneider RJ. Mitotic Raptor Promotes mTORC1 Activity, G2/M Cell Cycle Progression, and Internal Ribosome Entry Site-Mediated mRNA Translation. Molecular and Cellular Biology. 2010;30: 3151–3164. doi:10.1128/MCB. 00322-09
- 41. Proteolytic cleavage of p70 ribosomal S6 kinase by caspase-3 during DNA damage-induced apoptosis. Biochemistry. 2009;48: 1474–1480. doi:10.1021/bi801840s
- 42. Ma XM, Blenis J. Molecular mechanisms of mTOR-mediated translational control. Nat Rev Mol Cell Bio. 2009;10: 307–318. doi:10.1038/nrm2672
- 43. Bushell M, McKendrick L, Jänicke RU, Clemens MJ, Morley SJ. Caspase-3 is necessary and sufficient for cleavage of protein synthesis eukaryotic initiation factor 4G during apoptosis. FEBS Letters. 1999;451: 332–336.
- 44. Ding Q, Xia W, Liu J-C, Yang J-Y, Lee D-F, Xia J, et al. Erk associates with and primes GSK-3beta for its inactivation resulting in upregulation of beta-catenin. Mol Cell. 2005;19: 159–170. doi: 10.1016/j.molcel.2005.06.009
- 45. Zhang HH, Lipovsky AI, Dibble CC, Sahin M, Manning BD. S6K1 regulates GSK3 under conditions of mTOR-dependent feedback inhibition of Akt. Mol Cell. 2006;24: 185–197. doi:10.1016/j.molcel. 2006.09.019
- 46. Gervais JL, Seth P, Zhang H. Cleavage of CDK inhibitor p21(Cip1/Waf1) by caspases is an early event during DNA damage-induced apoptosis. J Biol Chem. 1998;273: 19207–19212.
- 47. Lu Z, Hunter T. Ubiquitylation and proteasomal degradation of the p21(Cip1), p27(Kip1) and p57(Kip2) CDK inhibitors. Cell Cycle. 2010;9: 2342–2352. doi:10.4161/cc.9.12.11988
- 48. Overton KW, Spencer SL, Noderer WL, Meyer T, Wang CL. Basal p21 controls population heterogeneity in cycling and quiescent cell cycle states. Proceedings of the National Academy of Sciences. National Acad Sciences; 2014;111: 201409797–E4393. doi:10.1073/pnas.1409797111
- 49. Fattman CL, Delach SM, Dou QP, Johnson DE. Sequential two-step cleavage of the retinoblastoma protein by caspase-3/-7 during etoposide-induced apoptosis. Oncogene. 2001;20: 2918–2926. doi:10.1038/sj.onc.1204414
- 50. Kato J, Matsushime H, Hiebert SW, Ewen ME, Sherr CJ. Direct binding of cyclin D to the retinoblastoma gene product (pRb) and pRb phosphorylation by the cyclin D-dependent kinase CDK4. Genes & Development. 1993;7: 331–342.
- 51. Novak B, Tyson JJ. A model for restriction point control of the mammalian cell cycle. J Theor Biol. 2004;230: 563–579. doi:10.1016/j.jtbi.2004.049
- 52. Zarkowska T, Mittnacht S. Differential phosphorylation of the retinoblastoma protein by G1/S cyclin-dependent kinases. J Biol Chem. 1997;272: 12738–12746.
- 53. Coqueret O. New roles for p21 and p27 cell-cycle inhibitors: a function for each cell compartment? Trends Cell Biol. 2003;13: 65–70.
- 54. Hinds PW, Mittnacht S, Dulic V, Arnold A, Reed SI, Weinberg RA. Regulation of retinoblastoma protein functions by ectopic expression of human cyclins. Cell. 1992;70: 993–1006.

- 55. Eymin B, Sordet O, Droin N, Munsch B, Haugg M, Van de Craen M, et al. Caspase-induced proteolysis of the cyclin-dependent kinase inhibitor p27Kip1 mediates its anti-apoptotic activity. Oncogene. 1999;18: 4839–4847. doi:10.1038/sj.onc.1202860
- 56. Levkau B, Koyama H, Raines EW, Clurman BE, Herren B, Orth K, et al. Cleavage of p21Cip1/Waf1 and p27Kip1 mediates apoptosis in endothelial cells through activation of Cdk2: role of a caspase cascade. Mol Cell. 1998;1: 553–563.
- 57. Montagnoli A, Fiore F, Eytan E, Carrano AC, Draetta GF, Hershko A, et al. Ubiquitination of p27 is regulated by Cdk-dependent phosphorylation and trimeric complex formation. Genes & Development. 1999;13: 1181–1189.
- 58. Ishida N, Hara T, Kamura T, Yoshida M, Nakayama K, Nakayama KI. Phosphorylation of p27Kip1 on serine 10 is required for its binding to CRM1 and nuclear export. J Biol Chem. 2002;277: 14355–14358. doi:10.1074/jbc.C100762200
- 59. Müller D, Bouchard C, Rudolph B, Steiner P, Stuckmann I, Saffrich R, et al. Cdk2-dependent phosphorylation of p27 facilitates its Myc-induced release from cyclin E/cdk2 complexes. Oncogene. 1997;15: 2561–2576. doi:10.1038/sj.onc.1201440
- 60. Sheaff RJ, Groudine M, Gordon M, Roberts JM, Clurman BE. Cyclin E-CDK2 is a regulator of p27Kip1. Genes & Development. 1997;11: 1464–1478.
- 61. Medema RH, Kops GJ, Bos JL, Burgering BM. AFX-like Forkhead transcription factors mediate cell-cycle regulation by Ras and PKB through p27kip1. Nature. 2000;404: 782–787. doi: 10.1038/35008115
- 62. Faure A, Naldi A, Chaouiya C, Thieffry D. Dynamical analysis of a generic Boolean model for the control of the mammalian cell cycle. Bioinformatics 2006;22: e124–31. doi:10.1093/bioinformatics/btl210
- 63. Sears R, Nuckolls F, Haura E, Taya Y, Tamai K, Nevins JR. Multiple Ras-dependent phosphorylation pathways regulate Myc protein stability. Genes & Development. 2000;14: 2501–2514. doi:10.1101/gad.836800
- 64. Lutterbach B, Hann SR. Hierarchical phosphorylation at N-terminal transformation-sensitive sites in c-Myc protein is regulated by mitogens and in mitosis. Molecular and Cellular Biology. 1994;14: 5510–5522.
- 65. Welcker M, Orian A, Jin J, Grim JE, Grim JA, Harper JW, et al. The Fbw7 tumor suppressor regulates glycogen synthase kinase 3 phosphorylation-dependent c-Myc protein degradation. Proc Natl Acad Sci U S A. 2004;101: 9085–9090. doi:10.1073/pnas.0402770101
- 66. Lin C-J, Malina A, Pelletier J. c-Myc and eIF4F constitute a feedforward loop that regulates cell growth: implications for anticancer therapy. Cancer Research. 2009;69: 7491–7494. doi: 10.1158/0008-5472.CAN-09-0813
- 67. Oswald F, Lovec H, Möröy T, Lipp M. E2F-dependent regulation of human MYC: trans-activation by cyclins D1 and A overrides tumour suppressor protein functions. Oncogene. 1994;9: 2029–2036. Available: http://europepmc.org/abstract/med/8208548
- 68. Thalmeier K, Synovzik H, Mertz R, Winnacker EL, Lipp M. Nuclear factor E2F mediates basic transcription and trans-activation by E1a of the human MYC promoter. Genes & Development. 1989;3: 527–536.
- 69. Batsché E, Lipp M, Cremisi C. Transcriptional repression and activation in the same cell type of the human c-MYC promoter by the retinoblastoma gene protein: antagonisation of both effects by SV40 T antigen. Oncogene. 1994;9: 2235–2243.
- 70. Roussel MF, Davis JN, Cleveland JL, Ghysdael J, Hiebert SW. Dual control of myc expression through a single DNA binding site targeted by ets family proteins and E2F-1. Oncogene. 1994;9: 405–415.
- 71. Stacey DW. Three Observations That Have Changed Our Understanding of Cyclin D1 and p27 in Cell Cycle Control. Genes Cancer. 2010;1: 1189–1199. doi:10.1177/1947601911403475

- 72. Diehl JA, Zindy F, Sherr CJ. Inhibition of cyclin D1 phosphorylation on threonine-286 prevents its rapid degradation via the ubiquitin-proteasome pathway. Genes & Development. 1997;11: 957–972.
- 73. Harper JW, Elledge SJ, Keyomarsi K, Dynlacht B, Tsai LH, Zhang P, et al. Inhibition of cyclin-dependent kinases by p21. Molecular Biology of the Cell. 1995;6: 387–400.
- 74. Xiong Y, Hannon GJ, Zhang H, Casso D, Kobayashi R, Beach D. p21 is a universal inhibitor of cyclin kinases. Nature. 1993;366: 701–704. doi:10.1038/366701a0
- 75. Guo Z-Y, Hao X-H, Tan F-F, Pei X, Shang L-M, Jiang X-L, et al. The elements of human cyclin D1 promoter and regulation involved. Clin Epigenetics. 2011;2: 63–76. doi:10.1007/s13148-010-0018-v
- 76. Fan J, Bertino JR. Functional roles of E2F in cell cycle regulation. Oncogene. 1997;14: 1191–1200. doi:10.1038/sj.onc.1200940
- 77. Chellappan SP, Hiebert S, Mudryj M, Horowitz JM, Nevins JR. The E2F transcription factor is a cellular target for the RB protein. Cell. 1991;65: 1053–1061. doi:10.1016/0092-8674(91)90557-F
- 78. Hitomi M, Stacey DW. Cyclin D1 production in cycling cells depends on ras in a cell-cycle-specific manner. Curr Biol. 1999;9: 1075–1084. doi:10.1016/S0960-9822(99)80476-X
- 79. Aktas H, Cai H, Cooper GM. Ras links growth factor signaling to the cell cycle machinery via regulation of cyclin D1 and the Cdk inhibitor p27KIP1. Molecular and Cellular Biology. 1997;17: 3850–3857.
- 80. Daksis JI, Lu RY, Facchini LM, Marhin WW, Penn LJ. Myc induces cyclin D1 expression in the absence of de novo protein synthesis and links mitogen-stimulated signal transduction to the cell cycle. Oncogene. 1994;9: 3635–3645.
- 81. Mateyak M, Obaya A, Sedivy J. c-Myc regulates cyclin D-Cdk4 and-Cdk6 activity but affects cell cycle progression at multiple independent points. Mol Cell Biol. 1999;19: 4672–4683.
- 82. Matsumura I, Tanaka H, Kanakura Y. E2F1 and c-Myc in cell growth and death. Cell Cycle. 2003;2: 333–338.
- 83. Diehl JA, Cheng M, Roussel MF, Sherr CJ. Glycogen synthase kinase-3beta regulates cyclin D1 proteolysis and subcellular localization. Genes & Development. 1998;12: 3499–3511. doi:10.1101/gad.12.22.3499
- 84. Leung JY, Ehmann GL, Giangrande PH, Nevins JR. A role for Myc in facilitating transcription activation by E2F1. Oncogene. 2008;27: 4172–4179. doi:10.1038/onc.2008.55
- 85. Krek W, Ewen ME, Shirodkar S, Arany Z, Kaelin WG, Livingston DM. Negative regulation of the growth-promoting transcription factor E2F-1 by a stably bound cyclin A-dependent protein kinase. Cell. 1994;78: 161–172.
- 86. Xu M, Sheppard KA, Peng CY, Yee AS, Piwnica-Worms H. Cyclin A/CDK2 binds directly to E2F-1 and inhibits the DNA-binding activity of E2F-1/DP-1 by phosphorylation. Mol Cell Biol. 1994;14: 8420–8431.
- 87. Weintraub SJ, Prater CA, Dean DC. Retinoblastoma protein switches the E2F site from positive to negative element. Nature. 1992;358: 259–261. doi:10.1038/358259a0
- 88. Johnson DG, Ohtani K, Nevins JR. Autoregulatory control of E2F1 expression in response to positive and negative regulators of cell cycle progression. Genes & Development. 1994;8: 1514–1525.
- 89. Leone G, DeGregori J, Sears R, Jakoi L, Nevins JR. Myc and Ras collaborate in inducing accumulation of active cyclin E/Cdk2 and E2F. Nature. 1997;387: 422–426. doi:10.1038/387422a0
- 90. Tanaka H, Matsumura I, Ezoe S, Satoh Y, Sakamaki T, Albanese C, et al. E2F1 and c-Myc potentiate apoptosis through inhibition of NF-kappaB activity that facilitates MnSOD-mediated ROS elimination. Mol Cell. 2002;9: 1017–1029.
- 91. Dong P, Maddali MV, Srimani JK, Thélot F, Nevins JR, Mathey-Prevot B, et al. Division of labour between Myc and G1 cyclins in cell cycle commitment and pace control. Nat Commun. 2014;5: 4750. doi:10.1038/ncomms5750

- 92. Ohtani K, DeGregori J, Nevins JR. Regulation of the cyclin E gene by transcription factor E2F1. Proc Natl Acad Sci U S A. 1995;92: 12146–12150. doi:10.1073/pnas.92.26.12146
- 93. Lunn CL, Chrivia JC, Baldassare JJ. Activation of Cdk2/Cyclin E complexes is dependent on the origin of replication licensing factor Cdc6 in mammalian cells. Cell Cycle. 2010;9: 4533–4541. doi: 10.4161/cc.9.22.13789
- 94. Geng Y, Lee Y-M, Welcker M, Swanger J, Zagozdzon A, Winer JD, et al. Kinase-independent function of cyclin E. Mol Cell. 2007;25: 127–139. doi:10.1016/j.molcel.2006.11.029
- 95. Helin K. Regulation of cell proliferation by the E2F transcription factors. Curr Opin Genet Dev. 1998;8: 28–35.
- 96. Sørensen CS, Syljuåsen RG, Falck J, Schroeder T, Rönnstrand L, Khanna KK, et al. Chk1 regulates the S phase checkpoint by coupling the physiological turnover and ionizing radiation-induced accelerated proteolysis of Cdc25A. Cancer Cell. 2003;3: 247–258.
- 97. Mazumder S, Gong B, Chen Q, Drazba JA, Buchsbaum JC, Almasan A. Proteolytic cleavage of cyclin E leads to inactivation of associated kinase activity and amplification of apoptosis in hematopoietic cells. Molecular and Cellular Biology. 2002;22: 2398–2409. doi:10.1128/MCB. 22.7.2398-2409.2002
- 98. Knudsen KE, Fribourg AF, Strobeck MW, Blanchard JM, Knudsen ES. Cyclin A is a functional target of retinoblastoma tumor suppressor protein-mediated cell cycle arrest. J Biol Chem. 1999;274: 27632–27641.
- 99. Kim I-M, Ackerson T, Ramakrishna S, Tretiakova M, Wang I-C, Kalin TV, et al. The Forkhead Box m1 transcription factor stimulates the proliferation of tumor cells during development of lung cancer. Cancer Res. 2006;66: 2153–2161. doi:10.1158/0008-5472.CAN-05-3003
- 100. Kalin TV, Wang I-C, Ackerson TJ, Major ML, Detrisac CJ, Kalinichenko VV, et al. Increased levels of the FoxM1 transcription factor accelerate development and progression of prostate carcinomas in both TRAMP and LADY transgenic mice. Cancer Res. 2006;66: 1712–1720. doi: 10.1158/0008-5472.CAN-05-3138
- 101. Laoukili J, Kooistra MRH, Brás A, Kauw J, Kerkhoven RM, Morrison A, et al. FoxM1 is required for execution of the mitotic programme and chromosome stability. Nature Cell Biology. 2005;7: 126–136. doi:10.1038/ncb1217
- 102. Alvarez-Fernández M, Halim VA, Krenning L, Aprelia M, Mohammed S, Heck AJ, et al. Recovery from a DNA-damage-induced G2 arrest requires Cdk-dependent activation of FoxM1. EMBO Rep. 2010;11: 452–458. doi:10.1038/embor.2010.46
- 103. Tategu M, Nakagawa H, Sasaki K, Yamauchi R, Sekimachi S, Suita Y, et al. Transcriptional regulation of human polo-like kinases and early mitotic inhibitor. J Genet Genomics. 2008;35: 215–224. doi:10.1016/S1673-8527(08)60030-2
- 104. Hsu JY, Reimann JDR, Sørensen CS, Lukas J, Jackson PK. E2F-dependent accumulation of hEmi1 regulates S phase entry by inhibiting APC(Cdh1). Nature Cell Biology. 2002;4: 358–366. doi:10.1038/ncb785
- 105. Lee J, Kim JA, Barbier V, Fotedar A, Fotedar R. DNA damage triggers p21WAF1-dependent Emi1 down-regulation that maintains G2 arrest. Mol Biol Cell. 2009;20: 1891–1902. doi:10.1091/mbc.e08-08-0818
- 106. Hansen DV, Loktev AV, Ban KH, Jackson PK. Plk1 regulates activation of the anaphase promoting complex by phosphorylating and triggering SCFbetaTrCP-dependent destruction of the APC Inhibitor Emi1. Molecular Biology of the Cell. 2004;15: 5623–5634. doi:10.1091/mbc.e04-07-0598
- 107. Pan H, Zhu Y, Wei W, Shao S, Rui X. Transcription factor FoxM1 is the downstream target of c-Myc and contributes to the development of prostate cancer. World J Surg Oncol. 2018;16: 59. doi: 10.1186/s12957-018-1352-3
- 108. Lüscher-Firzlaff JM, Lilischkis R, Lüscher B. Regulation of the transcription factor FOXM1c by Cyclin E/CDK2. FEBS Letters. 2006;580: 1716–1722. doi:10.1016/j.febslet.2006.02.021

- 109. Laoukili J, Alvarez M, Meijer LAT, Stahl M, Mohammed S, Kleij L, et al. Activation of FoxM1 during G2 requires cyclin A/Cdk-dependent relief of autorepression by the FoxM1 N-terminal domain. Mol Cell Biol. 2008;28: 3076–3087. doi:10.1128/MCB.01710-07
- 110. Sullivan C, Liu Y, Shen J, Curtis A, Newman C, Hock JM, et al. Novel interactions between FOXM1 and CDC25A regulate the cell cycle. PLoS ONE. 2012;7: e51277. doi:10.1371/journal.pone.0051277
- 111. Major ML, Lepe R, Costa RH. Forkhead box M1B transcriptional activity requires binding of Cdk-cyclin complexes for phosphorylation-dependent recruitment of p300/CBP coactivators.

 Molecular and Cellular Biology. 2004;24: 2649–2661. doi:10.1128/MCB.24.7.2649-2661.2004
- 112. Fu Z, Malureanu L, Huang J, Wang W, Li H, van Deursen JM, et al. Plk1-dependent phosphorylation of FoxM1 regulates a transcriptional programme required for mitotic progression. Nature Cell Biology. 2008;10: 1076–1082. doi:10.1038/ncb1767
- 113. Vigo E, Müller H, Prosperini E, Hateboer G, Cartwright P, Moroni MC, et al. CDC25A phosphatase is a target of E2F and is required for efficient E2F-induced S phase. Molecular and Cellular Biology. 1999;19: 6379–6395.
- 114. Wu L, Goodwin EC, Naeger LK, Vigo E, Galaktionov K, Helin K, et al. E2F-Rb complexes assemble and inhibit cdc25A transcription in cervical carcinoma cells following repression of human papillomavirus oncogene expression. Molecular and Cellular Biology. 2000;20: 7059–7067.
- 115. Donzelli M, Squatrito M, Ganoth D, Hershko A, Pagano M, Draetta GF. Dual mode of degradation of Cdc25 A phosphatase. EMBO J. 2002;21: 4875–4884. doi:10.1093/emboj/cdf491
- 116. Kang T, Wei Y, Honaker Y, Yamaguchi H, Appella E, Hung M-C, et al. GSK-3 beta targets Cdc25A for ubiquitin-mediated proteolysis, and GSK-3 beta inactivation correlates with Cdc25A overproduction in human cancers. Cancer Cell. 2008;13: 36–47. doi:10.1016/j.ccr.2007.12.002
- 117. Chen M-S, Ryan CE, Piwnica-Worms H. Chk1 kinase negatively regulates mitotic function of Cdc25A phosphatase through 14-3-3 binding. Molecular and Cellular Biology. 2003;23: 7488–7497. doi:10.1128/MCB.23.21.7488-7497.2003
- 118. Hoffmann I, Draetta G, Karsenti E. Activation of the phosphatase activity of human cdc25A by a cdk2-cyclin E dependent phosphorylation at the G1/S transition. EMBO J. 1994:13: 4302–4310.
- 119. Mazzolini L, Broban A, Froment C, Burlet-Schiltz O, Besson A, Manenti S, et al. Phosphorylation of CDC25A on SER283 in late S/G2 by CDK/cyclin complexes accelerates mitotic entry. Cell Cycle. 2016;15: 2742–2752. doi:10.1080/15384101.2016.1220455
- 120. Mailand N, Podtelejnikov AV, Groth A, Mann M, Bartek J, Lukas J. Regulation of G(2)/M events by Cdc25A through phosphorylation-dependent modulation of its stability. EMBO J. 2002;21: 5911–5920. doi:10.1093/emboj/cdf567
- 121. Geley S, Kramer E, Gieffers C, Gannon J, Peters JM, Hunt T. Anaphase-promoting complex/cyclosome-dependent proteolysis of human cyclin A starts at the beginning of mitosis and is not subject to the spindle assembly checkpoint. The Journal of Cell Biology. 2001;153: 137–148.
- 122. Elzen den N, Pines J. Cyclin A is destroyed in prometaphase and can delay chromosome alignment and anaphase. The Journal of Cell Biology. 2001;153: 121–136.
- 123. Di Fiore B, Pines J. How cyclin A destruction escapes the spindle assembly checkpoint. J Cell Biol. 2010;190: 501–509. doi:10.1083/jcb.201001083
- 124. Blomberg I, Hoffmann I. Ectopic expression of Cdc25A accelerates the G(1)/S transition and leads to premature activation of cyclin E- and cyclin A-dependent kinases. Mol Cell Biol. 1999;19: 6183–6194.
- 125. Timofeev O, Cizmecioglu O, Hu E, Orlik T, Hoffmann I. Human Cdc25A phosphatase has a non-redundant function in G2 phase by activating Cyclin A-dependent kinases. FEBS Letters. 2009;583: 841–847. doi:10.1016/j.febslet.2009.01.044
- 126. Rape M, Kirschner MW. Autonomous regulation of the anaphase-promoting complex couples mitosis to S-phase entry. Nature. 2004;432: 588–595. doi:10.1038/nature03023

- 127. Machida YJ, Dutta A. The APC/C inhibitor, Emi1, is essential for prevention of rereplication. Genes & Development. 2007;21: 184–194. doi:10.1101/gad.1495007
- 128. Reimann JD, Freed E, Hsu JY, Kramer ER, Peters JM, Jackson PK. Emi1 is a mitotic regulator that interacts with Cdc20 and inhibits the anaphase promoting complex. Cell. 2001;105: 645–655.
- 129. Reimann JD, Gardner BE, Margottin-Goguet F, Jackson PK. Emi1 regulates the anaphase-promoting complex by a different mechanism than Mad2 proteins. Genes & Development. 2001;15: 3278–3285. doi:10.1101/gad.945701
- 130. Zhou BB, Li H, Yuan J, Kirschner MW. Caspase-dependent activation of cyclin-dependent kinases during Fas-induced apoptosis in Jurkat cells. Proc Natl Acad Sci U S A. 1998;95: 6785–6790.
- 131. Lee J, Kumagai A, Dunphy WG. Positive regulation of Wee1 by Chk1 and 14-3-3 proteins. Molecular Biology of the Cell. 2001;12: 551–563. doi:10.1091/mbc.12.3.551
- 132. Domínguez-Kelly R, Martín Y, Koundrioukoff S, Tanenbaum ME, Smits VAJ, Medema RH, et al. Wee1 controls genomic stability during replication by regulating the Mus81-Eme1 endonuclease. J Cell Biol. 2011;194: 567–579. doi:10.1083/jcb.201101047
- 133. Watanabe N, Arai H, Iwasaki J-I, Shiina M, Ogata K, Hunter T, et al. Cyclin-dependent kinase (CDK) phosphorylation destabilizes somatic Wee1 via multiple pathways. Proc Natl Acad Sci U S A. 2005;102: 11663–11668. doi:10.1073/pnas.0500410102
- 134. Deibler RW, Kirschner MW. Quantitative reconstitution of mitotic CDK1 activation in somatic cell extracts. Mol Cell. 2010;37: 753–767. doi:10.1016/j.molcel.2010.02.023
- 135. Leung TW, Lin SS, Tsang AC, Tong CS, Ching JC, Leung WY, et al. Over-expression of FoxM1 stimulates cyclin B1 expression. FEBS Letters. 2001;507: 59–66.
- 136. Alvarez B, Martínez-A C, Burgering BM, Carrera AC. Forkhead transcription factors contribute to execution of the mitotic programme in mammals. Nature. 2001;413: 744–747. doi: 10.1038/35099574
- 137. Harper JW, Burton JL, Solomon MJ. The anaphase-promoting complex: it's not just for mitosis any more. Genes & Development. 2002;16: 2179–2206. doi:10.1101/gad.1013102
- 138. Wang I-C, Chen Y-J, Hughes D, Petrovic V, Major ML, Park HJ, et al. Forkhead box M1 regulates the transcriptional network of genes essential for mitotic progression and genes encoding the SCF (Skp2-Cks1) ubiquitin ligase. Molecular and Cellular Biology. 2005;25: 10875–10894. doi: 10.1128/MCB.25.24.10875-10894.2005
- 139. Dutertre S, Cazales M, Quaranta M, Froment C, Trabut V, Dozier C, et al. Phosphorylation of CDC25B by Aurora-A at the centrosome contributes to the G2-M transition. J Cell Sci. 2004;117: 2523–2531. doi:10.1242/jcs.01108
- 140. Dutertre S, Descamps S, Prigent C. On the role of aurora-A in centrosome function. Oncogene. 2002;21: 6175–6183. doi:10.1038/sj.onc.1205775
- 141. Lindon C, Pines J. Ordered proteolysis in anaphase inactivates Plk1 to contribute to proper mitotic exit in human cells. The Journal of Cell Biology. 2004;164: 233–241. doi:10.1083/jcb. 200309035
- 142. Kotani S, Tugendreich S, Fujii M, Jorgensen PM, Watanabe N, Hoog C, et al. PKA and MPF-activated polo-like kinase regulate anaphase-promoting complex activity and mitosis progression. Mol Cell. 1998;1: 371–380.
- 143. Qian YW, Erikson E, Li C, Maller JL. Activated polo-like kinase Plx1 is required at multiple points during mitosis in Xenopus laevis. Molecular and Cellular Biology. 1998;18: 4262–4271.
- 144. Thomas Y, Cirillo L, Panbianco C, Martino L, Tavernier N, Schwager F, et al. Cdk1 Phosphorylates SPAT-1/Bora to Promote Plk1 Activation in C. elegans and Human Cells. Cell Rep. 2016;15: 510–518. doi:10.1016/j.celrep.2016.03.049
- 145. Gheghiani L, Loew D, Lombard B, Mansfeld J, Gavet O. PLK1 Activation in Late G2 Sets Up Commitment to Mitosis. Cell Rep. 2017;19: 2060–2073. doi:10.1016/j.celrep.2017.05.031

- 146. Lobjois V, Jullien D, Bouché J-P, Ducommun B. The polo-like kinase 1 regulates CDC25B-dependent mitosis entry. Biochim Biophys Acta. 2009;1793: 462–468. doi:10.1016/j.bbamcr. 2008.12.015
- 147. Ouyang B, Li W, Pan H, Meadows J, Hoffmann I, Dai W. The physical association and phosphorylation of Cdc25C protein phosphatase by Prk. Oncogene. 1999;18: 6029–6036. doi: 10.1038/sj.onc.1202983
- 148. Cogswell JP, Brown CE, Bisi JE, Neill SD. Dominant-negative polo-like kinase 1 induces mitotic catastrophe independent of cdc25C function. Cell Growth Differ. 2000;11: 615–623.
- 149. De Souza CP, Ellem KA, Gabrielli BG. Centrosomal and cytoplasmic Cdc2/cyclin B1 activation precedes nuclear mitotic events. Experimental Cell Research. 2000;257: 11–21. doi:10.1006/excr. 2000.4872
- 150. Jackman M, Lindon C, Nigg EA, Pines J. Active cyclin B1-Cdk1 first appears on centrosomes in prophase. Nature Cell Biology. 2003;5: 143–148. doi:10.1038/ncb918
- 151. Lindqvist A, Källström H, Lundgren A, Barsoum E, Rosenthal CK. Cdc25B cooperates with Cdc25A to induce mitosis but has a unique role in activating cyclin B1-Cdk1 at the centrosome. The Journal of Cell Biology. 2005;171: 35–45. doi:10.1083/jcb.200503066
- 152. Lopez-Girona A, Furnari B, Mondesert O, Russell P. Nuclear localization of Cdc25 is regulated by DNA damage and a 14-3-3 protein. Nature. 1999;397: 172–175. doi:10.1038/16488
- 153. Schmitt E, Boutros R, Froment C, Monsarrat B, Ducommun B, Dozier C. CHK1 phosphorylates CDC25B during the cell cycle in the absence of DNA damage. J Cell Sci. 2006;119: 4269–4275. doi:10.1242/jcs.03200
- 154. Krämer A, Mailand N, Lukas C, Syljuåsen RG, Wilkinson CJ, Nigg EA, et al. Centrosome-associated Chk1 prevents premature activation of cyclin-B-Cdk1 kinase. Nature Cell Biology. 2004;6: 884–891. doi:10.1038/ncb1165
- 155. Hoffmann I, Clarke PR, Marcote MJ, Karsenti E, Draetta G. Phosphorylation and activation of human cdc25-C by cdc2--cyclin B and its involvement in the self-amplification of MPF at mitosis. EMBO J. 1993;12: 53–63.
- 156. Jackman MR, Pines JN. Cyclins and the G2/M transition. Cancer Surv. 1997;29: 47-73.
- 157. Heald R, McLoughlin M, McKeon F. Human wee1 maintains mitotic timing by protecting the nucleus from cytoplasmically activated Cdc2 kinase. Cell. 1993;74: 463–474.
- 158. Shteinberg M, Protopopov Y, Listovsky T, Brandeis M, Hershko A. Phosphorylation of the cyclosome is required for its stimulation by Fizzy/cdc20. Biochemical and Biophysical Research Communications. 1999;260: 193–198. doi:10.1006/bbrc.1999.0884
- 159. Golan A, Yudkovsky Y, Hershko A. The cyclin-ubiquitin ligase activity of cyclosome/APC is jointly activated by protein kinases Cdk1-cyclin B and Plk. J Biol Chem. 2002;277: 15552–15557. doi: 10.1074/jbc.M111476200
- 160. Manchado E, Eguren M, Malumbres M. The anaphase-promoting complex/cyclosome (APC/C): cell-cycle-dependent and -independent functions. Biochem Soc Trans. 2010;38: 65–71. doi: 10.1042/BST0380065
- 161. Peters J-M. The anaphase promoting complex/cyclosome: a machine designed to destroy. Nat Rev Mol Cell Biol. 2006;7: 644–656. doi:10.1038/nrm1988
- 162. D'Angiolella V, Mari C, Nocera D, Rametti L, Grieco D. The spindle checkpoint requires cyclin-dependent kinase activity. Genes & Development. 2003;17: 2520–2525. doi:10.1101/gad.267603
- 163. Hein JB, Nilsson J. Interphase APC/C-Cdc20 inhibition by cyclin A2-Cdk2 ensures efficient mitotic entry. Nat Commun. 2016;7: 10975. doi:10.1038/ncomms10975
- 164. Avram S, Mernea M, Mihailescu DF, Seiman CD, Seiman DD, Putz MV. Mitotic checkpoint proteins Mad1 and Mad2 structural and functional relationship with implication in genetic diseases. Curr Comput Aided Drug Des. 2014;10: 168–181.
- 165. Nezi L, Musacchio A. Sister chromatid tension and the spindle assembly checkpoint. Current opinion in cell biology. 2009;21: 785–795. doi:10.1016/j.ceb.2009.09.007

- 166. Thornton BR, Toczyski DP. Precise destruction: an emerging picture of the APC. Genes & Development. 2006;20: 3069–3078. doi:10.1101/gad.1478306
- 167. Kramer ER, Scheuringer N, Podtelejnikov AV, Mann M, Peters JM. Mitotic regulation of the APC activator proteins CDC20 and CDH1. Molecular Biology of the Cell. 2000;11: 1555–1569. doi: 10.1091/mbc.11.5.1555
- 168. Wang W, Kirschner MW. Emi1 preferentially inhibits ubiquitin chain elongation by the anaphase-promoting complex. Nat Cell Biol. 2013;15: 797–806. doi:10.1038/ncb2755
- 169. Ohtani K, DeGregori J, Leone G, Herendeen DR, Kelly TJ, Nevins JR. Expression of the HsOrc1 gene, a human ORC1 homolog, is regulated by cell proliferation via the E2F transcription factor. Molecular and Cellular Biology. 1996;16: 6977–6984.
- 170. DePamphilis ML, Blow JJ, Ghosh S, Saha T, Noguchi K, Vassilev A. Regulating the licensing of DNA replication origins in metazoa. Curr Opin Cell Biol. 2006;18: 231–239. doi:10.1016/j.ceb. 2006.04.001
- 171. Pelizon C, d'Adda di Fagagna F, Farrace L, Laskey RA. Human replication protein Cdc6 is selectively cleaved by caspase 3 during apoptosis. EMBO reports. 2002;3: 780–784. doi:10.1093/embo/reports/kvf161
- 172. Petersen BO, Lukas J, Sørensen CS, Bartek J, Helin K. Phosphorylation of mammalian CDC6 by cyclin A/CDK2 regulates its subcellular localization. EMBO J. 1999;18: 396–410. doi:10.1093/emboj/18.2.396
- 173. Yan Z, DeGregori J, Shohet R, Leone G, Stillman B, Nevins JR, et al. Cdc6 is regulated by E2F and is essential for DNA replication in mammalian cells. Proc Natl Acad Sci U S A. 1998;95: 3603–3608.
- 174. Yim H, Erikson RL. Cell division cycle 6, a mitotic substrate of polo-like kinase 1, regulates chromosomal segregation mediated by cyclin-dependent kinase 1 and separase. Proc Natl Acad Sci U S A. 2010;107: 19742–19747. doi:10.1073/pnas.1013557107
- 175. Valovka T, Schönfeld M, Raffeiner P, Breuker K, Dunzendorfer-Matt T, Hartl M, et al. Transcriptional control of DNA replication licensing by Myc. Sci Rep. 2013;3: 3444. doi:10.1038/srep03444
- 176. Yoshida K, Inoue I. Regulation of Geminin and Cdt1 expression by E2F transcription factors. Oncogene. 2004;23: 3802–3812. doi:10.1038/sj.onc.1207488
- 177. Yanagi K-I, Mizuno T, You Z, Hanaoka F. Mouse geminin inhibits not only Cdt1-MCM6 interactions but also a novel intrinsic Cdt1 DNA binding activity. J Biol Chem. 2002;277: 40871–40880. doi: 10.1074/jbc.M206202200
- 178. García-Higuera I, Manchado E, Dubus P, Cañamero M, Méndez J, Moreno S, et al. Genomic stability and tumour suppression by the APC/C cofactor Cdh1. Nat Cell Biol. 2008;10: 802–811. doi:10.1038/ncb1742
- 179. Clijsters L, Ogink J, Wolthuis R. The spindle checkpoint, APC/C(Cdc20), and APC/C(Cdh1) play distinct roles in connecting mitosis to S phase. J Cell Biol. 2013;201: 1013–1026. doi:10.1083/jcb. 201211019
- 180. Fragkos M, Ganier O, Coulombe P, Méchali M. DNA replication origin activation in space and time. Nat Rev Mol Cell Bio. 2015;16: 360–374. doi:10.1038/nrm4002
- 181. Ren B, Cam H, Takahashi Y, Volkert T, Terragni J, Young RA, et al. E2F integrates cell cycle progression with DNA repair, replication, and G(2)/M checkpoints. Genes & Development. 2002;16: 245–256. doi:10.1101/gad.949802
- 182. Cam H, Dynlacht BD. Emerging roles for E2F: beyond the G1/S transition and DNA replication. Cancer Cell. 2003;3: 311–316.
- 183. Coverley D, Laman H, Laskey RA. Distinct roles for cyclins E and A during DNA replication complex assembly and activation. Nature Cell Biology. 2002;4: 523–528. doi:10.1038/ncb813
- 184. Mailand N, Falck J, Lukas C, Syljuâsen RG, Welcker M, Bartek J, et al. Rapid destruction of human Cdc25A in response to DNA damage. Science. 2000;288: 1425–1429.

- 185. Donzelli M, Draetta GF. Regulating mammalian checkpoints through Cdc25 inactivation. EMBO reports. 2003;4: 671–677. doi:10.1038/sj.embor.embor887
- 186. Katsuno Y, Suzuki A, Sugimura K, Okumura K, Zineldeen DH, Shimada M, et al. Cyclin A-Cdk1 regulates the origin firing program in mammalian cells. Proc Natl Acad Sci U S A. 2009;106: 3184–3189. doi:10.1073/pnas.0809350106
- 187. Petermann E, Maya-Mendoza A, Zachos G, Gillespie DAF, Jackson DA, Caldecott KW. Chk1 requirement for high global rates of replication fork progression during normal vertebrate S phase. Molecular and Cellular Biology. 2006;26: 3319–3326. doi:10.1128/MCB.26.8.3319-3326.2006
- 188. Petronczki M, Glotzer M, Kraut N, Peters J-M. Polo-like kinase 1 triggers the initiation of cytokinesis in human cells by promoting recruitment of the RhoGEF Ect2 to the central spindle. Dev Cell. 2007;12: 713–725. doi:10.1016/j.devcel.2007.03.013
- 189. Burkard ME, Randall CL, Larochelle S, Zhang C, Shokat KM, Fisher RP, et al. Chemical genetics reveals the requirement for Polo-like kinase 1 activity in positioning RhoA and triggering cytokinesis in human cells. Proc Natl Acad Sci U S A. 2007;104: 4383–4388. doi:10.1073/pnas. 0701140104
- 190. Uhlmann F, Lottspeich F, Nasmyth K. Sister-chromatid separation at anaphase onset is promoted by cleavage of the cohesin subunit Scc1. Nature. 1999;400: 37–42. doi:10.1038/21831
- 191. Listovsky T, Sale JE. Sequestration of CDH1 by MAD2L2 prevents premature APC/C activation prior to anaphase onset. J Cell Biol. 2013;203: 87–100. doi:10.1083/jcb.201302060
- 192. Sánchez I, Dynlacht BD. New insights into cyclins, CDKs, and cell cycle control. Semin Cell Dev Biol. 2005;16: 311–321. doi:10.1016/j.semcdb.2005.02.007
- 193. Petronczki M, Lénárt P, Peters J-M. Polo on the Rise—from Mitotic Entry to Cytokinesis with Plk1. Dev Cell. 2008;14: 646–659. doi:10.1016/j.devcel.2008.04.014
- 194. Schmit TL, Zhong W, Setaluri V, Spiegelman VS, Ahmad N. Targeted depletion of Polo-like kinase (Plk) 1 through lentiviral shRNA or a small-molecule inhibitor causes mitotic catastrophe and induction of apoptosis in human melanoma cells. J Invest Dermatol. 2009;129: 2843–2853. doi: 10.1038/jid.2009.172
- 195. Vecchione A, Baldassarre G, Ishii H, Nicoloso MS, Belletti B, Petrocca F, et al. Fez1/Lzts1 absence impairs Cdk1/Cdc25C interaction during mitosis and predisposes mice to cancer development. Cancer Cell. 2007;11: 275–289. doi:10.1016/j.ccr.2007.01.014
- 196. Yuan Z, Becker EBE, Merlo P, Yamada T, DiBacco S, Konishi Y, et al. Activation of FOXO1 by Cdk1 in cycling cells and postmitotic neurons. Science. 2008;319: 1665–1668. doi:10.1126/science.1152337
- 197. Floyd S, Pines J, Lindon C. APC/C Cdh1 targets aurora kinase to control reorganization of the mitotic spindle at anaphase. Curr Biol. 2008;18: 1649–1658. doi:10.1016/j.cub.2008.09.058
- 198. Guicciardi ME, Gores GJ. Life and death by death receptors. The FASEB Journal. Federation of American Societies for Experimental Biology; 2009;23: 1625–1637. doi:10.1096/fj.08-111005
- 199. Pennarun B, Meijer A, de Vries EGE, Kleibeuker JH, Kruyt F, de Jong S. Playing the DISC: turning on TRAIL death receptor-mediated apoptosis in cancer. Biochim Biophys Acta. 2010;1805: 123–140. doi:10.1016/j.bbcan.2009.11.004
- 200. Slee EA, Harte MT, Kluck RM, Wolf BB, Casiano CA, Newmeyer DD, et al. Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. The Journal of Cell Biology. 1999;144: 281–292.
- 201. Cowling V, Downward J. Caspase-6 is the direct activator of caspase-8 in the cytochrome cinduced apoptosis pathway: absolute requirement for removal of caspase-6 prodomain. Cell Death and Differentiation. 2002;9: 1046–1056. doi:10.1038/sj.cdd.4401065
- 202. Li H, Bergeron L, Cryns V, Pasternack MS, Zhu H, Shi L, et al. Activation of caspase-2 in apoptosis. J Biol Chem. 1997;272: 21010–21017.

- 203. Slee EA, Adrain C, Martin SJ. Executioner caspase-3, -6, and -7 perform distinct, non-redundant roles during the demolition phase of apoptosis. J Biol Chem. 2001;276: 7320–7326. doi:10.1074/jbc.M008363200
- 204. Shah RB, Thompson R, Sidi S. A mitosis-sensing caspase activation platform? New insights into the PIDDosome. Molecular & Cellular Oncology. 2016;3: e1059921. doi: 10.1080/23723556.2015.1059921
- 205. Vogel C, Kienitz A, Müller R, Bastians H. The mitotic spindle checkpoint is a critical determinant for topoisomerase-based chemotherapy. J Biol Chem. 2005;280: 4025–4028. doi:10.1074/jbc.C400545200
- 206. Masuda A, Maeno K, Nakagawa T, Saito H, Takahashi T. Association between mitotic spindle checkpoint impairment and susceptibility to the induction of apoptosis by anti-microtubule agents in human lung cancers. Am J Pathol. 2003;163: 1109–1116. doi:10.1016/S0002-9440(10)63470-0
- 207. Andersen JL, Johnson CE, Freel CD, Parrish AB, Day JL, Buchakjian MR, et al. Restraint of apoptosis during mitosis through interdomain phosphorylation of caspase-2. The EMBO Journal. 2009;28: 3216–3227. doi:10.1038/emboj.2009.253
- 208. Sung E-S, Park K-J, Choi H-J, Kim C-H, Kim Y-S. The proteasome inhibitor MG132 potentiates TRAIL receptor agonist-induced apoptosis by stabilizing tBid and Bik in human head and neck squamous cell carcinoma cells. Experimental Cell Research. 2012;318: 1564–1576. doi:10.1016/j.yexcr.2012.04.003
- 209. Lin C-F, Tsai C-C, Huang W-C, Wang Y-C, Tseng P-C, Tsai T-T, et al. Glycogen Synthase Kinase-3β and Caspase-2 Mediate Ceramide- and Etoposide-Induced Apoptosis by Regulating the Lysosomal-Mitochondrial Axis. PLoS ONE. 2016;11: e0145460. doi:10.1371/journal.pone. 0145460
- 210. Wertz IE, Kusam S, Lam C, Okamoto T, Sandoval W, Anderson DJ, et al. Sensitivity to antitubulin chemotherapeutics is regulated by MCL1 and FBW7. Nature. 2011;471: 110–114. doi:10.1038/nature09779
- 211. Harley ME, Allan LA, Sanderson HS, Clarke PR. Phosphorylation of McI-1 by CDK1-cyclin B1 initiates its Cdc20-dependent destruction during mitotic arrest. The EMBO Journal. 2010;29: 2407–2420. doi:10.1038/emboj.2010.112
- 212. Croxton R, Ma Y, Song L, Haura EB, Cress WD. Direct repression of the Mcl-1 promoter by E2F1. Oncogene. 2002;21: 1359–1369. doi:10.1038/sj.onc.1205157
- 213. Ding Q, Huo L, Yang J-Y, Xia W, Wei Y, Liao Y, et al. Down-regulation of myeloid cell leukemia-1 through inhibiting Erk/Pin 1 pathway by sorafenib facilitates chemosensitization in breast cancer. Cancer Research. 2008;68: 6109–6117. doi:10.1158/0008-5472.CAN-08-0579
- 214. Townsend KJ, Trusty JL, Traupman MA, Eastman A, Craig RW. Expression of the antiapoptotic MCL1 gene product is regulated by a mitogen activated protein kinase-mediated pathway triggered through microtubule disruption and protein kinase C. Oncogene. 1998;17: 1223–1234. doi:10.1038/sj.onc.1202035
- 215. Maurer U, Charvet C, Wagman AS, Dejardin E, Green DR. Glycogen synthase kinase-3 regulates mitochondrial outer membrane permeabilization and apoptosis by destabilization of MCL-1. Mol Cell. 2006;21: 749–760. doi:10.1016/j.molcel.2006.02.009
- 216. Clem RJ, Cheng EH, Karp CL, Kirsch DG, Ueno K, Takahashi A, et al. Modulation of cell death by Bcl-XL through caspase interaction. Proc Natl Acad Sci U S A. 1998;95: 554–559.
- 217. Yang E, Zha J, Jockel J, Boise LH, Thompson CB, Korsmeyer SJ. Bad, a heterodimeric partner for Bcl-XL and Bcl-2, displaces Bax and promotes cell death. Cell. 1995;80: 285–291.
- 218. Gélinas C, White E. BH3-only proteins in control: specificity regulates MCL-1 and BAK-mediated apoptosis. Genes & Development. 2005;19: 1263–1268. doi:10.1101/gad.1326205
- 219. Bah N, Maillet L, Ryan J, Dubreil S, Gautier F, Letai A, et al. Bcl-xL controls a switch between cell death modes during mitotic arrest. Cell Death Dis. 2014;5: e1291–e1291. doi:10.1038/cddis. 2014.251

- 220. Weiß LM, Hugle M, Romero S, Fulda S. Synergistic induction of apoptosis by a polo-like kinase 1 inhibitor and microtubule-interfering drugs in Ewing sarcoma cells. Int J Cancer. 2016;138: 497–506. doi:10.1002/ijc.29725
- 221. Terrano DT, Upreti M, Chambers TC. Cyclin-dependent kinase 1-mediated Bcl-xL/Bcl-2 phosphorylation acts as a functional link coupling mitotic arrest and apoptosis. Mol Cell Biol. 2010;30: 640–656. doi:10.1128/MCB.00882-09
- 222. Zhou L, Cai X, Han X, Xu N, Chang DC. CDK1 switches mitotic arrest to apoptosis by phosphorylating Bcl-2/Bax family proteins during treatment with microtubule interfering agents. Cell Biol Int. 2014;38: 737–746. doi:10.1002/cbin.10259
- 223. Kirsch DG, Doseff A, Chau BN, Lim DS, de Souza-Pinto NC, Hansford R, et al. Caspase-3-dependent cleavage of Bcl-2 promotes release of cytochrome c. J Biol Chem. 1999;274: 21155–21161.
- 224. O'Connor L, Strasser A, O'Reilly LA, Hausmann G, Adams JM, Cory S, et al. Bim: a novel member of the Bcl-2 family that promotes apoptosis. EMBO J. 1998;17: 384–395. doi:10.1093/emboj/17.2.384
- 225. Elangovan B, Chinnadurai G. Functional dissection of the pro-apoptotic protein Bik. Heterodimerization with anti-apoptosis proteins is insufficient for induction of cell death. J Biol Chem. 1997;272: 24494–24498.
- 226. Eichhorn JM, Sakurikar N, Alford SE, Chu R, Chambers TC. Critical role of anti-apoptotic Bcl-2 protein phosphorylation in mitotic death. Cell Death Dis. 2013;4: e834–e834. doi:10.1038/cddis. 2013.360
- 227. Condorelli F, Salomoni P, Cotteret S, Cesi V, Srinivasula SM, Alnemri ES, et al. Caspase cleavage enhances the apoptosis-inducing effects of BAD. Molecular and Cellular Biology. 2001;21: 3025–3036. doi:10.1128/MCB.21.9.3025-3036.2001
- 228. Taghiyev AF, Guseva NV, Harada H, Knudson CM, Rokhlin OW, Cohen MB. Overexpression of BAD potentiates sensitivity to tumor necrosis factor-related apoptosis-inducing ligand treatment in the prostatic carcinoma cell line LNCaP. Mol Cancer Res. 2003;1: 500–507.
- 229. Datta SR, Dudek H, Tao X, Masters S, Fu H, Gotoh Y, et al. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. Cell. 1997;91: 231–241.
- 230. Fang X, Yu S, Eder A, Mao M, Bast RC, Boyd D, et al. Regulation of BAD phosphorylation at serine 112 by the Ras-mitogen-activated protein kinase pathway. Oncogene. 1999;18: 6635–6640. doi:10.1038/si.onc.1203076
- 231. Tan Y, Demeter MR, Ruan H, Comb MJ. BAD Ser-155 phosphorylation regulates BAD/Bcl-XL interaction and cell survival. J Biol Chem. 2000;275: 25865–25869. doi:10.1074/jbc.M004199200
- 232. Boyd JM, Gallo GJ, Elangovan B, Houghton AB, Malstrom S, Avery BJ, et al. Bik, a novel death-inducing protein shares a distinct sequence motif with Bcl-2 family proteins and interacts with viral and cellular survival-promoting proteins. Oncogene. 1995;11: 1921–1928.
- 233. Gillissen B, Essmann F, Hemmati PG, Richter A, Richter A, Oztop I, et al. McI-1 determines the Bax dependency of Nbk/Bik-induced apoptosis. J Cell Biol. 2007;179: 701–715. doi:10.1083/jcb. 200703040
- 234. Dijkers PF, Medema RH, Lammers JW, Koenderman L, Coffer PJ. Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the forkhead transcription factor FKHR-L1. Curr Biol. 2000;10: 1201–1204.
- 235. Hongisto V, Smeds N, Brecht S, Herdegen T, Courtney MJ, Coffey ET. Lithium blocks the c-Jun stress response and protects neurons via its action on glycogen synthase kinase 3. Molecular and Cellular Biology. 2003;23: 6027–6036. doi:10.1128/MCB.23.17.6027-6036.2003
- 236. Hughes R, Gilley J, Kristiansen M, Ham J. The MEK-ERK pathway negatively regulates bim expression through the 3' UTR in sympathetic neurons. BMC Neurosci. 2011;12: 69. doi: 10.1186/1471-2202-12-69

- 237. Gomez-Bougie P, Bataille R, Amiot M. The imbalance between Bim and McI-1 expression controls the survival of human myeloma cells. Eur J Immunol. 2004;34: 3156–3164. doi:10.1002/eji. 200424981
- 238. Yamada H, Tada-Oikawa S, Uchida A, Kawanishi S. TRAIL causes cleavage of bid by caspase-8 and loss of mitochondrial membrane potential resulting in apoptosis in BJAB cells. Biochemical and Biophysical Research Communications. 1999;265: 130–133. doi:10.1006/bbrc.1999.1641
- 239. Huang K, Zhang J, O'Neill KL, Gurumurthy CB, Quadros RM, Tu Y, et al. Cleavage by Caspase 8 and Mitochondrial Membrane Association Activate the BH3-only Protein Bid during TRAIL-induced Apoptosis. Journal of Biological Chemistry. 2016;291: 11843–11851. doi:10.1074/jbc.M115.711051
- 240. Li H, Zhu H, Xu CJ, Yuan J. Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. Cell. 1998;94: 491–501.
- 241. Upton J-P, Austgen K, Nishino M, Coakley KM, Hagen A, Han D, et al. Caspase-2 cleavage of BID is a critical apoptotic signal downstream of endoplasmic reticulum stress. Mol Cell Biol. 2008;28: 3943–3951. doi:10.1128/MCB.00013-08
- 242. Kim H, Rafiuddin-Shah M, Tu H-C, Jeffers JR, Zambetti GP, Hsieh JJ-D, et al. Hierarchical regulation of mitochondrion-dependent apoptosis by BCL-2 subfamilies. Nature Cell Biology. 2006;8: 1348–1358. doi:10.1038/ncb1499
- 243. Wei MC, Lindsten T, Mootha VK, Weiler S, Gross A, Ashiya M, et al. tBID, a membrane-targeted death ligand, oligomerizes BAK to release cytochrome c. Genes & Development. 2000;14: 2060–2071.
- 244. Sarosiek KA, Chi X, Bachman JA, Sims JJ, Montero J, Patel L, et al. BID preferentially activates BAK while BIM preferentially activates BAX, affecting chemotherapy response. Mol Cell. 2013;51: 751–765. doi:10.1016/j.molcel.2013.08.048
- 245. Mathai JP, Germain M, Shore GC. BH3-only BIK regulates BAX,BAK-dependent release of Ca2+ from endoplasmic reticulum stores and mitochondrial apoptosis during stress-induced cell death. J Biol Chem. 2005;280: 23829–23836. doi:10.1074/jbc.M500800200
- 246. Lee EF, Grabow S, Chappaz S, Dewson G, Hockings C, Kluck RM, et al. Physiological restraint of Bak by Bcl-xL is essential for cell survival. Genes & Development. 2016;30: 1240–1250. doi: 10.1101/gad.279414.116
- 247. Zhai D, Jin C, Huang Z, Satterthwait AC, Reed JC. Differential regulation of Bax and Bak by anti-apoptotic Bcl-2 family proteins Bcl-B and Mcl-1. J Biol Chem. 2008;283: 9580–9586. doi:10.1074/ibc.M708426200
- 248. Willis SN, Chen L, Dewson G, Wei A, Naik E, Fletcher JI, et al. Proapoptotic Bak is sequestered by Mcl-1 and Bcl-xL, but not Bcl-2, until displaced by BH3-only proteins. Genes & Development. 2005;19: 1294–1305. doi:10.1101/gad.1304105
- 249. Dai H, Meng XW, Lee S-H, Schneider PA, Kaufmann SH. Context-dependent Bcl-2/Bak interactions regulate lymphoid cell apoptosis. Journal of Biological Chemistry. 2009;284: 18311–18322. doi:10.1074/jbc.M109.004770
- 250. Germain M, Milburn J, Duronio V. MCL-1 inhibits BAX in the absence of MCL-1/BAX Interaction. J Biol Chem. 2008;283: 6384–6392. doi:10.1074/jbc.M707762200
- 251. Oltvai Z, MILLIMAN C, KORSMEYER S. Bcl-2 Heterodimerizes in-Vivo with a Conserved Homolog, Bax, That Accelerates Programmed Cell-Death. 1993;74: 609–619.
- 252. Manon S, Chaudhuri B, Guérin M. Release of cytochrome c and decrease of cytochrome c oxidase in Bax-expressing yeast cells, and prevention of these effects by coexpression of Bcl-xL. FEBS Letters. 1997;415: 29–32.
- 253. Wei MC, Zong WX, Cheng EH, Lindsten T, Panoutsakopoulou V, Ross AJ, et al. Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction and death. Science. 2001;292: 727–730. doi:10.1126/science.1059108

- 254. Zhou L, Chang DC. Dynamics and structure of the Bax-Bak complex responsible for releasing mitochondrial proteins during apoptosis. J Cell Sci. 2008;121: 2186–2196. doi:10.1242/jcs. 024703
- 255. Hu P, Han Z, Couvillon AD, Exton JH. Critical role of endogenous Akt/IAPs and MEK1/ERK pathways in counteracting endoplasmic reticulum stress-induced cell death. J Biol Chem. 2004;279: 49420–49429. doi:10.1074/jbc.M407700200
- 256. Du C, Fang M, Li Y, Li L, Wang X. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. Cell. 2000;102: 33–42.
- 257. Deveraux Q, Roy N, Stennicke H, Van Arsdale T, Zhou Q, Srinivasula S, et al. IAPs block apoptotic events induced by caspase-8 and cytochrome c by direct inhibition of distinct caspases. The EMBO Journal. 1998;17: 2215–2223. doi:10.1093/emboj/17.8.2215
- 258. Srinivasula SM, Ahmad M, Fernandes-Alnemri T, Alnemri ES. Autoactivation of procaspase-9 by Apaf-1-mediated oligomerization. Mol Cell. 1998;1: 949–957.
- 259. Riedl SJ, Renatus M, Schwarzenbacher R, Zhou Q, Sun C, Fesik SW, et al. Structural basis for the inhibition of caspase-3 by XIAP. 2001;104: 791–800.
- 260. Li P, Nijhawan D, Budihardjo I, Srinivasula S, Ahmad M, Alnemri E, et al. Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. 1997;91: 479–489.
- 261. Stennicke HR, Jürgensmeier JM, Shin H, Deveraux Q, Wolf BB, Yang X, et al. Pro-caspase-3 is a major physiologic target of caspase-8. J Biol Chem. 1998;273: 27084–27090.
- 262. Spencer SL, Gaudet S, Albeck JG, Burke JM, Sorger PK. Non-genetic origins of cell-to-cell variability in TRAIL-induced apoptosis. Nature. 2009;459: 428–432. doi:10.1038/nature08012
- 263. Wolf BB, Schuler M, Echeverri F, Green DR. Caspase-3 is the primary activator of apoptotic DNA fragmentation via DNA fragmentation factor-45/inhibitor of caspase-activated DNase inactivation. J Biol Chem. 1999;274: 30651–30656.