
Fungal orders within dominant phyla
Ascomycota
Pleosporales (16.16%)
Hypocreales (11.00%)
Unclassified order (9.96%)
Eurotiales (6.59%)
Microascales (4.54%)
Unclassified order (2.65%)
Mycosphaerellales (1.83%)
Other orders (1.62%)
Unclassified order (1.51%)
Chaetothyriales (1.26%)
Neophaeothecales (0.96%)
Cladosporiales (0.61%)
Thelebolales (0.57%)
Glomerellales (0.56%)
Xylariales (0.56%)
Basidiomycota
Unclassified order (10.90%)
Agaricales (4.23%)
Cystobasidiomycetes_ord_Incertae_sedis (2.94%)
Other orders (1.68%)
Agaricostilbales (0.88%)
Unclassified order (0.58%)
Microbotryomycetes_ord_Incertae_sedis (0.57%)
Ustilaginales (0.39%)
Tremellales (0.36%)
0
5
10
15
Unclassified fungi
Unclassified fungi (16.63%)
0
5
10
15
Mean relative abundance (%)
Class
Agaricomycetes
Agaricostilbomycetes
Cystobasidiomycetes
Dothideomycetes
Eurotiomycetes
Leotiomycetes
Microbotryomycetes
Other
Sordariomycetes
Tremellomycetes
Unclassified class
Ustilaginomycetes