



TRAUMA
Center


FALL RISK ASSESSMENT EVENT

Your Individual Results



Share these results with your
primary care provider, family
and friends.

Use them as a reference to see
how you are doing.



AGENDA

☐ Initial Surveys and Orthostatic Blood Pressure

First we will ask you to sit and fill out a few surveys. While you complete these surveys, we will start your first station, checking orthostatic blood pressures.

☐ Strength and Balance Stations

Next you will complete 3 strength and balance stations. The 4-stage balance test, the 30-second chair stand, and the timed up and go.

☐ Follow up Surveys

We will then ask you to make a fall prevention plan and complete an evaluation of the event. This is important so you remember all the great ideas you had and we can learn how to improve our events in the future.

☐ Optional Activities

- speak to an expert about your individual fall risks
 - enroll in our study about falls
 - visit our table for a snack, fall prevention supplies and more
 - sometimes we have guest stations available!
- 



STEADI Community Assessment Day SURVEY: PRE-assessment

Before you start today, please answer the following questions:

A. How would you rate your risk of falling? Please circle.

1. Your body's ability to maintain balance at all times at home:

1 2 3 4 5 6 7 8 9 10

Low Risk

High Risk

2. Your body's ability to maintain balance at all times outside of the home:

1 2 3 4 5 6 7 8 9 10

Low Risk

High Risk

B. What things have you done in the past to reduce your fall risks? Please circle.

1. I've talked to healthcare professionals about fall prevention in the past year.

Yes No Don't Remember

2. I've participated in fall prevention classes in the past.

Yes No Don't Remember

3. I've participated in fall prevention research in the past.

Yes No Don't Remember

4. I've participated in home safety evaluation in the past.

Yes No Don't Remember

C. What do you currently do now to reduce your fall risks? Please circle

1. I currently do exercises to improve my balance at least once a week (walking alone will not improve balance). Yes No

2. I walk or do other exercises or activities to keep me active.

Yes No

If yes, please describe:


What questions do you have about fall prevention today?

Short FES-I

Now we would like to ask some questions about how concerned you are about the possibility of falling. Please reply thinking about how you usually do the activity. If you currently don't do the activity, please answer to show whether you think you would be concerned about falling IF you did the activity. For each of the following activities, please tick the box which is closest to your own opinion to show how concerned you are that you might fall if you did this activity.

		Not at all concerned 1	Somewhat concerned 2	Fairly concerned 3	Very concerned 4
1	Getting dressed or undressed	☐ 1	☐ 2	☐ 3	☐ 4
2	Taking a bath or shower	☐ 1	☐ 2	☐ 3	☐ 4
3	Getting in or out of a chair	☐ 1	☐ 2	☐ 3	☐ 4
4	Going up or down stairs	☐ 1	☐ 2	☐ 3	☐ 4
5	Reaching for something above your head or on the ground	☐ 1	☐ 2	☐ 3	☐ 4
6	Walking up or down a slope	☐ 1	☐ 2	☐ 3	☐ 4
7	Going out to a social event (e.g. religious service, family gathering or club meeting)	☐ 1	☐ 2	☐ 3	☐ 4

STAY INDEPENDENT QUESTIONNAIRE



PLEASE CHECK "YES" OR "NO" FOR EACH BOLD STATEMENT BELOW. WHY IT MATTERS IS EXPLAINED AFTER THE QUESTION.

01	I HAVE FALLEN IN THE PAST YEAR. PEOPLE WHO HAVE FALLEN ONCE ARE LIKELY TO FALL AGAIN.	YES ☐	NO ☐
----	--	---------------------------------	--------------------------------

02	I USE OR HAVE BEEN ADVISED TO USE A CANE OR WALKER TO GET AROUND SAFELY. PEOPLE WHO HAVE BEEN ADVISED TO USE A CANE OR WALKER MAY ALREADY BE MORE LIKELY TO FALL.	YES ☐	NO ☐
----	---	---------------------------------	--------------------------------


03	SOMETIMES I FEEL UNSTEADY WHEN I AM WALKING. UNSTEADINESS OR NEEDING SUPPORT WHILE WALKING ARE SIGNS OF POOR BALANCE.	YES ☐	NO ☐
----	---	---------------------------------	--------------------------------

04	I STEADY MYSELF BY HOLDING ONTO FURNITURE WHEN WALKING AT HOME. THIS IS ALSO A SIGN OF POOR BALANCE.	YES ☐	NO ☐
----	--	---------------------------------	--------------------------------

05	I AM WORRIED ABOUT FALLING. PEOPLE WHO ARE WORRIED ABOUT FALLING ARE MORE LIKELY TO FALL.	YES ☐	NO ☐
----	--	---------------------------------	--------------------------------

06	I NEED TO PUSH WITH MY HANDS TO STAND UP FROM A CHAIR. THIS IS A SIGN OF WEAK LEG MUSCLES, A MAJOR REASON FOR FALLING.	YES ☐	NO ☐
----	---	---------------------------------	--------------------------------

07	I HAVE SOME TROUBLE STEPPING UP ONTO A CURB. THIS IS ALSO A SIGN OF WEAK LEG MUSCLES.	YES ☐	NO ☐
----	--	---------------------------------	--------------------------------



08

I OFTEN HAVE TO RUSH TO THE TOILET.

RUSHING TO THE BATHROOM,
ESPECIALLY AT NIGHT, INCREASES YOUR
CHANCE OF FALLING.

YES

☐

NO

☐

09

I HAVE LOST SOME FEELING IN MY FEET.

NUMBNESS IN YOUR FEET CAN CAUSE
STUMBLES AND LEAD TO FALLS.

YES

☐

NO

☐

10

**I TAKE MEDICINE THAT SOMETIMES
MAKES ME FEEL LIGHT-HEADED OR
MORE TIRED THAN USUAL.**

SIDE EFFECTS FROM MEDICINES CAN
SOMETIMES INCREASE YOUR CHANCE
OF FALLING.

YES

☐

NO

☐

11

**I TAKE MEDICINE TO HELP ME SLEEP
OR IMPROVE MY MOOD.**

THESE MEDICINES CAN SOMETIMES
INCREASE YOUR CHANCE OF FALLING.

YES

☐

NO

☐

12

I OFTEN FEEL SAD OR DEPRESSED.

SYMPTOMS OF DEPRESSION, SUCH AS
NOT FEELING WELL OR FEELING
SLOWED DOWN, ARE LINKED TO FALLS.

YES

☐

NO

☐

SCORE

EACH "YES" IS ONE POINT. TOTAL SCORE:

<4: MINIMAL OR NO RISK OF FALLING

4-8: MODERATE TO HIGH RISK OF FALLING

>8: HIGH TO SEVERE RISK OF FALLING

**PLEASE SHARE THIS QUESTIONNAIRE
WITH YOUR PRIMARY CARE PHYSICIAN.**

ASSESSMENT

Measuring Orthostatic Blood Pressure

1. Have the patient sit down for 5 minutes. Measure blood pressure and pulse rate.
2. Have the patient stand for 1 minute. Take blood pressure and pulse rate measurements after 1 minute.
3. Repeat blood pressure and pulse measurements at 3 minutes.

A drop in BP of ≥ 20 mm Hg, or in diastolic BP of ≥ 10 mm Hg, or experiencing lightheadedness or dizziness is considered abnormal.

POSITION	TIME	BP	ASSOCIATED SYMPTOMS
Sitting 	5 Mins.	BP ____ / ____ HR _____	
Standing 	1 Min.	BP ____ / ____ HR _____	
Standing 	3 Mins.	BP ____ / ____ HR _____	

CDC's STEADI tools and resources can help you screen, assess, and intervene to reduce your patient's fall risk. For more information, visit www.cdc.gov/steadi



**Centers for Disease
Control and Prevention**
National Center for Injury
Prevention and Control

2017

STEADI Stopping Elderly Accidents,
Deaths & Injuries

ASSESSMENT CONTINUED

The 4-Stage Balance Test

Instructions to the patient:

- I'm going to show you four positions.
- Try to stand in each position for 10 seconds.
- You can hold your arms out, or move your body to help keep your balance, but don't move your feet.
- For each position I will say, "Ready, begin." Then, I will start timing. After 10 seconds, I will say, "Stop."

	1 Stand with your feet side-by-side.	Time: _____ seconds
	2 Place the instep of one foot so it is touching the big toe of the other foot.	Time: _____ seconds
	3 Tandem stand: Place one foot in front of the other, heel touching toe.	Time: _____ seconds
	4 Stand on one foot.	Time: _____ seconds

Notes:

CDC's STEADI tools and resources can help you screen, assess, and intervene to reduce your patient's fall risk.
For more information, visit www.cdc.gov/steady

ASSESSMENT

30-Second Chair Stand


Purpose: To test leg strength and endurance

Equipment: A chair with a straight back without arm rests (seat 17" high), and a stopwatch.

1 Instruct the patient:

1. Sit in the middle of the chair.
2. Place your hands on the opposite shoulder crossed, at the wrists.
3. Keep your feet flat on the floor.
4. Keep your back straight, and keep your arms against your chest.
5. On "Go," rise to a full standing position, then sit back down again.
6. Repeat this for 30 seconds.

NOTE:
Stand next to the patient for safety.



2 On the word "Go," begin timing.

If the patient must use his/her arms to stand, stop the test.

Record "0" for the number and score.

3 Count the number of times the patient comes to a full standing position in 30 seconds.

If the patient is over halfway to a standing position when 30 seconds have elapsed, count it as a stand.

4 Record the number of times the patient stands in 30 seconds.

Number: _____

Score: _____

SCORING

Chair Stand Below Average Scores

AGE	MEN	WOMEN
60-64	< 14	< 12
65-69	< 12	< 11
70-74	< 12	< 10
75-79	< 11	< 10
80-84	< 10	< 9
85-89	< 8	< 8
90-94	< 7	< 4

A below average score indicates a risk for falls.

CDC's STEADI tools and resources can help you screen, assess, and intervene to reduce your patient's fall risk. For more information, visit www.cdc.gov/steadi



**Centers for Disease
Control and Prevention**
National Center for Injury
Prevention and Control

2017

STEADI Stopping Elderly Accidents,
Deaths & Injuries

ASSESSMENT

Timed Up & Go (TUG)

Purpose: To assess mobility

Equipment: A stopwatch

Directions: Patients wear their regular footwear and can use a walking aid, if needed. Begin by having the patient sit back in a standard arm chair and identify a line 3 meters, or 10 feet away, on the floor.

1 Instruct the patient:

When I say “Go,” I want you to:

1. Stand up from the chair.
2. Walk to the line on the floor at your normal pace.
3. Turn.
4. Walk back to the chair at your normal pace.
5. Sit down again.

NOTE:
Always stay by
the patient for
safety.

2 On the word “Go,” begin timing.

3 Stop timing after patient sits back down.

4 Record time.

Time in Seconds: _____

An older adult who takes ≥ 12 seconds to complete the TUG is at risk for falling.

CDC’s STEADI tools and resources can help you screen, assess, and intervene to reduce your patient’s fall risk. For more information, visit www.cdc.gov/steady

OBSERVATIONS

Observe the patient’s postural stability, gait, stride length, and sway.

Check all that apply:

- ☐ Slow tentative pace
- ☐ Loss of balance
- ☐ Short strides
- ☐ Little or no arm swing
- ☐ Steadying self on walls
- ☐ Shuffling
- ☐ En bloc turning
- ☐ Not using assistive

device properly


These changes may signify neurological problems that require further evaluation.



Centers for Disease
Control and Prevention
National Center for Injury
Prevention and Control

2017


STEADI Stopping Elderly Accidents,
Deaths & Injuries



EVENT NOTES

Please use this space to write down your fall prevention plans and reminders for yourself





STEADI Community Assessment Day SURVEY: POST-assessment

Thank you for participating today!

1. Please let us know which offerings were most helpful to you. **Please circle.**

- Fall risk assessment surveys
- Orthostatic blood pressure check
- 4-stage balance test
- 30-second chair stand
- Timed up & go
- Vision screening (if available)
- Pharmacy education (if available)
- Safety items for purchase: grab bars, nightlights
- Snacks

2. What is your fall prevention plan?

What?

How?

When?

3. What additional resources or changes would you like to see in a future event like this?

4. How did you hear about today's event?

5. Would you be interested in receiving more information about fall prevention and completing follow-up surveys to help us learn more about fall prevention?

Yes No

6. Would you like to be on the mailing list for future OHSU Trauma Program events?

Yes No

If yes, please provide your contact information. Thank you!

Name:

Date of birth:

Phone number:

Email:

Address:

Please tear off and return this page to event staff