

# Project 4, FYS 3150 / 4150, fall 2013

Odd Petter Sand and Nathalie Bonatout

November 2, 2013

All our source code can be found at our GitHub repository for this project:  
<https://github.com/NathalieB/Project4/>

## 1 Introduction

x

## 2 Theory and Technicalities

### 2.1 Closed form solution

We substitute

$$v(x, t) = u(x, t) - u_s(x) = u(x, t) + x - 1$$

where  $u_s(x) = 1 - x$  is the steady state solution that satisfies our boundary and initial conditions. We then set up the diffusion equation for  $v(x, t)$ :

$$\frac{\partial^2 v(x, t)}{\partial x^2} = \frac{\partial v(x, t)}{\partial t}$$

with known boundary conditions

$$v(0, t) = v(1, t) = 0 \quad t \geq 0.$$

The initial condition  $u(x, 0) = 0$  then becomes

$$v(x, 0) = u(x, 0) - u_s(x) = 0 - (1 - x) = x - 1 \quad 0 < x < 1.$$

The solution of this equation is known from pp. 313-314 in the lecture notes, using  $L = 1$ :

$$v(x, t) = \sum_{n=1}^{\infty} A_n \sin(n\pi x) e^{-n^2 \pi^2 t}.$$

We now find the Fourier series coefficients by partwise integration

$$\begin{aligned} A_n &= 2 \int_0^1 v(x, 0) \sin(n\pi x) dx = 2 \int_0^1 (x - 1) \sin(n\pi x) dx \\ &= 2 \left( \left[ -(x - 1) \frac{1}{n\pi} \cos(n\pi x) \right]_0^1 - \int_0^1 -\frac{1}{n\pi} \cos(n\pi x) dx \right) \\ &= \frac{2}{n\pi} \left( [(1 - x) \cos(n\pi x)]_0^1 + \left[ \frac{1}{n\pi} \sin(n\pi x) \right]_0^1 \right) \\ &= \frac{2}{n\pi} (1 + 0) = \frac{2}{n\pi} \end{aligned}$$

and finally, by substitution, our closed form solution is

$$u(x, t) = v(x, t) + u_s(x) = 1 - x + \sum_{n=1}^{\infty} \frac{2}{n\pi} \sin(n\pi x) e^{-n^2 \pi^2 t}.$$

## 2.2 Algorithms

### 2.2.1 Explicit scheme

input: nSteps (# of interior points), time

```
deltaX = 1.0 / (nSteps + 1)
alpha = 0.5
deltaT = alpha * deltaX ^ 2
tSteps = 1.0 / deltaT
```

define v, vNext

```

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    v[i] = v(x, 0)

for( t = 1; t <= tSteps)
    for( i = 1 —> nSteps )
        vNext[i] = (1 - 2 * alpha) * v[i]
        if( i > 0 ) : vNext[i] += alpha * v[i-1] // else += 0
        if( i < nSteps ) : vNext[i] += alpha * v[i+1] // else += 0
    v = vNext

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    u[i] = 1 - x + v[i]

output: u

```

### 2.2.2 Implicit scheme

```

input: nSteps (# of interior points), time, tSteps

deltaX = 1.0 / (nSteps + 1)
deltaT = 1.0 / tSteps
alpha = deltaT / (deltaX ^ 2)

define v, vNext

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    v[i] = v(x, 0)

a = -alpha // diagonal element
b = 1 + 2*alpha // off-diagonal element

for( t = 1; t <= tSteps)
    tridiagonalSolver(a, b, v, vNext)
    v = vNext

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    u[i] = 1 - x + v[i]

output: u

```

### 2.2.3 Crank-Nicholson scheme

```

input: nSteps (# of interior points), time, tSteps

deltaX = 1.0 / (nSteps + 1)
deltaT = 1.0 / tSteps
alpha = deltaT / (deltaX ^ 2)

define v, w

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    v[i] = v(x, 0)

a = 2 * (1 + alpha)      // diagonal element
b = -alpha               // off-diagonal element

for( t = 1; t <= tSteps)
    for( i = 0; i < nSteps )
        w[i] = 2 * (1 - alpha) * v[i]
        if( i > 0 ) : w[i] += alpha * v[i - 1] // else += 0
        if( i < nSteps - 1 ) : w[i] += alpha * v[i + 1] // else += 0
    tridiagonalSolver(a, b, w, v)

for( i = 0; i < nSteps)
    x = (i + 1) * deltaX
    u[i] = 1 - x + v[i]

output: u

```

### 2.3 Tridiagonal form of the implicit schemes

The methods to reformulate the problem into a tridiagonal matrix equation is described in great detail in pp. 308-312 in the lecture notes, so we will not reproduce them here.

In the case of the Crank-Nicholson scheme, it is easily seen by the definition of matrix addition that the sum of a diagonal matrix and a tridiagonal matrix  $(2\mathbf{I} + \alpha\mathbf{B})$  is also tridiagonal. We can then note that the known vector  $\mathbf{w}_{j-1} \equiv (2\mathbf{I} - \alpha\mathbf{B})\mathbf{v}_{j-1}$  is easily calculated in every step by

$$w_i = 2(1 - \alpha)v_i + \alpha(v_{i-1} + v_{i+1})$$

hence there is no reason to calculate the inverse matrix  $(2\mathbf{I} - \alpha\mathbf{B})^{-1}$  or demand that it be tridiagonal. The operation above is  $\mathcal{O}(n)$ , just as the tridiagonal solver itself, so this does not affect the scaling of the algorithm.

## **2.4 Truncation errors and stability**

Simply use the table from p. 350 in the slides.

If time: Do the Taylor expand for explicit scheme and find the actual error, not the order: p. 346-347 in the slides. (Can use just the result in the C-N case.)

For explanations of the stability criteria, we again refer to the lecture notes (p. 307-309, 312).

## **3 Results and analysis**

### **3.1 Explicit scheme**

x

### **3.2 Implicit scheme**

x

### **3.3 Cranck-Nicholson scheme**

x

## **4 Conclusion**

What we learned.

### **4.1 Critique**

x