

Econ 675 Assignment 1

Nathan Mather

October 1, 2018

Contents

1 Kernal Density Estimation	1
1.1 Part 1	1

1 Kernal Density Estimation

1.1 Part 1

Start by noting that

$$\hat{f}^{(r)}(x) = \frac{(-1)^s}{nh^{1+s}} \sum_{i=1}^n k^{(s)}\left(\frac{x_i - x}{h}\right)$$

Now taking the expectation of $\hat{f}^{(r)}(x)$ that we can apply the linearity of expectations to move the expectation inside the sum. Then we can use the i.i.d. assumption to show the sum is just n times the expectation. This leaves us with

$$E[\hat{f}^{(r)}(x)] = E\left[\frac{(-1)^s}{h^{1+s}} k^{(s)}\left(\frac{x_i - x}{h}\right)\right] = \int_{-\infty}^{\infty} \frac{(-1)^s}{h^{1+s}} k^{(s)}\left(\frac{z - x}{h}\right) f(z) dz$$

Where the second equality is just by the definition of the expectation. Next we use integration by parts. Note that

$$\int_{-\infty}^{\infty} \frac{(-1)^s}{h^{1+s}} k^{(s)}\left(\frac{z - x}{h}\right) f(z) dz = - \int_{-\infty}^{\infty} \frac{(-1)^s}{h^s} k^{(s-1)}\left(\frac{z - x}{h}\right) f^{(1)}(z) dz$$

Iterating this s times gives us

$$\int_{-\infty}^{\infty} \frac{(-1)^s}{h^{1+s}} k^{(s)}\left(\frac{z - x}{h}\right) f(z) dz = (-1)^s \int_{-\infty}^{\infty} \frac{(-1)^s}{h} k\left(\frac{z - x}{h}\right) f^{(s)}(z) dz = \int_{-\infty}^{\infty} \frac{1}{h} k\left(\frac{z - x}{h}\right) f^{(s)}(z) dz$$

Next we apply change of variables. let $u = \frac{z-x}{h}$ Note that $du = \frac{1}{h} dz$ so we get

$$\int_{-\infty}^{\infty} k(u) f^{(s)}(x + hu) du$$

Next we Taylor expand $f^{(s)}(x + hu)$ to the P^{th} order about x . Recall from properties of the kernal estimator that $\int_{-\infty}^{\infty} k(u) du = 1$ and that $\int_{-\infty}^{\infty} k(u) u^j du = 0$ for all $j \neq p$ This gives us

$$f^{(r)}(x) + \frac{1}{P!} f^{(s+P)}(x) h^P \int_{-\infty}^{\infty} k(u) u^p du + o(h^P) = f^{(r)}(x) + \frac{1}{P!} f^{(s+P)}(x) h^p \mu_P(k) + o(h^P)$$

which is the desired result.

Now for the variance recall again that

$$\hat{f}^{(r)}(x) = \frac{(-1)^s}{nh^{1+s}} \sum_{i=1}^n k^{(s)} \left(\frac{x_i - x}{h} \right)$$

So by the i.i.d. assumption we can get that

$$V \left(\hat{f}^{(r)}(x) \right) = \frac{1}{nh^{2+2s}} V \left(k^{(s)} \left(\frac{x_i - x}{h} \right) \right)$$