

## MATH 576 Homework 10

---

**Problem 1** Let  $G = \{10 \mid \{10 \mid -7\}\}$ . Compute  $m(G)$  and  $t(G)$ .

---

We can write

$$G = \{10 \mid \{10 \mid -7\}\} = \{10 \mid 10 + \{0 \mid -17\}\} = 10 + \{0 \mid \{0 \mid -17\}\} = 10 + +_{17}.$$

Then, we have

$$m(G) = m(10 + +_{17}) = m(10) + m(+_{17}) = 10 + 0 = 10,$$

and since  $t(10) = -1 \neq 0 = t(+_{17})$  we can write

$$t(G) = \max\{t(10), t(+_{17})\} = \max\{-1, 0\} = 0.$$

**Problem 2** Find the mean value and the temperature of the game

$$n + a \cdot \uparrow + * m \pm x_1 \dots \pm x_r,$$

where  $n$  is a number,  $a$  is an integer,  $m$  is a nonnegative integer, and  $x_1 > \dots > x_r > 0$  are numbers.

---

We have

$$\begin{aligned} m(n + a \cdot \uparrow + * m \pm x_1 \dots \pm x_r) &= m(n) + m(a \cdot \uparrow) + m(*m) + m(\pm x_1) + \dots + m(\pm x_r) \\ &= n + 0 + 0 + 0 + \dots + 0 \\ &= n. \end{aligned}$$

Also, we have  $t(a \cdot \uparrow + * m) = 0$  since  $a \cdot \uparrow + * m$  is an infinitesimal, so we have

$$\begin{aligned} t(n + a \cdot \uparrow + * m \pm x_1 \dots \pm x_r) &= \max\{t(n), t(a \cdot \uparrow + * m), t(\pm x_1), \dots, t(\pm x_r)\} \\ &= \max\{-1, 0, x_1, \dots, x_r\} \\ &= x_1. \end{aligned}$$

**Problem 3** Let  $x$  and  $y$  be two numbers with  $x > y \geq 0$ . Find examples of games  $G$  and  $H$  such that  $t(G) = t(H) = x$  and  $t(G + H) = y$ .

---

Consider  $G = \pm x$  and  $H = \pm x \pm y$ . Then  $t(G) = x$ , and since  $t(\pm x) = x \neq y = t(\pm y)$  we have

$$t(H) = t(\pm x \pm y) = \max\{t(\pm x), t(\pm y)\} = \max\{x, y\} = x.$$

Also, we have

$$t(G + H) = t(\pm x \pm y \pm x) = t(\pm y) = y.$$

□

**Problem 4** Let  $G_n$  be the position in Grundy's game with  $n$  tokens in a single heap (recall that in Grundy's game, the only legal move is to split the tokens in a heap into two nonempty heaps of unequal size). Prove that in misère play,  $G_3$ ,  $G_6$  and  $G_9$  are all  $\mathcal{P}$ -positions. (Hint: Proceed iteratively as you would in normal play, using the misère mex and misère nim addition rules.)

We can use the mex rule to write:

- $G_1 = G_2 = \{\} = *0$
- $G_3 = \{G_1 + G_2\} = \{*0 + *0\} = \{*0\} = *$
- $G_4 = \{G_1 + G_3\} = \{*0 + *\} = \{*\} = *0$
- $G_5 = \{G_1 + G_4, G_2 + G_3\} = \{*0 + *0, *0 + *\} = \{*0, *\} = *2$
- $G_6 = \{G_1 + G_5, G_2 + G_4\} = \{*0 + *2, *0 + *0\} = \{*0, *2\} = *$
- $G_7 = \{G_1 + G_6, G_2 + G_5, G_3 + G_4\} = \{*0 + *, *0 + *2, * + *0\} = \{*, *2, *\} = *0$
- $G_8 = \{G_1 + G_7, G_2 + G_6, G_3 + G_5\} = \{*0 + *0, *0 + *, * + *2\} = \{*0, *, *3\} = *2$
- $G_9 = \{G_1 + G_8, G_2 + G_7, G_3 + G_6, G_4 + G_5\} = \{*0 + *2, *0 + *0, * + *, *0 + *2\} = \{*2, *0, *0, *2\} = *.$

Since  $* = \{*0\}$  has all its options in  $\mathcal{N}$ ,  $* \in \mathcal{P}$ . Therefore, we have  $G_3 = G_6 = G_9 = * \in \mathcal{P}$ .  $\square$

**Problem 5** Let  $S_n$  be the position in Subtraction- $\{1, 4, 9\}$  with  $n$  tokens in a single heap. Compute the values in misère play of  $S_i$  for  $0 \leq i \leq 10$ .

We can use the mex rule to write:

- $S_0 = \{\} = *0$
- $S_1 = \{S_0\} = \{*0\} = *$
- $S_2 = \{S_1\} = \{*\} = *0$
- $S_3 = \{S_2\} = \{*0\} = *$
- $S_4 = \{S_0, S_3\} = \{*0, *\} = *2$
- $S_5 = \{S_1, S_4\} = \{*, *2\} = *0$
- $S_6 = \{S_2, S_5\} = \{*0, *0\} = \{*0\} = *$
- $S_7 = \{S_3, S_6\} = \{*, *0\} = *0$
- $S_8 = \{S_4, S_7\} = \{*2, *0\} = *$
- $S_9 = \{S_0, S_5, S_8\} = \{*0, *0, *\} = \{*0, *\} = *2$
- $S_{10} = \{S_1, S_6, S_9\} = \{*, *, *2\} = \{*, *2\} = *0.$