Weather Effects on the Efficiency of Photovoltaic Systems in Medellín, Colombia





Nathalia Correa Sánchez (1,2), Oscar J. Mesa Sánchez (2), Carlos D. Hoyos Ortíz (1,2)

1. Sistema de Alerta Temprana de Medellín y el Valle de Aburrá


2. Universidad Nacional de Colombia, sede Medellín

Introduction

6.3°N 6.2°N 0.2 1.2 2.2 3.2

Altitude [km]


 Daily average Irradiance in Medellín exceeds that of most of the cities worldwide with photovoltaic (PV) systems over buildings.




Colombia's energy matrix is 68% hydraulic, being vulnerable to periods with negative precipitation anomalies such as during El Niño

Urban areas consume most of the electricity but they could also generate a significant amount using PV systems in a household or industries, reducing the vulnerability of the energetic system


Reduction of surface solar radiation









Clouds and irradiance anomalies



-140.00 -120.00 -100.00 -80.00 -60.00 -40.00 -20.00 0.00 20.00 6 1 8 9 1012122314151611 6 1 8 9 1012122314151611







Irradiance anomalies composites

Amount of the hourly solar radiation that is being lost under the worst conditions.

Data & methods





GOES Band 2 Pixel Selection




Reflectances Thresholds


in contrasting locations of the Valley










Clearness (Kt) & clear sky (Kt*) indexes









Conclusions

- Clouds are the main limiting factor for solar radiation and are more frequent during April and May in the morning hours. The typical cloud forcing magnitude is approximately 200 W/m2.
- The slope of the adjustment line represents the efficiency of the solar panels at each point; according to this, the best performance is achieved at the west-center location.
- In all cases, the highest rates of variability of the Kt * index are between 0 and 0.3.

References

- [Data/information/map] obtained from the "Global Solar Atlas 2.0, a free, web-based application is developed and operated by the company Solargis s.r.o. on behalf of the World Bank Group, utilizing Solargis data, with funding provided by the Energy Sector Management Assistance Program (ESMAP). For additional information: https://globalsolaratlas.info
- Guzman Echavarria, G (2018). Análisis de la influencia del diseño urbano en la meteorología del Valle de
- Aburrá(Master Thesis). Universidad Nacional de Colombia, Medellín.
- Igbal, M. (2012). An introduction to solar radiation. Elsevier.
- Rossow, W. B., & Garder, L. C. (1993). Cloud detection using satellite measurements of infrared and visible radiances for ISCCP. Journal of climate, 6(12), 2341-2369.
- Weide Luiz, E (2018). Analysis of Cloud Cover Variability Using Geostationary Satellite Data(Doctoral Thesis). INPE, São José dos Campos.

Acknowledgments

This work was supported by Area Metropolitana de Medellín y del Valle de Aburrá, Municipio de Medellín, Grupo EPM, and ISAGEN under the contract CCT504 of 2019. Likewise, the Universidad Nacional de Colombia provided support to this work











nacorrreasa@unal.edu.co Nathalia Correa ncorreas.siata@gmail.com