

An Asymptotic Preserving scheme for the diffusion limit of a stochastic linear kinetic equation

Nathalie Ayi ⁽¹⁾ Erwan Faou ⁽²⁾

⁽¹⁾ Sorbonne Université

⁽²⁾ MINGuS Project, INRIA Bretagne Atlantique

6 Novembre 2018

Physical Context

Framework = Kinetic Theory.

Physical Context


Framework = **Kinetic Theory**.

Physical context = Models with *several levels of description* (example : gas).

Physical Context

Framework = **Kinetic Theory**.


Physical context = Models with *several levels of description* (example : gas).



Physical Context

Framework = Kinetic Theory.

Physical context = Models with *several levels of description* (example : gas).



Statistical point of view : we are interested in the evolution of *the density of particles* $f(t, x, v)$ where

Statistical point of view : we are interested in the evolution of *the density of particles* $f(t, x, v)$ where

t = time

x = position

v = velocity

Statistical point of view : we are interested in the evolution of *the density of particles* $f(t, x, v)$ where

t = time

x = position

v = velocity

For all infinitesimal volume $dx dv$ around the point (x, v) :

$f(t, x, v) dx dv =$ *number of particles* which have position x and velocity v at time t .

Statistical point of view : we are interested in the evolution of *the density of particles* $f(t, x, v)$ where

t = time

x = position

v = velocity

For all infinitesimal volume $dx dv$ around the point (x, v) :

$f(t, x, v) dx dv$ = *number of particles* which have position x and velocity v at time t .

Aim = **Numerical scheme** adapted to the passage from the **mesoscopic scale** to the **macroscopic one** by a diffusive limit for a **stochastic linear kinetic equation**.

Asymptotic Preserving schemes

Aim = A **numerical approximation** of the **kinetic equation** which **converges** to a **numerical approximation** of the **macroscopic equation** when passing to the limit.

Asymptotic Preserving schemes

Aim = A **numerical approximation** of the **kinetic equation** which **converges** to a **numerical approximation** of the **macroscopic equation** when passing to the limit.

$$\begin{array}{ccc} P^{\varepsilon,h} & \xrightarrow{h \rightarrow 0} & P^{\varepsilon} \\ \varepsilon \rightarrow 0 \downarrow & & \downarrow \varepsilon \rightarrow 0 \\ P^{0,h} & \xrightarrow{h \rightarrow 0} & P^0 \end{array}$$

Figure: Property of AP Schemes

Asymptotic Preserving schemes

Aim = A **numerical approximation** of the **kinetic equation** which **converges** to a **numerical approximation** of the **macroscopic equation** when passing to the limit.

$$\begin{array}{ccc} P^{\varepsilon,h} & \xrightarrow{h \rightarrow 0} & P^{\varepsilon} \\ \varepsilon \rightarrow 0 \downarrow & & \downarrow \varepsilon \rightarrow 0 \\ P^{0,h} & \xrightarrow{h \rightarrow 0} & P^0 \end{array}$$

Figure: Property of AP Schemes

Property = Good scheme for both the **kinetic** (i.e. $\varepsilon \approx 1$) and **diffusion regimes** (i.e. ε small).

Asymptotic Preserving schemes

Aim = A **numerical approximation** of the **kinetic equation** which **converges** to a **numerical approximation** of the **macroscopic equation** when passing to the limit.

$$\begin{array}{ccc} P^{\varepsilon,h} & \xrightarrow{h \rightarrow 0} & P^{\varepsilon} \\ \varepsilon \rightarrow 0 \downarrow & & \downarrow \varepsilon \rightarrow 0 \\ P^{0,h} & \xrightarrow{h \rightarrow 0} & P^0 \end{array}$$

Figure: Property of AP Schemes

Property = Good scheme for both the **kinetic** (i.e. $\varepsilon \approx 1$) and **diffusion regimes** (i.e. ε small).

Difficulty = Standard schemes $\rightsquigarrow h = \mathcal{O}(\varepsilon) \implies$ Computationally too expensive in “fluid” zone (i.e. ε small).

Asymptotic Preserving schemes

Aim = A **numerical approximation** of the **kinetic equation** which **converges** to a **numerical approximation** of the **macroscopic equation** when passing to the limit.

$$\begin{array}{ccc} P^{\varepsilon,h} & \xrightarrow{h \rightarrow 0} & P^{\varepsilon} \\ \varepsilon \rightarrow 0 \downarrow & & \downarrow \varepsilon \rightarrow 0 \\ P^{0,h} & \xrightarrow{h \rightarrow 0} & P^0 \end{array}$$

Figure: Property of AP Schemes

Property = Good scheme for both the **kinetic** (i.e. $\varepsilon \approx 1$) and **diffusion regimes** (i.e. ε small).

Difficulty = Standard schemes $\rightsquigarrow h = \mathcal{O}(\varepsilon) \implies$ Computationally too expensive in “fluid” zone (i.e. ε small).

\implies Construct a **scheme** for which h is **independent of ε** .

Context

A stochastic linear kinetic equation

$$df + \frac{1}{\varepsilon} v \partial_x f dt = \frac{\sigma}{\varepsilon^2} \mathcal{L} f dt + f \circ Q dW_t$$

whith $t \in [0, T]$, $x \in \mathbf{T}$, $v \in [-1, 1]$ and dW_t a **cylindrical Wiener process**.

Context

A stochastic linear kinetic equation

$$df + \frac{1}{\varepsilon} v \partial_x f dt = \frac{\sigma}{\varepsilon^2} \mathcal{L} f dt + f \circ Q dW_t$$

with $t \in [0, T]$, $x \in \mathbf{T}$, $v \in [-1, 1]$ and dW_t a **cylindrical Wiener process**.

Example = equation for the **radiative transfer**.

Left-hand side = **free transport**, linear operator \mathcal{L} = **interaction of the particles with the medium**, noise term = **random emission/absorption of particles**.

Context

A stochastic linear kinetic equation

$$df + \frac{1}{\varepsilon} v \partial_x f dt = \frac{\sigma}{\varepsilon^2} \mathcal{L} f dt + f \circ Q dW_t$$

with $t \in [0, T]$, $x \in \mathbf{T}$, $v \in [-1, 1]$ and dW_t a **cylindrical Wiener process**.

Example = equation for the **radiative transfer**.

Left-hand side = **free transport**, linear operator \mathcal{L} = **interaction of the particles with the medium**, noise term = **random emission/absorption of particles**.

$$dW_t = \sum_{k \geq 0} e_k d\beta_k(t)$$

with $(\beta_k)_{k \geq 0}$ **independent real-valued Brownian motions**, $(e_k)_{k \geq 0}$ an **orthonormal basis** of the Hilbert space $L^2(\mathbf{T})$, Q a linear **self-adjoint operator** on $L^2(\mathbf{T})$ such that

$$\sum_{k \geq 0} \|Q e_k\|_{L_x^\infty}^2 < +\infty,$$

σ a **function** which satisfies $0 < \sigma_m \leq \sigma(x) \leq \sigma_M$ for every x .

The linear operator

$$\mathcal{L}f(v) = \int_{-1}^1 s(v, v')(f(v') - f(v))dv',$$

with s such that $0 < s_m \leq s(v, v') \leq s_M$ for every $v, v' \in [-1, 1]$, s satisfies $\int_{-1}^1 s(v, v')dv' = 1$ and is symmetric: $s(v, v') = s(v', v)$.

The linear operator

$$\mathcal{L}f(v) = \int_{-1}^1 s(v, v')(f(v') - f(v))dv',$$

with s such that $0 < s_m \leq s(v, v') \leq s_M$ for every $v, v' \in [-1, 1]$, s satisfies $\int_{-1}^1 s(v, v')dv' = 1$ and is symmetric: $s(v, v') = s(v', v)$.

We define the **operator Π** such that

$$\Pi\phi = \frac{1}{2} \int_{-1}^1 \phi(v)dv.$$

The linear operator

$$\mathcal{L}f(v) = \int_{-1}^1 s(v, v')(f(v') - f(v))dv',$$

with s such that $0 < s_m \leq s(v, v') \leq s_M$ for every $v, v' \in [-1, 1]$, s satisfies $\int_{-1}^1 s(v, v')dv' = 1$ and is symmetric: $s(v, v') = s(v', v)$.

We define the **operator** Π such that

$$\Pi\phi = \frac{1}{2} \int_{-1}^1 \phi(v)dv.$$

\mathcal{L} satisfies the following **properties**:

- $\Pi(\mathcal{L}\phi) = 0$ for every $\phi \in L^2([-1, 1])$.
- The null space of \mathcal{L} is $\mathcal{N}(\mathcal{L}) = \{\phi = \Pi\phi\}$.
- The rank of \mathcal{L} is $\mathcal{R}(\mathcal{L}) = \mathcal{N}^\perp(\mathcal{L}) = \{\phi \text{ s.t. } \Pi\phi = 0\}$.
- \mathcal{L} admits a pseudo inverse from $\mathcal{N}^\perp(\mathcal{L})$ onto $\mathcal{N}^\perp(\mathcal{L})$ denoted by \mathcal{L}^{-1} .

Examples

The one-group transport equation

For

$$\mathcal{L}f = \int_{-1}^1 \frac{1}{2}(f(v') - f(v))dv' = \Pi f - f,$$

the **equation** becomes

$$df + \frac{1}{\varepsilon} v \partial_x f dt = \frac{\sigma}{\varepsilon^2} (\Pi f - f) dt + f \circ Q dW_t.$$

Examples

The one-group transport equation

For

$$\mathcal{L}f = \int_{-1}^1 \frac{1}{2}(f(v') - f(v))dv' = \Pi f - f,$$

the **equation** becomes

$$df + \frac{1}{\varepsilon} v \partial_x f dt = \frac{\sigma}{\varepsilon^2} (\Pi f - f) dt + f \circ Q dW_t.$$

The telegraph equation

With the velocity set $\{-1, 1\}$, dv the discrete Lebesgue measure, we denote $f(t, x, 1) := p(t, x)$ and $f(t, x, -1) := q(t, x)$. For $\sigma = 1$, the **equation** becomes

$$\begin{cases} dp + \frac{1}{\varepsilon} \partial_x p dt = \frac{1}{\varepsilon^2} \left(\frac{p+q}{2} - p \right) dt + p \circ Q dW_t \\ dq - \frac{1}{\varepsilon} \partial_x q dt = \frac{1}{\varepsilon^2} \left(\frac{p+q}{2} - q \right) dt + q \circ Q dW_t. \end{cases}$$

The diffusion limit

The macroscopic equation

$$d\rho + \partial_x(\kappa \partial_x \rho) dt = \rho \circ Q dW_t$$

with $\kappa(x) = \frac{\Pi(v\mathcal{L}^{-1}v)}{\sigma(x)}.$

The diffusion limit

The macroscopic equation

$$d\rho + \partial_x(\kappa \partial_x \rho) dt = \rho \circ Q dW_t$$

with $\kappa(x) = \frac{\Pi(v\mathcal{L}^{-1}v)}{\sigma(x)}$.

Hilbert expansion:

$$f(t, x, v) = f_0(t, x, v) + \varepsilon f_1(t, x, v) + \varepsilon^2 f_2(t, x, v) + \dots$$

The diffusion limit

The macroscopic equation

$$d\rho + \partial_x(\kappa \partial_x \rho) dt = \rho \circ Q dW_t$$

$$\text{with } \kappa(x) = \frac{\Pi(v\mathcal{L}^{-1}v)}{\sigma(x)}.$$

Hilbert expansion:

$$f(t, x, v) = f_0(t, x, v) + \varepsilon f_1(t, x, v) + \varepsilon^2 f_2(t, x, v) + \dots$$

Identifying the **same power** of ε :

$$\begin{aligned}\sigma \mathcal{L} f_0 &= 0 \quad \Rightarrow \quad f_0 = \Pi f_0 = \rho_0, \\ v \partial_x f_0 &= \sigma \mathcal{L} f_1 \quad \Rightarrow \quad f_1 = \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho_0 \right)\end{aligned}$$

$$df_0 = \sigma \mathcal{L} f_2 dt + f_0 \circ Q dW_t - v \partial_x \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho_0 \right) dt.$$

The diffusion limit

The macroscopic equation

$$d\rho + \partial_x(\kappa \partial_x \rho) dt = \rho \circ Q dW_t$$

$$\text{with } \kappa(x) = \frac{\Pi(v\mathcal{L}^{-1}v)}{\sigma(x)}.$$

Hilbert expansion:

$$f(t, x, v) = f_0(t, x, v) + \varepsilon f_1(t, x, v) + \varepsilon^2 f_2(t, x, v) + \dots$$

Identifying the **same power** of ε :

$$\begin{aligned}\sigma \mathcal{L} f_0 &= 0 \quad \Rightarrow \quad f_0 = \Pi f_0 = \rho_0, \\ v \partial_x f_0 &= \sigma \mathcal{L} f_1 \quad \Rightarrow \quad f_1 = \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho_0 \right)\end{aligned}$$

$$df_0 = \sigma \mathcal{L} f_2 dt + f_0 \circ Q dW_t - v \partial_x \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho_0 \right) dt.$$

We apply Π and obtain

$$d\rho_0 = \rho_0 \circ Q dW_t - \partial_x \left(\Pi \left[v \mathcal{L}^{-1} \left(\frac{v}{\sigma} \right) \right] \partial_x \rho_0 \right) dt.$$

The micro-macro decomposition

We introduce g such that $f = \rho + \varepsilon g$ with $\rho = \Pi(f) \Rightarrow g$ satisfies $\Pi(g) = 0$.

The micro-macro decomposition

We introduce g such that $f = \rho + \varepsilon g$ with $\rho = \Pi(f) \Rightarrow g$ satisfies $\Pi(g) = 0$.

The equation becomes

$$d(\rho + \varepsilon g) + \frac{1}{\varepsilon} v \partial_x (\rho + \varepsilon g) dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}(\rho + \varepsilon g) dt + (\rho + \varepsilon g) \circ Q dW_t. \quad (1)$$

The micro-macro decomposition

We introduce g such that $f = \rho + \varepsilon g$ with $\rho = \Pi(f) \Rightarrow g$ satisfies $\Pi(g) = 0$.

The equation becomes

$$d(\rho + \varepsilon g) + \frac{1}{\varepsilon} v \partial_x (\rho + \varepsilon g) dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}(\rho + \varepsilon g) dt + (\rho + \varepsilon g) \circ Q dW_t. \quad (1)$$

We apply Π and obtain

$$d\rho + \partial_x \Pi(vg) dt = \rho \circ Q dW_t. \quad (2)$$

The micro-macro decomposition

We introduce g such that $f = \rho + \varepsilon g$ with $\rho = \Pi(f) \Rightarrow g$ satisfies $\Pi(g) = 0$.

The equation becomes

$$d(\rho + \varepsilon g) + \frac{1}{\varepsilon} v \partial_x (\rho + \varepsilon g) dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}(\rho + \varepsilon g) dt + (\rho + \varepsilon g) \circ Q dW_t. \quad (1)$$

We apply Π and obtain

$$d\rho + \partial_x \Pi(vg) dt = \rho \circ Q dW_t. \quad (2)$$

We do (1) - (2) and obtain

$$dg + \frac{1}{\varepsilon} (I - \Pi)(v \partial_x g) dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}g dt + g \circ Q dW_t - \frac{1}{\varepsilon^2} v \partial_x \rho dt.$$

The micro-macro decomposition

The micro-macro formulation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2}\mathcal{L}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt, \end{cases}$$

\Rightarrow system which is equivalent to our kinetic equation.

The micro-macro decomposition

The micro-macro formulation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2} v\partial_x \rho dt, \end{cases}$$

⇒ system which is equivalent to our kinetic equation.

Advantages = The decomposition **only uses basic properties** of the collision operator common to most of kinetic equations

The micro-macro decomposition

The micro-macro formulation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2}\mathcal{L}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt, \end{cases}$$

⇒ system which is equivalent to our kinetic equation.

Advantages = The decomposition **only uses basic properties** of the collision operator common to most of kinetic equations, approach which works for **both diffusive and hydrodynamical limits**.

Examples

The one group transport equation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = -\frac{\sigma}{\varepsilon^2}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt. \end{cases}$$

Examples

The one group transport equation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = -\frac{\sigma}{\varepsilon^2}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt. \end{cases}$$

The telegraph equation

We denote $g = (\alpha, \gamma)$.

$$\begin{cases} d\rho + \partial_x \frac{\alpha - \gamma}{2}dt = \rho \circ QdW_t \\ d\alpha + \frac{1}{\varepsilon}\partial_x \frac{\alpha + \gamma}{2}dt = -\frac{1}{\varepsilon^2}\alpha dt + \alpha \circ QdW_t - \frac{1}{\varepsilon^2}\partial_x \rho dt \\ d\gamma - \frac{1}{\varepsilon}\partial_x \frac{\gamma + \alpha}{2}dt = -\frac{1}{\varepsilon^2}\gamma dt + \gamma \circ QdW_t + \frac{1}{\varepsilon^2}\partial_x \rho dt. \end{cases}$$

Formal analytical limit

$$\bullet \, dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2}\mathcal{L}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt$$

$$\Rightarrow g = \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho \right) + \mathcal{O}(\varepsilon).$$

Formal analytical limit

$$\bullet dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2}\mathcal{L}gdt + g \circ QdW_t - \frac{1}{\varepsilon^2}v\partial_x \rho dt$$

$$\Rightarrow g = \mathcal{L}^{-1} \left(\frac{1}{\sigma} v \partial_x \rho \right) + \mathcal{O}(\varepsilon).$$

$$\bullet d\rho + \partial_x \Pi(vg)dt = \rho \circ QdW_t$$

$$\Rightarrow d\rho + \partial_x (\kappa \partial_x \rho)dt = \rho \circ QdW_t + \mathcal{O}(\varepsilon)$$

$$\text{with } \kappa(x) = \frac{\Pi(v\mathcal{L}^{-1}v)}{\sigma(x)}.$$

Construction of the numerical scheme

Let us deal with the stiff term for the one-group transport equation with $\sigma = 1$.
We are interested in

$$\partial_t g = -\frac{1}{\varepsilon^2} g.$$

Construction of the numerical scheme

Let us deal with the stiff term for the one-group transport equation with $\sigma = 1$. We are interested in

$$\partial_t g = -\frac{1}{\varepsilon^2} g.$$

With **Explicit Euler**, we obtain

$$\frac{g^{n+1} - g^n}{\Delta t} = -\frac{1}{\varepsilon^2} g^n$$

$$\Rightarrow g^{n+1} = \left(1 - \frac{\Delta t}{\varepsilon^2}\right) g^n = \left(1 - \frac{\Delta t}{\varepsilon^2}\right)^{n+1} g^0.$$

Stability $\Rightarrow \Delta t \leq \varepsilon^2 \Rightarrow$ **Problem for ε small.**

Construction of the numerical scheme

Let us deal with the stiff term for the one-group transport equation with $\sigma = 1$.
We are interested in

$$\partial_t g = -\frac{1}{\varepsilon^2} g.$$

With **Explicit Euler**, we obtain

$$\frac{g^{n+1} - g^n}{\Delta t} = -\frac{1}{\varepsilon^2} g^n$$

$$\Rightarrow g^{n+1} = \left(1 - \frac{\Delta t}{\varepsilon^2}\right) g^n = \left(1 - \frac{\Delta t}{\varepsilon^2}\right)^{n+1} g^0.$$

Stability $\Rightarrow \Delta t \leq \varepsilon^2 \Rightarrow$ **Problem for ε small.**

With **Implicit Euler**, we obtain

$$\frac{g^{n+1} - g^n}{\Delta t} = -\frac{1}{\varepsilon^2} g^{n+1}$$

$$\Rightarrow g^{n+1} = \left(\frac{\varepsilon^2}{\varepsilon^2 + \Delta t}\right) g^n \Rightarrow \text{Stability issues independent of the size } \varepsilon.$$

The micro-macro formulation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho QdW_t + \frac{1}{2}\rho \sum_{k \geq 0} (Qe_k)^2 dt \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}gdt + gQdW_t + \frac{1}{2}g \sum_{k \geq 0} (Qe_k)^2 dt - \frac{1}{\varepsilon^2} v\partial_x \rho dt. \end{cases}$$

Implicitation of the collision term, **Upwind discretization** of $(I - \Pi)(v\partial_x g)$, **centered approximations** of $\partial_x \Pi(vg)$ and $v\partial_x \rho$.

The micro-macro formulation

$$\begin{cases} d\rho + \partial_x \Pi(vg)dt = \rho QdW_t + \frac{1}{2}\rho \sum_{k \geq 0} (Qe_k)^2 dt \\ dg + \frac{1}{\varepsilon}(I - \Pi)(v\partial_x g)dt = \frac{\sigma}{\varepsilon^2} \mathcal{L}gdt + gQdW_t + \frac{1}{2}g \sum_{k \geq 0} (Qe_k)^2 dt - \frac{1}{\varepsilon^2} v\partial_x \rho dt. \end{cases}$$

Implicitation of the collision term, **Upwind discretization** of $(I - \Pi)(v\partial_x g)$, **centered approximations** of $\partial_x \Pi(vg)$ and $v\partial_x \rho$.

Notations:

- Time: $t_n = n\Delta t$ with Δt time step.
- Space: Two staggered grids of step Δx with nodes $x_i = i\Delta x$ and $x_{i+\frac{1}{2}} = (i + \frac{1}{2})\Delta x$ extended by periodicity.

$$\rho_i^n \approx \rho(t_n, x_i) \text{ and } g_{i+\frac{1}{2}}^n(v) \approx g(t_n, x_{i+\frac{1}{2}}, v)$$

Numerical scheme

$$\left\{ \begin{array}{l} \rho_i^{n+1} = \rho_i^n - \Delta t \Pi \left(v \frac{g_{i+\frac{1}{2}}^{n+1} - g_{i-\frac{1}{2}}^{n+1}}{\Delta x} \right) + \rho_i^n \left(\frac{\Delta t}{2} \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right) \\ g_{i+\frac{1}{2}}^{n+1} = g_{i+\frac{1}{2}}^n - \frac{\Delta t}{\varepsilon \Delta x} (I - \Pi) \left(v^+ \left(g_{i+\frac{1}{2}}^n - g_{i-\frac{1}{2}}^n \right) + v^- \left(g_{i+\frac{3}{2}}^n - g_{i+\frac{1}{2}}^n \right) \right) \\ \quad - \frac{\sigma_{i+\frac{1}{2}}}{\varepsilon^2} \mathcal{L} g_{i+\frac{1}{2}}^{n+1} \Delta t + g_{i+\frac{1}{2}}^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{i+\frac{1}{2},k})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{i+\frac{1}{2},k} \xi_k^{n+1} \right) \\ \quad - \frac{1}{\varepsilon^2} v \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} \Delta t \end{array} \right.$$

with $v^+ = \max(v, 0)$ and $v^- = \min(v, 0)$, $(\xi_k^n)_{n \geq 1, k \geq 0}$ i.i.d. with a normal distribution, $b_{ik} := Qe_k(x_i)$ and $b_{i+\frac{1}{2},k} := Qe_k(x_{i+\frac{1}{2}})$.

Formal numerical limit

$$\bullet g_{i+\frac{1}{2}}^{n+1} = \frac{1}{\sigma_{i+\frac{1}{2}}} \mathcal{L}^{-1} \left(v \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} \right) + \mathcal{O}(\varepsilon)$$

Formal numerical limit

$$\bullet g_{i+\frac{1}{2}}^{n+1} = \frac{1}{\sigma_{i+\frac{1}{2}}} \mathcal{L}^{-1} \left(v \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} \right) + \mathcal{O}(\varepsilon)$$

$$\begin{aligned} \Rightarrow \rho_i^{n+1} = \rho_i^n - \frac{\Delta t}{\Delta x} & \left(\kappa_{i+\frac{1}{2}} \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} - \kappa_{i-\frac{1}{2}} \frac{\rho_i^n - \rho_{i-1}^n}{\Delta x} \right) \\ & + \rho_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right) + \mathcal{O}(\varepsilon) \end{aligned}$$

Formal numerical limit

$$\bullet g_{i+\frac{1}{2}}^{n+1} = \frac{1}{\sigma_{i+\frac{1}{2}}} \mathcal{L}^{-1} \left(v \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} \right) + \mathcal{O}(\varepsilon)$$

$$\begin{aligned} \Rightarrow \rho_i^{n+1} = \rho_i^n - \frac{\Delta t}{\Delta x} & \left(\kappa_{i+\frac{1}{2}} \frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} - \kappa_{i-\frac{1}{2}} \frac{\rho_i^n - \rho_{i-1}^n}{\Delta x} \right) \\ & + \rho_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right) + \mathcal{O}(\varepsilon) \end{aligned}$$

\Rightarrow the usual **3-points stencil explicit scheme** for **the diffusion equation**

$$\text{with } \kappa_{i+\frac{1}{2}} = - \frac{\Pi(v \mathcal{L}^{-1} v)}{\sigma_{i+\frac{1}{2}}}.$$

The one-group transport equation

Numerical scheme for the **kinetic equation**:

$$f_i^{n+1} = f_i^n - \frac{\Delta t}{\varepsilon \Delta x} (v^+ (f_i^n - f_{i-1}^n) + v^- (f_{i+1}^n - f_i^n)) \\ + \frac{1}{\varepsilon^2} (\Pi f_i^n - f_i^n) + f_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right)$$

The one-group transport equation

Numerical scheme for the **kinetic equation**:

$$f_i^{n+1} = f_i^n - \frac{\Delta t}{\varepsilon \Delta x} (v^+ (f_i^n - f_{i-1}^n) + v^- (f_{i+1}^n - f_i^n)) \\ + \frac{1}{\varepsilon^2} (\Pi f_i^n - f_i^n) + f_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right)$$

Numerical (Crank-Nicholson) scheme for the **diffusion equation**:

$$\rho_i^{n+1} = \rho_i^n - \frac{\kappa}{2} \frac{\Delta t}{\Delta x^2} (\rho_{i+1}^{n+1} - 2\rho_i^{n+1} + \rho_{i-1}^{n+1}) - \frac{\kappa}{2} \frac{\Delta t}{\Delta x^2} (\rho_{i+1}^n - 2\rho_i^n + \rho_{i-1}^n) \\ + \rho_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right)$$

with $\kappa = \Pi(v\mathcal{L}^{-1}v) = \Pi(-v^2) = -1/3$.

The one-group transport equation

Numerical scheme for the **kinetic equation**:

$$f_i^{n+1} = f_i^n - \frac{\Delta t}{\varepsilon \Delta x} (v^+ (f_i^n - f_{i-1}^n) + v^- (f_{i+1}^n - f_i^n)) \\ + \frac{1}{\varepsilon^2} (\Pi f_i^n - f_i^n) + f_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right)$$

Numerical (Crank-Nicholson) scheme for the **diffusion equation**:

$$\rho_i^{n+1} = \rho_i^n - \frac{\kappa}{2} \frac{\Delta t}{\Delta x^2} (\rho_{i+1}^{n+1} - 2\rho_i^{n+1} + \rho_{i-1}^{n+1}) - \frac{\kappa}{2} \frac{\Delta t}{\Delta x^2} (\rho_{i+1}^n - 2\rho_i^n + \rho_{i-1}^n) \\ + \rho_i^n \left(\frac{1}{2} \Delta t \sum_{k \geq 0} (b_{ik})^2 + \sqrt{\Delta t} \sum_{k \geq 0} b_{ik} \xi_k^{n+1} \right)$$

with $\kappa = \Pi(v\mathcal{L}^{-1}v) = \Pi(-v^2) = -1/3$.

- For the simulation, discretization in velocities $\Rightarrow \kappa \approx -1/2 \sum_{j=1}^N v_j^2 \Delta v$.

Simulations

- **Initial data:** $f_0(x, v) = (1 + \cos(2\pi x + \pi))$.
- **Space domain** $[0, 1]$ discretized with $N = 200$ points, periodic boundary conditions.
- **Noise** of the form $\sum_{k \in \mathbf{Z}} \frac{1}{|k| + 1} (\cos(kx) + \sin(kx)) d\beta_k$.
- **Comparison of three regimes:** kinetic for $\varepsilon = 1$, intermediate for $\varepsilon = 10^{-2}$, diffusion for $\varepsilon = 10^{-8}$.
- 100 **realizations** for each time \Rightarrow comparison of the mean.

Kinetic regime





Figure: one-group transport equation $\varepsilon = 1$: comparison between SMM and explicit schemes (200 grid points): $t = 0.1$ (left), $t = 0.3$ (right).

Kinetic regime





Figure: one-group transport equation $\varepsilon = 1$: comparison between SMM and explicit schemes (200 grid points): $t = 0.6$ (left), $t = 1$ (right).

Intermediate regime





Figure: one-group transport equation $\varepsilon = 10^{-2}$: comparison between SMM and CN schemes (200 grid points): $t = \varepsilon/10$ (left), $t = 4\varepsilon/10$ (right).

Intermediate regime





Figure: one-group transport equation $\varepsilon = 10^{-2}$: comparison between SMM and CN schemes (200 grid points): $t = 0.05$ (left), $t = 0.1$ (right).

Diffusion regime





Figure: one-group transport equation $\varepsilon = 10^{-8}$: comparison between SMM and CN schemes (200 grid points): $t = 0.01$ (left), $t = 0.1$ (right).

Diffusion regime





Figure: one-group transport equation $\varepsilon = 10^{-8}$: comparison between SMM and CN schemes (200 grid points): $t = 0.25$ (left), $t = 0.4$ (right).

Stability analysis

The telegraph equation with a **one dimensional Brownian motion** for noise:

$$\left\{ \begin{array}{l} d\rho + \partial_x \frac{\alpha - \gamma}{2} dt = \rho \circ d\beta(t) \\ d\alpha + \frac{1}{\varepsilon} \partial_x \frac{\alpha + \gamma}{2} dt = -\frac{1}{\varepsilon^2} \alpha dt + \alpha \circ d\beta(t) - \frac{1}{\varepsilon^2} \partial_x \rho dt \\ d\gamma - \frac{1}{\varepsilon} \partial_x \frac{\gamma + \alpha}{2} dt = -\frac{1}{\varepsilon^2} \gamma dt + \gamma \circ d\beta(t) + \frac{1}{\varepsilon^2} \partial_x \rho dt. \end{array} \right.$$

Stability analysis

The telegraph equation with a **one dimensional Brownian motion** for noise:

$$\left\{ \begin{array}{l} d\rho + \partial_x \frac{\alpha - \gamma}{2} dt = \rho \circ d\beta(t) \\ d\alpha + \frac{1}{\varepsilon} \partial_x \frac{\alpha + \gamma}{2} dt = -\frac{1}{\varepsilon^2} \alpha dt + \alpha \circ d\beta(t) - \frac{1}{\varepsilon^2} \partial_x \rho dt \\ d\gamma - \frac{1}{\varepsilon} \partial_x \frac{\gamma + \alpha}{2} dt = -\frac{1}{\varepsilon^2} \gamma dt + \gamma \circ d\beta(t) + \frac{1}{\varepsilon^2} \partial_x \rho dt. \end{array} \right.$$

We denote $j := \frac{1}{2\varepsilon}(p - q) = \frac{1}{2}(\alpha - \gamma)$. The **micro-macro scheme** rewrites

$$\left\{ \begin{array}{l} \rho_i^{n+1} = \rho_i^n - \frac{\Delta t}{\Delta x} \left(j_{i+\frac{1}{2}}^{n+1} - j_{i-\frac{1}{2}}^{n+1} \right) + \rho_i^n \left(\frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) \\ j_{i+\frac{1}{2}}^{n+1} = j_{i+\frac{1}{2}}^n + \frac{\Delta t}{2\varepsilon\Delta x} \left[j_{i+\frac{3}{2}}^n - 2j_{i+\frac{1}{2}}^n + j_{i-\frac{1}{2}}^n \right] \\ \quad - \frac{\Delta t}{\varepsilon^2} j_{i+\frac{1}{2}}^{n+1} + j_{i+\frac{1}{2}}^n \left(\frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) - \frac{1}{\varepsilon^2} \Delta t \left(\frac{\rho_{i+1}^n - \rho_i^n}{\Delta x} \right) \end{array} \right.$$

Stability analysis

Theorem

*There exist constants $C(T)$, Δt_0 , Δx_0 and ε_0 such that for all $\Delta t \leq \Delta t_0$, $\Delta x \leq \Delta x_0$ and $\varepsilon \leq \varepsilon_0$ satisfying the **CFL condition***

$$\Delta t \leq \frac{1}{2} \left(\frac{\Delta x^2}{2} + \varepsilon \Delta x \right)$$

then we have

$$\mathbb{E} \left[\sum_i (\rho_i^n)^2 + (\varepsilon j_{i+\frac{1}{2}}^n)^2 \right] \leq C(T) \mathbb{E} \left[\sum_i (\rho_i^0)^2 + (\varepsilon j_{i+\frac{1}{2}}^0)^2 \right]$$

for every n .

Proof:

We denote $J_{i+\frac{1}{2}}^n = \varepsilon j_{i+\frac{1}{2}}^n$, $\mu = \frac{\Delta t}{\varepsilon \Delta x}$ and $\lambda = \frac{1}{1+\Delta t/\varepsilon^2}$.

The scheme becomes

$$\begin{cases} \rho_j^{n+1} = \rho_j^n - \mu \left(J_{j+\frac{1}{2}}^{n+1} - J_{j-\frac{1}{2}}^{n+1} \right) + \rho_j^n \left(\frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) \\ J_{j+\frac{1}{2}}^{n+1} = \lambda \left(J_{j+\frac{1}{2}}^n \left(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) \right. \\ \qquad \qquad \qquad \left. + \frac{\mu}{2} \left[J_{j+\frac{3}{2}}^n - 2J_{j+\frac{1}{2}}^n + J_{j-\frac{1}{2}}^n \right] - \mu \left(\rho_{j+1}^n - \rho_j^n \right) \right) \end{cases}$$

Proof:

We denote $J_{i+\frac{1}{2}}^n = \varepsilon j_{i+\frac{1}{2}}^n$, $\mu = \frac{\Delta t}{\varepsilon \Delta x}$ and $\lambda = \frac{1}{1+\Delta t/\varepsilon^2}$.

The scheme becomes

$$\begin{cases} \rho_j^{n+1} = \rho_j^n - \mu \left(J_{j+\frac{1}{2}}^{n+1} - J_{j-\frac{1}{2}}^{n+1} \right) + \rho_j^n \left(\frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) \\ J_{j+\frac{1}{2}}^{n+1} = \lambda \left(J_{j+\frac{1}{2}}^n \left(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) \right. \\ \quad \left. + \frac{\mu}{2} \left[J_{j+\frac{3}{2}}^n - 2J_{j+\frac{1}{2}}^n + J_{j-\frac{1}{2}}^n \right] - \mu (\rho_{j+1}^n - \rho_j^n) \right) \end{cases}$$

We take ρ_j^n and $J_{j+\frac{1}{2}}^n$ on the form of elementary waves $\rho_j^n = \rho^n(\varphi)e^{ij\varphi}$ and $J_{j+\frac{1}{2}}^n = J^n(\varphi)e^{i(j+\frac{1}{2})\varphi}$.

$$\begin{cases} \rho^{n+1} = \rho^n \left(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} \right) - 2i\mu J^{n+1} \sin \theta \\ J^{n+1} = \lambda \left(J^n \left(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta \right) - 2i\mu \sin \theta \rho^n \right) \end{cases}$$

with $\theta = \frac{\varphi}{2}$.

This rewrites as $\begin{pmatrix} \rho^{n+1} \\ J^{n+1} \end{pmatrix} = A_{n+1} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix}$ with

$$\begin{pmatrix} 1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 4\mu^2 \lambda \sin^2 \theta & -i(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) \lambda \end{pmatrix}$$

$$= A_{n+1}.$$

This rewrites as $\begin{pmatrix} \rho^{n+1} \\ J^{n+1} \end{pmatrix} = A_{n+1} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix}$ with

$$\begin{pmatrix} 1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 4\mu^2 \lambda \sin^2 \theta & -i(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) \lambda \end{pmatrix}$$

$$= A_{n+1}.$$

A_{n+1} is a **stochastic perturbation** of the matrix \tilde{A} appearing in the deterministic version :

$$\tilde{A} = \begin{pmatrix} 1 - 4\mu^2 \lambda \sin^2 \theta & -i(1 - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 - 2\mu \sin^2 \theta) \lambda \end{pmatrix}.$$

$$\Rightarrow A_{n+1} = \tilde{A} + \sqrt{\Delta t} \xi^{n+1} \tilde{B} + \Delta t \tilde{C},$$

with $\tilde{B} = \tilde{B}(\mu\lambda, \mu^2\lambda, \sin \theta)$ and $\tilde{C} = \tilde{C}(\mu\lambda, \mu^2\lambda, \sin \theta)$.

This rewrites as $\begin{pmatrix} \rho^{n+1} \\ J^{n+1} \end{pmatrix} = A_{n+1} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix}$ with

$$\begin{pmatrix} 1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 4\mu^2 \lambda \sin^2 \theta & -i(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) \lambda \end{pmatrix}$$

$$= A_{n+1}.$$

A_{n+1} is a **stochastic perturbation** of the matrix \tilde{A} appearing in the deterministic version :

$$\tilde{A} = \begin{pmatrix} 1 - 4\mu^2 \lambda \sin^2 \theta & -i(1 - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 - 2\mu \sin^2 \theta) \lambda \end{pmatrix}.$$

$$\Rightarrow A_{n+1} = \tilde{A} + \sqrt{\Delta t} \xi^{n+1} \tilde{B} + \Delta t \tilde{C},$$

with $\tilde{B} = \tilde{B}(\mu\lambda, \mu^2\lambda, \sin \theta)$ and $\tilde{C} = \tilde{C}(\mu\lambda, \mu^2\lambda, \sin \theta)$.

Under the **CFL assumption**, $\lambda\mu \leq C$, $\lambda\mu^2 \leq C$.

This rewrites as $\begin{pmatrix} \rho^{n+1} \\ J^{n+1} \end{pmatrix} = A_{n+1} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix}$ with

$$\begin{pmatrix} 1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 4\mu^2 \lambda \sin^2 \theta & -i(1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 + \frac{\Delta t}{2} + \sqrt{\Delta t} \xi^{n+1} - 2\mu \sin^2 \theta) \lambda \end{pmatrix}$$

$$= A_{n+1}.$$

A_{n+1} is a **stochastic perturbation** of the matrix \tilde{A} appearing in the deterministic version :

$$\tilde{A} = \begin{pmatrix} 1 - 4\mu^2 \lambda \sin^2 \theta & -i(1 - 2\mu \sin^2 \theta) 2\lambda \mu \sin \theta \\ -2i\mu \lambda \sin \theta & (1 - 2\mu \sin^2 \theta) \lambda \end{pmatrix}.$$

$$\Rightarrow A_{n+1} = \tilde{A} + \sqrt{\Delta t} \xi^{n+1} \tilde{B} + \Delta t \tilde{C},$$

with $\tilde{B} = \tilde{B}(\mu\lambda, \mu^2\lambda, \sin \theta)$ and $\tilde{C} = \tilde{C}(\mu\lambda, \mu^2\lambda, \sin \theta)$.

Under the **CFL assumption**, $\lambda\mu \leq C$, $\lambda\mu^2 \leq C$.

$\Rightarrow \tilde{B}(\lambda\mu, \lambda\mu^2, \sin \theta)$ and $\tilde{C}(\lambda\mu, \lambda\mu^2, \sin \theta)$ are **uniformly bounded**.

We denote by \mathcal{F}_n the σ -algebra generated by

$$\rho^n, J^n, \xi^n, \rho^{n-1}, J^{n-1}, \xi^{n-1}, \dots, \rho^0, J^0.$$

$\Rightarrow \xi^{n+1}$ independent of \mathcal{F}_n and ρ^n, J^n are \mathcal{F}_n -measurable.

$\Rightarrow \mathbb{E}(\xi^{n+1}|\mathcal{F}_n) = 0, \mathbb{E}(\rho^n|\mathcal{F}_n) = \rho^n, \mathbb{E}(J^n|\mathcal{F}_n) = J^n.$

$$\begin{aligned} \Rightarrow \mathbb{E} \left[\left\| \begin{pmatrix} \rho^{n+1} \\ J^{n+1} \end{pmatrix} \right\|_2^2 \middle| \mathcal{F}_n \right] &= \mathbb{E} \left[\left\| A_{n+1} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix} \right\|_2^2 \middle| \mathcal{F}_n \right] \\ &\leq \left\| (\tilde{A} + \Delta t \tilde{C}) \begin{pmatrix} \rho^n \\ J^n \end{pmatrix} \right\|_2^2 + \Delta t \left\| \tilde{B} \begin{pmatrix} \rho^n \\ J^n \end{pmatrix} \right\|_2^2 \leq (\|\tilde{A}\|_2^2 + L\Delta t) \left\| \begin{pmatrix} \rho^n \\ J^n \end{pmatrix} \right\|_2^2, \end{aligned}$$

Let us prove that $\|\tilde{A}\|_2^2 \leq 1$ under **the CFL condition**.

Let us prove that $\|\tilde{A}\|_2^2 \leq 1$ under **the CFL condition**.

$\|\tilde{A}\|_2^2$ is the **largest eigenvalue** of the matrix $\tilde{A}^* \tilde{A}$. Denoting by \tilde{T} and \tilde{D} the trace and determinant of this latter matrix, the largest eigenvalue is

$$\frac{\tilde{T} + \sqrt{\tilde{T}^2 - 4\tilde{D}}}{2}$$

and the condition $\|\tilde{A}\|_2^2 \leq 1$ is thus equivalent to $1 - \tilde{T} + \tilde{D} \geq 0$.

Let us prove that $\|\tilde{A}\|_2^2 \leq 1$ under **the CFL condition**.

$\|\tilde{A}\|_2^2$ is the **largest eigenvalue** of the matrix $\tilde{A}^* \tilde{A}$. Denoting by \tilde{T} and \tilde{D} the trace and determinant of this latter matrix, the largest eigenvalue is

$$\frac{\tilde{T} + \sqrt{\tilde{T}^2 - 4\tilde{D}}}{2}$$

and the condition $\|\tilde{A}\|_2^2 \leq 1$ is thus equivalent to $1 - \tilde{T} + \tilde{D} \geq 0$.

$$1 - \tilde{T} + \tilde{D} = 8\lambda\mu^2 X(1 - \lambda + 2\lambda\mu X - 2\lambda\mu^2 X^2 - 2\lambda\mu^2 X)$$

with $X = (\sin \theta)^2$.

Let us prove that $\|\tilde{A}\|_2^2 \leq 1$ under **the CFL condition**.

$\|\tilde{A}\|_2^2$ is the **largest eigenvalue** of the matrix $\tilde{A}^* \tilde{A}$. Denoting by \tilde{T} and \tilde{D} the trace and determinant of this latter matrix, the largest eigenvalue is

$$\frac{\tilde{T} + \sqrt{\tilde{T}^2 - 4\tilde{D}}}{2}$$

and the condition $\|\tilde{A}\|_2^2 \leq 1$ is thus equivalent to $1 - \tilde{T} + \tilde{D} \geq 0$.

$$1 - \tilde{T} + \tilde{D} = 8\lambda\mu^2 X(1 - \lambda + 2\lambda\mu X - 2\lambda\mu^2 X^2 - 2\lambda\mu^2 X)$$

with $X = (\sin \theta)^2$.

We denote $Q(X) := 1 - \lambda + 2\lambda\mu X - 2\lambda\mu^2 X^2 - 2\lambda\mu^2 X$.

Q **concave** and satisfies $Q(0) = 1 - \lambda > 0$, $Q(1) = 1 - \lambda + 2\lambda\mu - 4\lambda\mu^2 \geq 0$ under **the CFL condition**.

$$\Rightarrow 1 - \tilde{T} + \tilde{D} \geq 0 \Rightarrow \|\tilde{A}\|_2^2 \leq 1.$$



Stability in the general case

Theorem

If Δt satisfies the following CFL condition

$$\Delta t \leq \frac{2s_m\sigma_m\Delta x^2}{2(2+\varepsilon)} + \frac{\varepsilon\Delta x}{2+\varepsilon},$$

then the sequence ρ^n and g^n defined by the scheme satisfy the energy estimate

$$\begin{aligned} \mathbb{E} \left[\sum_i (\rho_i^n)^2 \right] + \varepsilon^2 \mathbb{E} \left[\sum_i \Pi \left((g_{i+\frac{1}{2}}^n)^2 \right) \right] \\ \leq C(T) \left(\mathbb{E} \left[\sum_i (\rho_i^0)^2 \right] + \varepsilon^2 \mathbb{E} \left[\sum_i \Pi \left((g_{i+\frac{1}{2}}^0)^2 \right) \right] \right) \end{aligned}$$

for every n with $C(T)$ a constant which only depends on T .

Conclusion and Perspectives

- Diffusion equation recovered when $\varepsilon \rightarrow 0 \Rightarrow$ **AP scheme**.
- CFL conditions ensuring the **stability of our scheme**.
- **Numerical results** confirming the **good performances** of the scheme, in particular in the diffusion regime.
- **Perspectives**: Similar technics to study stochastic perturbations of
 - ◇ the non-linear case of the radiative transfer equation,
 - ◇ the Boltzmann equation,
 - ◇ the Vlasov equation,
 - ◇ hydrodynamical limits . . .

Thank you for your attention.