

STAT 210A: INTRODUCTION TO MATHEMATICAL STATISTICS

Problem Set 3

Student: Ivana Malenica

Due: Tuesday, Sep. 27

Note: All measure-theoretic niceties about conditioning on measure-zero sets, almost-sure equality vs. actual equality, “all functions” vs. “all measurable functions,” etc. are disregarded for the time being.

1. U-estimable functions

Assume that the model $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$ has complete sufficient statistic $T(X)$, which can take exactly d distinct values (all with positive probability, $d < \infty$).

- (a) Let \mathcal{G} denote the set of real-valued U -estimable functions $g(\theta)$. Show that \mathcal{G} is a vector space of dimension d . (Hint: use completeness)
- (b) If $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \text{Bern}(\theta)$, show that $g(\theta)$ is U -estimable if and only if it is a polynomial of degree at most n .

2. Poisson UMVU estimation

Let $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \text{Pois}(\theta)$ and consider estimating

$$g(\theta) = e^{-\theta} = \mathbb{P}_\theta(X_1 = 0)$$

- (a) Find the UMVU estimator for $g(\theta)$ by Rao-Blackwellizing $1\{X_1 = 0\}$. (Hint: start by showing that $(X_1, \dots, X_n) \sim \text{Multinom}(t, (n^{-1}, \dots, n^{-1}))$ given $\sum_{i=1}^n X_i = t$).
- (b) Find the UMVU estimator for $g(\theta)$ directly, using the power series method from class.
- (c) If $\theta = 1$, numerically compute the efficiency of the UMVU (relative to the CRLB) for $n = 1, 10, 100$.

3. UMVUE for uniforms

Let $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} U[0, \theta]$ and consider estimating $\beta = g(\theta)$ where $g(\cdot)$ is a differentiable function on $(0, \infty)$. Show that the UMVUE of β is given by

$$h(X) = g(X_{(n)}) + \frac{X_{(n)}g'(X_{(n)})}{n},$$

where $X_{(n)} = \max\{X_1, \dots, X_n\}$.

4. Fisher information for scale families

Consider a scale family

$$p_\theta(x) = \frac{1}{\theta} p\left(\frac{x}{\theta}\right), \quad \theta > 0.$$

where p is some fixed density function with respect to the Lebesgue measure.

- (a) Show that the Fisher information of a single observation X is given by

$$J(\theta) = \frac{1}{\theta^2} \int_{-\infty}^{\infty} \left[\frac{yp'(y)}{p(y)} + 1 \right]^2 p(y) dy.$$

- (b) Show that the Fisher information of a single observation X about $\xi = \log \theta$ is independent of θ .

5. HCR inequality for exponential location family

Suppose that X_1, \dots, X_n are i.i.d. with a single observation having density

$$p_{\theta}^{(1)}(x) = e^{\theta-x} 1\{x \geq \theta\}$$

with respect to the Lebesgue measure on \mathbb{R} .

Note that this family does not have common support and the log-likelihood is certainly not differentiable w.r.t. θ at $\theta = x_{(1)}$.

- (a) Show that (in general) the HCR inequality still applies when the support of $p_{\theta+\varepsilon}$ is contained in the support of p_{θ} , and conclude that we can therefore still apply the HCR inequality in this problem for $\varepsilon \geq 0$ (Note: to alleviate any existential angst about dividing 0 by 0, we can formally define the function $L(x) = p_{\theta+\varepsilon}(x)/p_{\theta}(x)$ if the denominator is positive and $L(x) = 1$ if the numerator and denominator are both 0)
- (b) If $\delta(X)$ is an unbiased estimator of θ , optimize the HCR bound to show that

$$\text{Var}_{\theta}(\delta(X)) \geq \frac{a^2}{(e^a - 1)n^2},$$

where a solves

$$\frac{2}{a} - \frac{e^a}{e^a - 1} = 0.$$

- (c) Prove that $\delta(X) = \min_i X_i - \frac{1}{n}$ is unbiased and has variance $1/n^2$. (Note: this is much better than the $1/n$ scaling predicted by the CRLB, which does not apply here)