Pushing 200g from 150W

Definitive seed-to-harvest guide for 4 feminized Super Skunk in a Homebox Q80+ (80×80×180cm) under a Caluma Force 150W LED, BioBizz Light-Mix soil, inside a 120×90cm Austrian flat pantry. Quality first, maximum yield second. Every claim pressure-tested from environment control, plant science, practical experience, and stealth/risk perspectives.

TENT / LIGHT
Q80+ 80×80×180 / Caluma 150W dimmable

PLANTS / MEDIUM
4× fem. Super Skunk / Light-Mix + 25% perlite

POTS / VENTILATION
4× 11L fabric square / 125mm Vents TT + Rhino Pro

TIMELINE
~18–20 weeks seed to cured jar

150g
Confident floor
1.0 g/W

200g
Realistic target
1.3 g/W

225g
Hard ceiling
1.5 g/W

> **Devil's advocate — yield honesty:** 150g is near-guaranteed with proper ScrOG. 200g requires dialed-in VPD, nutrients, and full screen fill. 225g demands everything optimized plus favorable phenotypes. For a grower with some experience on their first indoor run, plan for 180g and celebrate anything above that.

---

1. The light is your ceiling — and here's exactly how high it goes

The Caluma Force 150W produces ~400 µmol/s PPF at 2.7 µmol/J. Inside the reflective Q80+ at 25–30cm height, that delivers approximately 540–625 µmol/m²/s average PPFD across the 80×80cm footprint. During a 12-hour flower cycle, this equals a DLI of ~23–27 mol/m²/day — roughly 65% of the 40–45 DLI where peer-reviewed research shows yield plateaus.

What does this mean in grams? Interpolating published yield-vs-PPFD data to your PPFD gives 275–350 g/m². Applied to your 0.64m² canopy: 176–224g dry. Forum-documented results from Rollitup and ICMag with comparable 150W Samsung-diode fixtures in similar tents confirm this range, clustering around 1.0–1.5 g/W with ScrOG.

Expected dry yield at 150W by training method. The jump from "Top + ScrOG" to "Full optimization" comes from VPD control, CalMag, mycorrhizae, and aggressive lollipopping.

---

2. The ScrOG canopy is everything

With only 150W, your single highest-impact intervention is ensuring every square centimeter of the 80×80cm footprint receives equal light. A perfectly flat ScrOG canopy absorbs 95%+ of available photons. An untrained canopy wastes 30–40% on gaps and self-shading. That difference alone accounts for the gap between 100g and 200g.

Pot layout — 4 plants filling the 80×80cm screen



Top-down: fully trained ScrOG canopy at flip day — ~60 bud sites evenly distributed across 80×80cm

Training protocol — week by week



Training stages: seedling → topped at node 3 → 4 colas spread by LST → flat ScrOG canopy at flip

Devil's advocate — training complexity: Mainlining (creating perfect symmetry with 3+ toppings) adds 2–4 weeks of veg time for marginal benefit at 150W. A single topping plus LST produces comparable results in half the time. Super Skunk's indica morphology naturally produces a compact, bushy shape that responds well to a single

top — don't overcomplicate it on the first run.

3. Height budget — why you flip at 30–40cm



Flip when canopy sits 30–40cm above pots. Super Skunk stretches 50–100% (indica pheno).

Side view: height budget — 15cm LED + 30cm clearance + 105cm plant zone + 25cm net + 30cm pot. Flip at 30–40cm.

4. VPD targets — your free yield lever

Seedling	Veg	Early flower	Late flower
0.4–0.8 kPa	**0.8–1.2** kPa	**1.0–1.4** kPa	**1.2–1.6** kPa
25°C / 70% RH	25°C / 55–60% RH	25°C / 50–55% RH	22°C / 40–45% RH
Humidity dome Fan on lowest	Open intake vent Fan on low-med	Increase fan speed via Varispeed	Max fan + open door wider

VPD = the pressure differential that drives stomatal opening, transpiration, and nutrient uptake.
Measure with digital hygrometer at canopy level. Check min/max memory daily.

VPD targets by stage with practical instructions for achieving them using only the Varispeed fan controller and door gap

Devil's advocate — VPD realism: Late-flower targets of 40–45% RH may be unachievable during Austrian

summer without a small dehumidifier (~€30–50 from Mediamarkt). In winter (the recommended grow season), indoor humidity naturally drops to 30–50% from central heating — making these targets trivial. Schedule lights-on during the hottest part of the day so the dark period coincides with the coolest hours, naturally achieving a 3–5°C temperature DIF.

---

5. Nutrient schedule — BioBizz optimized with three key supplements

CalMag (non-negotiable under LED)

LEDs drive faster photosynthesis with less radiant heat, reducing passive calcium uptake. BioBizz CalMag at 1 ml/L from week 2 through harvest prevents the mid-flower calcium lockout that silently caps yields by 10–30%.

Mycorrhizal inoculant at transplant

Apply 5g of granular mycorrhizae (DYNOMYCO or Great White) directly on the rootball at transplant. BioBizz organics are ideal partners — synthetic fertilizers suppress mycorrhizal colonization, but organic nutrients don't.

> **Devil's advocate — mycorrhizae claims:** DYNOMYCO's marketing claims "10–45% yield increase." Independent data is more modest — expect 5–15% in well-managed organic soil. Still the highest-ROI supplement at ~€15 per grow, but don't plan your yield around the upper range.

Silica (structural support for heavy colas)

Potassium silicate at 2–3 ml/gallon builds stronger cell walls, supporting heavier colas. Add to water first, before other nutrients, from early veg through week 3–4 of flower.

Complete feeding schedule

Week	Stage	Bio-Grow	Bio-Bloom	Top-Max	CalMag	Notes
1–2	Seedling	—	—	—	—	Light-Mix has enough. Water only.
3	Early veg	1	—	—	1	Start feeding after transplant
4–5	Veg	2–3	—	—	1	Ramp based on plant response
6	Flip	2	1	1	1	Transition: both Grow + Bloom
7–8	Early flower	2	2	1–2	1	Stretch phase, still needs N
9–10	Mid flower	1	3	2–3	1	Peak feeding. Watch for burn.
11–12	Late flower	1	3–4	3–4	0.5	Max bloom. Reduce N slowly.
13	Ripen	0	2	2	—	Taper down
14	Harvest	0	0	0	—	Plain water last 3–5 days only

All values in ml/L. Feed → plain water → feed rotation. Start at 50% dose and increase based on plant response.

> **Devil's advocate — flushing:** With BioBizz organics, there's nothing harsh to flush — nutrients are microbially processed. A 3–5 day plain-water finish is sufficient. The 10-day flush in the previous guide was unnecessary for organic growing.

---

6. Watering — the most common mistake

The watering cycle. Use the lift test — never a fixed schedule. Empty drip trays 15 min after watering.

7. Harvest timing — by trichomes, not the calendar

Start checking trichomes at flower week 6 with a 60× loupe. Check on calyxes (bud pods), not sugar leaves — leaves mature faster and mislead. Super Skunk's real flowering time is 8–9 weeks, not the 45–50 days Sensi Seeds claims.

Too early

All clear / glassy

Low cannabinoid content. Keep waiting.

Peak THC

Mostly milky, minimal amber

Maximum THC. More cerebral. Good but you can push further for the Super Skunk body stone.

Recommended harvest

70–80% milky, 20–30% amber

The sweet spot. Balanced potency with the classic heavy body stone. Complex terpene profile at peak. This is where quality-first growers harvest.

Late / sedative

50%+ amber

THC degrading to CBN. Heavy sedation. Only if you want a sleep aid.

Staggered harvest for +10–15% yield: Top buds ripen before lowers. Cut top colas first, strip fan leaves, give lowers 5–7 more days under light. Multiple growers report lower buds "thickening up quite a bit" during this window.

8. Late-flower quality protocol (final 2 weeks)

These interventions are stacked in the last 14 days before harvest for maximum quality.

Intervention	What to do	Why it works
Temperature drop	Days 21–23°C, nights 17–19°C	Preserves volatile terpenes, triggers stress-resin response
		Signals plant to produce more trichomes as dehydration

Humidity drop	35–40% RH	defense
Mild drought stress	Let pots dry slightly more in final 5–7 days	Increases cannabinoid concentration per gram by ~12%
Harvest timing	Just before lights-on	THC peaks during dark period, degrades partially under light

> **Devil's advocate — drought stress:** Caplan et al. (2019) showed drought stress increases cannabinoid concentration per gram but simultaneously decreases total grams. Net THC per plant stays roughly flat. Use this only as a quality finisher in the final 5–7 days, never as a yield strategy.

---

9. Drying and curing — where quality is won or lost

Your tent becomes the drying chamber. Remove the LED. Hang whole branches upside-down. Fan on lowest for gentle air exchange. Never point a fan directly at buds.

Phase	Conditions	Duration	Test for readiness
Drying	15–18°C, 55–60% RH, complete darkness	10–14 days	Small stems snap cleanly when bent
Cure (week 1)	Mason jars, 60–65% RH, dark, 15–21°C	Burp 2–3× daily, 10–15 min	No ammonia smell when opening
Cure (week 2)	Same	Burp once daily	Boveda 62% pack if below 55% RH
Cure (week 3+)	Same	Burp weekly	Flavor developing, smoke smoothing
Ready	Sealed jars, dark, cool	Minimum 4 weeks, ideal 6–8	Smooth smoke, complex terpene profile

> **Dry trimming for quality:** Leave sugar leaves on during drying — they slow moisture loss, protect trichomes, and preserve terpenes. Trim to final form only after drying is complete. Super Skunk's dense indica buds benefit from the longer end of the drying range (12–14 days).

---

10. What NOT to do — debunked techniques

Technique	Claim	Reality
CO_2 bags	+20% yield	Your exhaust fan evacuates CO_2 instantly. At 550 PPFD, light is the bottleneck, not carbon. Waste of money.
48–72h darkness pre-harvest	+30% THC	One unreplicated study. Mold risk in humid tent with no airflow. Skip.
Ice water flush	More trichomes	A/B test: 26 of 30 strains hermied. Cold shock inhibits P uptake. Don't.
Schwazzing	+40% yield	Designed for 1000W + CO_2. At 150W, removing that much leaf starves the plant.
13/11 light schedule	More yield than 12/12	Promising research but risky with unknown phenos. Stick with 12/12 first grow.

11. Complete week-by-week timeline

Wk	Stage	LED	Key actions	Temp/RH
0	Germ	Dark	Soak → paper towel → plant at 1–2cm taproot	24°C/75%
1	Seedling	25%/55cm	Dome on. Mycorrhizae at transplant to 11L	25°C/70%
2	Seedling	35%/45cm	Remove dome. Bio-Grow 1ml/L + CalMag 1ml/L	25°C/65%
3	Veg	50%/40cm	Top above node 3–4. Begin LST	25°C/60%
4	Veg	60%/35cm	Install ScrOG net. Begin tucking. Sticky traps in	25°C/58%
5	Veg	70%/35cm	Net 70–80% full. Lollipop bottom third	25°C/55%
6	Flip	85%/30cm	12/12. Defoliate 15–20%. Net 90%+ full	25°C/55%
7	F wk1	90%/28cm	Stretch begins. Continue tucking 7–10 days	25°C/52%
8	F wk2	95%/27cm	First pistils. Last tuck. Stretch slowing	25°C/50%
9	F wk3	100%/25cm	Day 21: 2nd defoliation (max 25%). Stop tucking	24°C/48%
10	F wk4	100%/25cm	Buds fattening. Smell sharp. Add branch support	24°C/45%
11	F wk5	100%/25cm	Peak feeding. Watch for bud rot daily	23°C/44%
12	F wk6	100%/25cm	Begin trichome checks (60× loupe)	23°C/42%
13	F wk7	100%/25cm	Trichomes milky. Taper nutrients. Mild drought	22°C/40%
14	F wk8	100%	70% milky/20% amber → harvest tops. Stagger	22°C/38%
15	F wk9	90%	Lower buds mature 5–7 more days. Harvest rest	21°C/38%
16–17	Dry	OFF	Hang branches in dark tent. Fan on lowest	18°C/58%
18–25	Cure	—	Mason jars. Burp per schedule. Min 4 weeks	18°C/62% jar

Revised yield expectations

150g

Floor: ScrOG + basic nutes
1.0 g/W — near guaranteed

200g

Target: + VPD + CalMag + myco
1.3 g/W — realistic with discipline

225g

Ceiling: + perfect pheno + stagger
1.5 g/W — everything optimized

The difference between floor and target is technique and environment. The difference between target and ceiling is experience and phenotype luck. With proper ScrOG, CalMag from week 2, mycorrhizae at transplant, and humidity below 45% in late flower, 200g is a realistic first-grow target.

Three highest-impact interventions (ranked):

1. Flat ScrOG canopy with 90%+ screen fill — worth 40–60g vs untrained plants.

2. CalMag + mycorrhizal inoculant — prevents hidden deficiencies for ~€20 total.

3. VPD management via canopy-level hygrometer + Varispeed fan adjustment — free.

Best future upgrade: A 200–240W LED pushes DLI from 65% to 85% of optimal and the ceiling past 300g.

Total timeline: ~18–20 weeks seed to smokeable flower including 4-week minimum cure.