

# Titanic

2022-11

## Titanic

Trabajaremos con el Dataset **Titanic**

Table 1: Primeros registros de la base de datos Titanic

PassengerId	Survived	Pclass	Name	Sex	Age	SibSp	Parch	Ticket	Fare	Cabin	Embarked
1	0	3	Braund, Mr. Owen Harris	male	22	1	0	A/5 21171	7.2500	NA	S
2	1	1	Cumings, Mrs. John Bradley (Florence Briggs Thayer)	female	38	1	0	PC 17599	71.2833	C85	C
3	1	3	Heikkinen, Miss. Laina	female	26	0	0	STON/O2. 3101282	7.9250	NA	S
4	1	1	Futrelle, Mrs. Jacques Heath (Lily May Peel)	female	35	1	0	113803	53.1000	C123	S
5	0	3	Allen, Mr. William Henry	male	35	0	0	373450	8.0500	NA	S
6	0	3	Moran, Mr. James	male	NA	0	0	330877	8.4583	NA	Q

## Data Frame Summary

**TitanicT Dimensions:** 891 x 12

**Duplicates:** 0


No	Variable	Stats / Values	Freqs (% of Valid)	Valid	Missing
1	PassengerId [numeric]	Mean (sd) : 446 (257.4) min < med < max: 1 < 446 < 891 IQR (CV) : 445 (0.6)	891 distinct values	891 (100.0%)	0 (0.0%)
2	Survived [numeric]	Min : 0 Mean : 0.4 Max : 1	0 : 549 (61.6%) 1 : 342 (38.4%)	891 (100.0%)	0 (0.0%)
3	Pclass [numeric]	Mean (sd) : 2.3 (0.8) min < med < max: 1 < 3 < 3 IQR (CV) : 1 (0.4)	1 : 216 (24.2%) 2 : 184 (20.7%) 3 : 491 (55.1%)	891 (100.0%)	0 (0.0%)
4	Name [character]	1. Abbing, Mr. Anthony 2. Abbott, Mr. Rossmore Edwa 3. Abbott, Mrs. Stanton (Ros 4. Abelson, Mr. Samuel 5. Abelson, Mrs. Samuel (Han [ 886 others ]	1 ( 0.1%) 1 ( 0.1%) 1 ( 0.1%) 1 ( 0.1%) 1 ( 0.1%) 886 (99.4%)	891 (100.0%)	0 (0.0%)
5	Sex [character]	1. female 2. male	314 (35.2%) 577 (64.8%)	891 (100.0%)	0 (0.0%)

No	Variable	Stats / Values	Freqs (% of Valid)	Valid	Missing
6	Age [numeric]	Mean (sd) : 29.7 (14.5) min < med < max: 0.4 < 28 < 80 IQR (CV) : 17.9 (0.5)	88 distinct values	714 (80.1%)	177 (19.9%)
7	SibSp [numeric]	Mean (sd) : 0.5 (1.1) min < med < max: 0 < 0 < 8 IQR (CV) : 1 (2.1)	7 distinct values	891 (100.0%)	0 (0.0%)
8	Parch [numeric]	Mean (sd) : 0.4 (0.8) min < med < max: 0 < 0 < 6 IQR (CV) : 0 (2.1)	7 distinct values	891 (100.0%)	0 (0.0%)
9	Ticket [character]	1. 1601 2. 347082 3. CA. 2343 4. 3101295 5. 347088 [ 676 others ]	7 ( 0.8%) 7 ( 0.8%) 7 ( 0.8%) 6 ( 0.7%) 6 ( 0.7%) 858 (96.3%)	891 (100.0%)	0 (0.0%)
10	Fare [numeric]	Mean (sd) : 32.2 (49.7) min < med < max: 0 < 14.5 < 512.3 IQR (CV) : 23.1 (1.5)	248 distinct values	891 (100.0%)	0 (0.0%)
11	Cabin [character]	1. B96 B98 2. C23 C25 C27 3. G6 4. C22 C26 5. D [ 142 others ]	4 ( 2.0%) 4 ( 2.0%) 4 ( 2.0%) 3 ( 1.5%) 3 ( 1.5%) 186 (91.2%)	204 (22.9%)	687 (77.1%)
12	Embarked [character]	1. C 2. Q 3. S	168 (18.9%) 77 ( 8.7%) 644 (72.4%)	889 (99.8%)	2 (0.2%)

1. ¿La varianza de las edades de quienes sobrevivieron es diferente para ambos grupos?
2. Con base en la respuesta anterior, prueba si las edades promedio son iguales o diferentes para quienes sobrevivieron o no?


```
TitanicT %>% ggplot(aes(x=Age,colour=as.factor(Survived),group=Survived)) +  
  geom_density()
```

```
## Warning: Removed 177 rows containing non-finite values (`stat_density()`).
```



```
TitanicT %>% ggplot(aes(x=as.factor(Survived), y=Age)) +  
  geom_boxplot()
```

```
## Warning: Removed 177 rows containing non-finite values (`stat_boxplot()`).
```



```
TitanicT %>% group_by(Survived) %>% summarise(varAge = var(Age, na.rm = T))
```

```
## # A tibble: 2 x 2
##   Survived varAge
##   <dbl>   <dbl>
## 1       0    201.
## 2       1    224.
```

```
stats::var.test( Age ~ as.factor(Survived), data = TitanicT, conf.level = 0.95)
```

```
##
## F test to compare two variances
##
## data: Age by as.factor(Survived)
## F = 0.89853, num df = 423, denom df = 289, p-value = 0.317
## alternative hypothesis: true ratio of variances is not equal to 1
## 95 percent confidence interval:
##  0.7253979 1.1082231
## sample estimates:
## ratio of variances
##      0.8985274
```

$H_0: \mu_{age0} = \mu_{age1}, \mu_{age0} - \mu_{age1} = 0$

```
t.test(Age ~ Survived, data = TitanicT, conf.level = 0.95)
```

```
##
## Welch Two Sample t-test
##
```

```
## data: Age by Survived
## t = 2.046, df = 598.84, p-value = 0.04119
## alternative hypothesis: true difference in means between group 0 and group 1 is not equal to 0
## 95 percent confidence interval:
## 0.09158472 4.47339446
## sample estimates:
## mean in group 0 mean in group 1
## 30.62618 28.34369
TitanicT %>% t_test(Age ~ Survived)
```

```
## Warning: The statistic is based on a difference or ratio; by default, for
## difference-based statistics, the explanatory variable is subtracted in the order
## "0" - "1", or divided in the order "0" / "1" for ratio-based statistics. To
## specify this order yourself, supply `order = c("0", "1")`.
```

```
## # A tibble: 1 x 7
##   statistic t_df p_value alternative estimate lower_ci upper_ci
##   <dbl> <dbl> <dbl> <chr> <dbl> <dbl> <dbl>
## 1 2.05 599. 0.0412 two.sided 2.28 0.0916 4.47
```

Se dice que aproximadamente una tercera parte de la gente sobrevivió en el Titanic, estos datos respaldan esta afirmación

```
prop.test(sum(TitanicT$Survived),891,p=1/3)
```

```
##
## 1-sample proportions test with continuity correction
##
## data: sum(TitanicT$Survived) out of 891, null probability 1/3
## X-squared = 10.001, df = 1, p-value = 0.001564
## alternative hypothesis: true p is not equal to 0.3333333
## 95 percent confidence interval:
## 0.3519194 0.4167722
## sample estimates:
## p
## 0.3838384
```

```
TitanicT %>% mutate(Survived=as.factor(Survived)) %>% prop_test(Survived ~ NULL,p=1/3)
```

```
## # A tibble: 1 x 4
##   statistic chisq_df p_value alternative
##   <dbl> <int> <dbl> <chr>
## 1 319. 1 1.90e-71 two.sided
```

```
prop.test(sum(TitanicT$Sex=="female"),891,p=.5)
```

```
##
## 1-sample proportions test with continuity correction
##
## data: sum(TitanicT$Sex == "female") out of 891, null probability 0.5
## X-squared = 77.042, df = 1, p-value < 2.2e-16
## alternative hypothesis: true p is not equal to 0.5
## 95 percent confidence interval:
## 0.3211923 0.3849235
## sample estimates:
## p
## 0.352413
```

```

TitanicT %>% prop_test(Sex ~ NULL,p=.5, order = c("female","male"))

## # A tibble: 1 x 4
##   statistic chisq_df p_value alternative
##   <dbl>      <int>   <dbl> <chr>
## 1      77.0         1 1.67e-18 two.sided

TitanicT %>% prop_test(Sex ~ NULL,p=.5, success = "male", z=TRUE)

## # A tibble: 1 x 3
##   statistic p_value alternative
##   <dbl>     <dbl> <chr>
## 1      8.81 1.24e-18 two.sided

(tabla<- with(TitanicT,addmargins(table(Sex,Survived))))

##           Survived
## Sex           0    1 Sum
## female      81 233 314
## male       468 109 577
## Sum        549 342 891

with(TitanicT,prop.table(table(Sex,Survived),margin = 1))

##           Survived
## Sex           0          1
## female 0.2579618 0.7420382
## male   0.8110919 0.1889081

prop.test(x=c(233,109),n=c(314,577),alternative = "greater")

##
## 2-sample test for equality of proportions with continuity correction
##
## data:  c(233, 109) out of c(314, 577)
## X-squared = 260.72, df = 1, p-value < 2.2e-16
## alternative hypothesis: greater
## 95 percent confidence interval:
##  0.5020113 1.0000000
## sample estimates:
##   prop 1    prop 2
## 0.7420382 0.1889081

TitanicT %>%
  group_by(Sex) %>%
  summarise(
    p_hat=mean(Survived),
    n=n()
  )

## # A tibble: 2 x 3
##   Sex    p_hat    n
##   <chr> <dbl> <int>
## 1 female 0.742   314
## 2 male  0.189   577

TitanicT %>% mutate(Survived=as.factor(Survived)) %>%
  prop_test(Survived ~ Sex,

```

```

    order = c("female", "male"),
    success = "1",
    alternative = "greater",
    correct = F)

## # A tibble: 1 x 6
##   statistic chisq_df p_value alternative lower_ci upper_ci
##   <dbl>      <dbl>   <dbl> <chr>          <dbl>    <dbl>
## 1      263.        1 1.86e-59 greater        0.485      1
(tabla<- with(TitanicT, addmargins(table(Pclass, Survived))))

##           Survived
## Pclass    0    1 Sum
##    1      80 136 216
##    2      97  87 184
##    3     372 119 491
##    Sum   549 342 891

TitanicT %>%
  mutate(Survived=as.factor(Survived),
         Pclass = as.factor(Pclass)) %>%
  chisq_test(Survived ~ Pclass)


## # A tibble: 1 x 3
##   statistic chisq_df p_value
##   <dbl>      <int>   <dbl>
## 1      103.        2 4.55e-23

TitanicT %>%
  mutate(Survived=as.factor(Survived),
         Pclass = as.factor(Pclass)) %>%
  chisq_test(Pclass ~ Survived)

## # A tibble: 1 x 3
##   statistic chisq_df p_value
##   <dbl>      <int>   <dbl>
## 1      103.        2 4.55e-23

TitanicT %>%
  ggplot(aes(x=as.factor(Pclass), y = Age)) +
  geom_boxplot()

## Warning: Removed 177 rows containing non-finite values (`stat_boxplot()`).
```



```
TitanicT %>% group_by(Pclass) %>% summarise(meanAge = mean(Age, na.rm = T))
```

```
## # A tibble: 3 x 2
##   Pclass meanAge
##   <dbl>   <dbl>
## 1     1     38.2
## 2     2     29.9
## 3     3     25.1
```

```
mdl_Age_Pclass <- lm(Age ~ Pclass, data = TitanicT) # variable_num ~ variable_cat
anova(mdl_Age_Pclass)
```

```
## Analysis of Variance Table
```

```
##
```

```
## Response: Age
```

```
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Pclass      1  20511  20511.4   112.39 < 2.2e-16 ***
## Residuals 712  129945    182.5
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
pairwise.t.test(TitanicT$Age, TitanicT$Pclass, p.adjust.method = "none")
```

```
##
```

```
## Pairwise comparisons using t tests with pooled SD
```

```
##
```

```
## data: TitanicT$Age and TitanicT$Pclass
```


```
##
```

```
##      1      2
```



```
## 2 7e-09 -
## 3 < 2e-16 0.00017
##
## P value adjustment method: none
```

```
TitanicT %>%
  ggplot(aes(Pclass, fill = as.factor(Survived)))+
  geom_bar(position = "fill")+
  ylab("proportion")
```



```
TitanicT %>% group_by(Pclass) %>% summarise(meanSurv = mean(Survived, na.rm = T))
```


```
## # A tibble: 3 x 2
##   Pclass meanSurv
##   <dbl>   <dbl>
## 1     1     0.630
## 2     2     0.473
## 3     3     0.242
```

```
pairwise.t.test(TitanicT$Survived, TitanicT$Pclass, p.adjust.method = "none")
```

```
##
## Pairwise comparisons using t tests with pooled SD
##
## data: TitanicT$Survived and TitanicT$Pclass
##
##    1      2
## 2 0.00068 -
## 3 < 2e-16 8.2e-09
```

```
##
## P value adjustment method: none
```

```
TitanicT %>%
  ggplot(aes(Pclass, fill = Sex ))+
  geom_bar(position = "fill")+
  ylab("proporcion")
```



```
TitanicT %>% mutate(Sex=(Sex=="female")) %>% group_by(Pclass) %>% summarise(meanSex = mean(Sex, na.rm = TRUE))
```

```
## # A tibble: 3 x 2
##   Pclass meanSex
##   <dbl>   <dbl>
## 1     1     0.435
## 2     2     0.413
## 3     3     0.293
```

```
TitanicT %>% mutate(Sex=(Sex=="female")) %>% with(pairwise.t.test(Sex,Pclass, p.adjust.method = "none"))
```

```
##
## Pairwise comparisons using t tests with pooled SD
##
## data: Sex and Pclass
##
##    1      2
## 2 0.64156 -
## 3 0.00026 0.00355
##
## P value adjustment method: none
```