Deviation

Emphasise variations (+/-) from a fixed reference point. Typically the reference point is zero but it can also be a target or a long-term average. Can also be used to show sentiment (positive/neutral/negative).

Example FT uses Trade surplus/deficit, climate change

Diverging bar




chart that can handle both negative and positive magnitude

Diverging stacked bar



Splits a single value into two contrasting components (eg male/female).



The shaded area of these charts allows a balance to be shown either against a baseline or between two series.

Visual

money with the Financial Times.

Inspired by the Graphic Continuum by Jon Schwabish and Severino Ribecca


a login by scanning here.

Correlation

Show the relationship between two or more variables. Be mindful that, unless you tell them otherwise, many readers will assume the relationships you show them to be causal (i.e. one causes the


Example FT uses Inflation and unemployment, income and life expectancy

Scatterplot




show the relationship between two continuous variables, each of which

Column + line timeline




an amount (columns) and a rate (line).

Connected scatterplot




Usually used to show how the relationship between 2 variables has changed over time.




Like a scatterplot, but adds additional detail by sizing the circles according to a third


XY heatmap





A good way of showing the patterns between 2 effective at showing fine




The standard way to









categories of data, less differences in amounts.

Ranking

Use where an item's position in an ordered list is more important than its absolute or relative value. Don't be afraid to highlight the points of interest.

Example FT uses Wealth, deprivation, league tables constituency election results

Ordered bar




Standard bar charts display the ranks of values much more easily when sorted into order.

Ordered column


See above.

Ordered proportional symbol



Use when there are big variations between values and/or seeing fine differences between data is not so


Dot strip plot



Dots placed in order on a strip are a space-efficient ... method of laying out ranks across multiple categories.



Perfect for showing how ranks have changed over time or vary between categories.



Lollipops draw more attention to the data value than standard bar/column and can also show rank and value effectively.




Effective for showing changing rankings across multiple dates. For large datasets, consider grouping lines using colour.

Distribution

Show values in a dataset and how often they occur. The shape (or 'skew') of a distribution can be a memorable way of highlighting the lack of uniformity or equality in the data.


Example FT uses Income distribution, population (age/sex) distribution, revealing inequality

Histogram




The standard way to show a statistical distribution - keep the gaps between columns small to highlight the shape' of the data.

Dot plot



Dot strip plot




Good for showing individual values in a distribution, can be a problem when too many dots have the same value.

Barcode plot



Like dot strip plots, good for displaying all the data in a table, they work best when highlighting individual

Boxplot




Summarise multiple distributions by showing the median (centre) and range of the data

Violin plot




Similar to a box plot but more effective with complex distributions (data that cannot be summarised with simple average).

Population pyramid




A standard way for showing the age and sex breakdown of a population distribution; effectively, back to back

Cumulative curve



A good way of showing how unequal a distribution is: y axis is always cumulative frequency, x axis is always a measure.

Frequency polygons



For displaying multiple distributions of data. Like a regular chart, best limited to a maximum of 3 or 4 datasets.

Beeswarm




Use to emphasise individual points in a distribution. Points can be sized to an additional variable. Best with mediumsized datasets

Change over Time

Give emphasis to changing trends These can be short (intra-day) movements or extended series traversing decades or centuries: Choosing the correct time period is important to provide suitable context

Example FT uses Share price movements, economic time series, sectoral changes in a market

for the reader.




The standard way to show a changing time series. If data are irregular, consider markers to represent


Column

Columns work well for showing change over time - but usually best with only one series of

Column + line timeline



A good way of showing the relationship over time between an amount (columns) and a rate




Good for showing changing data as long as the data can be simplified into 2 or 3 points without missing a key part of story.

Area chart




Use with care – these are good at showing changes to total, but seeing change in components can be very difficult.

Candlestick




Usually focused on day-to-day activity, these charts show opening/closing and high/low points of each day.

Fan chart (projections)




Use to show the uncertainty in future projections - usually this grows the further forward to projection.

Connected scatterplot



A good way of showing changing data for two variables whenever there is a relatively clear pattern of

Calendar heatmap




A great way of showing temporal patterns (daily, weekly, monthly) – at the expense of showing precision in

Priestley timeline



Great when date and duration are key elements of the story in the data.

Circle timeline



Good for showing discrete values of varying size across multiple categories (eg earthquakes by

Presents time on the Y

axis. Good for


displaying detailed

time series that work

especially well when


scrolling on mobile.

Vertical timeline



Another alternative to the circle timeline for showing series where there are big variations in the data.

Streamgraph



changes in proportions over time is more important than individual values


Magnitude

Show size comparisons. These can be relative (just being able to see larger/bigger) or absolute (need to see fine differences). Usually these show a 'counted' number (for example, barrels dollars or people) rather than a

Example FT uses Commodity production, market capitalisation, volumes in general

calculated rate or per cent.

Column



See above. Good when

the data are not time


series and labels have

long category names.

compare the size of

things. Must always

Paired column



As per standard column but allows for multiple series. Can become tricky to read with more than 2

See above.

Paired bar




Marimekko




showing the size and proportion of data at the same time – as long as the data are not too complicated

Proportional symbol



Use when there are big variations between values and/or seeing fine differences between data is not so important.

Isotype (pictogram)



Lollipop charts draw more attention to the data value than standard bar/column -


does not have to start a

zero (but preferable).



A space-efficient way of showing value of multiple variables – but make sure they are organised in a way that makes sense to reader.

Parallel coordinates An alternative to radar



Good for showing a measurement against the context of a targe or performance range


arrangement of the variables is important.

Usually benefits from

highlighting values.

charts – again, the arrangement of the variables is importan

Grouped symbol



useful.

An alternative to

being able to count

individual elements is

data or highlight

bar/column charts when


Part-to-whole

Show how a single entity can be broken down into its component elements. If the reader's interest is solely in the size of the components, consider a magnitude-type chart instead.

Example FT uses Fiscal budgets, company structures,

Stacked column/bar

national election results



A good way of showing the size and proportion of data at

the same time – as


ong as the data are not too complicated

Marimekko



data – but be aware that it's difficult to accurately compare the size of the

Donut



space to include more information about the data (eg total).

relationships; can be

difficult to read when

there are many small

Use for hierarchical


part-to-whole

good way of making

Treemap



A way of turning points into areas – any point within each area is closer to the central point than any other



A hemicycle, often used for visualising parliamentary composition by number of seats.


Good for showing %

best when used on

whole numbers and

work well in small

information, they work



Generally only used for schematic representation.

Can be useful for

some of the


negative.

components are

showing part-to-whole

relationships where

Waterfall




Spatial


Aside from locator maps only used when precise locations or geographical patterns in data are more important to the reader than anything else.

Example FT uses Population density, natural resource locations, natural disaster risk/impact, catchment areas, variation in election


Basic choropleth (rate/ratio)



Proportional symbol (count/magnitude)



Flow map



Contour map For showing areas of



Equalised cartogram

good for representing voting regions with equal value.

Converting each unit on

a map to a regular and

equally-sized shape –

Stretching and


Used to show the

any patterns the

reader should see.

location of individual

make sure to annotate



Dot density

Heat map




Grid-based data values mapped with an intensity colour scale.

As choropleth map -

but not snapped to an

admin/political unit.



Flow

Show the reader volumes or intensity of


movement between two or more states

or conditions. These might be logical


sequences or geographical locations.

Sankey

Example FT uses




Waterfall






Chord



Network




Used for showing the

FINANCIAL TIMES







and geographical learning

Royal

© Financial Times 2016-2019. This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

FT graphic: Alan Smith; Chris Campbell; Ian Bott; Liz Faunce; Graham Parrish; Billy Ehrenberg-Shannon; Paul McCallum; Martin Stabe

ft.com/schoolsarefree

Vocabulary

Read the FT for free

Add fresh examples and insights to your work, make more

informed career choices, and feel more confident managing

Teachers can register your school and students can create