

Review of the paper:

Mathematical modelling of cancer cell invasion of tissue: Local and non-local models and the effect of adhesion

(BY A. GERISCH, M.A.J. CHAPLAIN)

Julia N. Navarro

School of Mathematics and Statistics

University of St. Andrews

2022-2023

Candlemas Semsester



University of
St Andrews

Motivation

(and previous work done)

- ▶ Cancer spread depends on a multitude of factors such as type of cancer (e.g. after a mutation)
- ▶ Invasion rate facilitated by enzymes that destroy the surrounding tissue
- ▶ Previous papers (such as by Anderson et al. (2000)) have modelled the migration of cancer cells by haptotaxis
- ▶ Armstrong et al. (2006) has introduced cell motility by inclusion of non-local terms
- ▶ Authors choose non-motile ECM population (**only** changes due to degradation by enzymes) and a cancer population driven by adhesive effects
- ▶ Researchers of this paper take the (so-called local) model and modify it and compare it to a non-local model with more parameters (adhesion interactions between cells and cells and the matrix)
- ▶ Look at how rate of migration is affected via numerical simulations

Motivation

(and previous work done)

- ▶ Cancer spread depends on a multitude of factors such as type of cancer (e.g. after a mutation)
- ▶ Invasion rate facilitated by enzymes that destroy the surrounding tissue
- ▶ Previous papers (such as by Anderson et al. (2000)) have modelled the migration of cancer cells by haptotaxis
- ▶ Armstrong et al. (2006) has introduced cell motility by inclusion of non-local terms
- ▶ Authors choose non-motile ECM population (**only** changes due to degradation by enzymes) and a cancer population driven by adhesive effects
- ▶ Researchers of this paper take the (so-called local) model and modify it and compare it to a non-local model with more parameters (adhesion interactions between cells and cells and the matrix)
- ▶ Look at how rate of migration is affected via numerical simulations

Motivation

(and previous work done)

- ▶ Cancer spread depends on a multitude of factors such as type of cancer (e.g. after a mutation)
- ▶ Invasion rate facilitated by enzymes that destroy the surrounding tissue
- ▶ Previous papers (such as by Anderson et al. (2000)) have modelled the migration of cancer cells by haptotaxis
- ▶ Armstrong et al. (2006) has introduced cell motility by inclusion of non-local terms
- ▶ Authors choose non-motile ECM population (**only** changes due to degradation by enzymes) and a cancer population driven by adhesive effects
- ▶ Researchers of this paper take the (so-called local) model and modify it and compare it to a non-local model with more parameters (adhesion interactions between cells and cells and the matrix)
- ▶ Look at how rate of migration is affected via numerical simulations



Motivation

(and previous work done)

- ▶ Cancer spread depends on a multitude of factors such as type of cancer (e.g. after a mutation)
- ▶ Invasion rate facilitated by enzymes that destroy the surrounding tissue
- ▶ Previous papers (such as by Anderson et al. (2000)) have modelled the migration of cancer cells by haptotaxis
- ▶ Armstrong et al. (2006) has introduced cell motility by inclusion of non-local terms
- ▶ Authors choose non-motile ECM population (**only** changes due to degradation by enzymes) and a cancer population driven by adhesive effects
- ▶ Researchers of this paper take the (so-called local) model and modify it and compare it to a non-local model with more parameters (adhesion interactions between cells and cells and the matrix)
- ▶ Look at how rate of migration is affected via numerical simulations

Motivation

(and previous work done)

- ▶ Cancer spread depends on a multitude of factors such as type of cancer (e.g. after a mutation)
- ▶ Invasion rate facilitated by enzymes that destroy the surrounding tissue
- ▶ Previous papers (such as by Anderson et al. (2000)) have modelled the migration of cancer cells by haptotaxis
- ▶ Armstrong et al. (2006) has introduced cell motility by inclusion of non-local terms
- ▶ Authors choose non-motile ECM population (**only** changes due to degradation by enzymes) and a cancer population driven by adhesive effects
- ▶ Researchers of this paper take the (so-called local) model and modify it and compare it to a non-local model with more parameters (adhesion interactions between cells and cells and the matrix)
- ▶ Look at how rate of migration is affected via numerical simulations

Models

The local and non-local models

- ▶ Use the following notation for the PDEs:
 - * $c(t, x)$ - the cancer cell density
 - * $v(t, x)$ - extra-cellular matrix density
 - * $m(t, x)$ - matrix-degrading enzyme density
- ▶ After non-dimensionalizing the equations (for both models) we have the following systems:
- ▶ Some changes to the local model: the $(1 - c - v)$ terms ensures no overcrowding (**volume filling term**)
- ▶ Note the similarities and differences: main change is the flux for the cancer density equation $\chi_{12}(1 - c - v)$ replaced by \mathcal{A} and in the left model we have slightly different expressions to previous papers (underlined in orange)



Models

The local and non-local models

- ▶ Use the following notation for the PDEs:
 - * $c(t, x)$ - the cancer cell density
 - * $v(t, x)$ - extra-cellular matrix density
 - * $m(t, x)$ - matrix-degrading enzyme density
- ▶ After non-dimensionalizing the equations (for both models) we have the following systems:

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - \chi_{12}(1 - c - v)c \nabla v] + \mu_1 c(1 - c - v), \quad (2a)$$

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - c \mathcal{A}\{\underline{u}(t, \cdot)\}] + \mu_1 c(1 - c - v), \quad (4a)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (2b)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (4b)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (2c)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (4c)$$

- ▶ Some changes to the local model: the $(1 - c - v)$ terms ensures no overcrowding (volume filling term)
- ▶ Note the similarities and differences: main change is the flux for the cancer density equation $\chi_{12}(1 - c - v)$ replaced by \mathcal{A} and in the left model we have slightly different expressions to previous papers (underlined in orange)

Models

The local and non-local models

- ▶ Use the following notation for the PDEs:
 - * $c(t, x)$ - the cancer cell density
 - * $v(t, x)$ - extra-cellular matrix density
 - * $m(t, x)$ - matrix-degrading enzyme density
- ▶ After non-dimensionalizing the equations (for both models) we have the following systems:

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - \chi_{12}(1 - c - v)c \nabla v] + \mu_1 c(1 - c - v), \quad (2a)$$

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - c \mathcal{A}(\underline{u}(t, \cdot))] + \mu_1 c(1 - c - v), \quad (4a)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (2b)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (4b)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (2c)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (4c)$$

- ▶ Some changes to the local model: the $(1 - c - v)$ terms ensures no overcrowding (**volume filling term**)
- ▶ Note the similarities and differences: main change is the flux for the cancer density equation $\chi_{12}(1 - c - v)$ replaced by \mathcal{A} and in the left model we have slightly different expressions to previous papers (underlined in orange)

Models

The local and non-local models

- ▶ Use the following notation for the PDEs:
 - * $c(t, x)$ - the cancer cell density
 - * $v(t, x)$ - extra-cellular matrix density
 - * $m(t, x)$ - matrix-degrading enzyme density
- ▶ After non-dimensionalizing the equations (for both models) we have the following systems:

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - \chi_{12}(1 - c - v)c \nabla v] + \mu_1 c(1 - c - v), \quad (2a)$$

$$\frac{\partial c}{\partial t} = \nabla \cdot [D_1 \nabla c - c \mathcal{A}\{\underline{u(t, \cdot)}\}] + \mu_1 c(1 - c - v), \quad (4a)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (2b)$$

$$\frac{\partial v}{\partial t} = -\gamma m v + \mu_2(1 - c - v), \quad (4b)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (2c)$$

$$\frac{\partial m}{\partial t} = \nabla \cdot [D_3 \nabla m] + \alpha c - \lambda m. \quad (4c)$$

- ▶ Some changes to the local model: the $(1 - c - v)$ terms ensures no overcrowding (**volume filling term**)
- ▶ Note the similarities and differences: main change is the flux for the cancer density equation $\chi_{12}(1 - c - v)$ replaced by \mathcal{A} and in the left model we have slightly different expressions to previous papers (underlined in orange)



Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) \, dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(\cdot)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)
- ▶ Two types of Ω :
 - * Ω_1 : all points in sensing disk have equal share in direction of movements
 - * Ω_2 : sensitivity decreases the further away points are from centre
- ▶ Ω_1 in 1D and 2D is: **(independent of r)**
- ▶ And Ω_2 is: **(decays linearly)**

Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

$$g(c, v) = (S_{cc}c + S_{cv}v) \cdot (1 - c - v)^+$$

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(\cdot)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)
- ▶ Two types of Ω :
 - * Ω_1 : all points in sensing disk have equal share in direction of movements
 - * Ω_2 : sensitivity decreases the further away points are from centre
- ▶ Ω_1 in 1D and 2D is: (independent of r)
- ▶ And Ω_2 is: (decays linearly)



Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

$$g(c, v) = (S_{cc}c + S_{cv}v) \cdot (1 - c - v)^+$$

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(\cdot)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)
- ▶ Two types of Ω :
 - * Ω_1 : all points in sensing disk have equal share in direction of movements
 - * Ω_2 : sensitivity decreases the further away points are from centre
- ▶ Ω_1 in 1D and 2D is: (independent of r)
- ▶ And Ω_2 is: (decays linearly)



Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

$$g(c, v) = (S_{cc}c + S_{cv}v) \cdot (1 - c - v)^+$$

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(\cdot)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)
- ▶ Two types of Ω :
 - * Ω_1 : all points in sensing disk have equal share in direction of movements
 - * Ω_2 : sensitivity decreases the further away points are from centre
- ▶ Ω_1 in 1D and 2D is: (independent of r)
- ▶ And Ω_2 is: (decays linearly)



Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

$$g(c, v) = (S_{cc}c + S_{cv}v) \cdot (1 - c - v)^+$$

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(\cdot)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)
- ▶ Two types of Ω :
 - * Ω_1 : all points in sensing disk have equal share in direction of movements
 - * Ω_2 : sensitivity decreases the further away points are from centre
- ▶ Ω_1 in 1D and 2D is: $\Omega_1(r) = \frac{1}{2R}$ and $\Omega_1(r) = \frac{1}{\pi R^2}$, **(independent of r)**
- ▶ And Ω_2 is: **(decays linearly)**



Model parameters

The functions $g(u)$ and $\Omega(r)$

- ▶ In the non-local model, \mathcal{A} (adhesion velocity) is an integral of a sum of a product between a function $g(u)$ and $\Omega(r)$ defined as:

$$\mathcal{A}\{\underline{u}(t, \cdot)\}(\underline{x}) := \frac{1}{R} \int_0^R \sum_{k=0}^1 \underline{\eta}(k) \cdot \Omega(r) g(\underline{u}(t, \underline{x} + r \underline{\eta}(k))) dr,$$

where $\eta(k) = (-1)^k$

- ▶ The $g(u)$ equation is defined as

$$g(c, v) = (S_{cc}c + S_{cv}v) \cdot (1 - c - v)^+$$

where S_{cc} and S_{cv} are adhesion parameters

- ▶ The second bracket $(.)^+$ ensures that densely packed areas do not influence in direction of invasion (to avoid aggregation)

- ▶ Two types of Ω :

- * Ω_1 : all points in sensing disk have equal share in direction of movements
- * Ω_2 : sensitivity decreases the further away points are from centre

- ▶ Ω_1 in 1D and 2D is: $\Omega_1(r) = \frac{1}{2R}$ and $\Omega_1(r) = \frac{1}{\pi R^2}$, **(independent of r)**

- ▶ And Ω_2 is: $\Omega_2(r) = \frac{1}{R} \left(1 - \frac{r}{R}\right)$ and $\Omega_2(r) = \frac{3}{\pi R^2} \left(1 - \frac{r}{R}\right)$ **(decays linearly)**

Numerical Results

Travelling wave cluster

- ▶ Researchers show results for the two coefficients (non-local model)
- ▶ Also switch on MDE degradation term to avoid build-up of the enzyme concentration ($\mu_2 > 0$)
- ▶ Choose Ω_2 (more realistic), $R = 0.1$ (larger R leads to slower invasion) and $S_{cv} = S_{cc} = 0.1$ (same value):
- ▶ True detachment of cells travelling in wave-like manner along matrix edge
- ▶ Also have a population around $x = 0$



Numerical Results

Travelling wave cluster


- ▶ Researchers show results for the two coefficients (non-local model)
- ▶ Also switch on MDE degradation term to avoid build-up of the enzyme concentration ($\mu_2 > 0$)
- ▶ Choose Ω_2 (more realistic), $R = 0.1$ (larger R leads to slower invasion) and $S_{cv} = S_{cc} = 0.1$ (same value):
- ▶ True detachment of cells travelling in wave-like manner along matrix edge
- ▶ Also have a population around $x = 0$



Numerical Results

Travelling wave cluster

- ▶ Researchers show results for the two coefficients (non-local model)
- ▶ Also switch on MDE degradation term to avoid build-up of the enzyme concentration ($\mu_2 > 0$)
- ▶ Choose Ω_2 (more realistic), $R = 0.1$ (larger R leads to slower invasion) and $S_{cv} = S_{cc} = 0.1$ (same value):




- ▶ True detachment of cells travelling in wave-like manner along matrix edge
- ▶ Also have a population around $x = 0$

Numerical Results

Travelling wave cluster

- ▶ Researchers show results for the two coefficients (non-local model)
- ▶ Also switch on MDE degradation term to avoid build-up of the enzyme concentration ($\mu_2 > 0$)
- ▶ Choose Ω_2 (more realistic), $R = 0.1$ (larger R leads to slower invasion) and $S_{cv} = S_{cc} = 0.1$ (same value):




- ▶ True detachment of cells travelling in wave-like manner along matrix edge
- ▶ Also have a population around $x = 0$

Numerical Results

Travelling wave cluster

- ▶ Researchers show results for the two coefficients (non-local model)
- ▶ Also switch on MDE degradation term to avoid build-up of the enzyme concentration ($\mu_2 > 0$)
- ▶ Choose Ω_2 (more realistic), $R = 0.1$ (larger R leads to slower invasion) and $S_{cv} = S_{cc} = 0.1$ (same value):




- ▶ True detachment of cells travelling in wave-like manner along matrix edge
- ▶ Also have a population around $x = 0$

Numerical Results

One parameter fixed, one varied

- ▶ Paper now fixes one of the parameters to 0.5 and varies the other on $[0.02, 0.5]$




- ▶ The left plot has $S_{cc} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cv} < 0.5$ most of the tumour cells remain at $x = 0$ with few migrating
 - ▶ for $S_{cv} = 0.5$ all at centre ($x = 0$)
 - ▶ for $S_{cv} > 0.5$ gives rise to only migrating cells
- ▶ The right plot has $S_{cv} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cc} < 0.5$ cells mostly invade matrix tissue with few at $x = 0$ at constant speed
 - ▶ for $S_{cc} > 0.5$ no cells migrating, all remain at $x = 0$
- ▶ Summary: If $S_{cv} > S_{cc}$ we have an invasive tumour. If $S_{cv} < S_{cc}$ the tumour is compact (see plot)

Numerical Results

One parameter fixed, one varied

- ▶ Paper now fixes one of the parameters to 0.5 and varies the other on $[0.02, 0.5]$




- ▶ The left plot has $S_{cc} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cv} < 0.5$ most of the tumour cells remain at $x = 0$ with few migrating
 - ▶ for $S_{cv} = 0.5$ all at centre ($x = 0$)
 - ▶ for $S_{cv} > 0.5$ gives rise to only migrating cells
- ▶ The right plot has $S_{cv} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cc} < 0.5$ cells mostly invade matrix tissue with few at $x = 0$ at constant speed
 - ▶ for $S_{cc} > 0.5$ no cells migrating, all remain at $x = 0$
- ▶ Summary: If $S_{cv} > S_{cc}$ we have an invasive tumour. If $S_{cv} < S_{cc}$ the tumour is compact (see plot)

Numerical Results

One parameter fixed, one varied

- ▶ Paper now fixes one of the parameters to 0.5 and varies the other on $[0.02, 0.5]$




- ▶ The left plot has $S_{cc} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cv} < 0.5$ most of the tumour cells remain at $x = 0$ with few migrating
 - ▶ for $S_{cv} = 0.5$ all at centre ($x = 0$)
 - ▶ for $S_{cv} > 0.5$ gives rise to only migrating cells
- ▶ The right plot has $S_{cv} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cc} < 0.5$ cells mostly invade matrix tissue with few at $x = 0$ at constant speed
 - ▶ for $S_{cc} > 0.5$ no cells migrating, all remain at $x = 0$
- ▶ Summary: If $S_{cv} > S_{cc}$ we have an invasive tumour. If $S_{cv} < S_{cc}$ the tumour is compact (see plot)

Numerical Results

One parameter fixed, one varied

- ▶ Paper now fixes one of the parameters to 0.5 and varies the other on $[0.02, 0.5]$



- ▶ The left plot has $S_{cc} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cv} < 0.5$ most of the tumour cells remain at $x = 0$ with few migrating
 - ▶ for $S_{cv} = 0.5$ all at centre ($x = 0$)
 - ▶ for $S_{cv} > 0.5$ gives rise to only migrating cells
- ▶ The right plot has $S_{cv} = 0.5$ and varies the other, and we observe:
 - ▶ for $S_{cc} < 0.5$ cells mostly invade matrix tissue with few at $x = 0$ at constant speed
 - ▶ for $S_{cc} > 0.5$ no cells migrating, all remain at $x = 0$
- ▶ Summary: If $S_{cv} > S_{cc}$ we have an invasive tumour. If $S_{cv} < S_{cc}$ the tumour is compact (see plot)

Numerical Results in 2D


Steady-state cancer

- ▶ Now look at non-local model in 2D with a heterogeneous matrix density
- ▶ Note top row is cancer cell density, middle row is matrix density and bottom row is the enzyme concentration
- ▶ Reaches a steady-state at around $t = 30$
- ▶ Note elongation at $t = 10$ (cell-cell adhesion pulls cells together afterwards)
- ▶ Not really a lot of diffusion of the enzymes
- ▶ Remains a steady-state non-invasive tumour ... unless **key** parameters change

Numerical Results in 2D

Steady-state cancer

- ▶ Now look at non-local model in 2D with a heterogeneous matrix density
- ▶ Note top row is cancer cell density, middle row is matrix density and bottom row is the enzyme concentration




- ▶ Reaches a steady-state at around $t = 30$
- ▶ Note elongation at $t = 10$ (cell-cell adhesion pulls cells together afterwards)
- ▶ Not really a lot of diffusion of the enzymes
- ▶ Remains a steady-state non-invasive tumour ... unless **key** parameters change

Numerical Results in 2D

Steady-state cancer

- ▶ Now look at non-local model in 2D with a heterogeneous matrix density
- ▶ Note top row is cancer cell density, middle row is matrix density and bottom row is the enzyme concentration




- ▶ Reaches a steady-state at around $t = 30$
- ▶ Note elongation at $t = 10$ (cell-cell adhesion pulls cells together afterwards)
- ▶ Not really a lot of diffusion of the enzymes
- ▶ Remains a steady-state non-invasive tumour ... unless **key** parameters change

Numerical Results in 2D

Mutation: Invasion of matrix by cancer cells

- ▶ To, again, “imitate” a mutation we change the parameters for proliferation and decrease cell-cell adhesion at $t = 30$:




- ▶ Observe how the cancer, its enzymes diffuse **asymmetrically**
- ▶ At $t = 50$ see the top row, cancer does not invade low density area in matrix and at $t = 60$ it avoids high density regions
- ▶ Matrix degrades and cancer moves into it

Numerical Results in 2D

Mutation: Invasion of matrix by cancer cells

- ▶ To, again, “imitate” a mutation we change the parameters for proliferation and decrease cell-cell adhesion at $t = 30$:



- ▶ Observe how the cancer, its enzymes diffuse **asymmetrically**
- ▶ At $t = 50$ see the top row, cancer does not invade low density area in matrix and at $t = 60$ it avoids high density regions
- ▶ Matrix degrades and cancer moves into it

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)

Summary

- ▶ This paper presented a modified local model and a non-local model (with adhesion parameters and replaced haptotaxis migration by \mathcal{A})
- ▶ Have briefly discussed the non-local term \mathcal{A}
- ▶ Non-local model gave rise to cells travelling in wave-like manner (where $S_{cc} = S_{cv}$)
- ▶ Demonstrated these are important in determining if tumour is aggressive (relative magnitudes of the adhesion parameters)
 - * e.g., spreads when $S_{cv} > S_{cc}$
 - * and remains compact for $S_{cv} < S_{cc}$
- ▶ Considered 2D case with an imitated mutation (parameter change)
- ▶ Conclude that adhesion terms influence both the cancer migration rate (not discussed here) and the cancer type (benign or aggressive)