Wildlife Ecology

Week 10 Assignment Solutions (2026)

1 A community already affected by disease, weeds, pollutants, or livestock pressure is best described as which type of community?

(a) Pristine community

(b) Climax community

(c) Resistant community

(d) **Disturbed community**

2 When an organism absorbs a toxic substance into its tissues faster than it loses it, the process is called:

(a) Biomagnification

(b) Bioavailability

(c) **Bioaccumulation**

(d) Biodegradation

3 What percentage of marine plastic debris is stated to sink to the ocean bottom?

(a) 15%

(b) 30%

(c) 50%

(d) **70%**

4 Hydrocarbons derived directly from mineral oils are called:

(a) Pyrogenic hydrocarbons

(b) Biogenic hydrocarbons

(c) Aromatic hydrocarbons

(d) **Petrogenic hydrocarbons**

5 “Any changes in natural or human systems that inadvertently increase vulnerability to climatic stimuli; an adaptation that does not succeed in reducing vulnerability but increases it instead” is a definition for

a. adaptation

b. mitigation

c. **maladaptation**

d. malmitigation

6 A normal ecological community faces one large and infrequent disturbance and then returns close to its earlier condition. Which outcome is best represented?

(a) Long-term alteration

(b) **Recovery**

(c) Bioaccumulation

(d) Habitat displacement

7 Macrodebris in the context of plastic debris has fragments of size

a. **> 20 mm**

b. 5 - 20 mm

c. < 5 mm

d. < 1 mm

8 The Deepwater Horizon incident is classified by origin as which kind of oil spill?

(a) Natural

(b) **Accidental**

(c) Intentional

(d) Biogenic

9 Harmful effects caused by a single or short-term exposure are called:

(a) Chronic toxicity

(b) **Acute toxicity**

(c) Sub-lethal exposure

(d) Bioavailability

10 Mesodebris in the context of plastic debris has fragments of size

a. > 20 mm

b. **5 - 20 mm**

c. < 5 mm

d. < 1 mm