An Introduction to Higher Algebraic K-Theory

Naveen Maurya

November 12, 2026

Contents

Chapter 1

Projective Modules

The subject of K-Theory concerns with studying vector bundles on spaces. Algebraic K-Theory studies the algebraic analogue of the same, projective modules over associative rings.

1.1 Projective Modules

1.1.1 Idempotence Criteria for Projective Modules

1.1.2 Constructing Projective Modules

- Patching along localizations
- Milnor Patching

1.1.3 Classification of Projective Modules

- Projective modules ¡-¿ vector bundles -

Chapter 2

The Grothendieck Group K_0

In this chapter, we will develop the notion of the Grothendieck Group K_0 accross a variety of contexts. First, we discuss the grothendieck group of a ring, then symmetric monoidal categories.

2.1 K_0 of a ring

Let R be a (unital) ring. The interesting objects of study will be finitely generated projective modules over R. We denote by $\mathbf{P}(R)$ the set of isomorphism classes of fintely generated projective modules over R. Note the $\mathbf{P}(R)$ derives a monoid structure through $\oplus$ because direct sums yield an associative operation that preserves isomorphisms. This monoid is also abelian due to the isomorphisms $M \oplus N \cong N \oplus M$ for any R-modules.

The idea is to consider the group naturally derived by implementing inverses for all elements in this monoid.

2.1.1 Group Completion of a Monoid

Given an (abelian) monoid $(M, +)$, we can derive an (abelian) group associated to M, which we shall call $\tilde{M}$, via the following procedure:

Define an equivalence relation on $M \times M$ by

$$(a, b) \sim (c, d) \Leftrightarrow a + d + e = b + c + e \text{ for some } e \text{ in } M.$$

This indeed gives us an equivalence relation. Let $\tilde{M}$ be the set of equivalence classes of $M \times M$ under this relation. This set admits a natural (abelian) group operation defined by


$$[(a, b)] + [(c, d)] := [(a + c, b + d)]$$

This is a well-defined, abelian and associative operation on $\tilde{M}$ with the identity element $0 := [(0,0)]$ represented by any (l,l). Each element $[(a,b)]$ admits an inverse given by $[(b,a)]$ as $[(a,b)] + [(b,a)] = [(a+b, b+a)] = 0$.

A very familiar example of this construction is the abelian group $\mathbb{Z}$, which is the group completion of the abelian monoid $\mathbb{N}$ under $+$.

This construction is natural in M and follows a suitable universal property which we shall state.

Proposition 2.1 (Universal property of Group Completion)**.** We have a monoid homomorphism $i : M \to \tilde{M}$ such that for any (abelian) group G, and a monoid homomorphism $\varphi : M \to G$, we have that φ must factor uniquely through i. That is, there is a unique map $\tilde{\varphi} : \tilde{M} \to G$ such that the following diagram commutes:



Proof. The monoid homomorphism $i : M \to \tilde{M}$ is simply given by mapping a to $[(a,0)]$. This is indeed a homomorphism since $[(a,0)] + [(b,0)] = [(a+b,0)]$. Now suppose we are given $\varphi : M \to G$. Then $\tilde{\varphi} : \tilde{M} \to G$ is given by $[(a,b)] \mapsto \varphi(a) - \varphi(b)$, making the diagram commute. This map is unique since for any $a \in M$, we have $\tilde{\varphi}([(a,0)]) = \varphi(a)$, and consequently $\tilde{\varphi}([(0,a)]) = -\tilde{\varphi}([(a,0)]) = -\varphi(a)$, which altogether determines the map $\tilde{\varphi}$. □

Furthermore, any abelain group satisfying the given universal property is isomorphic to $\tilde{M}$ via a unique isomorphism.

There is an alternative (but equivalent) characterization of $\tilde{M}$ via the quotienting the free group on M by the monoidal relations.

Proposition 2.2. Let $K(M)$ be the abelian group given by taking the free abelian group $F(M)$ on the set M and quotienting by all relations of the form $[a] + [b] - [a+b]$ where $a, b \in M$ and $[x]$ represents the element in $F(M)$ corresponding to the element $x \in M$. Then $K(M) \cong \tilde{M}$.

Proof. We will show that $K(M)$ satisfies the universal property of $\tilde{M}$. Let $F(M)$ represent the free abelian group on the set M. The map $i : M \to K(M)$ on x is given by taking the equivalence class of the element x. Suppose $\varphi : M \to H$ is a monoid homomorphism to an abelian group H. Then define $\tilde{\varphi}([a])$ as $\varphi(a)$. Suppose $[a] = [b]$, then $[a] + [-b] = [0]$ so $a - b = 0$ and consequently $\varphi(a) - \varphi(b) = 0$. Thus the map $\tilde{\varphi}$ is well-defined

and clearly a group homomorphism. It is also unique since $\tilde{\varphi}$ is specified uniquely on the basis of $F(M)$. □

Example 2.1.1. Let R be a (unital) ring satisfying the *Invariant Basis Property (IBP)*. That is, $R^m \cong R^n$ implies $m = n$. Then the monoid of isomorphism classes of free R-modules with the operation $\oplus$ is isomorphic to $\mathbb{N}$, the isomorphism given by the map $R^n \mapsto n$. Consequently, its group completion is isomorphic to the abelian group $\mathbb{Z}$.

Note that all fields (and division rings) F satisfy IBP since dimension of an F-module is independent of the choice of basis. Further, any ring R with a map $R \to F$, a field or a division ring F satisfies IBP. In particular, commutative rings all satisfy IBP.

Example 2.1.2. Let X be a topological space. Then the set $[X, \mathbb{N}]$ of continuous maps $X \to \mathbb{N}$ forms an abelian monoid under pointwise addition of maps. Then the group completion of this monoid is isomorphic to the abelian group of continuous maps $X \to \mathbb{Z}$ denoted $[X, \mathbb{Z}]$. If X is quasi-compact, this is in fact a free abelian group. Notably, $[X, \mathbb{N}]$ also admits a commutative semiring structure via pointwise multiplication in addtion to the monoid operation. This makes $[X, \mathbb{Z}]$ into a commutative ring.

2.1.2 Defining $K_0(R)$

Now we can define the group $K_0(R)$ as the group completion of the monoid $\mathbf{P}(R)$ consisting of finitely generated projective R-modules with the monoid operation given by $\oplus$.

Example 2.1.3. Let F be a field or a division ring. Then all finitely generated projective F-modules are free. This gives us $K_0(F) \cong \mathbb{Z}$. More generally, for any local ring R, all finitely generated projective R-modules are free. This gives us $K_0(R) \cong \mathbb{Z}$.

Example 2.1.4. Note that finitely generated projectives over $\mathbb{Z}$ are free. Thus $K_0(\mathbb{Z}) \cong \mathbb{Z}$. More generally, for any ring R with the IBP

2.1.3 Morita Equivalence and Invariance

Suppose we have (unital) rings R and S such that we have an equivalence of the respective categories of right modules:

$$\textbf{mod-}R \leftrightharpoons \textbf{mod-}S$$

Then we say that R and S are **Morita equivalent**.

A consequence of Morita Equivalence is that R and S yield isomorphic K_0 groups.

2.2 K_0 of a Symmetric Monoidal Category

2.3 K_0 of an Exact Category

Chapter 3

Groups K_1 and K_2 of a ring

Chapter 4

Constructions of Higher K-Theory

In this chapter, we shall explore the definitions of higher K-theory. The primary goal of higher K-theory is to (functorially) define a 'K-Theory space' $K(R)$ associated to a unital ring R. We then define the higher K-groups to be $K_n(R) = \pi_n K(R)$, and verify that this description matches with the classical theory for $n = 0, 1, 2$.

A number of approaches can be taken in different contexts, and we have some results showing that in some restricted contexts, these various constructions are equivalent, i.e., yield homotopy-equivalent K-theory spaces.

Finally, this allows us to define and relate the higher K-theory of finitely generated projective modules over a ring, among other applications.

In Section §1, we first recall some homotopy-theoretic notions. In Section §2, we describe the +-construction, introduced in [] by ¡¿ and used by Quillen in [] to describe higher K-groups. Section §3 describes higher K-groups of a ring using the +-construction. In Section §4, we describe the $S^{-1}S$ construction on a symmetric monoidal category, and show its equivalence with the +-construction for $(\mathbf{P}(R), \oplus)$. In Section §4, we describe Quillen's Q-construction for exact categories. Section §5 is devoted to the $+ = Q$ theorem, which implies equivalence of the + and Q-construction for $\mathbf{P}(R)$.

4.1 Homotopy Theory Preliminaries

A based space is a topological space X along with a distinguished basepoint which we denote $*_X$. A map of based spaces $X \to Y$ is a map of the underlying topological spaces which preserves basepoints.

We can functorially associate homotopy groups π_n to a based space X, denoted $\pi_n(X)$. We assume familiarity with the definitions of homotopy and homology groups. $H_*(X; M)$ will refer to the singular homology of X with coefficients in some abelian group M, or alternatively local coefficients, i.e. M being a $\pi_1(X)$-module. When $M = \mathbb{Z}$, this is referred to as integral homology. See [] for a detailed treatment.

A CW complex (or cellular complex) X is a colimit of a filtration of n-skeleta, denoted $\emptyset = X_{-1} \subseteq X_0 \subseteq X_1 \subseteq X_2 \subseteq \cdots$ where X_0 is a discrete set of points and X_n is a pushout of X_{n-1} via a disjoint union $\cup_\alpha \varphi_\alpha$ of attaching maps $\varphi_\alpha : S^{n-1}_\alpha \to X_{n-1}$ and the disjoint union $\cup_\alpha j_\alpha$ of inclusions $j_\alpha : S^{n-1}_\alpha \to D^n_\alpha$. (Note S^{-1} is taken to be $\emptyset$.)

That is, $X_n = X_{n-1} \cup_{S^{n-1}_\alpha} D^n_\alpha$. The images of the interiors of the disks D^n_α in X_n are called n-cells. Note that 0-cells correspond to a set of points, 1-cells correspond to line segments, 2-cells to disks, and so on.

The colimit carries the weak topology. The homology of CW complexes is particularly tractable. See [Hatcher] for a detailed treatment.

4.1.1 Geometric Realisation of a small category

Given a small category C, there is a way to interpret C as a CW complex by the following procedure:

Definition 4.1 (Simplicial sets)**.** A simplicial set X is a contravariant functor $\mathbf{\Delta} \to \mathbf{Sets}$, where $\mathbf{\Delta}$ is the simplex category consisting of objects $\mathbf{n} = \{0, 1, \ldots, n-1\}$ and morphisms $\mathbf{i} \to \mathbf{j}$ given by weakly-order preserving maps. Or alternatively, a sequence of sets $X_0, X_1, X_2, \ldots$ with face maps $\partial^n_i : X_n \to X_{n-1}$ and degeneracy maps $\sigma^n_j : X_n \to X_{n+1}$ subject to certain identities $(0 \leq i, j, \leq n)$.

The category of simplicial sets is denoted **sSet**. It is clear **sSet** is the functor category $[\mathbf{\Delta}^{op}, \mathbf{Sets}]$.

Example 4.1.1. The category Δ[1] of standard n-simplices Δ^n with morphisms being the standard inclusion maps $\Delta^i \to \Delta^j$ for $0 \leq i \leq j$ gives a simplicial set by $\mathbf{n} \to \Delta^n$ and $f : \mathbf{i} \to \mathbf{j}$ associated to $\Delta^i \to \Delta^j$. In fact, this is an isomorphism of categories $\Phi : \mathbf{\Delta} \cong \Delta$.

Example 4.1.2. Any space X determines a simplicial set by the representable functor $\mathbf{Top}(\Phi(-), X)$. This construction is clearly a (covariant) functor $S : \mathbf{Top} \to \mathbf{sSet}$. It is called the *singular functor*.

A simplicial set X may be functorially turned into a topological space, and this functor is called Geometric Realization.

[1] not identical to $\mathbf{\Delta}$

Proposition 4.2 (Geometric Realization)**.** There exists a functor $|-|$: **sSet** $\to$ **Top** which is left-adjoint to the singular functor.

$$\mathbf{Top}(|X|, Y) \cong \mathbf{sSet}(X, SY)$$

naturally in X and Y.

We can define $|-|$ to be the functor taking X to the coend $\int_{\mathbf{n}\in\mathbf{\Delta}} X_n \times \Delta^n$ or equivalently, the colimit $\varinjlim_{\Delta^n \to X} \Delta^n$ in the slice category $\Delta \downarrow X$. The result follows due to Yoneda's Lemma. See [Maclane]. Furthermore, $|X|$ has a natural CW structure induced by CW structures on Δ^n.

Definition 4.3 (Nerve of a small category)**.** Let C be a small category. Then its nerve NC is a simplicial set defined by taking the n-simplices as functors $c : \mathbf{n} \to C$, i.e. diagrams of the form

$$c_0 \to c_1 \to \cdots \to c_{n-1}$$

for objects $c_0, c_1, \ldots, c_{n-1}$ in C. The i-th face map $\partial_i^n(c)$ on this simplex is obtained by deleting c_i in the evident way. Similarly, the j-th degeneracy map $\sigma_j^n(c)$ is obtained by replacing c_j by $c_j \to c_j$ in the obvious manner.

We define the *classifying space* of a small category C, denoted BC, as the geometric realization of NC.

Definition 4.4 (Classifying space of a small category)**.** $BC := |NC|$.

Example 4.1.3. For the categories $\mathbf{n}$, the geometric realization $B\mathbf{n} = \Phi\mathbf{n}$.

Furthermore, for any small category C, BC is the colimit (with the weak topology) of the directed system of the functors $B\mathbf{n} \to BC$ coming from $\mathbf{n} \to C$ and the morphisms coming from factorizations $\mathbf{i} \to \mathbf{n} \to C$.

Proposition 4.5. A functor $F : C \to D$ of small categories determines a cellular map $BF : BC \to BD$. Further, $B(\mathrm{id}_C) : BC \to BC$ is the identity map on BC, and for functors $F : C \to D, G : D \to E$, we have $B(G \circ F) = BG \circ BF$.

Proof. Due to the functoriality of geometric realization on simplicial sets, it suffices to show that the association $N : \mathbf{Cat} \to \mathbf{sSet}$ is functorial. But this follows directly from axioms of a category. □

Thus we have a functor $B : \mathbf{Cat} \to \mathbf{Top}$.

Example 4.1.4. A group G may be interpreted as a small category with a single object and all invertible arrows. We call BG the classifying space of the group G.

Proposition 4.6. The fundamental group $\pi_1(BG)$ is naturally isomorphic to G.

4.1.2 More Homotopy Theory

We will note that the following homotopy-theoretic generalization of a topological group will be relevant when considering upcoming constructions.

Definition 4.7 (H-space)**.** A H-space [2] X is a topological space with a binary operation $\mu : X \times X \to X$ and a distinguished element e such that

1. $\mu(e, e) = e$
2. $\mu(e, -) \simeq \mathrm{id}_X$
3. $\mu(-, e) \simeq \mathrm{id}_X$

Additionally, we say that a H-space is homotopy-associative or homotopy-commutative if those respective properties hold up to homotopy. A detailed description is available in [].

Topological groups are trivially examples of H-spaces. A class of illuminating nontrivial examples come from n-th loop spaces $\Omega^n X$ of some based space X, where the multiplication is given by concatenation of paths.

The (path) connected components of a homotopy-associative H-space determine a monoid.

Homotopy Fiber and Fiber Sequences

We now describe the important notion of a *homotopy fiber*, which is useful for organizing how homotopy groups behave across a map $f : E \to B$.

Definition 4.8 (Homotopy Fiber)**.** Let $f : E \to B$ be a map of based spaces. The homotopy fiber $F(f)$ of this map is the homotopy-pullback of the cospan $E \to B \leftarrow PB$, where PB is the (based) space of paths in Y.

More concretely, we can define $F(f)$ to be consisting of pairs (e, γ) where $e \in E$ and γ is a path in B starting at the basepoint such that $f(e) = \gamma(1)$. $F(f)$ inherits the subspace topology from $E \times PB$ where PB itself carries the compact-open topology. We may declare $F(f)$ to have the basepoint $(*_E, *_{PB})$.

The key property of the homotopy fiber is that the sequence $F(f) \to E \to B$ of based space maps which yields a long exact sequence:

$$\cdots \to \Omega^{n+1} B \xrightarrow{\partial} \Omega^n F(f) \to \Omega^n E \to \Omega^n B \xrightarrow{\partial} \cdots$$
$$\cdots \xrightarrow{\partial} \Omega F(f) \to \Omega E \to \Omega B \xrightarrow{\partial} F(f) \to E \to B$$

[2] The H stands for Hopf

which in turn yields a long exact sequence of groups/pointed sets:

$$\cdots \to \pi_{n+1}B \xrightarrow{\partial} \pi_n F(f) \to \pi_n E \to \pi_n B \xrightarrow{\partial} \cdots$$
$$\cdots \xrightarrow{\partial} \pi_1 F(f) \to \pi_1 E \to \pi_1 B \xrightarrow{\partial} \pi_0 F(f) \to \pi_0 E \to \pi_0 B$$

Any other such sequence $F \to E \to B$ which results in the evident map $F \to F(f)$ (use $*_{PB}$ in the image) being a homotopy equivalence, is called a *homotopy fibration sequence*, and consequently also yields such a long exact sequence.

If $f : E \to B$ were a homotopy equivalence, then the homotopy fiber $F(f)$ is weakly contractible, i.e., it has the homotopy groups of a point, $\pi_* F(f) \cong \pi_*(\star)$. If E and B have homotopy type of CW complexes, then so is $F(f)$ and by Whitehead's Theorem [], weak homotopy equivalence implies homotopy equivalence.

Conversely, if the fiber $F(f)$ is contractible, then f is a weak homotopy equivalence, i.e., the induced maps $\pi_n(E) \to \pi_n(B)$ are isomorphisms. If E and B have homotopy type of CW complexes, then f is a homotopy equivalence.

4.1.3 +-construction on a connected CW complex

The +-construction, devised by ¡¿ in [], is a universal way to kill a perfect normal subgroup of the fundamental group of a connected CW complex, while preserving homology theory. The discussion of the +-construction fits into the framework of acyclic maps, which we explore here.

Definition 4.9. A space X is said to be acyclic if it has the homology of a point, i.e., $H_*(X;\mathbb{Z}) \cong H_*(\star;\mathbb{Z}) = \underline{\mathbb{Z}}$.

In particular, contractible spaces are acyclic. Acyclic spaces serve an analogous purpose to homology as contractible spaces do to homotopy.

Definition 4.10 (Acyclic maps)**.** Let $f : X \to Y$ be a map of based connected CW complexes. We say f is acyclic if the homotopy fiber $F(f)$ is acyclic, i.e., has the homology of a point.

In particular, a homotopy equivalence is acyclic since the former implies that the homotopy fiber is contractible. Note an acyclic map induces isomorphisms on integral homology $f_* : H_*(X;\mathbb{Z}) \cong H_*(Y;\mathbb{Z})$. Furthermore, we have the following result characterising acyclic maps:

Lemma 4.11. Let X, Y be based connected CW complexes. A map $f : X \to Y$ is acyclic if and only if for every $\pi_1(Y)$-module M, it induces an isomorphism $H_*(X;M) \cong H_*(Y;M)$.

The proof for this uses Serre Spectral sequences, which we skip and refer the reader to [Weibel].

Definition 4.12 (relative +-construction)**.** Let X be a based connected CW complex and P be a perfect normal subgroup of $\pi_1(X)$. Then we say a based acyclic map of CW-complexes $f : X \to Y$ is a +-construction relative to P if P is the kernel of the induced map $\pi_1(X) \to \pi_1(Y)$.

Since f was assumed to be acyclic, we can deduce from the exact sequence $\pi_1 F(f) \to \pi_1(X) \to \pi_1(Y) \to \pi_0 F(f)$ that the map $\pi_1(X) \to \pi_1(Y)$ is onto. (because $\pi_0 F(f) = \{\star\}$) And $\pi_1(Y) \cong \pi_1(X)/P$.

Theorem 4.13 (Quillen)**.** *Let X be a based connected CW complex and P be a perfect normal subgroup of $\pi_1(X)$.*

1. *There exists a +-construction $X \to Y$ relative to P.*
2. *A +-construction $f : X \to Y$ is universal among maps which kill P in the fundamental group. That is, for any $g : X \to Z$ such that the image of P in $\pi_1(Z)$ vanishes, we have a map $h : Y \to Z$, unique up to homotopy, such that $g = hf$.*
3. *In particular, if g is another +-construction relative to P, then there is a homotopy equivalence $h : Y \xrightarrow{\sim} Z$ compatible with the maps.*

The proof of this uses topological obstruction theory. We describe a construction for 1 in §**??** and refer the reader to [Berrick] for a full proof of 1 and 2. 3 follows immediately from 2 since we have maps both way which must compose to identity up to homotopy.

Remark 4.1.1. Any group G has a maximal perfect normal subgroup, which we call the *perfect radical*. In fact, it is the subgroup generated by all perfect normal subgroups, hence their maximum. From now on, for any based CW complex X, we will denote by X^+ any +-construction on X relative to the perfect radical $P \lhd \pi_1(X)$.

4.1.4 Explicit Construction

A number of explicit *models* exist for the +-construction. Here, we present a construction due to [Kervaire] which constructs X^+ by attaching 2-cells to kill the maximal perfect subgroup of $\pi_1(X)$, and then 3-cells to recover the integral homology of X, making sure that the inclusion $X \hookrightarrow X^+$ is acyclic. We refer the reader to [Gersten] and [Suslin] for a variety of constructions. We will describe more models for K-theory in Sections §3 and §4.

Construction

Let X be a connected CW complex with a perfect normal subgroup P of the fundamental group $\pi_1(X)$. Identify the 1-cells in X corresponding to of P as e_p^1 for each $p \in P$. Now construct a pushout Y by attaching disks D_p^2 with the boundary S_p^2 along every e_p^1. Call these new cells e_p^2.

That is, X' is the pushout of the diagram

$$\begin{array}{ccc} \coprod_{p\in P} S_p^1 & \hookrightarrow & \coprod_{p\in P} D_p^2 \\ \downarrow & & \downarrow \\ X & \hookrightarrow & Y \end{array}$$

Using the van-Kampen Theorem, we have the pushout through π_1 in **Grp**, which gives us $\pi_1(Y) = \pi_1(X) *_P 1 = \pi_1(X)/P$. (amalgamated product)

Consequently, H_1 is the abelianization. Since P is perfect, it is killed by abelianization and we have $H_1(Y) = H_1(X)$.

Consider the pair (Y, X). Since only 2-cells were attached, the homology of the pair is

$$H_2(Y, X) = \bigoplus_{[e_p^2]} \mathbb{Z}$$

and 0 elsewhere. We wish to cleverly attach 3-cells to kill relative homology without affecting H_3.

Consider the long exact sequence of the pair (Y, X)

$$\cdots \to H_{n+1}(Y, X) \xrightarrow{\partial} H_n(X) \to H_n(Y) \to H_n(Y, X) \xrightarrow{\partial} \cdots$$

At $n = 1$, since the map $H_1(X) \to H_1(Y)$ is an isomorphism, we have that $H_2(Y, X) \to H_1(X)$ is 0. Consequently, the map $H_2(Y) \to H_2(Y, X)$ is onto. We can lift each representative in $H_2(Y, X)$ to a representative in $H_2(Y)$, say σ_p for each class $[e_p^2]$.

Note each σ_p is represented by a map $h_p : S_p^2 \to Y$. Now attach 3-cells e_p^3 corresponding to each σ_p along the maps $h_p : S_p^2 \to Y$.

This yields another pushout which we call Z

$$Z = Y \cup \bigcup_{[e_p^2]} e_p^3$$

The triple $X \subseteq Y \subseteq Z$ induces a long exact sequence in relative homology

$$\cdots \to H_{n+1}(Z, Y) \xrightarrow{\partial} H_n(Y, X) \to H_n(Z, X) \to H_n(Z, Y) \xrightarrow{\partial} \cdots$$

At $n = 1$, since $H_2(Y, X) = 0$ and $H_0(Z, Y) = 0$, this gives us

$$\cdots \xrightarrow{\partial} 0 \to H_1(Z, X) \xrightarrow{\cong} H_1(Z, Y) \xrightarrow{\partial} 0 \to \cdots$$

Similarly, for $n \geq 4$, since $H_{n+1}(Z, Y) = 0$ and $H_{n-1}(Y, X) = 0$, we get

$$\cdots \xrightarrow{\partial} 0 \to H_n(Z, X) \xrightarrow{\cong} H_n(Z, Y) \xrightarrow{\partial} 0 \to \cdots$$

So we conclude $H_n(Z, X) \cong H_n(Z, Y) = 0$ for $n \neq 1, 2, 3$.

The only nontrivial terms in the sequence that remain are

$$\cdots \xrightarrow{\partial} 0 \to H_3(Z, X) \to H_3(Z, Y) \xrightarrow{\partial} H_2(Y, X) \to 0 \to \cdots$$

But $H_3(Z, Y)$ and $H_2(Y, X)$ are computable precisely due to construction. Z was constructed so as to kill

$$H_3(Z, Y) = \bigoplus_{[\sigma_p]} \mathbb{Z}$$

which is isomorphic via the evident maps to $H_2(Y, X) = \bigoplus_{[e_p]} \mathbb{Z}$.

So finally, we have $H_3(Z, X) = 0$, and altogether $H_n(Z, X) = 0$ for all n. Consequently, in the long exact sequence of the pair (Z, X), we get

$$\cdots \to H_{n+1}(Z, X) \xrightarrow{\partial} H_n(X) \to H_n(Z) \to H_n(Z, X) \xrightarrow{\partial} \cdots$$

which gives us that $H_n(X) \cong H_n(Z)$ induced via the inclusions are isomorphisms for all n.

It is clear that P is the kernel of the induced map $\pi_1(X) \to \pi_1(Y)$. This concludes the result that the stated construction preserves integral homology.

If this result were to extend to local coefficient systems, this would mean that the map $X \to Z$ is acyclic and forms a model for X^+. We do not pursue the full proof here and instead refer the reader to [Rosenberg]. Indeed, there is an alternate result in Proposition **??** which shows that what we have proven suffices for the case we are interested in, $BGL(R)^+$, which is what we shall discuss now.

4.2 The $BGL(R)^+$ construction for a ring R

?? We apply the +-construction on the classifying space $BGL(R)$ of the stable general linear group over R. The maximal perfect subgroup of $GL(R) = \pi_1 BGL(R)$ is the stable group of elementary matrices, $E(R)$.

Theorem 4.14. *With this definition of $BGL(R)^+$, we obtain:*

1. *$\pi_1 BGL(R)^+ \cong K_1(R)$ with the kernel of the induced map $GL(R) = \pi_1 BGL(R) \to \pi_1 BGL(R)^+$ being $E(R)$.*
2. *We have induced isomoprhisms $H_*(BGL(R); M) \cong H_*(BGL(R)^+; M)$ for every $K_1(R)$-module M.*

Proof. Since $E(R)$ is the maximal perfect normal subgroup of $GL(R)$, 1 follows immediately from Theorem **??**. 2 follows from the characterization of acyclic maps in Lemma **??**. □

Due to Theorem **??**, $BGL^+(R)$ is uniquely defined up to homotopy equivalence via a canonical equivalence. Consequently, we may define $K_n(R) = \pi_n BGL(R)^+$ for $n \geq 1$. It is obvious that $K_1(R)$ defined here matches the description in Chapter 3. It remains to verify that this agrees with the classical theory for $n = 2$.

Proposition 4.15. Let P be a perfect normal subgroup of G with a $+$-construction $f : BG \to BG^+$ relative to P. Let $F(f)$ be the homotopy fiber of f. Then $\pi_1 F(f)$ is the universal central extension of P, and consequently $\pi_2(BG^+) \cong H_2(P; \mathbb{Z})$.

Proof. We have the fibration long exact sequence

$$\cdots \to \pi_2(BG) \to \pi_2(BG^+) \to \pi_1 F(f) \to G \to G/P \to 1$$

Since BG is an Eilenberg-Maclane space $K(G, 1)$, we have $\pi_2(BG) = 0$. So $\pi_2(BG^+)$ injects into $\pi_1 F(f)$, which in turns surjects onto the kernel of $G \to G/P$ which is P. Since $\pi_2(BG^+)$ is abelian, it must inject into the center of $\pi_1 F(f)$. This tells us that $\pi_1 F(f) \to P$ is a central extension. Since f is acylic, we have $\pi_1 F(f)$ is perfect and so $H_2(\pi_1 F(f), \mathbb{Z}) = 0$. By the Recognition Theorem [Chapter 3], we have that $\pi_1 F(f)$ is the universal central extension of P. Consequently, the kernel $\pi_2(BG^+)$ is given by $H_2(P; \mathbb{Z})$.

□

Corollary 4.16. We have $\pi_2(BGL(R))^+ \cong H_2(E(R); \mathbb{Z})$, and thus our definition matches with the classical theory since $H_2(E(R); \mathbb{Z}) \cong K_2(R)$ from Chapter 3.

However, at this point, it is not clear that the associations $R \mapsto BGL(R)^+$ or $R \mapsto K_n(R)$ are functorial. This invites some discussion.

4.2.1 Functoriality of the $BGL(R)^+$-construction

Apart from the upcoming models via $S^{-1}S$ and Q-constructions, we can make the following argument to show that the constructions of $BGL(R)^+$ can be performed functorially.

We can first define $BGL(\mathbb{Z})^+$ and take a homotopy pushout (using the mapping cylinder) via the canonical map $\mathbb{Z} \to R$ which induces $BGL(\mathbb{Z}) \to BGL(R)$. Note that the homotopy pushout

$$\begin{array}{ccc} BGL(\mathbb{Z}) & \longrightarrow & BGL(\mathbb{Z})^+ \\ \downarrow & & \downarrow \\ BGL(R) & \longrightarrow & BGL(R)^+ \end{array}$$

is a plus construction for $BGL(R)$.

4.2.2 H-Space Properties of $BGL(R)^+$

It's not hard to see that we can put a H-space structure on $BGL(R)^+$.

Proposition 4.17. $BGL(R)^+$ is a homotopy-associative H-space.

Proof. We have the group map $\oplus : GL_m(R) \times GL_n(R) \to GL_{m+n}(R) \subset GL(R)$ which induces $BGL_m(R) \times BGL_n(R) \to BGL(R)$. Since we have show that the plus construction can be taken to be functorial, we have maps

$$(BGL_m(R) \times BGL_n(R))^+ \to BGL(R)^+$$

. [Finish this] □

There is a recognition criteria for +-constructions of $BGL(R)$ in terms of homology and H-spaces, due to Quillen. The proof again uses obstruction theory. See [Gersten].

Theorem 4.18. *The plus construction $i : BGL(R) \to BGL(R)^+$ is universal for maps into H-spaces. That is, for any map $f : BGL(R) \to H$ for some H-space H, we have a map $g : BGL(R)^+ \to H$ so that $f = gi$, such that g is unique up to weak homotopy, i.e. the induced maps $\pi_n(BGL(R)^+) \to \pi_n(H)$ is independent of g.*

Proposition 4.19. If further $f_* : H_*(BGL(R); \mathbb{Z}) \xrightarrow{\cong} H_*(H; \mathbb{Z})$ is an isomorphism, then f is acyclic and g is a homotopy equivance.

Proof. It suffices to prove that f is acyclic. □

4.3 $S^{-1}S$ construction

For a symmetric monoidal category $(S, \square)$, the construction of Chapter 2 for K_0 may be extended to a K-theory space $K^{\square}(S)$ by constructing an appropriate group-completion of S with respect to the monoidal operation $\square$.

Note that the geometric realization of a symmetric monoidal category is a homotopy-associative homotopy-commutative H-space with the multiplication coming from the monoidal product $\square$. This is because [] We wish to construct a symmetric monoidal category $S^{-1}S$ which describes a "group completion" of BS so as to make $BS^{-1}S$ be a group-like H-space. [Grayson]

But first, for either of BS or $BS^{-1}S$ to be interesting, we must reduce our scope.

Definition 4.20 (symmetric monoidal groupoid)**.** A *symmetric monoidal groupoid* is a symmetric monoidal category $(S, \square)$ in which all morphisms are isomorphisms.

Further, we say S is *faithful* if all translations $s\square : \mathrm{Aut}(t) \to \mathrm{Aut}(s\square t)$ are injective.

Definition 4.21. Given a symmetric monoidal category S, $\mathrm{iso}(S)$ is the symmetric monoidal groupoid which is a subcategory of S consisting of the same objects but inheriting only isomorphisms.

A variety of symmetric monoidal groupoids yield interesting geometric realizations.

Example 4.3.1. The category $\mathbf{F}(R)$ with $\square = \oplus$ is contractible since it has an initial object 0. However $\mathrm{iso}(\mathbf{F}(R))$ is more interesting and in fact equivalent to the disjoint union $\coprod_n GL_n(R)$ where n runs through the integers.

Example 4.3.2. Similarly, 0 is also an initial object for the category $\mathbf{P}(R)$ with $\square = \oplus$. But $\mathrm{iso}(\mathbf{P}(R))$ is equivalent to $\coprod_P \mathrm{Aut}(P)$ where P runs over representatives of isomorphism clases of finitely generated projective modules.

Let S be a faithful symmetric monoidal groupoid.

Definition 4.22 ($S^{-1}S$)**.** Define $S^{-1}S$ to be a category where the objects are pairs of objects (m, n) for $m, n \in S$. Morphisms in $S^{-1}S$ from $(m_1, m_2) \to (n_1, n_2)$ are equivalence classes of triples (s, f_1, f_2) for some object $s \in S$ and $(f_1, f_2) : (s\square m_1, s\square m_2) \to (n_1, n_2)$ being a morphism in $S \times S$.

That is,

$$(m_1, m_2) \xrightarrow{s\square} (s\square m_1, s\square m_2) \xrightarrow{(f_1,f_2)} (n_1, n_2)$$

which we consider equivalent to

$$(m_1, m_2) \xrightarrow{t\square} (t\square m_1, t\square m_2) \xrightarrow{(g_1,g_2)} (n_1, n_2)$$

if there is an isomorphism $\alpha : s \xrightarrow{\cong} t$ such that $\alpha\square g_1 = f_1$ and $\alpha\square g_2 = f_2$. The composition $[(s, f_1, f_2)] \circ [(t, g_1, g_2)]$ is defined as $[(s\square t, f_1 \circ g_1, f_2 \circ g_2)]$. One may verify this is well-defined and indeed forms a category.

Remark 4.3.1. Consider two distinguished classes of morphisms in $S^{-1}S$. One is simultaneous translations, which we formally write $s\square : (m_1, m_2) \to (s\square m_1, s\square m_2)$, represented by $(s, \mathrm{id}, \mathrm{id})$. The other being pairs of maps $(f_1, f_2) : (m_1, m_2) \to (n_1, n_2)$ coming from morphisms in $S \times S$ where the translation composite is trivial. That is, the morphism is represented by (e, f_1, f_2).

Proposition 4.23. Let S be a faithful symmetric monoidal groupoid. Then any morphism $[(s, f_1, f_2)]$ in $S^{-1}S$ determines s, f_1, f_2 uniquely up to unique isomorphisms.

Proof. It is clear s, f_1, f_2 are determined uniquely up to isomorphism. Suppose (s, f_1, f_2) is equivalent to (t, g_1, g_2) via isomorphisms $\alpha, \beta : s \xrightarrow{\cong} t$ Since $\alpha\square g_i = f_i = \beta\square g_i$, we have $\alpha\square m_i = \beta\square m_i$ and since S was faithful, we recover $\alpha = \beta$. □

Thus, we can think of morphisms $[(s, f_1, f_2)]$ in $S^{-1}S$ as comprising of a *translation component* $[s\square]$ and *mapping component* $[(f_1, f_2)]$.

Remark 4.3.2. $S^{-1}S$ is a symmetric monoidal category with the multiplication given by $(m_1, m_2)\square(n_1, n_2) := (m_1\square n_1, m_2\square n_2)$. It is clear that the functor $S \to S^{-1}S$, which sends $s \mapsto (s, e)$ is a monoidal functor. Further, a monoidal functor $S \to T$ describes a (monoidal) natural transformation

$$\begin{array}{ccc} S & \longrightarrow & T \\ \downarrow & & \downarrow \\ S^{-1}S & \longrightarrow & T^{-1}T \end{array}$$

We also have an induced H-space map $BS \to B(S^{-1}S)$, and a map $\pi_0(S) \to \pi_0(S^{-1}S)$ of abelian monoids. (The component of s maps to the component of (s, e))

Further, $\pi_0(S^{-1}S)$ is an abelian group since for the connected coponent of (m,n) we have we have inverse (n,m) due to the isomorphism $(e,e) \xrightarrow{\cong} (m,n)\square(n,m)$ given by the mapping component $[(c_{mn}, c_{mn}^{-1})]$ where c_{mn} is the symmetric braiding given by isomorphism $m\square n \to n\square m$ in S. It is abelian due to S being symmetric.

Definition 4.24. Let S be a symmetric monoidal groupoid. We define

$$K^{\square}(S) := B(S^{-1}S)$$

and consequently we can define $K_n^{\square}(S) = \pi_n B(S^{-1}S)$.

Note that a (strict) monoidal functor $S \to T$ induces $S^{-1}S \to T^{-1}T$ which in turn induces $K^{\square}(S) \to K^{\square}(T)$, Thus the $S^{-1}S$ construction is functorial!

Proposition 4.25. The group $K_0^{\square}(S) = \pi_0(S^{-1}S)$ is indeed the group completion of the monoid given by isomorphism classes of S via the inclusion $i_* : \pi_0(S) \to \pi_0(S^{-1}S)$.

Proof. Let $\varphi : \pi_0(S) \to G$ be a group completion of $\pi_0(S)$. We will show that the universal homomorphism $\widetilde{i}_* : G \to \pi_0(S^{-1}S)$ is invertible. Let j be the set map $S^{-1}S \to G$ as $j(m,n) := \varphi(m) - \varphi(n)$. If we have $s \in S$ then $j(m_1, m_2) = j(s\square m_1, s\square m_2)$ and if $f_i : m_i \to n_i$, then $j(m_1, m_2) = \varphi(m_1) - \varphi(m_2) = \varphi(n_1) - \varphi(n_2) = j(n_1, n_2)$. This shows j induces a well-defined homomorphism $j_* : \pi_0(S^{-1}S) \to G$. By construction, it is the inverse to $G \to \pi_0(S^{-1}S)$. □

This shows our definition matches the definition of $K_0(S)$ in Chapter 2.

4.3.1 Group Completions of BS

We turn our attention to

Theorem 4.26. *If S*

Theorem 4.27 (Quillen)**.** *Let $S = \coprod GL_n(R)$ which is equivalent to $\mathbf{F}(R)$. Then $B(S^{-1}S)$ is the group-completion of BS and*

$$B(S^{-1}S) \simeq \mathbb{Z} \times BGL(R)^+$$

Proof. □

Theorem 4.28. *Let S be a symmetric monoidal groupoid with faithful transitions satisfying cofinality.*

We have then

$$K^{\square}(S) \simeq K_0^{\square}(S) \times B\,\mathrm{Aut}(S)^+ A$$

4.4 Quillen's Q-construction

In [Quillen, 19XX], Quillen proposed another construction for the K-theory space. Given an exact category $\mathcal{A}$, [] we will define a category $Q\mathcal{A}$, and then define $K_n(\mathcal{A}) = \pi_{n+1}(BQ\mathcal{A})$ for all $n \geq 0$.

Definition 4.29 (Q-construction)**.** Let $\mathcal{A}$ be an exact category. We define $Q\mathcal{A}$ as the category with the same objects as $\mathcal{A}$, and the morphisms between $C \to C'$ are equivalence classes of diagrams

$$C \xleftarrow[j]{} D \xrightarrow{i} C'$$

where two such diagrams are considered isomorphic when there is an isomorphism of these diagrams which is identity on C and C'.

The composition of the above arrow with $C' \twoheadleftarrow D' \rightarrowtail C''$ is given by $C \twoheadleftarrow D \times_{C'} D' \rightarrowtail C''$

$$\begin{array}{ccccc} D \times_{C'} D' & \xrightarrow{\pi_2} & D' & \xrightarrow{i'} & C'' \\ {\scriptstyle \pi_1}\downarrow & & \downarrow{\scriptstyle j'} & & \\ D & \xrightarrow[i]{} & C' & & \\ {\scriptstyle j}\downarrow & & & & \\ C & & & & \end{array}$$

It is clear that composition is well-defined and associative and so $Q\mathcal{A}$ forms a well-defined category.

[Set theoretic remark]

Remark 4.4.1. Admissible monomorphisms $C \rightarrowtail C'$ determine a morphism $C \to C'$ by taking $D = C'$ and $D \twoheadleftarrow C'$ as identity. We will call these morphisms *injective*.

Similarly, admissible epimorphisms $C \twoheadleftarrow C'$ determine a morphism $C \to C'$ by taking $D = C$ and $C \rightarrowtail D$ as identity. We will call these morphisms *surjective*.

Note that these distinguished families of morphisms are closed under composition, and generate all morphisms. Furthermore, any morphism $C \twoheadleftarrow D \rightarrowtail C'$ factors as $i_* j^*$ where i_* is the injective morphism determined by $D \rightarrowtail C'$ and j^* is the surjective morphism determined by $C \twoheadleftarrow D$. This factorization is unique up to unique isomorphism.

4.4.1 Alternative Description of Morphisms in $Q\mathcal{A}$

There is an alternative way to describe the morphisms $C \to D$ in $Q\mathcal{A}$ in terms of *subobjects* and *subquotients*. Fix some object C' in $\mathcal{A}$.


Definition 4.30. An *admissible subobject* of C' is an equivalence class of admissible monomorphisms $C \rightarrowtail C'$ with equivalence given by whether they factor through each other, or in other words as isomorphism in $\mathcal{A} \downarrow C$.

An *admissible subquotient* is defined in the dual manner, using admissible epimorphisms.

One can define a partial order on the set of admissible monomorphisms into C'. We declare $C_1 \leq C_2$ if there is an admissible monomorphism $C_1 \rightarrowtail C_2$, which if it exists has to be unique. We then say (C_1, C_2) is an *admissible layer* in C' and the cokernel C_2/C_1 is an *admissible subquotient* of C'.

Proposition 4.31. The morphisms from $C \to C'$ may be identified with pairs $((C_1, C_2), \varphi)$, where (C_1, C_2) is an admissible layer in C' and $\varphi : C \xrightarrow{\cong} C_2/C_1$ is an isomorphism in $\mathcal{A}$.

Proof. For $C \twoheadleftarrow D \rightarrowtail C'$, take the admissible layer (C_1, C_2) where $C_2 = D$ and C_1 is the kernel of $C \twoheadleftarrow D$. Clearly, since C_1 gives a subobject of $D = C_2$ which gives a subobject of C', the given pair is indeed an admissible layer. A unique isomorphism follows since both C and the C_2/C_1 are cokernels of the kernel of an admissible epimorphism.



Conversely, an admissible layer (C_1, C_2) in C' evidently determines a morphism $C_2/C_1 \twoheadleftarrow C_2 \rightarrowtail C$ in $Q\mathcal{A}$ where the isomorphism may be taken as identity.

Furthermore, composition of morphisms $C \to C'$ and $C' \to C''$ may be described this way and arises from the fact that subquotient of a subquotient is a subquotient.

Let morphisms $C \to C'$ and $C' \to C''$ correspond to $((C_1, C_2), \varphi)$ and $((C_1', C_2'), \psi)$ where (C_1, C_2) and (C_1', C_2') are admissible layers in C and C'' respectively with isomorphisms $\varphi : C \xrightarrow{\cong} C_2/C_1$ and $\psi : C' \xrightarrow{\cong} C_2'/C_1'$. Let the map $C_2' \to C_2'/C_1'$ be denoted q.

Then the composite $C \to C' \to C''$ corresponds to the layer (D_1, D_2) where $D_1 = C_1 \times_{C'} C_2'$ and $D_2 = C_2 \times_{C'} C_2'$ with the isomorphism $\Phi : C \xrightarrow{\cong} D_2/D_1$ given by

$$C \cong C_2/C_1$$

This follows from the evident isomorphisms and the fact that D_2 is the pullback of $C_2 \rightarrowtail C' \twoheadleftarrow C_2'$. □

This characterization is particularly helpful to eludicate certain properties of the category $Q\mathcal{A}$.

Example 4.4.1. Using this characterization, we can describe the morphisms with source 0 in $Q\mathcal{A}$. Morphisms $0 \to C$ are in 1-1 correspondence with the subobjects of C, thus injective. There is a canonoical morphism $C \to 0$ since 0 has a unique subquotient.

Example 4.4.2. Isomorphisms $C \xrightarrow{\cong} D$ in $Q\mathcal{A}$ are in 1-1 correspondence with isomorphisms $C \xrightarrow{\cong} D$ in $\mathcal{A}$.

4.4.2 K-Theory using the Q-construction

We now prove a few homotopy-theoretic properties of the Q-construction, before we eventually define K_n-groups corresponding to a given (skeletally small) exact category.

Proposition 4.32. An exact functor $F : \mathcal{A} \to \mathcal{B}$ induces a functor $QF : Q\mathcal{A} \to Q\mathcal{B}$.

Proof. On the level of objects, it suffices to define QF precisely as F. Suppose $g : C \to D$ is a morphism in $Q\mathcal{A}$. We may identify g with an admissible layer (D_1, D_2) of D. Define $QF(g)$ to be the morphism $FC \to FD$ in $Q\mathcal{B}$ corresponding to the admissible layer (FD_1, FD_2) of FD, which is indeed an admissible layer since the exactness relations in $\mathcal{A}$ carry over to $\mathcal{B}$ by the exact functor F. Functoriality follows since compositions are defined naturally across $Q\mathcal{A}$ and $Q\mathcal{B}$. □

We present a different proof of the theorem proven by Quillen in [].


Theorem 4.33 (Quillen)**.** *Let $\mathcal{A}$ be a small exact category. The fundmental group of $BQ\mathcal{A}$ at the basepoint 0 is canonically isomorphic to the $K_0(\mathcal{A})$ group defined in Chapter 2.*

Proof. Note that $\pi_1(BQ\mathcal{A})$ is generated by loops given by the morphisms $0 \to C \leftarrow 0$ for each C in $Q\mathcal{A}$, where $0 \to C$ is the injective morphism given by the subobject $0 \rightarrowtail C$, and $C \leftarrow 0$ is the surjective morphism given by $0 \twoheadleftarrow C$. This also shows $BQ\mathcal{A}$ is (path) connected.

It is clear that the loops given by $0 \to C \leftarrow 0$ determine C up to isomorphism, and loops $0 \to C \leftarrow 0$ generate $\pi_1(BQ\mathcal{A})$.

Given any short exact sequence $C' \rightarrowtail C \twoheadrightarrow C''$ in $\mathcal{A}$, we wish to show that the loop given by the concatenated diagram $0 \to C' \leftarrow 0 \to C'' \leftarrow 0$ in $Q\mathcal{A}$

is homotopic to the loop given by $0 \to C \leftarrow 0$. Drawing out the morphisms of these diagrams in $Q\mathcal{A}$, we have the following diagram in $\mathcal{A}$



Note that there are four triangles with vertices and morphisms in $Q\mathcal{A}$ which divide this diagram as displayed here. To show the desired claim, it suffices to show that each of the four traingles commute in $Q\mathcal{A}$, which would imply each corresponds to a 2-simplex in $\pi_1(BQ\mathcal{A})$, making the loop $0 \to C' \leftarrow 0 \to C'' \leftarrow 0$ homotopic to $0 \to C \leftarrow 0$ due to shared edges.

But in each of the triangles the computation of the pullback in the composition is trivial and yields either the injective morphism $0 \to C$ (by $0 \twoheadleftarrow 0 \rightarrowtail C$) or the surjective morphism $0 \to C$ (by $0 \twoheadleftarrow C \rightarrowtail C$). In any case, it is equal to the composite edge.

It also follows that $\pi_1(BQ\mathcal{A})$ is abelian.

We can now define a group homomorphism $\Theta : K_0(\mathcal{A}) \to \pi_1(BQ\mathcal{A})$ by mapping $[C]$ to the loop $0 \to C \leftarrow 0$ as mentioned. This is a (surjective) group homomorphism since exact sequences lead to homotopic loops.

To show Θ is injective, We construct an Φ from morphisms in $Q\mathcal{A}$ to $K_0(\mathcal{A})$ by mapping the morphism determined by the layer (D_1, D_2) of C' to D_1. This is additive since □

[Set Theory remark]

Definition 4.34. Let $\mathcal{A}$ be a small exact category. Then $K\mathcal{A}$ is defined to be the space $\Omega BQ\mathcal{A}$ pointed at 0 and further

$$K_n(\mathcal{A}) := \pi_n(K\mathcal{A}) = \pi_{n+1}(BQ\mathcal{A})$$

This definition agrees with $K_0(\mathcal{A})$ defined in Chapter 2. Further, any exact functor $F : \mathcal{A} \to \mathcal{B}$ induces a functor $Q\mathcal{A} \to Q\mathcal{B}$, hence $BQ\mathcal{A} \to BQ\mathcal{B}$ showing that $K_n(\mathcal{A}) \to K_n(\mathcal{B})$ showing that the Q-construction leads to a functorial K-theory.

Definition 4.35. If R is a unital ring, we define $K(R) := K(\mathbf{P}(R))$ and consequently

$$K_n(R) = K_n(\mathbf{P}(R)).$$

This matches our original definition for $n = 0$. We prove in §?? the $+ = Q$ Theorem, which implies that this matches with the description of the K_n groups defined using the $+$-construction in §??.

Definition 4.36. If R is a Noetherian ring with unity, we define $G(R) := K(\mathbf{M}(R))$ and consequently

$$G_n(R) := K_n(\mathbf{M}(R)).$$

This gives us G-Theory of unital Noetherian rings and matches the classical definition for $n = 0$.

4.4.3 Key Properties of the Q-construction

Morita Invariance

Since K-theory from the Q-construction is functorial, for Morita-equivalent rings R, S, i.e., there is an equivalence $\mathbf{P}(R) \simeq \mathbf{P}(S)$ of categories, from which it follows that $K_n(R) \cong K_n(S)$.

Cofinality

4.5 The "$+ = Q$" Theorem

Given an additive category $\mathcal{A}$, there are two approaches to defining K-theory over $\mathcal{A}$. One is through

Theorem 4.37 (Quillen)**.** *Let $\mathcal{A}$ be a split-exact category and $S = \operatorname{iso}\mathcal{A}$ be the symmetric monoidal groupoid of isomorphisms in $\mathcal{A}$ with $\square = \oplus$, then*

$$\Omega BQ\mathcal{A} \simeq B(S^{-1}S)$$

Corollary 4.38 ($+ = Q$)**.** For any unital ring R,

$$\Omega BQ\mathbf{P}(R) \simeq K_0(R) \times BGL(R)^+$$

This means that the K-Theory of $\mathbf{P}(R)$ from the Q-construction is equivalent to the K-Theory defined using the $+$-construction.

4.5.1 Proof of the $+ = Q$ Theorem

For the proof, we wish to produce a fibration sequence

$$S^{-1}S \to ?? \to Q\mathcal{A}$$

where **??** is chosen appropriately and $B(??)$ is contractible. Then upon taking the homotopy long exact sequence of the fibration, we would have our desired homotopy equivalence $\Omega BQ\mathcal{A} \simeq BS^{-1}S$. This sequence of leading up to the theorem is taken from [Weibel].

Definition 4.39. Given an exact category $\mathcal{A}$, we define $\mathcal{EA}$ where the objects of $\mathcal{EA}$ are admissible exact sequences in $\mathcal{A}$, and a morphism between sequences $C' \rightarrowtail C \twoheadrightarrow C''$ and $D' \rightarrowtail D \twoheadrightarrow D''$ is an equivalence class of the following diagrams admssible exact sequences $D' \rightarrowtail C \twoheadrightarrow E$ such that the following diagram commutes

$$\begin{array}{ccccc} C' & \rightarrowtail & C & \twoheadrightarrow & C'' \\ {\scriptstyle\alpha}\uparrow & & \| & & \uparrow \\ D' & \rightarrowtail & C & \twoheadrightarrow & E \\ \| & & {\scriptstyle\beta}\downarrow & & \downarrow \\ D' & \rightarrowtail & D & \twoheadrightarrow & D'' \end{array}$$

Two such diagrams are considered equivalent when there is an isomorphism between them which is identity at all vertices except for E.

Note since the right edge in the diagram describes a morphism in $Q\mathcal{A}$, we have a functor $f : \mathcal{EA} \to Q\mathcal{A}$ given by picking C'' from the exact sequence $C' \rightarrowtail C \twoheadrightarrow C''$.

We now consider the fiber category $f^{-1}(E)$ for an object E which we denote $\mathcal{E}_E$.

Remark 4.5.1. If we consider the identity morphism to be fixed at E , then the fiber category $\mathcal{E}_E$ consists of exact sequences with target E, and the morphisms from $C' \rightarrowtail C \twoheadrightarrow E$ to $D' \rightarrowtail D \twoheadrightarrow E$ are diagrams of the form

$$\begin{array}{ccccc} C' & \rightarrowtail & C & \twoheadrightarrow & E \\ {\scriptstyle\alpha}\uparrow{\scriptstyle\cong} & & {\scriptstyle\beta}\downarrow{\scriptstyle\cong} & & \| \\ D' & \rightarrowtail & D & \twoheadrightarrow & E \end{array}$$

It follows automatically that each morphism in $\mathcal{E}_E$ is an isomorphism which we may describe as a pair (α, β) where $\alpha : D' \cong C'$ and $\beta : C \cong D$.

The following result allows us to relate the fiber category with the symmetric monoidal groupoid $S = \operatorname{iso}\mathcal{A}$.

Proposition 4.40. The fiber category $\mathcal{E}_0$ is homotopy equivalent to $S = \operatorname{iso}\mathcal{A}$.

Proof. Let $F : S \to \mathcal{E}_0$ be the functor given by mapping an object C to $C \rightarrowtail C \twoheadrightarrow 0$, and an isomorphism $f : C \to C$ to the morphism (f, f^{-1}) with the diagram

$$\begin{array}{ccccc} C & \rightarrowtail & C & \twoheadrightarrow & E \\ {\scriptstyle f}\uparrow & & {\scriptstyle f^{-1}}\downarrow & & \| \\ C & \rightarrowtail & C & \twoheadrightarrow & E \end{array}$$

In fact, since every object in $\mathcal{E}_0$ is naturally isomorphic to such a sequence, the functor F is defined naturally. It is also a full embedding. This gives us the claim: □

Proposition 4.41. For any object E in $\mathcal{A}$, $\mathcal{E}_E$ is a symmetric monoidal category, and there is a (faithful) monoidal functor $\eta_E : S \to \mathcal{E}_E$ sending C to the sequence $C \rightarrowtail C \oplus E \twoheadrightarrow E$.

Proof. We define the monoidal structure on $\mathcal{E}_E$ as follows. Let $e = (0 \rightarrowtail E \twoheadrightarrow E)$ be the identity. Let x denote $C' \rightarrowtail C \twoheadrightarrow E$ and y denote $D' \rightarrowtail D \twoheadrightarrow E$. Then we define $x \square y$ as

$$C' \oplus D' \rightarrowtail C \times_E D \twoheadrightarrow E$$

It follows that this obeys the necessary unit, commutatitivity and coherence laws and does define a symmetric monoidal structure on $\mathcal{E}_E$. The symmetric monoidal functor $S \to \mathcal{E}_E$ follows naturally by mapping C to the split-exact sequence $C \rightarrowtail C \oplus E \twoheadrightarrow E$. [] It is also faithful. □

Now, here is a wonderful observation which reduces the complexity of the fiber category $\mathcal{E}_E$ in a restricted case.

Remark 4.5.2. If $\mathcal{A}$ were split-exact, then every sequence in $\mathcal{E}_E$ is of the form $C \rightarrowtail C \oplus E \twoheadrightarrow E$, hence comes from S via the functor η_E. Consequently, the category $\langle S, \mathcal{E}_E \rangle$ is connected.

Proposition 4.42. If $\mathcal{A}$ is split exact, each $S^{-1}S \to S^{-1}\mathcal{E}_E$ is a homotopy equivalence.

Proof. Note the sequence $S^{-1}S \to S^{-1}\mathcal{E}_E \to \langle S, \mathcal{E}_E \rangle$ is a fibration. It suffices to show that $L = \langle S, \mathcal{E}_E \rangle$ is contractible. Note that the monoidal operation on $\mathcal{E}_E$ induces a monoidal operation on L, making BL a H-space. Since L is connected, by [Whitehead] it implies BL is group-like and has a homotopy inverse.

Now for any exact sequence $x = (C \rightarrowtail C \oplus E \twoheadrightarrow E)$ we have a diagonal morphism $\Delta_x : x \to x \square x$ in L,

$$\begin{array}{ccccc} C & \rightarrowtail & D & \twoheadrightarrow & C \\ \downarrow & & \downarrow & & \| \\ C \oplus C & \rightarrowtail & D \times_C D & \twoheadrightarrow & C \end{array}$$

The diagonal Δ_x is natural in x. This natural transformation induces a homotopy between $\mathrm{id} : BL \to BL$ and $(-)^2 : BL \to BL$. Using the homotopy inverse to subtract away id, we get $\mathrm{id} : BL \to BL$ is null-homotopic.

Hence BL is contractible. □

Lemma 4.43. For any morphism $f : E' \to E$ in $Q\mathcal{A}$, there is a canonical "base-change" functor $f^* : \mathcal{E}_E \to \mathcal{E}_{E'}$ and a natural transformation $\eta : f^* \Rightarrow i$, where i is the inclusion $\mathcal{E}_E \hookrightarrow \mathcal{E}\mathcal{A}$.

Theorem 4.44. *If $\mathcal{A}$ is a split exact category and $S = \mathrm{iso}\,\mathcal{A}$, then the sequence*

$$S^{-1}S \to S^{-1}\mathcal{E}\mathcal{A} \to Q\mathcal{A}$$

is a homotopy fibration.

Proof. We use Quillen's Theorem B to prove this. For Theorem B to apply, we need to show that the base changes f^* are homotopy equivalences.

We show the same on properly injective and properly surjective morphisms $0 \to C$. For properly injective morphisms, the base change is given by $(C \rightarrowtail D \twoheadrightarrow E) \mapsto (C \rightarrowtail C \twoheadrightarrow 0)$ which can be identified with C in S. The composition $S^{-1}S \to S^{-1}\mathcal{E}_E \to Q\mathcal{A}$ is thus identity.

For properly surjective morphisms, the base change is $(C \rightarrowtail D \twoheadrightarrow E) \mapsto (D \rightarrowtail D \twoheadrightarrow 0)$ which can be identified with D in S. The composition $S^{-1}S \to S^{-1}\mathcal{E}_E \to Q\mathcal{A}$ in this case is $C \mapsto (D = C \oplus E)$, which is realized by the natural transformation given by $E\oplus$, hence is also a homotopy equivalence.

Hence we can conclude that in general, f^* is a homotopy equivalence. □

***Proof of Theorem* ??.** The proof follows from Theorem **??** if we show that $S^{-1}\mathcal{E}\mathcal{A}$ is contractible. Note that $\mathcal{E}\mathcal{A}$ is contractible. [] Any action of S on a contractible category must be invertible. [] So we have the homotopy equivalence $\mathcal{E}\mathcal{A} \to S^{-1}\mathcal{E}\mathcal{A}$, concluding that $S^{-1}\mathcal{E}\mathcal{A}$ is contractible. □

This concludes the section and gives us Theorem **??** and Corollary **??**.