



Department of Mathematics, IIT Delhi

MTL733: Assignment-2

---

**Q.1)** Let  $X(t) = \int_0^t B(s) ds$  be a stochastic process. Show that

$$\mathbb{E}[X^2(t)] = \frac{t^3}{3}, \quad \forall t \geq 0.$$

**Q.2)** Explain whether the stochastic process

$$X(t) = t^2 B(t) - 2 \int_0^t s B(s) ds$$

is a martingale or not (with respect to the filtration  $\mathcal{F}_t = \sigma(B(s) : 0 \leq s \leq t)$ ).

**Q.3)** Let  $m_k(t) = \mathbb{E}[B^k(t)]$ ,  $k = 0, 1, 2, 3, \dots$ . Use Ito-formula to prove that

$$m_k(t) = \frac{1}{2}k(k-1) \int_0^t m_{k-2}(s) ds, \quad k \geq 2.$$

Deduce that

$$\mathbb{E}[B^4(t)] = 3t^2; \quad \mathbb{E}[B^6(t)] = 15t^3.$$

**Q.4)** For  $c, \alpha$  constants, define a stochastic process  $X(t) := \exp\{ct + \alpha B(t)\}$ . Show that

$$dX(t) = \left(c + \frac{1}{2}\alpha^2\right)X(t) dt + \alpha X(t) dB(t).$$

Let  $m(t) := \mathbb{E}[X(t)]$ . Then show that  $m(t)$  satisfies the ODE

$$\begin{cases} m'(t) = \left(c + \frac{1}{2}\alpha^2\right)m(t) \\ m(0) = 1. \end{cases}$$

Show that if  $c < -\frac{1}{2}\alpha^2$ , then  $\lim_{t \rightarrow \infty} m(t) = 0$ .

**Q.5)** Let  $X(t)$  be an Ito process given by  $X(t) = X(0) + \int_0^t v(s) dB(s)$ . Then show that  $M(t) := X^2(t) - \int_0^t v^2(s) ds$  is a martingale.

**Q.6)** Let  $X(t)$  be an Ito process given by

$$dX(t) = u(t)dt + dB(t).$$

Define  $Y(t) = X(t)M(t)$  where  $M(t)$  is a stochastic process given as

$$M(t) = \exp \left\{ - \int_0^t u(s) dB(s) - \frac{1}{2} \int_0^t u^2(s) ds \right\}.$$

Use Ito formula to show that  $Y(t)$  is a martingale. In particular, show that

$$Z(t) = (t + B(t))e^{-B(t) - \frac{t}{2}}$$

is a martingale.

**Q.7)** Using Ito formula, show that the process  $Y(t) = e^{B(t) - \frac{t}{2}}$  is an Ito process with the differential form:

$$dY(t) = Y(t) dB(t)$$

**Q.8)** Write down the differential form of  $\sin(B(t))$ . Using Ito formula, show that the processes  $X(t) = e^{\frac{t}{2}} \sin(B(t))$  and  $Y(t) = e^{\frac{t}{2}} \cos(B(t))$  are martingale.

**Q.9)** Let  $S(t)$  be a positive stochastic process that satisfies the generalized geometric Brownian motion differential equation

$$dS(t) = \alpha(t)S(t)dt + \sigma(t)S(t) dB(t)$$

for some adapted processes  $\alpha(\cdot)$  and  $\sigma(\cdot)$ . Show that  $S(t)$  is given by the formula

$$S(t) = S(0) \exp \left\{ \int_0^t \sigma(s) dB(s) + \int_0^t \left( \alpha(s) - \frac{1}{2} \sigma^2(s) \right) ds \right\}.$$

**Q.10)** Let  $R(t)$  a stochastic process satisfying the Vasicek interest rate equation

$$dR(t) = (\alpha - \beta R(t)) dt + \sigma dB(t)$$

where  $\alpha, \beta$  and  $\sigma$  are positive constants and  $R(0)$  is nonrandom. Show that  $R(t)$  is given by

$$R(t) = e^{-\beta t} \left\{ R(0) + \frac{\alpha}{\beta} (e^{\beta t} - 1) + \sigma \int_0^t e^{\beta s} dB(s) \right\}.$$

Let  $m(t) = \mathbb{E}[R(t)]$ . Show that  $m(t)$  is decreasing if  $R(0) > \frac{\alpha}{\beta}$ , and increasing if  $R(0) < \frac{\alpha}{\beta}$ . Moreover, prove that  $\lim_{t \rightarrow \infty} m(t) = \frac{\alpha}{\beta}$ .

**Q.11)** Show that

$$(1-t) \int_0^t \frac{1}{1-r} dB(r) = B(t) - \int_0^t \left( \int_0^s \frac{1}{1-r} dB(r) \right) ds, \quad 0 \leq t < 1.$$

**Q.12)** Let  $u = u(t, x)$  be a smooth solution of the diffusion equation

$$u_t(t, x) + \frac{1}{2} u_{xx}(t, x) = 0.$$

Show that for each time  $t > 0$ ,  $\mathbb{E}[u(t, B(t))] = u(0, 0)$ .

**Q.13)** Let  $B(t)$  be a Brownian motion. Define a stochastic process

$$M(t) := \int_0^t \text{sign}(B(s)) dB(s), \text{ where } \text{sign}(x) = \begin{cases} 1, & x \geq 0 \\ -1, & x < 0. \end{cases}$$

- Show that  $M(t)$  is a Brownian motion and  $\mathbb{E}[M(t)B(t)] = 0$ .
- Show that  $\mathbb{E}[M(t)B^2(t)] \neq \mathbb{E}[M(t)]\mathbb{E}[B^2(t)]$ . Explain, whether  $M(t)$  and  $B(t)$  are independent or not.

**Q.14)** Let  $\mathbf{X}(t) = (X_1(t), X_2(t))$  be a 2-dimensional stochastic process given by the SDE

$$\begin{aligned} dX_1(t) &= X_2(t) dt \\ dX_2(t) &= \left( -\frac{R}{L}X_2(t) - \frac{1}{CL}X_1(t) + \frac{g(t)}{L} \right) dt + \frac{\alpha}{L} dB(t) \end{aligned}$$

where  $B(t)$  is a one-dimensional Brownian motion,  $R, L, C, \alpha$  are positive constants and  $g(t)$  is a given adapted process. Show that  $\mathbf{X}(t)$  is given by the following formula:

$$\mathbf{X}(t) = \exp(t\mathbf{A}) \left\{ \mathbf{X}(0) + \exp(-t\mathbf{A})\mathbf{K}B(t) + \int_0^t \exp(-s\mathbf{A})[\mathbf{H}(s) + \mathbf{A}\mathbf{K}B(s)] ds \right\},$$

where

$$\mathbf{A} = \begin{pmatrix} 0 & 1 \\ -\frac{1}{CL} & -\frac{R}{L} \end{pmatrix}, \quad \mathbf{K} = \begin{pmatrix} 0 \\ \frac{\alpha}{L} \end{pmatrix}, \quad \mathbf{H}(t) = \begin{pmatrix} 0 \\ \frac{g(t)}{L} \end{pmatrix}.$$

**Q.15)** Show that the process  $\mathbf{X}(t) = (X_1(t), X_2(t))$ , defined by

$$X_1(t) = a \cos(B(t)), \quad X_2(t) = b \sin(B(t)), \quad a, b > 0$$

is a solution of the SDE

$$d\mathbf{X}(t) = -\frac{1}{2}\mathbf{X}(t) dt + \mathbf{M}\mathbf{X}(t) dB(t)$$

where  $\mathbf{M} = \begin{pmatrix} 0 & -\frac{a}{b} \\ \frac{b}{a} & 0 \end{pmatrix}$  and  $B(t)$  is a one-dimensional Brownian motion.

**Q.16)** Let  $\mathbf{B}(t) = (B_1(t), B_2(t), \dots, B_m(t))$  be a  $m$ -dimensional Brownian motion. For any  $\mathbf{g} = (g_{ij}) \in \mathcal{Y}_{n \times m}$  with  $g_{ij} > 0$ , define a stochastic process.

$$g_i(t) := \left( \sum_{j=1}^m g_{ij}^2(t) \right)^{\frac{1}{2}} \quad 1 \leq i \leq n.$$

- For each  $i$ , show that the process  $M_i(t) := \sum_{j=1}^m \int_0^t \frac{g_{ij}(s)}{g_i(s)} dB_j(s)$  is a one-dimensional Brownian motion.
- Explain whether the process  $\mathbf{M}(t) = (M_1(t), M_2(t), \dots, M_n(t))$  is a  $n$ -dimensional Brownian motion or not.

**Q.17)** Find the process  $f(t, \omega) \in \mathcal{Y}(0, T)$  such that  $F = \mathbb{E}[F] + \int_0^T f(t, \omega) dB(t)$  for  $F = B^2(T)$  and  $F = e^{B(T)}$ .

**Q.18)** Find the Ito representation form for the martingales:

- i)  $X(t) := B^3(t) - 3tB(t), \ t \geq 0$
- ii)  $Y(t) := B^4(t) - 6tB^2(t) + 3t^2, \ t \geq 0$
- iii)  $Z(t) = \mathbb{E}[B^2(T)|\mathcal{F}_t], \ 0 \leq t \leq T.$