




Exported on: 21/02/2023

Clinical guidelines - Diagnosis and treatment manual

For curative programmes in hospitals and dispensaries
Guidance for prescribing



This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Médecins Sans Frontières. Clinical guidelines - Diagnosis and treatment manual.

December 2022

ISBN 978-2-37585-200-2

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Table of contents

[Authors/Contributors](#)

[Preface](#)

[Abbreviations and acronyms](#)

[Chapter 1: A few symptoms and syndromes](#)

[Shock](#)

[Seizures](#)

[Hypoglycaemia](#)

[Fever](#)

[Pain](#)

[Anaemia](#)

[Dehydration](#)

[Severe acute malnutrition](#)

[Chapter 2: Respiratory diseases](#)

[Acute upper airway obstruction](#)

[Rhinitis and rhinopharyngitis \(common cold\)](#)

[Acute sinusitis](#)

[Acute pharyngitis](#)

[Diphtheria](#)

[Other upper respiratory tract infections](#)

[Laryngotracheitis and laryngotracheobronchitis \(croup\)](#)

[Epiglottitis](#)

[Bacterial tracheitis](#)

[Otitis](#)

[Acute otitis externa](#)

[Acute otitis media \(AOM\)](#)

[Chronic suppurative otitis media \(CSOM\)](#)

[Whooping cough \(pertussis\)](#)

[Bronchitis](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Asthma

[Asthma attack \(acute asthma\)](#)

[Chronic asthma](#)

[Pulmonary tuberculosis](#)

Chapter 3: Gastrointestinal disorders

[Acute diarrhoea](#)

[Shigellosis](#)

[Amoebiasis](#)

[Disorders of the stomach and duodenum](#)

[Gastro-oesophageal reflux](#)

[Gastric and duodenal ulcers in adults](#)

[Dyspepsia](#)

[Stomatitis](#)

[Oral and oropharyngeal candidiasis](#)

[Oral herpes](#)

[Other infectious causes](#)

[Stomatitis from scurvy \(vitamin C deficiency\)](#)

[Other lesions resulting from a nutritional deficiency](#)

Chapter 4: Skin diseases

[Dermatology](#)

[Scabies](#)

[Lice \(pediculosis\)](#)

[Superficial fungal infections](#)

[Bacterial skin infections](#)

[Impetigo](#)

[Furuncles and carbuncles](#)

[Erysipelas and cellulitis](#)

[Cutaneous anthrax](#)

[Endemic treponematoses](#)

[Leprosy](#)

[Herpes simplex and herpes zoster](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

[Xerophthalmia \(vitamin A deficiency\)](#)

[Conjunctivitis](#)

[Neonatal conjunctivitis](#)

[Viral epidemic keratoconjunctivitis](#)

[Trachoma](#)

[Periorbital and orbital cellulitis](#)

[Other pathologies](#)

[Onchocerciasis \(river blindness\)](#)

[Loiasis](#)

[Pterygium](#)

[Cataract](#)

[Chapter 6: Parasitic diseases](#)

[Malaria](#)

[Human African trypanosomiasis \(sleeping sickness\)](#)

[American trypanosomiasis \(Chagas disease\)](#)

[Leishmaniasis](#)

[Intestinal protozoan infections \(parasitic diarrhoea\)](#)

[Flukes](#)

[Schistosomiasis](#)

[Cestodes](#)

[Nematode infections](#)

[Filariasis](#)

[Onchocerciasis \(river blindness\)](#)

[Loiasis](#)

[Lymphatic filariasis \(LF\)](#)

[Chapter 7: Bacterial diseases](#)

[Bacterial meningitis](#)

[Tetanus](#)

[Enteric \(typhoid and paratyphoid\) fevers](#)

[Brucellosis](#)

[Plague](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

[Rabies](#)

[Viral hepatitis](#)

[Dengue](#)

[Viral haemorrhagic fevers](#)

[HIV infection and AIDS](#)

[Chapter 9: Genito-urinary diseases](#)

[Nephrotic syndrome in children](#)

[Urolithiasis](#)

[Acute cystitis](#)

[Acute pyelonephritis](#)

[Acute prostatitis](#)

[Genital infections](#)

[Urethral discharge](#)

[Abnormal vaginal discharge](#)

[Genital ulcers](#)

[Lower abdominal pain in women](#)

[Upper genital tract infections \(UGTI\)](#)

[Venereal warts](#)

[Major genital infections \(summary\)](#)

[Abnormal uterine bleeding \(in the absence of pregnancy\)](#)

[Chapter 10: Medical and minor surgical procedures](#)

[Dressings](#)

[Treatment of a simple wound](#)

[Burns](#)

[Cutaneous abscess](#)

[Pyomyositis](#)

[Leg ulcers](#)

[Necrotising infections of the skin and soft tissues](#)

[Venomous bites and stings](#)

[Dental infections](#)

[Chapter 11: Mental disorders in adults](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

[Chronic psychoses](#)

[Bipolar disorder](#)

[Chapter 12: Other conditions](#)

[Sickle cell disease](#)

[Diabetes type 2 in adults](#)

[Essential hypertension in adults](#)

[Heart failure in adults](#)

[Chronic heart failure](#)

[Acute heart failure \(acute pulmonary oedema\)](#)

[Endemic goitre and iodine deficiency](#)

[Appendices](#)

[Appendix 1. Normal daily maintenance IV fluids in children > 1 month](#)

[Main references](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Authors/Contributors

The *Clinical guidelines* has been developed by Médecins Sans Frontières.

MSF would like to express its sincere gratitude to everyone who has contributed to developing these guidelines.

Co-authors: Grace Dubois, Blandine Vasseur-Binachon, Cedric Yoshimoto

Contributors: Gabriel Alcoba, Beatriz Alonso, Mohana Amirtharajah, Haydar Alwash, Catherine Bachy, Roberta Caboclo, Severine Caluwaerts, Cristina Carreno, Arlene Chua, Kate Clezy, Anne-Sophie Coutin, Marcio da Fonseca, Martin De Smet, Eva Deplecker, Carolina Echeverri, Sylvie Fagard-Sultan, Roopan Gill, Sonia Guinovart, Jarred Halton, Kerstin Hanson, Christian Heck, Caroline Henry-Ostian, Cathy Hewison, Yves-Laurent Jackson, Carolina Jimenez, John Johnson, Rupa Kanapathipillai, Mohamad Khalife, Nadia Lafferty, Amin Lamrous, James Lee, Helen McColl, Natasha Mlakar, Juno Min, Miguel Palma, Isabella Panunzi, Roberta Petrucci, Nicolas Peyraud, Ernestina Repetto, Jean Rigal, Koert Ritmeijer, Julia Sander, Raghda Sleit, Erin Stratta, Alex Telnov, Malcolm Townsend, Clara Van Gulik.

Specific support has been given by the **International Guidelines Publication team:**

Editor: Véronique Grouzard

Language editors: Mohamed Elsonbaty Ramadan, Carolina López, Anna Romero

Lay-out designer: Evelyne Laissu

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Preface

This guide is designed for use by medical professionals involved in curative care at the dispensary and primary hospital.

We have tried to respond in the simplest and most practical way possible to the questions and problems faced by field medical staff, using the accumulated field experience of Médecins Sans Frontières, the recommendations of reference organizations such as the World Health Organization (WHO) and specialized works in each field.

This edition touches on the curative and, to a lesser extent, the preventive aspects of the main diseases encountered in the field. The list is incomplete, but covers the essential needs.

This guide is used not only in programmes supported by Médecins Sans Frontières, but also in other programmes and in other contexts. It is notably an integral part of the WHO Emergency Health Kit.

Médecins Sans Frontières has also issued French, Spanish and Arabic editions. Editions in other languages have also been produced in the field.

This guide is a collaborative effort of medical professionals from many disciplines, all with field experience.

Despite all efforts, it is possible that certain errors may have been overlooked in this guide. Please inform the authors of any errors detected. It is important to remember, that if in doubt, it is the responsibility of the prescribing medical professional to ensure that the doses indicated in this manual conform to the manufacturer's specifications.

To ensure that this guide continues to evolve while remaining adapted to field realities, please send any comments or suggestions.

As treatment protocols are regularly revised, please check the monthly [updates](#).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Abbreviations and acronyms

Last update : November 2022

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

ACE	angiotensin converting enzyme
ACT	artemisinin-based combination therapy
AFB	acid-fast bacillus
ALT	alanine aminotransferase
ARV	antiretroviral
AST	aspartate aminotransferase
BCG	bacillus Calmette-Guérin
BMI	body mass index
BP	blood pressure
°C	degree Celsius
co-amoxiclav	amoxicillin + clavulanic acid
co-trimoxazole	sulfamethoxazole + trimethoprim
CRT	capillary refill time
CSF	cerebrospinal fluid
CMV	cytomegalovirus
D1 (D2, D3, etc.)	Day 1 or first day (Day 2 or 2 nd day, Day 3 or 3 rd day, etc.)
dl	decilitre
(e)FAST	(extended) focused assessment with sonography for trauma
g	gram
.....

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

IM	intramuscular
IO	intraosseous
IU	international unit
IV	intravenous
kcal	kilocalorie
kg	kilogram
LP	lumbar puncture
mg	milligram
MIU	million of international units
ml	millilitre
mmHg	millimetre of mercury
mmol	millimole
MSF	Médecins Sans Frontières
NSAID	nonsteroidal anti-inflammatory drug
ORS	oral rehydration solution or salts
PCP	pneumocystosis
PCR	polymerase chain reaction
PO	per os – oral administration
POCUS	point-of-care ultrasound
RR	respiratory rate

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

tab	tablet
TB	tuberculosis
TMP	trimethoprim
WHO	World Health Organization

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Chapter 1: A few symptoms and syndromes

[Shock](#)

[Seizures](#)

[Hypoglycaemia](#)

[Fever](#)

[Pain](#)

[Anaemia](#)

[Dehydration](#)

[Severe acute malnutrition](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Shock

Acute circulatory failure leading to inadequate tissue perfusion which, if prolonged, results in irreversible organ failure. Mortality is high without early diagnosis and treatment.

Aetiology and pathophysiology

Hypovolaemic shock

Absolute hypovolaemia due to significant intravascular fluid depletion:

- Internal or external haemorrhage: post-traumatic, peri or postoperative, obstetrical (ectopic pregnancy, uterine rupture, etc.), blood loss due to an underlying condition (gastrointestinal ulcer, etc.). A loss of greater than 30% of blood volume in adults will lead to haemorrhagic shock.
- Dehydration: severe diarrhoea and vomiting, intestinal obstruction, diabetic ketoacidosis or hyperosmolar coma, etc.
- Plasma leaks: extensive burns, crushed limbs, etc.

Relative hypovolaemia due to vasodilation without concomitant increase in intravascular volume:

- Anaphylactic reaction: allergic reaction to insect bites or stings; drugs, mainly neuromuscular blockers, antibiotics, acetylsalicylic acid, colloid solutions (dextran, modified gelatin fluid); equine sera; vaccines containing egg protein; food, etc.
- Acute haemolysis: severe malaria, drug poisoning (rare).

Septic shock

By a complex mechanism, often including vasodilation, heart failure and absolute hypovolaemia.

Cardiogenic shock

By decrease of cardiac output:

- Direct injury to the myocardium: infarction, contusion, trauma, poisoning.
- Indirect mechanism: arrhythmia, constrictive pericarditis, haemopericardium, pulmonary embolism, tension pneumothorax, valvular disease, severe anaemia, beri beri, etc.

Clinical features

Signs common to most forms of shock

- Pallor, mottled skin, cold extremities, sweating and thirst.
- Rapid and weak pulse often only detected on major arteries (femoral or carotid)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

The common signs of shock listed above are typical of hypovolaemic shock.

Do not underestimate hypovolaemia. Signs of shock may not become evident until a 50% loss of blood volume in adults.

Anaphylactic shock

- Significant and sudden drop in BP
- Tachycardia
- Frequent cutaneous signs: rash, urticaria, angioedema
- Respiratory signs: dyspnoea, bronchospasm

Septic shock

- High fever or hypothermia (< 36 °C), rigors, confusion
- BP may be initially maintained, but rapidly, same pattern as for hypovolaemic shock.

Cardiogenic shock

- Respiratory signs of left ventricular failure (acute pulmonary oedema) are dominant: tachypnoea, crepitations on auscultation.
- Signs of right ventricular failure: raised jugular venous pressure, hepatomegaly, sometimes alone, more often associated with signs of left ventricular failure.

The aetiological diagnosis is oriented by:

- The context: trauma, insect bite, ongoing medical treatment, etc.
- The clinical examination:
 - fever
 - skin pinch consistent with dehydration
 - thoracic pain from a myocardial infarction or pulmonary embolus
 - abdominal pain or rigidity of the abdominal wall from peritonitis, abdominal distension from intestinal obstruction
 - blood in stools, vomiting blood in intestinal haemorrhage
 - subcutaneous crepitations, likely anaerobic infection

Management

Symptomatic and aetiological treatment must take place simultaneously.

In all cases

- Emergency: immediate attention to the patient.
- Warm the patient, lay him flat, elevate legs (except in respiratory distress, acute pulmonary oedema).
- Insert a peripheral IV line using a large calibre catheter (16G in adults). If no IV access, use intraosseous route.
- ~~Oxygen therapy, assisted ventilation in the event of respiratory distress~~

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Priority: restore vascular volume as quickly as possible:
Insert 2 peripheral IV lines (2 catheters 16G in adults).
Ringer lactate or 0.9% sodium chloride: replace 3 times the estimated losses
and/or **plasma substitute:** replace 1.5 times the estimated losses
- Transfuse: classically once estimated blood loss represents approximately 30 to 40% of blood volume (25% in children). The blood must be tested (HIV, hepatitis B and C, syphilis, malaria in endemic areas, etc.)

Severe acute dehydration due to bacterial/viral gastroenteritis

- Urgently restore circulating volume using IV therapy with **Ringer lactate:**
Children under 5 years: 20 ml/kg over 15 minutes (to be repeated 2 times if necessary) then 70 ml/kg over 3 hours
Children 5 years and over and adults: 30 ml/kg over 30 minutes (to be repeated once if necessary) then 70 ml/kg over 3 hours
- As soon as the patient is able to drink (often within 2 hours), provide oral rehydration solution (ORS) as the patient tolerates.
- Closely monitor the patient; be careful to avoid fluid overload in young children and elderly patients.
- For aetiological treatment, see [Acute diarrhoea](#), Chapter 3.

Note: in severely malnourished children the IV rate is different than those for healthy children (see [Severe acute malnutrition](#), Chapter 1).

Severe anaphylactic reaction

- Determine the causal agent and remove it, e.g. stop ongoing injections or infusions, but if in place, maintain the IV line.
- Administer **epinephrine (adrenaline)** IM, into the anterolateral part of the thigh, in the event of hypotension, pharyngolaryngeal oedema, or breathing difficulties:
 - Use *undiluted* solution (1:1000 = 1 mg/ml) and a 1 ml syringe graduated in 0.01 ml:
 - Children under 6 years: 0.15 ml
 - Children from 6 to 12 years: 0.3 ml
 - Children over 12 years and adults: 0.5 ml
 - In children, if 1 ml syringe is not available, use a *diluted* solution, i.e. add 1 mg epinephrine to 9 ml of 0.9% sodium chloride to obtain a 0.1 mg/ml solution (1:10 000):
 - Children under 6 years: 1.5 ml
 - Children from 6 to 12 years: 3 ml
 - At the same time, administer rapidly **Ringer lactate or 0.9% sodium chloride:** 1 litre in adults (maximum rate); 20 ml/kg in children, to be repeated if necessary.
If there is no clinical improvement, repeat IM epinephrine every 5 to 15 minutes.
 - In shock persists after 3 IM injections, administration of IV epinephrine at a constant rate by a syringe pump is necessary:
Use a *diluted* solution, i.e. add 1 mg epinephrine (1:1000) to 9 ml of 0.9% sodium chloride to obtain a 0.1 mg/ml

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Septic shock

- Vascular fluid replacement with **Ringer lactate or 0.9% sodium chloride or plasma substitute.**
- Use of vasoconstrictors:
dopamine IV at a constant rate by syringe pump (see [box](#)): 10 to 20 micrograms/kg/minute
or
epinephrine IV at a constant rate by syringe pump:
Use a *diluted* solution, i.e. add 1 mg epinephrine (1:1000) to 9 ml of 0.9% sodium chloride to obtain a 0.1 mg/ml solution (1:10 000). Start with 0.1 microgram/kg/minute. Increase the dose progressively until a clinical improvement is seen.
If syringe pump is not available, see [box](#).
- Look for the origin of the infection (abscess; ENT, pulmonary, digestive, gynaecological or urological infection etc.). Antibiotic therapy according to the origin of infection:

Origin	Antibiotic therapy	Alternative
Cutaneous staphylococci, streptococci	cloxacillin + gentamicin	
Pulmonary pneumococci, <i>Haemophilus influenzae</i>	ampicillin or ceftriaxone +/- gentamicin	co-amoxiclav or ceftriaxone + ciprofloxacin
Intestinal or biliary enterobacteria, anaerobic bacteria, enterococci	co-amoxiclav + gentamicin	ceftriaxone + gentamicin + metronidazole
Gynaecological streptococci, gonococci, anaerobic bacteria, <i>E. coli</i>	co-amoxiclav + gentamicin	ceftriaxone + gentamicin + metronidazole
Urinary enterobacteria, enterococci	ampicillin + gentamicin	ceftriaxone + ciprofloxacin
Other or undetermined	ampicillin + gentamicin	ceftriaxone + ciprofloxacin

ampicillin IV

Children over 1 month: 50 mg/kg every 6 to 8 hours

Adults: 1 to 2 g every 6 to 8 hours

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

ceftriaxone slow IV^a (3 minutes)

Children: 100 mg/kg once daily

Adults: 2 g once daily

ciprofloxacin PO (by nasogastric tube)

Children: 15 mg/kg 2 times daily

Adults: 500 mg 2 times daily

gentamicin IM or slow IV (3 minutes)

Children ≥ 1 month and adults: 6 mg/kg once daily

metronidazole IV infusion (30 minutes)

Children over 1 month: 10 mg/kg every 8 hours (max. 1500 mg daily)

Adults: 500 mg every 8 hours

- Corticosteroids: not recommended, the adverse effects outweigh the benefits.

Cardiogenic shock

The objective is to restore efficient cardiac output. The treatment of cardiogenic shock depends on its mechanism.

- **Acute left heart failure with pulmonary oedema**

- Acute pulmonary oedema (for treatment, see [Heart failure in adults](#), Chapter 12).
- In the event of worsening signs with vascular collapse, use a strong cardiotonic:
dopamine IV at a constant rate by syringe pump (see [box](#)): 3 to 10 micrograms/kg/minute
- Once the haemodynamic situation allows (normal BP, reduction in the signs of peripheral circulatory failure), nitrates or morphine may be cautiously introduced.
- Digoxin should no longer be used for cardiogenic shock, except in the rare cases when a supraventricular tachycardia has been diagnosed by ECG. Correct hypoxia before using digoxin.

digoxin slow IV

Adults: one injection of 0.25 to 0.5 mg, then 0.25 mg 3 or 4 times per 24 hours if necessary

Children: one injection of 0.010 mg/kg (10 micrograms/kg), to be repeated up to 4 times per 24 hours if necessary

- **Cardiac tamponade:** restricted cardiac filling as a result of haemopericardium or pericarditis.

Requires immediate pericardial tap after restoration of circulating volume.

- **Tension pneumothorax:** drainage of the pneumothorax.

- **Symptomatic pulmonary embolism:** treat with an anticoagulant in a hospital setting.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

Administration of **dopamine** or **epinephrine** at a constant rate requires the following conditions:

close medical supervision in a hospital setting;

- use of a dedicated vein (no other infusion/injection in this vein), avoid the antecubital fossa if possible;
- use of an electric syringe pump;
- progressive increase and adaptation of doses according to clinical response;
- intensive monitoring of drug administration, particularly during syringe changes.

Example:

dopamine: 10 micrograms/kg/minute in a patient weighing 60 kg

Hourly dose: $10 \text{ (micrograms)} \times 60 \text{ (kg)} \times 60 \text{ (min)} = 36\,000 \text{ micrograms/hour} = 36 \text{ mg/hour}$

In a 50 ml syringe, dilute one 200 mg-ampoule of dopamine with 0.9% sodium chloride to obtain 50 ml of solution containing 4 mg of dopamine per ml.

For a dose of 36 mg/hour, administer the solution (4 mg/ml) at 9 ml/hour.

If there is no electric syringe pump, dilution in an infusion bag may be considered. However, it is important to consider the risks related to this type of administration (accidental bolus or insufficient dose). The infusion must be constantly monitored to prevent any, even small, change from the prescribed rate of administration.

Example for **epinephrine:**

- In adults:

Dilute 10 ampoules of 1 mg epinephrine (10 000 micrograms) in 1 litre of 5% glucose or 0.9% sodium chloride to obtain a solution containing 10 micrograms of epinephrine per ml.

Knowing that 1 ml = 20 drops, *in an adult weighing 50 kg:*

- 0.1 microgram/kg/minute = 5 micrograms/minute = 10 drops/minute
- 1 microgram/kg/minute = 50 micrograms/minute = 100 drops/minute, etc.

- In children:

Dilute 1 ampoule of 1 mg epinephrine (1000 micrograms) in 100 ml of 5% glucose or 0.9% sodium chloride to obtain a solution containing 10 micrograms of epinephrine per ml.

For administration, use a **paediatric infusion set**; knowing that 1 ml = 60 drops, *in a child weighing 10 kg:*

- 0.1 microgram/kg/minute = 1 microgram/minute = 6 drops/minute
- 0.2 microgram/kg/minute = 2 micrograms/minute = 12 drops/minute, etc.

Note: account for all infused volumes when recording ins and outs.

Footnotes

- (a) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must never be administered by IV route. For IV administration, water for injection must always be used.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Seizures

Involuntary movements of cerebral origin (stiffness followed by clonic movements), accompanied by a loss of consciousness, and often urinary incontinence (generalized tonic-clonic seizures).

In pregnant women, eclamptic seizures require specific medical and obstetrical care. Refer to the guide [Essential obstetric and newborn care](#), MSF.

Initial treatment

During a seizure

- Protect from trauma, maintain airway, place patient in 'recovery position', loosen clothing.
- Most seizures are quickly self-limited. Immediate administration of an anticonvulsant is not systematic. If generalized seizure lasts more than 5 minutes, use diazepam to stop it:

diazepam

Children: 0.5 mg/kg preferably rectally^a without exceeding 10 mg

IV administration is possible (0.3 mg/kg over 2 or 3 minutes), only if means of ventilation are available (Ambu bag and mask).

Adults: 10 mg rectally or by slow IV

In all cases:

- If seizure continues, repeat dose once after 10 minutes.
- In infants and elderly patients, monitor respiratory rate and blood pressure.
- If seizure continues after the second dose, treat as status epilepticus.

The patient is no longer seizing

- Look for the cause of the seizure and evaluate the risk of recurrence.
- Keep diazepam and glucose available in case the patient starts seizing again.

Status epilepticus

Several distinct seizures without complete restoration of consciousness in between or an uninterrupted seizure lasting more than 30 minutes.

- Protect from trauma, loosen clothing, maintain airway and administer oxygen as required.
- Insert an intravenous or intraosseous line.
- Treat for hypoglycaemia (see [Hypoglycaemia](#), Chapter 1).
- If 2 doses of diazepam have not stopped the seizures, use phenytoin or phenobarbital if phenytoin is not available or if seizures persist despite phenytoin.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

phenytoin slow IV infusion 250 mg in 5 ml ampoule (50 mg/ml)	<ul style="list-style-type: none"> Children 1 month and over and adults: one dose of 15 to 20 mg/kg administered over 20 minutes minimum and 60 minutes maximum The concentration of the diluted solution should be between 5 and 10 mg/ml. The infusion rate should not exceed 1 mg/kg/minute or 50 mg/minute (25 mg/minute in elderly patients or patients with cardiac disorders). <p>For example:</p> <p>Child weighing 8 kg: 160 mg (20 mg x 8 kg), i.e. 3.2 ml of phenytoin in 17 ml of 0.9% sodium chloride over 30 minutes</p> <p>Adult weighing 50 kg: 1 g (20 mg x 50 kg), i.e. 20 ml of phenytoin in a bag of 100 ml of 0.9% sodium chloride over 30 minutes</p> <p>Do not dilute phenytoin in glucose. Do not administer via a line used for glucose solution administration. Use a large catheter. Check the insertion site and for blood backflow (risk of necrosis in the event of extravasation). After each infusion, rinse with 0.9% sodium chloride to limit local venous irritation.</p>
phenobarbital slow IV infusion 200 mg in 1 ml ampoule (200 mg/ml)	<ul style="list-style-type: none"> Children 1 month to < 12 years: one dose of 15 to 20 mg/kg (max. 1 g) administered over 20 minutes minimum If necessary, a second dose of 10 mg/kg may be administered 15 to 30 minutes after the first dose. Children ≥ 12 years and adults: one dose of 10 mg/kg (max. 1 g) administered over 20 minutes minimum If necessary, a second dose of 5 to 10 mg/kg may be administered 15 to 30 minutes after the first dose. Do not administer more than 1 mg/kg/minute. <p>For example:</p> <p>Child weighing 8 kg: 120 mg (15 mg x 8 kg), i.e. 0.6 ml of phenobarbital in 20 ml of 0.9% sodium chloride over 20 minutes</p> <p>Adult weighing 50 kg: 500 mg (10 mg x 50 kg), i.e. 2.5 ml of phenobarbital in a bag of 100 ml of 0.9% sodium chloride over 20 minutes</p> <p>For doses less than 1 ml, use a 1 ml syringe graduated 0.01 ml to draw the phenobarbital.</p>

Further treatment

Febrile seizures

- Determine the cause of the fever. Give paracetamol (see [Fever](#), Chapter 1).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Hypoglycaemia: administer glucose by slow IV injection to all patients who do not regain consciousness, to patients with severe malaria and to newborns and malnourished children. When possible, confirm hypoglycaemia (reagent strip test).

Iatrogenic causes

Withdrawal of antiepileptic therapy in a patient being treated for epilepsy should be managed over a period of 4-6 months with progressive reduction of the doses. An abrupt stop of treatment may provoke severe recurrent seizures.

Epilepsy

- A first brief seizure does not need further protective treatment. Only patients with chronic repetitive seizures require further regular protective treatment with an antiepileptic drug, usually over several years.
- Once a diagnosis is made, abstention from treatment may be recommended due to the risks associated with treatment. However, these risks must be balanced with the risks of aggravation of the epilepsy, ensuing seizure-induced cerebral damage and other injury if the patient is not treated.
- It is always preferable to start with monotherapy. The effective dose must be reached progressively and symptoms and drug tolerance evaluated every 15 to 20 days.
- An abrupt interruption of treatment may provoke status epilepticus. The rate of dose reduction varies according to the length of treatment; the longer the treatment period, the longer the reduction period (see [Iatrogenic causes](#)). In the same way, a change from one antiepileptic drug to another must be made progressively with an overlap period of a few weeks.
- First line treatments for generalised tonic-clonic seizures in children under 2 years are carbamazepine or phenobarbital, in older children and adults sodium valproate or carbamazepine.

For information:

▫ **sodium valproate PO**

Adults: initial dose of 300 mg 2 times daily; increase by 200 mg every 3 days if necessary until the optimal dose has been reached (usually 500 mg to 1 g 2 times daily).

Children over 20 kg: initial dose of 200 mg 2 times daily irrespective of weight; increase the dose progressively if necessary until the optimal dose has been reached (usually 10 to 15 mg/kg 2 times daily).

▫ **carbamazepine PO**

Adults: initial dose of 100 to 200 mg once or 2 times daily; increase the dose every week by 100 to 200 mg, up to 400 mg 2 to 3 times daily (max. 1600 mg daily)

Children 1 month and over: initial dose of 5 mg/kg once daily or 2.5 mg/kg 2 times daily; increase the dose every week by 2.5 to 5 mg/kg, up to 5 mg/kg 2 to 3 times daily (max. 20 mg/kg daily)

▫ **phenobarbital PO**

Adults: initial dose of 2 mg/kg once daily (max. 100 mg); increase the dose progressively up to 6 mg/kg daily if necessary

Children: initial dose of 3 to 4 mg/kg once daily at bedtime; increase the dose progressively up to 8 mg/kg daily if necessary

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

Hypoglycaemia

Last update: November 2022

Hypoglycaemia is an abnormally low concentration of blood glucose. Severe hypoglycaemia can be fatal or lead to irreversible neurological damage.

Blood glucose levels should be measured whenever possible in patients presenting symptoms of hypoglycaemia. If hypoglycaemia is suspected but blood glucose measurement is not available, glucose (or another available sugar) should be given empirically.

Always consider hypoglycaemia in patients presenting impaired consciousness (lethargy, coma) or seizures.

For diagnosis and treatment of hypoglycaemia in neonates, refer to the guide [Essential obstetric and newborn care](#), MSF.

Clinical features

Rapid onset of non-specific signs, mild to severe depending on the degree of the hypoglycaemia: sensation of hunger and fatigue, tremors, tachycardia, pallor, sweats, anxiety, blurred vision, difficulty speaking, confusion, convulsions, lethargy, coma.

Diagnosis

Capillary blood glucose concentration (reagent strip test):

- Non-diabetic patients:
 - Hypoglycaemia: < 3.3 mmol/litre (< 60 mg/dl)
 - Severe hypoglycaemia: < 2.2 mmol/litre (< 40 mg/dl)
- Diabetic patients on home treatment: < 3.9 mmol/litre (< 70 mg/dl)^[1]

If blood glucose measurement is not available, diagnosis is confirmed when symptoms resolve after the administration of sugar or glucose.

Symptomatic treatment

- Conscious patients:

Children: a teaspoon of powdered sugar in a few ml of water or 50 ml of fruit juice, maternal or therapeutic milk or 10 ml/kg of 10% glucose by oral route or nasogastric tube.

Adults: 15 to 20 g of sugar (3 or 4 cubes) or sugar water, fruit juice, soda, etc.

Symptoms improve approximately 15 minutes after taking sugar by oral route.

- Patients with impaired consciousness or prolonged convulsions:

Children: 0.1 ml/kg of 10% glucose by slow intravenous infusion over 10 to 20 minutes^[2]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

If patient does not return to full alertness after an episode of severe hypoglycaemia, monitor blood glucose levels regularly.

Aetiological treatment

- Other than diabetes:
 - Treat severe malnutrition, neonatal sepsis, severe malaria, acute alcohol intoxication, etc.
 - End prolonged fast.
 - Replace drugs inducing hypoglycaemia (e.g. quinine IV, pentamidine, ciprofloxacin, enalapril, beta-blockers, high-dose aspirin, tramadol), or anticipate hypoglycaemia (e.g. administer quinine IV in a glucose infusion).
- In diabetic patients:
 - Avoid missing meals, increase intake of carbohydrates if necessary.
 - Adjust dosage of insulin according to blood glucose levels and physical activity.
 - Adjust dosage of oral antidiabetics, taking into account possible drug interactions.

Footnotes

- (a) If ready-made 10% glucose solution is not available: remove 100 ml of 5% glucose from a 500 ml bottle or bag, then add 50 ml of 50% glucose to the remaining 400 ml of 5% glucose to obtain 450 ml of 10% glucose solution.

References

1. American Diabetes Association Standards of Medical Care in Diabetes, 2017.
http://care.diabetesjournals.org/content/diacare/suppl/2016/12/15/40.Supplement_1.DC1/DC_40_S1_final.pdf [Accessed 24 May 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Fever

Fever is defined as an axillary temperature higher than or equal to 37.5 °C.

Fever is frequently due to infection. In a febrile patient, first look for signs of serious illness then, try to establish a diagnosis.

Signs of severity

- Severe tachycardia, tachypnoea, respiratory distress, SpO₂ ≤ 90%.
- Shock, altered mental status, petechial or purpuric rash, meningeal signs, seizures, heart murmur, severe abdominal pain, dehydration, critically ill appearance^a; a bulging fontanel in young children.

Infectious causes of fever according to localizing symptoms

Signs or symptoms	Possible aetiology
Meningeal signs, seizures	Meningitis/meningoencephalitis/severe malaria
Abdominal pain or peritoneal signs	Appendicitis/peritonitis/typhoid fever
Diarrhoea, vomiting	Gastroenteritis/typhoid fever
Jaundice, enlarged liver	Viral hepatitis
Cough	Pneumonia/measles/tuberculosis if persistent
Ear pain, red tympanic membrane	Otitis media
Sore throat, enlarged lymph nodes	Streptococcal pharyngitis, diphtheria
Dysuria, urinary frequency, back pain	Urinary tract infection
Red, warm, painful skin	Erysipelas, cellulitis, abscess
Limp, difficulty walking	Osteomyelitis/septic arthritis
Rash	Measles/dengue/haemorrhagic fever/chikungunya

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Laboratory and other examinations

- Children less than 2 months with a temperature higher than or equal to 37.5 °C without a focus:
 - Urinary dipstick;
 - Lumbar puncture (LP) if child less than 1 month or if any of the following: meningeal signs, coma, seizures, critically ill appearance^a, failure of prior antibiotic therapy, suspicion of staphylococcal infection;
 - Chest X-Ray (if available) in case of signs of respiratory disease.
- Children 2 months to 3 years with a temperature higher than or equal to 38 °C without a focus:
 - Urine dipstick;
 - White blood cell count (WBC) if available;
 - LP if meningeal signs.
- Children over 3 years and adults with a temperature higher than or equal to 39 °C:
According to clinical presentation.

Aetiological treatment

- Antibiotherapy according to the cause of fever.
- For patients with sickle cell disease, see [Sickle cell disease](#), Chapter 12.
- If no source of infection is found, hospitalise and treat the following children with empiric antibiotics:
 - Children less than 1 month;
 - Children 1 month to 3 years with WBC ≥ 15000 or ≤ 5000 cells/mm³;
 - All critically ill appearing^a patients or those with signs of serious illness;For antibiotic doses according to age, see [Acute pneumonia](#), Chapter 2.

Symptomatic treatment

- Undress the patient. Do not wrap children in wet towels or cloths (not effective, increases discomfort, risk of hypothermia).
- Antipyretics may increase the patient's comfort but they do not prevent febrile convulsions. Do not treat for more than 3 days with antipyretics.

paracetamol PO

Children less than 1 month: 10 mg/kg 3 to 4 times daily (max. 40 mg/kg daily)

Children 1 month and over: 15 mg/kg 3 to 4 times daily (max. 60 mg/kg daily)

Adults: 1 g 3 to 4 times daily (max. 4 g daily)

or

ibuprofen PO

Children over 3 months and < 12 years: 5 to 10 mg/kg 3 to 4 times daily (max. 30 mg/kg daily)

Children 12 years and over and adults: 200 to 400 mg 3 to 4 times daily (max. 1200 mg daily)

or

acetylsalicylic acid (ASA) PO

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- In case of haemorrhagic fever and dengue: acetylsalicylic acid and ibuprofen are contraindicated; use paracetamol with caution in the presence of hepatic dysfunction.

Footnotes

- (a) Critically ill appearing child: weak grunting or crying, drowsiness, difficult to arouse, does not smile, unconjugate or anxious gaze, pallor or cyanosis, general hypotonia.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Pain

Pain results from a variety of pathological processes. It is expressed differently by each patient depending on cultural background, age, etc. It is a subjective experience meaning that only the individual is able to assess his/her level of pain. Regular assessment of the intensity of pain is indispensable in establishing effective treatment.

Clinical features

Pain assessment

- Intensity: use a simple verbal scale in children over 5 years and adults, and NFCS or FLACC scales in children less than 5 years (see [Pain evaluation scales](#)).
- Pattern: sudden, intermittent, chronic; at rest, at night, on movement, during care procedures, etc.
- Character: burning, cramping, spasmodic, radiating, etc.
- Aggravating or relieving factors, etc.

Clinical examination

- Of the organ or area where the pain is located.
- Specific signs of underlying disease (e.g. bone or osteoarticular pain may be caused by a vitamin C deficiency) and review of all systems.
- Associated signs (fever, weight loss, etc.).

Synthesis

The synthesis of information gathered during history taking and clinical examination allows aetiological diagnosis and orients treatment. It is important to distinguish:

- **Nociceptive pain:** it presents most often as acute pain and the cause-effect relationship is usually obvious (e.g. acute post-operative pain, burns, trauma, renal colic, etc.). The pain may be present in different forms, but neurological exam is normal. Treatment is relatively well standardized.
- **Neuropathic pain**, due to a nerve lesion (section, stretching, ischaemia): most often chronic pain. On a background of constant, more or less localized pain, such as paraesthesia or burning, there are recurrent acute attacks such as electric shock-like pain, frequently associated with disordered sensation (anaesthesia, hypo or hyperaesthesia). This type of pain is linked to viral infections directly affecting the CNS (herpes simplex, herpes zoster), neural compression by tumors, post- amputation pain, paraplegia, etc.
- **Mixed pain** (cancer, HIV) for which management requires a broader approach.

Pain evaluation scales

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Observational evaluation scale - Children 2 months-5 years

FLACC scale (Face Limb Activity Cry Consolability)

Items	Scoring		
	0	1	2
Face	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested	Frequent to constant frown, clenched jaw, quivering chin
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid or jerking
Cry	No cry (awake or asleep)	Moans or whimpers, occasional complaint	Crying steadily, screams or sobs, frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging or being talked to, distractible	Difficult to console or comfort


Each category is scored from 0 to 2, giving a final score between 0 and 10.

0 to 3: mild pain, 4 to 7: moderate pain, 7 to 10: severe pain

Observational evaluation scale - Children under 2 months

NFCS scale (Neonatal Facial Coding System)

Items	Scoring	
	0	1
Brow bulge	no	yes
Eye squeeze	no	yes
Nasolabial furrow	no	yes



This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Nociceptive pain

The WHO classifies analgesics used for this type of pain on a three-step ladder:

- **Step 1:** non-opioid analgesics such as paracetamol and nonsteroidal anti-inflammatory drugs (NSAIDs).
- **Step 2:** weak opioid analgesics such as codeine and tramadol. Their combination with one or two Step 1 analgesics is recommended.
- **Step 3:** strong opioid analgesics, first and foremost morphine. Their combination with one or two Step 1 analgesics is recommended.

The treatment of pain is based on a few fundamental concepts:

- Pain can only be treated correctly if it is correctly evaluated. The only person who can evaluate the intensity of pain is the patient himself. The use of pain assessment scales is invaluable.
- The pain evaluation observations should be recorded in the patient chart in the same fashion as other vital signs.
- Treatment of pain should be as prompt as possible.
- It is recommended to administer analgesics in advance when appropriate (e.g. before painful care procedures).
- Analgesics should be prescribed and administered at fixed time intervals (not on demand).
- Oral forms should be used whenever possible.
- The combination of different analgesic drugs (multimodal analgesia) is advantageous.
- Start with an analgesic from the level presumed most effective: e.g., in the event of a fractured femur, start with a Step 3 analgesic.
- The treatment and dose chosen are guided by the assessment of pain intensity, but also by the patient's response which may vary significantly from one person to another.

Treatment of acute pain

Mild pain	Paracetamol + /- NSAID
Moderate pain	Paracetamol + /- NSAID + tramadol or codeine
Severe pain	Paracetamol + /- NSAID + morphine

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	Analgesics	Children	Adults (except pregnant/breast-feeding women)	Remarks
Level 1	paracetamol PO	< 1 month: 10 mg/kg every 6 to 8 hours (max. 40 mg/kg daily) ≥ 1 month: 15 mg/kg every 6 to 8 hours (max. 60 mg/kg daily)	1 g every 6 to 8 hours (max. 4 g daily)	The efficacy of IV paracetamol is not superior to the efficacy of oral paracetamol; the IV route is restricted to situations where oral administration is impossible.
	paracetamol IV	< 1 month: 7.5 mg/kg every 6 hours (max. 30 mg/kg daily) ≥ 1 month and < 10 kg: 10 mg/kg every 6 hours (max. 30 mg/kg daily) ≥ 10 kg: 15 mg/kg every 6 hours (max. 60 mg/kg daily)	< 50 kg: 15 mg/kg every 6 hours (max. 60 mg/kg daily) ≥ 50 kg: 1 g every 6 hours (max. 4 g daily)	
	acetylsalicylic acid (aspirin) PO	–	300 mg to 1 g every 4 to 6 hours (max. 4 g daily)	Avoid in children less than 16 years.
	diclofenac IM	–	75 mg once daily	Treatment must be as short as possible.
	ibuprofen PO	> 3 months: 5 to 10 mg/kg every 6 to 8 hours (max. 30 mg/kg daily) > 12 years: as for adults	200 to 400 mg every 6 to 8 hours (max. 1200 mg daily)	Respect contra-indications.
	codeine PO	> 12 years: 30 to 60 mg every 4 to 6 hours (max. 240 mg daily)	30 to 60 mg every 4 to 6 hours (max. 240 mg daily)	Add a laxative if treatment > 48 hours.
Level 2	tramadol PO	> 12 years: 50 to 100 mg every 4 to 6 hours (max.	50 to 100 mg every 4 to 6 hours (max. 400 mg daily)	25 to 50 mg every 12 hours in

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

			<ul style="list-style-type: none"> • Add a laxative if treatment > 48 hours.
morphine PO sustained release (MSR)	The daily dose is determined during the initial treatment with immediate release morphine (MIR). If treatment is initiated directly with MSR: ≥ 6 months: 0.5 mg/kg every 12 hours, to be adjusted in relation to pain intensity	The daily dose is determined during the initial treatment with immediate release morphine (MIR). If treatment is initiated directly with MSR: 30 mg every 12 hours, to be adjusted in relation to pain intensity	<ul style="list-style-type: none"> • Do not initiate treatment with the MSR in elderly patients and patients with renal or hepatic impairment. Begin treatment with MIR. • Add a laxative if treatment > 48 hours.
morphine SC, IM	> 6 months: 0.1 to 0.2 mg/kg every 4 hours	0.1 to 0.2 mg/kg every 4 hours	<ul style="list-style-type: none"> • Reduce doses by half and administer less frequently, according to clinical response, in elderly patients and patients with severe renal or hepatic impairment.
morphine IV	> 6 months: 0.1 mg/kg administered in fractionated doses (0.05 mg/kg every 10 minutes) every 4 hours if necessary	0.1 mg/kg administered in fractionated doses (0.05 mg/kg every 10 minutes) every 4 hours if necessary	<ul style="list-style-type: none"> • Add a laxative if treatment > 48 hours.

Notes on the use of morphine and derivatives:

- Morphine is an effective treatment for many types of severe pain. Its analgesic effect is dose-dependent. Its adverse effects have often been exaggerated and should not be an obstacle to its use.
- The most serious adverse effect of morphine is respiratory depression, which may be fatal. This adverse effect results from overdose. It is, therefore, important to increase doses gradually. Respiratory depression is preceded by drowsiness, which is a warning to monitor respiratory rate (RR).

The RR should remain equal to or greater than the thresholds indicated below:

Children 1 to 12 months	RR ≥ 25 respirations/minute
Children 1 to 2 years	RR ≥ 20 respirations/minute
Children 2 to 5 years	RR ≥ 15 respirations/minute
Children < 5 years and adults	RR ≥ 10 respirations/minute

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

If the patient's stools are soft, a stimulant laxative (**bisacodyl** PO: children > 3 years: 5 to 10 mg once daily; adults: 10 to 15 mg once daily) is preferred.

- Nausea and vomiting are common at the beginning of treatment.

Children:

ondansetron PO: 0.15 mg/kg (max. 4 mg per dose) up to 3 times daily

Do not use metoclopramide in children.

Adults:

haloperidol PO (2 mg/ml oral solution): 1 to 2 mg up to 6 times daily or **metoclopramide** PO: 5 to 10 mg 3 times daily with an interval of at least 6 hours between each dose

Do not combine haloperidol and metoclopramide.

- For chronic pain in late stage disease (cancer, AIDS etc.), morphine PO is the drug of choice. It may be necessary to increase doses over time according to pain assessment. Do not hesitate to give sufficient and effective doses.
- Morphine, tramadol and codeine have similar modes of action and should not be combined.
- Buprenorphine, nalbuphine and pentazocine must not be combined with morphine, pethidine, tramadol or codeine because they have competitive action.

Treatment of nociceptive pain in pregnant and breast-feeding women

Analgesics		Pregnancy		Breast-feeding
		0-5 months	From 6 th month	
Level 1	paracetamol	first choice	first choice	first choice
	aspirin	avoid	contra-indicated	avoid
	ibuprofen	avoid	contra-indicated	possible
Level 2	codeine	possible	The neonate may develop withdrawal symptoms, respiratory depression and drowsiness in the event of prolonged administration of large doses at the end of the third trimester. Closely monitor the neonate.	Use with caution, for a short period (2-3 days), at the lowest effective dose. Monitor the mother and the child: in the event of excessive drowsiness, stop treatment.
	tramadol	possible	The child may develop drowsiness when the mother receives tramadol at the end of the third trimester and during breast-feeding. Administer with caution, for a short period, at the lowest effective dose, and monitor the child.	

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Commonly used analgesics are often ineffective in treating this type of pain.

Treatment of neuropathic pain is based on a combination of two centrally acting drugs:

amitriptyline PO

Adults: 25 mg once daily at bedtime (Week 1); 50 mg once daily at bedtime (Week 2); 75 mg once daily at bedtime (as of Week 3); max.150 mg daily. Reduce the dose by half in elderly patients.

carbamazepine PO

Adults: 200 mg once daily at bedtime (Week 1); 200 mg 2 times daily (Week 2); 200 mg 3 times daily (as of Week 3)

Given its teratogenic risk, carbamazepine should only be used in women of childbearing age when covered by effective contraception (intrauterine device or injectable progestogen). It is not recommended in pregnant women.

Mixed pain

In mixed pain with a significant component of nociceptive pain, such as in cancer or AIDS, morphine is combined with antidepressants and antiepileptics.

Chronic pain

In contrast to acute pain, medical treatment alone is not always sufficient in controlling chronic pain. A multidisciplinary approach including medical treatment, physiotherapy, psychotherapy and nursing is often necessary to allow good pain relief and encourage patient selfmanagement.

Co-analgesics

The combination of certain drugs may be useful or even essential in the treatment of pain: antispasmodics, muscle relaxants, anxiolytics, corticosteroids, local anaesthesia, etc.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Anaemia

Anaemia is defined as a haemoglobin (Hb) level below reference values, which vary depending on sex, age and pregnancy status (see [Table 2](#)).

Anaemia may be caused by:

- Decreased production of red blood cells: iron deficiency, nutritional deficiencies (folic acid, vitamin B₁₂, vitamin A), depressed bone marrow function, certain infections (HIV, visceral leishmaniasis), renal failure;
- Loss of red blood cells: acute or chronic haemorrhage (ancylostomiasis, schistosomiasis, etc.);
- Increased destruction of red blood cells (haemolysis): parasitic (malaria), bacterial and viral (HIV) infections; haemoglobinopathies (sickle cell disease, thalassaemia); intolerance to certain drugs (primaquine, dapsone, co-trimoxazole, etc.) in patients with G6PD deficiency.

In tropical settings, the causes of anaemia are often interlinked.

Clinical features

- Common signs: pallor of the conjunctivae, mucous membranes, palms of hands and soles of feet; fatigue, dizziness, dyspnoea, tachycardia, heart murmur.
- Signs that anaemia may be immediately life threatening: sweating, thirst, cold extremities, oedema in the lower limbs, respiratory distress, angina, shock.
- Significant signs: cheilosis and glossitis (nutritional deficiency), jaundice, hepatosplenomegaly, dark coloured urine (haemolysis), bleeding (maelena, haematuria, etc.), signs of [malaria](#).

Laboratory

- Hb levels
- Rapid test or systematic thick and thin blood films in areas where malaria is endemic.
- Urinary dipstick: check for haemoglobinuria or haematuria.
- Emmel test if sickle cell disease is suspected.
- Blood cell count if available to guide diagnosis.

Table 1 - Possible diagnoses with blood cell count

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Characteristics	Possible diagnoses
Macrocytic	Deficiency (folic acid, vitamin B ₁₂), chronic alcoholism
Microcytic	Iron deficiency (malnutrition, chronic haemorrhage), chronic inflammation (HIV infection, cancer), thalassaemia
Normocytic	Acute haemorrhage, renal failure, haemolysis
Reduced number of reticulocytes	Deficiency (iron, folic acid, vitamin B ₁₂), spinal tumour, renal failure
Increased or normal number of reticulocytes	Haemolysis, sickle cell disease, thalassaemia
Eosinophilia	Ancylostomiasis, trichuriasis, schistosomiasis, HIV infection, malignant haemopathies

Aetiological treatment

Anaemia in itself is not an indication for transfusion. Most anaemias are well tolerated and can be corrected with simple aetiological treatment.

Aetiological treatment may be given alone or together with transfusion.

- Iron deficiency

ferrous salts PO for 3 months. Doses are expressed in elemental iron^a:

Neonates: 1 to 2 mg/kg 2 times daily

Children 1 month to < 6 years: 1.5 to 3 mg/kg 2 times daily

Children 6 to < 12 years: 65 mg 2 times daily

Children ≥ 12 years and adults: 65 mg 2 to 3 times daily

Age	Weight	Treatment	
		45 mg/5 ml syrup	65 mg tablet
< 1 month	< 4 kg	0.5 ml x 2	-
1 month to < 1 year	4 to < 10 kg	1.5 ml x 2	-

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Children < 1 year: 0.5 mg/kg once daily

Children ≥ 1 year and adults: 5 mg once daily

- Malaria: see [Malaria](#) (Chapter 6). In the event of associated iron deficiency, wait 4 weeks after malaria treatment before prescribing iron supplements.
- Suspected haemolytic anaemia: stop any drug that causes haemolysis in patients with (or that may possibly have) G6PD deficiency.

Blood transfusion

Indications

To decide whether to transfuse, several parameters should be taken into account:

- Clinical tolerance of anaemia
- Underlying conditions (cardiovascular disease, infection, etc.)
- Rate at which anaemia develops.
- Hb levels

If transfusion is indicated, it should be carried out without delay^b. For transfusion thresholds, see [Table 2](#).

Volume to be transfused

In the absence of hypovolaemia or shock:

Children < 20 kg: 15 ml/kg of red cell concentrate in 3 hours or 20 ml/kg of whole blood in 4 hours

Children ≥ 20 kg and adults: start with an adult unit of whole blood or red cell concentrate; do not exceed a transfusion rate of 5 ml/kg/hour

Repeat if necessary, depending on clinical condition.

Monitoring

Monitor the patient's condition and vital signs (heart rate, blood pressure, respiratory rate, temperature):

- During the transfusion: 5 minutes after the start of transfusion, then every 15 minutes during the first hour, then every 30 minutes until the end of the transfusion.
- After the transfusion: 4 to 6 hours after the end of the transfusion.

If signs of circulatory overload appear:

- Stop temporarily the transfusion.
- Sit the patient in an upright position.
- Administer oxygen.
- Administer **furosemide** by slow IV:

Children: 0.5 to 1 mg/kg

Adults: 20 to 40 mg

Repeat the injection (same dose) after 2 hours if necessary.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Children 1 month to < 12 years: 1 to 2 mg/kg once daily (max. 65 mg daily)
Children ≥ 12 years and adults: 65 mg once daily

Age	Weight	Prevention	
		45 mg/5 ml syrup	65 mg tablet
< 1 month	< 4 kg	0.5 ml	-
1 month to < 1 year	4 to < 10 kg	1 ml	-
1 to < 6 years	10 to < 20 kg	2.5 ml	-
6 to < 12 years	20 to < 40 kg	5 ml	-
≥ 12 years and adults	≥ 40 kg	-	1 tab

or preferably,

ferrous salts + folic acid PO based on elemental iron dosages.

- Nutritional supplements (if the basic diet is insufficient)
- In the event of sickle cell anaemia: see [Sickle cell disease](#) (Chapter 12).
- Early treatment of malaria, helminthic infections, etc.

Table 2 - Definition of anaemia and transfusion thresholds

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Patients	Hb levels defining anaemia	Transfusion threshold
Children 2-6 months	< 9.5 g/dl	Hb < 4 g/dl, even if there are no signs of decompensation Hb ≥ 4 g/dl and < 6 g/dl if there are signs of decompensation or sickle cell disease or severe malaria or serious bacterial infection or pre-existing heart disease
Children 6 months-5 years	< 11 g/dl	
Children 6-11 years	< 11.5 g/dl	
Children 12-14 years	< 12 g/dl	
Men	< 12 g/dl	Hb < 7 g/dl if there are signs of decompensation or sickle cell disease or severe malaria or serious bacterial infection or pre-existing heart disease
Women	< 13 g/dl	
Pregnant women	<p>< 11 g/dl (1st and 3rd trimester)</p> <p>< 10.5 g/dl (2nd trimester)</p>	<p>< 36 weeks</p> <p>Hb ≤ 5 g/dl, even if there are no signs of decompensation Hb > 5 g/dl and < 7 g/dl if there are signs of decompensation or sickle cell disease or severe malaria or serious bacterial infection or pre-existing heart disease</p> <p>≥ 36 weeks</p> <p>Hb ≤ 6 g/dl, even if there are no signs of decompensation Hb > 6 g/dl and < 8 g/dl if there are signs of decompensation or sickle cell disease or severe malaria or serious bacterial infection or pre-existing heart disease</p>

Footnotes

- (a) Available in 140 mg/5 ml syrup of ferrous fumarate containing approximately 45 mg/5 ml of elemental iron and 200 mg ferrous sulfate tablets or ferrous sulfate + folic acid tablets containing 65 mg of elemental iron. Tablets of 185 or 200 mg ferrous fumurate or sulfate + folic acid (60 or 65 mg of elemental iron) contain 400 micrograms folic acid.
- (b) Before transfusing: determine the recipient's and potential donors' blood groups/rhesus and carry out screening tests on the donor's blood for HIV-1 and 2, syphilis and, in endemic areas, malaria and Chagas disease.

References

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Dehydration

Dehydration results from excessive loss of water and electrolytes from the body. If prolonged, dehydration can compromise organ perfusion, resulting in shock.

It is principally caused by diarrhoea, vomiting and severe burns.

Children are particularly susceptible to dehydration due to frequent episodes of gastroenteritis, high surface area to volume ratio and inability to fully communicate, or independently meet their fluid needs.

The protocols below are focused on treatment of dehydration caused by diarrhoea and vomiting. Alternative treatment protocols should be used for children with malnutrition (see [Severe acute malnutrition](#), Chapter 1) or in patients with severe burns (see [Burns](#), Chapter 10).

Clinical features and assessment

- History of diarrhoea and/or vomiting and concomitant reduced urine output.
- Clinical features depend on the degree of dehydration (see table below). Features such as dry mouth, absence of tears may also be noted.
- Patients with severe dehydration should be assessed for shock (tachycardia, low blood pressure and delayed capillary refill time etc.).
- Electrolyte disorders may cause tachypnoea, muscle cramps or weakness, cardiac arrhythmia (irregular heart rate, palpitation), confusion and/or seizures.

Classification of degree of dehydration (adapted from the WHO)^{[1][2]}

	Severe dehydration At least 2 of the following signs:	Some dehydration At least 2 of the following signs:	No dehydration No signs of "severe" or "some" dehydration.
Mental status	Lethargic or unconscious	Restless or irritable	Normal
Radial pulse	Weak or absent	Palpable	Easily palpable
Eyes^(a)	Sunken	Sunken	Normal
Skin pinch^(b)	Goes back very slowly (> 2 seconds)	Goes back slowly (< 2 seconds)	Goes back quickly (< 1 second)

Drinks no fluids

Thirst

No thirst

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Severe dehydration

- Treat shock if present (see [Shock](#), Chapter 1).
- If able to drink, administer oral rehydration solution (ORS) PO whilst obtaining IV access. according to WHO Treatment Plan C, monitoring infusion rate closely:
- Insert peripheral IV line using large caliber catheter (22-24G in children or 18G in adults) or intraosseous needle.
- Administer **Ringer lactate** (RL)^a

WHO Treatment Plan C^{[1][2]}

Age	First, give 30 ml/kg over ^(c) :	Then, give 70 ml/kg over:
Children < 1 year	1 hour	5 hours
Children ≥ 1 year and adults	30 minutes	2 ½ hours

(c) Repeat once if radial pulse remains weak or absent after first bolus.

- In case of suspected severe anaemia, measure haemoglobin and treat accordingly (see [Anaemia](#), Chapter 1).^b
- As soon as the patient is able to drink safely (often within 2 hours), provide ORS as the patient tolerates. ORS contains glucose and electrolytes which prevent development of complications.
- Monitor ongoing losses closely. Assess clinical condition and degree of dehydration at regular intervals to ensure continuation of appropriate treatment.

If over the course of treatment the patient:

- remains or becomes lethargic: measure blood glucose level and/or treat hypoglycaemia (see [Hypoglycaemia](#), Chapter 1).
- develops muscle cramps/weakness and abdominal distension: treat for moderate hypokalaemia with **7.5% potassium chloride** syrup (1 mmol of K⁺/ml) PO for 2 days:
Children under 45 kg: 2 mmol/kg (2 ml/kg) daily (according to weight, the daily dose is divided into 2 or 3 doses)
Children 45 kg and over and adults: 30 mmol (30 ml) 3 times daily
This treatment should only be given as an inpatient^c.
- develops peri-orbital or peripheral oedema: reduce the infusion rate to a minimum, auscultate the lungs, re-evaluate the stage of dehydration and the necessity of continuing IV rehydration. If IV rehydration is still required, continue the infusion at a slower rate and observe the patient closely. If IV rehydration is no longer required, change to oral treatment with ORS.
- develops dyspnoea, cough and bibasal crepitations are heard on auscultation of the lungs: sit the patient up, reduce the infusion rate to a minimum and administer one dose of **furosemide** IV (1 mg/kg in children; 40 mg in adults). Monitor the patient closely over 30 minutes and assess for underlying cardiorespiratory or renal disease.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Age	< 4 months	4 to 11 months	12 to 23 months	2 to 4 years	5 to 14 years	≥ 15 years
Weight	< 5 kg	5 to 7.9 kg	8 to 10.9 kg	11 to 15.9 kg	16 to 29.9 kg	≥ 30 kg
Quantity of ORS over 4 hours	200 to 400 ml	400 to 600 ml	600 to 800 ml	800 to 1200 ml	1200 to 2200 ml	2200 to 4000 ml

- Encourage additional age-appropriate fluid intake, including breastfeeding in young children. Give additional ORS after each loose stool (see below).
- Monitor ongoing losses closely. Assess clinical condition and degree of dehydration at regular intervals to ensure continuation of appropriate treatment.

No dehydration

Prevent dehydration:

- Encourage age-appropriate fluid intake, including breastfeeding in young children.
- Administer ORS according to WHO Treatment Plan A after any loose stool.

WHO Treatment Plan A^{[1][2]}

Age	Quantity of ORS
Children < 2 years	50 to 100 ml (<i>10 to 20 teaspoons</i>)
Children 2 to 10 years	100 to 200 ml (<i>½ to 1 glass</i>)
Children > 10 years and adults	at least 250 ml (<i>at least 1 glass</i>)

Treatment of diarrhoea

In addition to the WHO treatment plan corresponding to patient's degree of dehydration:

- Administer aetiological treatment if required.
- Administer zinc sulfate to children under 5 years (see [Acute diarrhoea](#), Chapter 3).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

References

1. World Health Organization. The treatment of diarrhoea : a manual for physicians and other senior health workers, 4th rev. World Health Organization. 2005.
<https://apps.who.int/iris/handle/10665/43209>
2. World Health Organization. Pocket book of Hospital Care for children. Guidelines for the Management of Common Childhood Illnesses. 2013.
https://apps.who.int/iris/bitstream/handle/10665/81170/9789241548373_eng.pdf?sequence=1

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Severe acute malnutrition

Last update: December 2022

Severe acute malnutrition (SAM) results from insufficient energy (kilocalories), fat, protein and/or other nutrients (vitamins and minerals, etc.) to cover individual needs.

SAM is frequently associated with medical complications due to metabolic disturbances and compromised immunity. It is a major cause of morbidity and mortality in children globally.

The protocols below are focused on the diagnosis and management of SAM in children 6 to 59 months only. For further details regarding this age group, and guidance for other age groups, refer to national recommendations and/or specialised protocols.

Clinical assessment

Characteristic physical signs

- In marasmus: skeletal appearance resulting from significant loss of muscle mass and subcutaneous fat.
- In kwashiorkor:
 - Bilateral oedema of the lower limbs sometimes extending to other parts of the body (e.g. arms and hands, face).
 - Discoloured, brittle hair; shiny skin which may crack, weep, and become infected.

Diagnostic and admission criteria

Diagnostic criteria for SAM are both anthropometric and clinical:

- Mid-upper arm circumference (MUAC)^a measures the degree of muscle wasting. MUAC < 115 mm indicates SAM and significant mortality risk.
- Weight-for-height z-score (WHZ) indicates the degree of weight loss by comparing the weight of the child with the median weight of non-malnourished children of the same height and sex. SAM is defined as WHZ < -3 with reference to the WHO Child Growth Standards^b.
- The presence of bilateral pitting oedema of the lower limbs (when other causes of oedema have been ruled out) indicates SAM, regardless of MUAC and WHZ.

Admission criteria for SAM treatment programmes vary with context. Refer to national recommendations.

Medical complications

- Children with any of the following severe medical conditions should receive hospital-based medical management:
 - Pitting oedema extending from the lower limbs up to the face;
 - Anorexia (observed during appetite test);

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Nutritional treatment is based on the use of specialised nutritious foods enriched with vitamins and minerals: F-75 and F-100 therapeutic milks, and ready-to-use therapeutic food (RUTF).
- Nutritional treatment is organised into phases:
 - Phase 1 (inpatient) intends to restore metabolic functions and treat or stabilize medical complications. Children receive F-75 therapeutic milk. This phase may last 1 to 7 days, after which children usually enter transition phase. Children with medical complications generally begin with phase 1.
 - Transition phase (inpatient) intends to ensure tolerance of increased food intake and continued improvement of clinical condition. Children receive F-100 therapeutic milk and/or RUTF. This phase usually lasts 1 to 3 days, after which children enter phase 2.
 - Phase 2 (outpatient or inpatient) intends to promote rapid weight gain and catch-up growth. Children receive RUTF. This phase usually lasts 1 to 3 days when inpatient, after which children are discharged for outpatient care. Children without medical complications enter directly into this phase as outpatients. The outpatient component usually lasts several weeks.
- Breastfeeding should be continued in breastfed children.
- Drinking water should be given in addition to meals, especially if the ambient temperature is high, or the child has a fever or is receiving RUTF.

Routine medical management

The following should be provided to all inpatients and outpatients with SAM:

Antibiotic treatment	From D1, unless specific signs of infection are present: amoxicillin PO: 50 mg/kg (max. 1 g) 2 times daily for 5 to 7 days
Malaria	On D1, rapid diagnostic test in endemic areas and treatment for malaria according to results or if testing is not available (see <u>Malaria</u> , Chapter 6).
Intestinal parasites	In transition phase or upon outpatient admission, albendazole PO: Children 12 to 23 months: 200 mg single dose Children 24 months and over: 400 mg single dose
Vaccination	<ul style="list-style-type: none"> • In transition phase or upon outpatient admission, measles vaccine for children 6 months to 5 years, unless a document shows that the child received 2 doses of vaccine administered as follows: one dose at or after 9 months and one dose at least 4 weeks after the first dose. Children vaccinated between 6 and 8 months should be re-vaccinated as above (i.e. with 2 doses) once they reach 9 months of age, provided that an interval of 4 weeks from the first dose is respected. • Other vaccines included in the EPI: check vaccination status and refer the child to vaccination services at discharge.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Management of complications

Infections

- Respiratory, cutaneous and urinary infections are common. However, classic signs of infection, such as fever, may be absent [1].
- Severe infection or sepsis should be suspected in children that are lethargic or apathetic or suffering from an acute complication such as hypothermia, hypoglycaemia, seizures, difficulty breathing, or shock. Immediately administer **ampicillin** IV 50 mg/kg every 8 hours + **gentamicin** IV 7.5 mg/kg once daily. Continue this treatment unless the source of infection is identified and different antibiotic treatment is required.
- If circulatory impairment or shock, immediately administer **ceftriaxone** IV, one dose of 80 mg/kg, then assess the source of infection to determine further antibiotic treatment. See also [Shock](#), Chapter 1. Transfuse urgently as for severe anaemia (see below) if haemoglobin (Hb) is < 6 g/dl.
- In less severe infections, assess the source of infection (see [Fever](#), Chapter 1) and treat accordingly.
- If fever is present and causes discomfort, undress the child. If insufficient, administer **paracetamol** PO in low dose: 10 mg/kg, up to 3 times maximum per 24 hours. Encourage oral fluids (including breast milk).
- If hypothermia is present, place the child skin-to-skin against the mother's body and cover with a warm blanket. Treat for infection as above. Check blood glucose level and treat for hypoglycaemia if necessary (see [Hypoglycaemia](#), Chapter 1).
- In children with kwashiorkor, infection of cutaneous lesions is common and may progress to soft tissue or systemic infection. If cutaneous infection is present, stop amoxicillin and start **amoxicillin/clavulanic acid** PO. Use formulations in a ratio of 8:1 or 7:1. The dose is expressed in amoxicillin: 50 mg/kg 2 times daily for 7 days.

Severe anaemia

- Children with Hb < 4 or < 6 with signs of decompensation such as respiratory distress require transfusion within the first 24 hours (preferably with packed red blood cells, PRBC).
- Transfusion volume is based on presence or absence of fever at time of ordering blood [2]:
 - No fever (< 37.5 °C): 30 ml/kg of whole blood or 15 ml/kg of PRBC over 4 hours
 - Fever present (≥ 37.5 °C): 20 ml/kg of whole blood or 10 ml/kg of PRBC over 4 hours
- Monitor closely for transfusion reactions and for signs of fluid overload (see [box](#) below).
- After the transfusion is completed: measure Hb at 8, 24 and 48 hours.

Diarrhoea and dehydration

- Diarrhoea is common. Therapeutic foods facilitate the recovery of physiological function of the gastrointestinal tract. Amoxicillin administered as part of routine treatment reduces intestinal bacterial overgrowth. Diarrhoea generally resolves without additional treatment. If an aetiological treatment is necessary, see [Acute diarrhoea](#), Chapter 3.
- Zinc supplementation is not needed if children consume recommended amounts of therapeutic foods.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Clinical features (2 or more of the following signs)	No dehydration	Some dehydration	Severe dehydration
Mental status	Normal	Restless, irritability	Lethargic or unconscious
Thirst	No thirst, drinks normally	Thirsty, drinks eagerly	Unable to drink or drinks poorly
Urine output	Normal	Reduced	Absent for several hours
Recent frequent watery diarrhoea and/or vomiting	Yes	Yes	Yes
Recent obvious rapid weight loss	No	Yes	Yes

Acute diarrhoea with no dehydration (Plan A SAM)

- Stools are neither frequent nor abundant (outpatient): **oral rehydration solution (ORS)** PO: 5 ml/kg after each loose stool to prevent dehydration.
- Stools are frequent and/or abundant (inpatient): **ReSoMal^c** PO or by nasogastric tube (NGT): 5 ml/kg after each loose stool to prevent dehydration.
- In all cases, continue feeding and breastfeeding, encourage oral fluids.

Acute diarrhoea with some dehydration (Plan B SAM)

- Determine the target weight (weight before the onset of diarrhoea) before starting rehydration. If not feasible (e.g. new admission), estimate target weight as current weight x 1.06.
- **ReSoMal^c** PO or by NGT: 20 ml/kg/hour for 2 hours. In addition, administer 5 ml/kg of **ReSoMal** after each loose stool if tolerated.
- Assess after 2 hours (clinical evaluation and weight):
 - If improvement (diarrhoea and signs of dehydration regress):
 - ▷ Reduce **ReSoMal** to 10 ml/kg/hour until the signs of dehydration and/or weight loss (known or estimated) have been corrected.
 - ▷ Assess every 2 hours.
 - ▷ Once there are no signs of dehydration and/or the target weight is reached, change to Plan A SAM to prevent dehydration.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Measure blood glucose level and treat [hypoglycaemia](#) (Chapter 1) if necessary.
- Monitor vital signs and signs of dehydration every 15 to 30 minutes.
- Monitor urine output.
- Monitor for signs of fluid overload (see [box](#) below).
- If there is no circulatory impairment:
 - **ReSoMal** PO or by NGT: 20 ml/kg over 1 hour
 - If the child is alert, continue feeding including breastfeeding.
 - Assess after 1 hour:
 - ▷ If improvement: change to Plan B SAM, but keep the same target weight.
 - ▷ If rehydration PO/NGT not tolerated (e. g. vomiting):
 - ▷ Stop ReSoMal. Administer **glucose 5%-Ringer lactate (G5%-RL)**^d IV infusion: 10 ml/kg/hour for 2 hours.
 - ▷ Assess after 2 hours of IV fluids:
 - ▷ If improvement and/or not vomiting, stop G5%-RL IV infusion and change to Plan B SAM.
 - ▷ If no improvement or still vomiting, continue G5%-RL IV infusion: 10 ml/kg/hour for 2 hours.
 - ▷ If deterioration with circulatory impairment: see below.
 - If there is circulatory impairment:
 - Stabilize (see [Shock](#), Chapter 1).
 - Administer **ceftriaxone** IV, one dose of 80 mg/kg. Subsequent antibiotic treatment depends on assessment of underlying cause.
 - Administer **G5%-RL** IV infusion: 10 ml/kg/hour for 2 hours. Stop ReSoMal if the child was taking it.
 - Assess after 1 hour of IV fluids:
 - ▷ If improvement and no vomiting: stop IV fluid and change to Plan B SAM, but keep the same target weight.
 - ▷ If no improvement:
 - ▷ Continue **G5%-RL** IV infusion: 10 ml/kg/hour.
 - ▷ Prepare for blood transfusion.
 - Assess after 2 hours of IV fluids:
 - ▷ If improvement: change to Plan B SAM, but keep the same target weight.
 - ▷ If no improvement or deterioration:
 - ▷ Check Hb as baseline and administer whole blood: 10 ml/kg over at least 3 hours^[3] (max. 4 hours) using a separate IV line.
 - ▷ While transfusing, continue G5%-RL IV infusion 10 ml/kg/hour for another 2 hours.



During rehydration, monitor continuously for signs of fluid overload. These include:

- RR ≥ 10 breaths/minute compared to initial RR, or
- HR ≥ 20 beats/minute compared to initial HR

Plus any one of the following:

- New or worsening hypoxia (decrease in SpO₂ by > 5%)
- New onset of rales and/or fine crackles in lung fields
- New galloping heart rhythm
- Increased liver size (must have marked liver border with pen before rehydration)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Discharge criteria

In general:

- Children can be discharged from hospital and be treated as outpatients if the following criteria are met:
 - clinically well;
 - medical complications controlled;
 - able to eat RUTF (observed during appetite test);
 - reduction or absence of oedema;
 - caregiver feels able to provide care as outpatient;
 - vaccinations up to date or referral to vaccination service organised.
- Children can be discharged from nutritional treatment if the following criteria are met:
 - co-existing medical conditions stable and outpatient treatment organised if necessary (e.g. dressing changes, follow-up for chronic diseases);
 - vaccinations up to date or referral to vaccination service organised;
 - absence of oedema and WHZ > -2 or MUAC > 125 mm for at least 2 weeks.

Discharge criteria vary with context. Refer to national recommendations.

Footnotes

- (a) MUAC is measured at the mid-point of the left upper arm. The arm should be relaxed. The measuring tape should be in contact with the skin all around the arm, without exerting pressure.
- (b) For WHZ, see WHO simplified field tables in z-scores for girls and for boys:
<https://www.who.int/tools/child-growth-standards/standards/weight-for-length-height>
- (c) ReSoMal is a specific oral rehydration solution for malnourished children, containing less sodium and more potassium than standard ORS. It should be administered under medical supervision to avoid overdosing and hyponatremia.
- (d) Remove 50 ml of Ringer lactate (RL) from a 500 ml RL bottle or bag, then add 50 ml of 5% glucose to the remaining 450 ml of RL to obtain 500 ml of 5% glucose-RL solution.

References

1. Jones KDJ, Berkley JA. Severe acute malnutrition and infection. *Paediatrics and International Child Health* 2014; 34(sup1): S1-S29.
<https://www.tandfonline.com/doi/full/10.1179/2046904714Z.000000000218> [Accessed 24 August 2022]
2. Maitland K, Olupot-Olupot P, Kiguli S, Chagaluka G, Alaroker F, Opoka RO, Mpoya A, Engoru C, Nteziyaremye J, Mallewa M, Kennedy N, Nakuya M, Namayanja C, Kayaga J, Uyoga S, Kyeyune Byabazaire D, M'baya B, Wabwire B, Frost G, Bates I, Evans JA, Williams TN, Saramago Goncalves P, George EC, Gibb DM, Walker AS; TRACT Group. Transfusion Volume for Children with Severe Anemia in Africa. *N Engl J Med*. 2019 Aug 1;381(5):420-431.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6830210/>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 2: Respiratory diseases

[Acute upper airway obstruction](#)

[Rhinitis and rhinopharyngitis \(common cold\)](#)

[Acute sinusitis](#)

[Acute pharyngitis](#)

[Diphtheria](#)

[Other upper respiratory tract infections](#)

[Laryngotracheitis and laryngotracheobronchitis \(croup\)](#)

[Epiglottitis](#)

[Bacterial tracheitis](#)

[Otitis](#)

[Acute otitis externa](#)

[Acute otitis media \(AOM\)](#)

[Chronic suppurative otitis media \(CSOM\)](#)

[Whooping cough \(pertussis\)](#)

[Bronchitis](#)

[Acute bronchitis](#)

[Chronic bronchitis](#)

[Bronchiolitis](#)

[Acute pneumonia](#)

[Pneumonia in children under 5 years of age](#)

[Pneumonia in children over 5 years and adults](#)

[Persistent pneumonia](#)

[Staphylococcal pneumonia](#)

[Asthma](#)

[Asthma attack \(acute asthma\)](#)

[Chronic asthma](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Acute upper airway obstruction

Acute upper airway obstruction can be caused by foreign body aspiration, viral or bacterial infections (croup, epiglottitis, tracheitis), anaphylaxis, burns or trauma.

Initially stable and partial obstruction may worsen and develop into a life-threatening emergency, especially in young children.

Clinical features

Clinical signs of the severity of obstruction:

Obstruction	Signs	Danger signs
Complete	<ul style="list-style-type: none">Respiratory distress followed by cardiac arrest	
Imminent complete	<ul style="list-style-type: none">Severe respiratory distress with cyanosis or $\text{SpO}_2 < 90\%$Agitation or lethargyTachycardia, capillary refill time > 3 seconds	
Severe	<ul style="list-style-type: none">Stridor (abnormal high pitched sound on inspiration) at restSevere respiratory distress:<ul style="list-style-type: none">Severe intercostal and subcostal retractionsNasal flaringSubsternal retractions (inward movement of the breastbone during inspiration)Severe tachypnoea	Yes
Moderate	<ul style="list-style-type: none">Stridor with agitationModerate respiratory distress:<ul style="list-style-type: none">Mild intercostal and subcostal retractionsModerate tachypnoea	No
Mild	<ul style="list-style-type: none">Cough, hoarse voice, no respiratory distress	

Management in all cases

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Management of foreign body aspiration

Acute airway obstruction (the foreign body either completely obstructs the pharynx or acts as a valve on the laryngeal inlet), no warning signs, most frequently in a child 6 months-5 years playing with a small object or eating. Conscience is initially maintained.

Perform maneuvers to relieve obstruction only if the patient cannot speak or cough or emit any sound:

- Children over 1 year and adults:

Heimlich manoeuvre: stand behind the patient. Place a closed fist in the pit of the stomach, above the navel and below the ribs. Place the other hand over fist and press hard into the abdomen with a quick, upward thrust. Perform one to five abdominal thrusts in order to compress the lungs from the below and dislodge the foreign body.

- Children under 1 year:

Place the infant face down across the forearm (resting the forearm on the leg) and support the infant's head with the hand. With the heel of the other hand, perform one to five slaps on the back, between shoulder plates.

If unsuccessful, turn the infant on their back. Perform five forceful sternal compressions as in cardiopulmonary resuscitation: use 2 or 3 fingers in the center of the chest just below the nipples. Press down approximately one-third the depth of the chest (about 3 to 4 cm).

Repeat until the foreign body is expelled and the patient resumes spontaneous breathing (coughing, crying, talking). If the patient loses consciousness ventilate and perform cardiopulmonary resuscitation. Tracheostomy if unable to ventilate.

Differential diagnosis and management of airway obstructions of infectious origin

Infections	Symptoms	Appearance	Timing of symptoms
Viral croup	Stridor, cough and moderate respiratory difficulty	Prefers to sit	Progressive
Epiglottitis	Stridor, high fever and severe respiratory distress	Prefers to sit, drooling (cannot swallow their own saliva)	Rapid
Bacterial tracheitis	Stridor, fever, purulent secretions and severe respiratory distress	Prefers to lie flat	Progressive
Retropharyngeal or tonsillar abscess	Fever, sore throat and painful swallowing, earache, trismus and hot potato voice	Prefers to sit, drooling	Progressive

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Footnotes

- (a) If possible it is better to treat all patients with a SpO₂ < 95% with oxygen.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Rhinitis and rhinopharyngitis (common cold)

Rhinitis (inflammation of the nasal mucosa) and rhinopharyngitis (inflammation of the nasal and pharyngeal mucosa) are generally benign, self-limited and most often of viral origin. However, they may be an early sign of another infection (e.g. measles or influenza) or may be complicated by a bacterial infection (e.g. otitis media or sinusitis).

Clinical features

- Nasal discharge or obstruction, which may be accompanied by sore throat, fever, cough, lacrimation, and diarrhoea in infants. Purulent nasal discharge is not indicative of a secondary bacterial infection.
- In children under 5 years, routinely check the tympanic membranes to look for an associated otitis media.

Treatment

- Antibiotherapy is not recommended: it does not promote recovery nor prevent complications.
- Treatment is symptomatic:
 - Clear the nose with 0.9% sodium chloride^a.
 - Fever, throat soreness: paracetamol PO for 2 to 3 days ([Fever](#), Chapter 1).

Footnotes

- (a) For a child: place him on his back, head turned to the side, and instil 0.9% sodium chloride into each nostril.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute sinusitis

Acute sinusitis is an inflammation of one or more of the sinus cavities, caused by an infection or allergy. Most acute sinus infections are viral and resolve spontaneously in less than 10 days. Treatment is symptomatic. Acute bacterial sinusitis may be a primary infection, a complication of viral sinusitis or of dental origin. The principal causative organisms are *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis*. It is essential to distinguish between bacterial sinusitis and common rhinopharyngitis (see [Rhinitis and rhinopharyngitis](#)). Antibiotic therapy is required in case of bacterial sinusitis only. Without treatment, severe sinusitis in children may cause serious complications due to the spread of infection to the neighbouring bony structures, orbits or the meninges.

Clinical features

Sinusitis in adults

- Purulent unilateral or bilateral discharge, nasal obstruction and
- Facial unilateral or bilateral pain that increases when bending over; painful pressure in maxillary area or behind the forehead.
- Fever is usually mild or absent.

Sinusitis is likely if symptoms persist for longer than 10 to 14 days or worsen after 5 to 7 days or are severe (severe pain, high fever, deterioration of the general condition).

Sinusitis in children

- Same symptoms; in addition, irritability or lethargy or cough or vomiting may be present.
- In the event of severe infection: deterioration of the general condition, fever over 39 °C, periorbital or facial oedema.

Treatment

Symptomatic treatment

- [Fever](#) and [pain](#) (Chapter 1).
- Clear the nose with 0.9% sodium chloride^a.

Antibiotherapy

- In adults:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Children: 30 mg/kg 3 times daily (max. 3 g daily)
 Adults: 1 g 3 times daily
- In the event of failure to respond within 48 hours of therapy:
amoxicillin/clavulanic acid PO for 7 to 10 days. Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin:
 - Children < 40 kg: 25 mg/kg 2 times daily
 - Children ≥ 40 kg and adults:
 - Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)
 - Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)
 - In penicillin-allergic patients:
erythromycin PO for 7 to 10 days:
 - Children: 30 to 50 mg/kg daily^b
 - Adults: 1 g 2 to 3 times daily
- In infants with ethmoiditis, see [Periorbital and orbital cellulitis](#) (Chapter 5).

Other treatments

- For sinusitis secondary to dental infection: dental extraction while under antibiotic treatment.
- In the event of ophthalmologic complications (ophthalmoplegia, mydriasis, reduced visual acuity, corneal anesthesia), refer for surgical drainage.

Footnotes

- (a) For a child: place him on his back, head turned to the side, and instil 0.9% sodium chloride into each nostril.
- (b) For dosage according to age or weight, see [erythromycin](#) in the guide Essential drugs, MSF.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Acute pharyngitis

Last updated: November 2020

Acute inflammation of the tonsils and pharynx. The majority of cases are of viral origin and do not require antibiotic treatment. Group A streptococcus (GAS) is the main bacterial cause, and mainly affects children aged 3 to 14 years. Acute rheumatic fever (ARF), a serious late complication of GAS pharyngitis, can be prevented with antibiotic treatment.

One of the main objectives of assessing acute pharyngitis is to identify patients requiring antibiotic treatment.

Clinical features

- Features common to all types of pharyngitis: throat pain, dysphagia (difficulty swallowing), inflammation of the tonsils and pharynx, tender anterior cervical lymph nodes, with or without fever.
- Specific features, depending on the cause:

Common forms:

- **Erythematous (red throat) or exudative (red throat and whitish exudate) pharyngitis:** this appearance is common to both viral and GAS pharyngitis. Centor criteria help assessment and decrease the empirical use of antibiotics in settings where rapid testing for GAS is not available. A Centor score of less than 2 rules out GAS infection^{[1][2]}. Nevertheless, in patients with risk factors (immunosuppression, personal or family history of ARF) for poststreptococcal complications, or for local or general complications, do not use Centor score and prescribe empirical antibiotic treatment.

Centor criteria

Criteria	Score
Temperature > 38 °C	1
Absence of cough	1
Tender anterior cervical lymph node(s)	1
Tonsillar swelling or exudate	1

In patients over 14 years, the probability of GAS pharyngitis is low. Infectious mononucleosis (IM) due to the Epstein-Barr virus should be suspected in adolescents and young adults with extreme fatigue, generalized adenopathy and often splenomegaly.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Spots on oral mucosa (Koplik's spots) accompanied by conjunctivitis and skin rash (see [Measles](#), Chapter 8).
- "Strawberry" (red and bumpy) tongue accompanied by a skin rash: scarlet fever caused by GAS.
- Local complications:
Peritonsillar, retropharyngeal or lateral pharyngeal abscess: fever, intense pain, dysphagia, hoarse voice, trismus (limitation of mouth opening), unilateral deviation of the uvula.
- General complications:
 - Complications due to the toxin: diphtheria (see [Diphtheria](#), Chapter 2).
 - Poststreptococcal complications: ARF, acute glomerulonephritis.
 - Signs of serious illness in children: severe dehydration, severe difficulty swallowing, upper airway compromise, deterioration of general condition.
- Differential diagnosis: epiglottitis (see [Epiglottitis](#), Chapter 2).

Treatment

- Symptomatic treatment (fever and pain): paracetamol or ibuprofen PO ([Fever](#), Chapter 1).
- Centor score ≤ 1 : viral pharyngitis, which typically resolves within a few days (or weeks, for IM): no antibiotic treatment.
- Centor score ≥ 2 or scarlet fever: antibiotic treatment for GAS infections ^[3] :
 - If single-use injection equipment is available, benzathine benzylpenicillin is the drug of choice as streptococcus A resistance to penicillin remains rare; it is the only antibiotic proven effective in reducing the incidence of rheumatic fever; and the treatment is administered as a single dose.
benzathine benzylpenicillin IM
Children under 30 kg (or under 10 years): 600 000 IU single dose
Children 30 kg and over (or 10 years and over) and adults: 1.2 MIU single dose
 - Penicillin V is the oral reference treatment, but poor adherence is predictable due to the length of treatment.
phenoxymethylenicillin (penicillin V) PO for 10 days
Children 1 to < 6 years: 250 mg 2 times daily
Children 6 to < 12 years: 500 mg 2 times daily
Children 12 years and over and adults: 1 g 2 times daily
Children under 1 year: 125 mg 2 times daily
 - Amoxicillin is an alternative and the treatment has the advantage of being relatively short. However, it can cause adverse skin reactions in patients with undiagnosed IM and thus should be avoided when IM has not been excluded
amoxicillin PO for 6 days
Children: 25 mg/kg 2 times daily
Adults: 1 g 2 times daily
 - Macrolides should be reserved for penicillin allergic patients as resistance to macrolides is frequent and their

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- If signs of serious illness or epiglottitis are present in children: hospitalise.

References

1. Fine AM, Nizet V, Mandl KD. Large-scale validation of the Centor and McIsaac scores to predict group A streptococcal pharyngitis. *Arch Intern Med.* 2012;172(11):847-852.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3627733/> [Accessed 20 October 2020]
2. National Institute for Health and Care Excellence. Sore throat (acute): antimicrobial prescribing. 2018.
<http://www.nice.org.uk/ng84> [Accessed 20 October 2020]
3. Group A Streptococcal Disease, Centers for Disease Control and Prevention. Atlanta (GA): CDC; 2020.
<https://www.cdc.gov/groupastrep/diseases-hcp/strep-throat.html> [Accessed 20 October 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Diphtheria

Last updated: October 2022

Diphtheria is a bacterial infection due to *Corynebacterium diphtheriae*, spread from person to person through inhalation of infected respiratory droplets of symptomatic or asymptomatic individuals, or direct contact with contaminated objects or diphtheria skin lesions^{[1][2]a}.

After infection, *C. diphtheriae* has an incubation period of 1 to 5 days (max. 10 days)^[1] during which time it multiplies in the upper respiratory tract. The bacteria secretes a toxin which causes severe local as well as systemic effects. Death can occur from airway obstruction or as a result of systemic complications, including damage to the myocardium and nervous system, caused by the toxin.

Cases can remain infectious up to 8 weeks after initial infection^[2]. Antibiotic treatment can reduce infectiousness to 6 days^[3].

Vaccination is the key to prevention and control of diphtheria. It protects individuals from severe disease (fewer and less severe symptoms) but does not prevent the spread of *C. diphtheriae*. Clinical disease does not confer protective immunity and vaccination is an integral part of case management.

Clinical features

- During clinical examination respect standard, contact, and droplet precautions (handwashing, gloves, gown, mask, etc.). Conduct a careful examination of the throat.
- Signs of respiratory diphtheria^a :
 - pharyngitis, rhinopharyngitis, tonsillitis or laryngitis with tough, greyish, firmly adherent pseudo-membranes of the pharynx, nasopharynx, tonsils, or larynx;
 - dysphagia and cervical adenitis, at times progressing to massive swelling of the neck;
 - airway obstruction and possible suffocation when the infection extends to the nasal passages, larynx, trachea and bronchi;
 - fever is generally low-grade^[2].
- Generalised signs due to effects of the toxin:
 - cardiac dysfunction (tachycardia, arrhythmias), severe myocarditis with heart failure and possibly cardiogenic shock (see [Shock](#), Chapter 1) 3 to 7 days or 2 to 3 weeks after onset of the disease;
 - neuropathies in 2 to 8 weeks after the onset of disease leading to nasal voice and difficulty with swallowing (paralysis of the soft palate), vision (ocular motor paralysis), breathing (paralysis of respiratory muscles) and ambulation (limb paralysis);
 - oliguria, anuria and acute renal failure.
- Differential diagnoses: [Epiglottitis](#) and [Acute pharyngitis](#), Chapter 2, [Stomatitis](#), Chapter 3.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- **Diphtheria antitoxin (DAT)^b** derived from horse serum:

Administer DAT as soon as possible after disease onset. Do not wait for bacteriological confirmation^[1]; administer DAT under close monitoring in a hospital setting, according to the Besredka method to assess possibility of allergy. Any delay can diminish efficacy.



There is a risk of anaphylactic reaction, especially in patients with asthma. Close monitoring of the patient is essential, with immediate availability of equipment for manual ventilation (Ambu bag, face mask) and intubation, Ringer lactate and epinephrine (see [Shock](#), Chapter 1).

Besredka method: inject 0.1 ml SC and wait 15 minutes. If there is no allergic reaction (no erythema at the injection site or a flat erythema of less than 0.5 cm in diameter), inject a further 0.25 ml SC. If there is no reaction after 15 minutes, inject the rest of the product IM or IV depending on the volume to be administered.

Doses are given as a function of the severity of illness, and the delay in treatment:

Clinical signs	Dose in units	Administration route
Laryngitis or pharyngitis or duration < 48 hours	20 to 40 000	
Rhinopharyngitis	40 to 60 000	IM or IV infusion in 250 ml of 0.9% sodium chloride in 2 to 4 hours for doses of more than 20 000 units.
Severe disease (respiratory distress, shock), cervical oedema or duration ≥ 48 hours	80 to 100 000	

- Antibiotic treatment (as soon as possible without waiting for bacteriological confirmation) for 14 days or according to length of treatment recommended by the national protocol:

- if the patient can swallow:

azithromycin PO (first-line)

Children: 10 to 12 mg/kg once daily (max. 500 mg daily)

Adults: 500 mg once daily

or

erythromycin PO

Children under 40 kg: 10 to 15 mg/kg (max. 500 mg) 4 times daily

Children 40 kg and over and adults: 500 mg 4 times daily

or

phenoxycephalothin (penicillin V) PO

Children under 40 kg: 10 to 15 mg/kg (max. 500 mg) 4 times daily

Children 40 kg and over and adults: 500 mg 4 times daily

- If the patient cannot swallow, start with one of the treatments below and change as soon as possible to oral route with one of the oral treatments above to complete 14 days of treatment:

~~metronidazole~~

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Management of close contacts

Close contacts include household members living under the same roof and people who were directly exposed (less than one metre) to nasopharyngeal secretions of the patient on a regular basis (e.g. family or close friends, children in the same class, medical personnel) during the 5 days or nights prior to onset of symptoms of the case^[4].

- Collect nasal and pharyngeal swabs for culture before starting antibiotic prophylaxis; temperature and throat examination daily (10 days); exclusion from school or work until 48 hours after starting antibiotic prophylaxis. If symptoms of respiratory infection appear: treat immediately as a case of diphtheria.
- Antibiotic prophylaxis:

benzathine benzylpenicillin IM

Children under 30 kg: 600 000 IU single dose

Children 30 kg and over and adults: 1.2 MIU single dose



Benzathine benzylpenicillin should never be administered by IV route.

or azithromycin PO or erythromycin PO as above for 7 days.

- Check vaccination status:
 - if less than 3 injections received: complete vaccination schedule (see [Prevention](#) below);
 - if 3 injections received, with the last injection over one year ago: administer a booster dose immediately;
 - if 3 injections received, with the last injection less than one year ago: a booster dose is not immediately necessary.

Outbreak surveillance measures

- A suspected case of diphtheria is defined as a person with:
 - pharyngitis, rhinopharyngitis, tonsillitis and/or laryngitisAND
 - an adherent pseudo-membrane of the pharynx, nose, tonsils and/or larynx^[1].
- Isolate and treat suspect cases without delay. Collect swab samples before starting antibiotic treatment. Submit case notification to the public health authorities within 24 hours^[1].

Prevention

- Routine vaccination (EPI), for information: 3 doses of conjugate vaccine containing the higher potency (D) formulation of diphtheria toxoid as soon as possible as of 6 weeks of age and at 4 week intervals; D booster between 12 and 23 months, then between 4 and 7 years; booster with a vaccine containing a reduced dose (d) of diphtheria toxoid between 9 and 15 years^[5].
- Catch-up vaccination (individuals who have not received routine vaccination), for information:
 - children 1 to 6 years: 3 doses of conjugate vaccine containing the higher potency (D) formulation of diphtheria toxoid at least 4 weeks apart;
 - children 7 years and over and adults (including medical staff): 3 doses of conjugate vaccine containing a reduced

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

(b) DAT reduces mortality and should be given to all diphtheria patients. However, as supply is very limited, it may be necessary to define criteria and reserve DAT for the treatment of patients who will benefit the most from it. DAT can be administered to pregnant women.

(c) **erythromycin IV infusion (60 minutes)**

Children: 12.5 mg/kg every 6 hours (max. 2 g daily); adults: 500 mg every 6 hours

Erythromycin powder (1 g) should be reconstituted in 20 ml of water for injection only. Then, dilute each dose of erythromycin in 10 ml/kg of 0.9% sodium chloride in children less than 20 kg and in a bag of 250 ml of 0.9% sodium chloride in children 20 kg and over and in adults. Do not dilute in glucose.

References

1. World Health Organization. Diphtheria. Vaccine-Preventable Diseases Surveillance Standards. 2018. https://www.who.int/immunization/monitoring_surveillance/burden/vpd/WHO_SurveillanceVaccinePreventable_04_Diphtheria_R2.pdf?ua=1 [Accessed 11 August 2020]
2. Tiwari TSP, Wharton M. Chapter 19: Diphtheria Toxoid. In: Plotkin SA, Orenstein WA, Offit PA, editors. Vaccines. 7th ed. Philadelphia, PA: Elsevier; 2018. p. 261–275.
3. Truelove SA, Keegan LT, Moss WJ, Chaisson LH, Macher E, Azman AS, Lessler J. Clinical and Epidemiological Aspects of Diphtheria: A Systematic Review and Pooled Analysis. Clin Infect Dis. 2020 Jun 24;71(1):89–97. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7312233/> [Accessed 24 November 2020]
4. Pan American Health Organization, World Health Organization. Diphtheria in the Americas - Summary of the situation 2018. Epidemiological Update Diphtheria. 16 April 2018. https://www.paho.org/hq/index.php?option=com_docman&view=download&category_slug=diphtheria-098968&alias=44497-16-april-2018-diphtheria-epidemiological-update-497&Itemid=270&lang=en [Accessed 11 August 2020]
5. World Health Organization. Diphtheria vaccine: WHO position paper - August 2017. Weekly epidemiological record 2017; 92(31):417–436. https://www.who.int/immunization/policy/position_papers/wer_31_diphtheria_updated_position_paper.pdf?ua=1 [Accessed 11 August 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Other upper respiratory tract infections

- [Laryngotracheitis and laryngotracheobronchitis \(croup\)](#)
- [Epiglottitis](#)
- [Bacterial tracheitis](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Laryngotracheitis and laryngotracheobronchitis (croup)

Viral infection in children aged 3 months to 4 years.

Clinical features

- Typical barking cough, hoarse voice or cry.
- Inspiratory stridor (abnormal high pitched sound on inspiration):
 - Croup is considered mild or moderate if the stridor only occurs with agitation;
 - Croup is considered severe if there is stridor at rest, especially when it is accompanied by respiratory distress.
- Wheezing may also be present if the bronchi are involved.

Treatment

- In the absence of inspiratory stridor or retractions, treat symptomatically: ensure adequate hydration, seek medical attention if symptoms worsen (e.g. respiratory difficulty, noisy breathing and inability to tolerate oral fluids).
- If stridor is only present with agitation (moderate croup):
 - Hospitalize for treatment and observation (risk of worsening).
 - Assure adequate hydration.
 - **dexamethasone^a** PO (use IV preparation flavoured with sugar water, 10% or 50% glucose or juice) or IM if child is vomiting: 0.6 mg/kg single dose (see [table](#)).
- If danger signs are present (stridor at rest, respiratory distress), admit in intensive care:
 - Oxygen continuously: at least 5 litres/minute or to maintain the SpO₂ between 94 and 98%.
 - Insert a peripheral IV line and provide IV hydration.
 - **epinephrine** (adrenaline) via nebulizer (1 mg/ml, 1 ml ampoule): 0.5 mg/kg (max. 5 mg) to be repeated every 20 minutes if danger signs persist.
Monitor heart rate during nebulization (if heart rate greater than 200, stop the nebulization).

Age	3 months	4-6 months	7-9 months	10-11 months	1-4 years
Weight	6 kg	7 kg	8 kg	9 kg	10-17 kg
Dose in mg	3 mg	3.5 mg	4 mg	4.5 mg	5 mg
Dose in ml	3 ml	3.5 ml	4 ml	4.5 ml	5 ml

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Age	3-11 months	1-2 years	3-4 years
Weight	6-9 kg	10-13 kg	14-17 kg
Dose in mg	4 mg	8 mg	10 mg
Dose in ml	1 ml	2 ml	2.5 ml

Suspect bacterial tracheitis in a critically ill appearing child^b with croup who does not improve with the above treatment.

- If wheezing is present:
salbutamol aerosol (using a spacer): 2 to 3 puffs every 20 to 30 minutes as needed
- If the patient has a complete airway obstruction, intubation if possible or emergency tracheotomy.

Footnotes

- (a) Administer orally if possible in order to avoid causing agitation in the child as this may worsen symptoms.
- (b) Critically ill appearing child: weak grunting or crying, drowsiness, difficult to arouse, does not smile, unconjugate or anxious gaze, pallor or cyanosis, general hypotonia.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Epiglottitis

Bacterial infection of the epiglottis in young children caused by *Haemophilus influenzae* (Hib), it is rare when Hib vaccine coverage is high. It can be caused by other bacteria and occur in adults.

Clinical features

- Rapid (less than 12-24 hours) onset of high fever.
- Typical “tripod or sniffing” position, preferring to sit, leaning forward with an open mouth, anxious appearing.
- Difficulty swallowing, drooling, and respiratory distress.
- Stridor may be present (as opposed to croup, hoarse voice and cough are usually absent).
- Critically ill appearing^a.



Allow the child to sit in a comfortable position or on the parent’s lap. Do not force them to lie down (may precipitate airway obstruction). Avoid any examination that will upset the child including examination of the mouth and throat.

Treatment

- In case of imminent airway obstruction, emergency intubation or tracheotomy is indicated. The intubation is technically difficult and should be performed under anaesthesia by a physician familiar with the procedure. Be prepared to perform a tracheotomy if intubation is unsuccessful.
- In all cases:
 - Insert a peripheral IV line and provide IV hydration.
 - Antibiotherapy:
ceftriaxone slow IV (3 minutes) or IV infusion (30 minutes)^b. Avoid IM route (may agitate the child and precipitate a respiratory arrest).
Children: 50 mg/kg once daily
Adults: 1 g once daily
The IV treatment is administered for at least 5 days then, if the clinical condition has improved^c and oral treatment can be tolerated, change to:
amoxicillin/clavulanic acid (co-amoxiclav) PO to complete a total of 7 to 10 days of treatment. Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin:
Children < 40 kg: 50 mg/kg 2 times daily
Children ≥ 40 kg and adult:
Ratio 8:1: 3000 mg daily (2 tablets of 500/62.5 mg 3 times daily)
Ratio 7:1: 2625 mg daily (1 tablet of 875/125 mg 3 times daily)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Bacterial tracheitis

Bacterial infection of the trachea in children, occurring as a complication of a previous viral infection (croup, influenza, measles, etc.).

Clinical features

- Fever in a critically ill appearing child^a.
- Stridor, cough and respiratory distress.
- Copious purulent secretions.
- As opposed to epiglottitis the onset of symptoms is gradual and the child prefers to lie flat.
- In severe cases there is a risk of complete airway obstruction, especially in very young children.

Treatment

- Suction purulent secretions.
- Insert a peripheral IV line and provide IV hydration.

- Antibiotherapy:

ceftriaxone slow IV^b (3 minutes) or IV infusion (30 minutes). Do not administer by IM route (may agitate the child and precipitate a respiratory arrest).

Children: 50 mg/kg once daily

Adults: 1 g once daily

+

cloxacillin IV infusion (60 minutes)

Children less than 12 years: 25 to 50 mg/kg every 6 hours

Children 12 years and over and adults: 2 g every 6 hours

The IV treatment is administered for at least 5 days then, if the clinical condition has improved^c and oral treatment can be tolerated, change to :

amoxicillin/clavulanic acid (co-amoxiclav) PO to complete 7 to 10 days of treatment, as in epiglottitis.

- If the event of complete airway obstruction, intubation if possible or emergency tracheotomy.

Footnotes

(a) Critically ill appearing child: weak grunting or crying, drowsiness, difficult to arouse, does not smile, unconjugate or anxious gaze, pallor or cyanosis, general hypotonia.

(b) For administration by IV route, ceftriaxone powder should be reconstituted in water for injection only. For administration by IV infusion, dilute each dose of ceftriaxone in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Otitis

- [Acute otitis externa](#)
- [Acute otitis media \(AOM\)](#)
- [Chronic suppurative otitis media \(CSOM\)](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Acute otitis externa

Diffuse inflammation of the external ear canal, due to bacterial or fungal infection. Common precipitants of otitis externa are maceration, trauma of the ear canal or presence of a foreign body or dermatologic diseases (such as eczema, psoriasis).

Clinical features

- Ear canal pruritus or ear pain, often severe and exacerbated by motion of the pinna; feeling of fullness in the ear; clear or purulent ear discharge or no discharge
- Otoscopy (remove skin debris and secretions from the auditory canal by gentle dry mopping (use a dry cotton bud or a small piece of dry cotton wool):
 - diffuse erythema and edema, or infected eczema, of the ear canal
 - look for a foreign body
 - if visible, the tympanic membrane is normal (swelling and pain very often prevent adequate visualization of the tympanic membrane)

Treatment

- Remove a foreign body, if present.
- Treatment of pain: paracetamol PO (Chapter 1, [Pain](#)).
- Local treatment:
 - Remove secretions from the auditory canal by gentle dry mopping (use a dry cotton bud or a small piece of dry cotton wool). Consider ear irrigation (0.9% sodium chloride, using a syringe) only if the tympanic membrane can be fully visualised and is intact (no perforation). Otherwise, ear irrigation is contra-indicated.
 - Apply **ciprofloxacin** ear drops in the affected ear(s) for 7 days:
Children \geq 1 year: 3 drops 2 times daily
Adults: 4 drops 2 times daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute otitis media (AOM)

Acute inflammation of the middle ear, due to viral or bacterial infection, very common in children under 3 years, but uncommon in adults.

The principal causative organisms of bacterial otitis media are *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis* and in older children, *Streptococcus pyogenes*.

Clinical features

- Rapid onset of ear pain (in infants: crying, irritability, sleeplessness, reluctance to nurse) and ear discharge (otorrhoea) or fever.
- Other signs such as rhinorrhoea, cough, diarrhoea or vomiting are frequently associated, and may confuse the diagnosis, hence the necessity of examining the tympanic membranes.
- Otoscopy: bright red tympanic membrane (or yellowish if rupture is imminent) and presence of pus, either externalised (drainage in ear canal if the tympanic membrane is ruptured) or internalised (opaque or bulging tympanic membrane). The combination of these signs with ear pain or fever confirms the diagnosis of AOM.

Note:

The following otoscopic findings are not sufficient to make the diagnosis of AOM:

- A red tympanic membrane alone, with no evidence of bulging or perforation, is suggestive of viral otitis in a context of upper respiratory tract infection, or may be due to prolonged crying in children or high fever.
- The presence of air bubbles or fluid behind an intact tympanic membrane, in the absence of signs and symptoms of acute infection, is suggestive of otitis media with effusion (OME).
- Complications, particularly in high-risk children (malnutrition, immunodeficiency, ear malformation) include chronic suppurative otitis media, and rarely, mastoiditis, brain abscess or meningitis.

Treatment

- In all cases:
 - Treatment of [fever](#) and [pain](#): paracetamol PO (Chapter 1).
 - Ear irrigation is contra-indicated if the tympanic membrane is ruptured, or when the tympanic membrane cannot be fully visualised. Ear drops are not indicated.
- Indications for antibiotic therapy:
 - Antibiotics are prescribed in children less than 2 years, children whose assessment suggests severe infection (vomiting, fever > 39 °C, severe pain) and children at risk of unfavourable outcome (malnutrition, immunodeficiency, ear malformation).
 - For other children:
 - ▷ If the child can be re-examined within 48 to 72 hours: it is preferable to delay antibiotic prescription. Spontaneous resolution is probable and a short symptomatic treatment of fever and pain may be sufficient.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Amoxicillin/clavulanic acid is used as second-line treatment, in the case of treatment failure. Treatment failure is defined as persistence of fever and/or ear pain after 48 hours of antibiotic treatment.

amoxicillin/clavulanic acid (co-amoxiclav) PO for 5 days

Use formulations in a ratio of 8:1 or 7:1. The dose is expressed in amoxicillin:

Children < 40 kg: 25 mg/kg 2 times daily

Children ≥ 40 kg and adult:

Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)

Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)

Persistence of a ear drainage alone, without fever and pain, in a child who has otherwise improved (reduction in systemic symptoms and local inflammation) does not warrant a change in antibiotic therapy. Clean ear canal by gentle dry mopping until no more drainage is obtained.

- Macrolides should be reserved for very rare penicillin-allergic patients, as treatment failure (resistance to macrolides) is frequent.

azithromycin PO

Children over 6 months: 10 mg/kg once daily for 3 days

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

Chronic suppurative otitis media (CSOM)

Chronic bacterial infection of the middle ear with persistent purulent discharge through a perforated tympanic membrane.

The principal causative organisms are *Pseudomonas aeruginosa*, *Proteus* sp, staphylococcus, other Gram negative and anaerobic bacteria.

Clinical features

- Purulent discharge for more than 2 weeks, often associated with hearing loss or even deafness; absence of pain and fever
- Otoscopy: perforation of the tympanic membrane and purulent exudate
- Complications:
 - Consider a superinfection (AOM) in the case of new onset of fever with ear pain, and treat accordingly.
 - Consider mastoiditis in the case of new onset of high fever, severe ear pain and/or tender swelling behind the ear, in a patient who appears significantly unwell.
 - Consider brain abscess or meningitis in the case of impaired consciousness, neck stiffness and focal neurological signs (e.g. facial nerve paralysis).

Treatment

- Remove secretions from the auditory canal by gentle dry mopping (use a dry cotton bud or a small piece of dry cotton wool).
- Apply **ciprofloxacin** ear drops until no more drainage is obtained (approximately 2 weeks, max. 4 weeks):
Children 1 year and over: 3 drops 2 times daily
Adults: 4 drops 2 times daily
- Complications:
 - Chronic mastoiditis is a medical emergency that requires prompt hospitalisation, prolonged antibiotic therapy that covers the causative organisms of CSOM (**ceftriaxone** IM for 10 days + **ciprofloxacin** PO for 14 days), atraumatic cleaning of the ear canal; surgical treatment may be required. Before transfer to hospital, if the patient needs to be transferred, administer the first dose of antibiotics.
 - [Meningitis](#) (Chapter 7).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Whooping cough (pertussis)

Whooping cough is a highly contagious bacterial infection of the lower respiratory tract, of prolonged duration, due to *Bordetella pertussis*.

B. pertussis is transmitted through inhalation of droplets spread by infected individuals (coughing, sneezing).

The majority of cases arise in non-vaccinated or incompletely vaccinated individuals. Whooping cough affects all age groups. Signs and symptoms are usually minor in adolescents and adults. As a result the infection may be ignored, thus contributing to the spread of *B. pertussis* and infection in infants and young children, in whom the illness is severe.

Clinical features

After an incubation period of 7 to 10 days, the illness evolves in 3 phases:

- Catarrhal phase (1 to 2 weeks): coryza and cough. At this stage, the illness is indistinguishable from a minor upper respiratory infection.
- Paroxysmal phase (1 to 6 weeks):
 - Typical presentation: cough of at least 2 weeks duration, occurring in characteristic bouts (paroxysms), followed by a laboured inspiration causing a distinctive sound (whoop), or vomiting. Fever is absent or moderate, and the clinical exam is normal between coughing bouts; however, the patient becomes more and more fatigued.
 - Atypical presentations:
 - ▷ Infants under 6 months: paroxysms are poorly tolerated, with apnoea, cyanosis; coughing bouts and whoop may be absent.
 - ▷ Adults: prolonged cough, often without other symptoms.
 - Complications:
 - ▷ Major: in infants, secondary bacterial pneumonia (new-onset fever is an indicator); malnutrition and dehydration triggered by poor feeding due to cough and vomiting; rarely, seizures, encephalopathy; sudden death.
 - ▷ Minor: subconjunctival haemorrhage, petechiae, hernias, rectal prolapse.
- Convalescent phase: symptoms gradually resolve over weeks or months.

Management and treatment

Suspect cases

- Routinely hospitalise infants less than 3 months, as well as children with severe cases. Infants under 3 months must be monitored 24 hours per day due to the risk of apnoea.
- When children are treated as outpatients, educate the parents about signs that should lead to re-consultation (fever, deterioration in general condition, dehydration, malnutrition, apnoea, cyanosis).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Antibiotic treatment is indicated in the first 3 weeks after onset of cough. Infectivity is virtually nil after 5 days of antibioticotherapy.

	Antibiotic	Children	Adults
First line	azithromycin PO for 5 days	10 mg/kg once daily (max. 500 mg daily)	D1 500 mg D2 to D5 250 mg once daily
Alternative^(a)	co-trimoxazole PO for 14 days (if macrolides contra-indicated or not tolerated)	20 mg/kg SMX + 4 mg/kg TMP 2 times daily (avoid in infant < 1 month, and in the last month of pregnancy)	800 mg SMX + 160 mg TMP 2 times daily

(a) Erythromycin (7 days) is a possible alternative but azithromycin is better tolerated and simpler to administrate (shorter treatment duration, fewer daily doses). For dosage according to age or weight, see [erythromycin](#) in the guide Essential drugs, MSF.

- For hospitalised children:
 - Place the child in a semi-reclining position ($\pm 30^\circ$).
 - Oro-pharyngeal suction if needed.

Post-exposure prophylaxis

- Antibiotic prophylaxis (same treatment as for suspect cases) is recommended for unvaccinated or incompletely vaccinated infants of less than 6 months, who have had contact with a suspect case.
- Isolation of contacts is not necessary.

Note: pertussis vaccination should be updated in all cases (suspects and contacts). If the primary series has been interrupted, it should be completed, rather than restarted from the beginning.

Prevention

Routine vaccination with polyvalent vaccines containing pertussis antigens (e.g. DTP, or DTP + Hep B, or DTP + Hib + Hep B) from the age of 6 weeks or according to national protocol.

Neither vaccination nor natural disease confers lasting immunity. Booster doses are necessary to reinforce immunity and reduce the risk of developing disease and transmitting it to young children.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Bronchitis

- [Acute bronchitis](#)
- [Chronic bronchitis](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Acute bronchitis

An acute inflammation of the bronchial mucosa, most commonly of viral origin. In older children it can be caused by *Mycoplasma pneumoniae*. In children over 2 years of age with repetitive acute bronchitis or 'wheezing' bronchitis, consider asthma (see [Asthma](#)). In children under 2 years of age, consider bronchiolitis (see [Bronchiolitis](#)).

Clinical features

Often begins with a rhinopharyngitis that descends progressively: pharyngitis, laryngitis, tracheitis.

- Heavy cough, dry at the beginning then becoming productive
- Low-grade fever
- No tachypnoea, no dyspnoea
- On pulmonary auscultation: bronchial wheezing

Treatment

- **Fever:** paracetamol PO (Chapter 1).
- Keep the patient hydrated, humidify air (with a bowl of water or a wet towel).
- Children: nasal irrigation with 0.9% sodium chloride or Ringer lactate, 4 to 6 times daily to clear the airway.
- Antibiotherapy is not useful for patients in good overall condition with rhinopharyngitis or influenza.
- Antibiotherapy is indicated only if:
 - the patient is in poor general condition: malnutrition, measles, rickets, severe anaemia, cardiac disease, elderly patient etc.
 - if the patient has dyspnoea, fever greater than 38.5 °C and purulent expectorations: a secondary infection with *Haemophilus influenzae* or with pneumococcus is probable.

amoxicillin PO

Children: 30 mg/kg 3 times daily (max. 3 g daily) for 5 days

Adults: 1 g 3 times daily for 5 days

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chronic bronchitis

A chronic inflammation of the bronchial mucosa due to irritation (tobacco, pollution), allergy (asthma) or infection (repetitive acute bronchitis). It may develop into chronic obstructive pulmonary disease.

Clinical features

- Productive cough for 3 consecutive months per year for 2 successive years.
- No dyspnoea at onset. Dyspnoea develops after several years, first on exertion, then becoming persistent.
- On pulmonary auscultation: bronchial wheeze (always exclude tuberculosis).

A patient with an acute exacerbation of chronic bronchitis presents with:

- Onset or increase of dyspnoea.
- Increased volume of sputum.
- Purulent sputum.

Treatment

- Antibiotic treatment is not useful in treating simple chronic bronchitis.
- Antibiotic treatment may be useful, for patients in a poor general condition only, for acute exacerbations of chronic bronchitis (see [Acute bronchitis](#)).
- Discourage smoking and other irritating factors.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Bronchiolitis

Bronchiolitis is an epidemic and seasonal viral infection of the lower respiratory tract in children less than 2 years of age, characterised by bronchiolar obstruction.

Respiratory syncytial virus (RSV) is responsible for 70% of cases of bronchiolitis. Transmission of RSV is direct, through inhalation of droplets (coughing, sneezing), and indirect, through contact with hands or materials contaminated by infected secretions.

In the majority of cases, bronchiolitis is benign, resolves spontaneously (relapses are possible), and can be treated on an outpatient basis.

Severe cases may occur, which put the child at risk due to exhaustion or secondary bacterial infection. Hospitalisation is necessary when signs/criteria of severity are present (10 to 20% of cases).

Clinical features

- Tachypnoea, dyspnoea, wheezing, cough; profuse, frothy, obstructive secretions.
- On auscultation: prolonged expiration with diffuse, bilateral wheezes; sometimes diffuse fine, end-inspiratory crackles.

Rhinopharyngitis, with dry cough, precedes these features by 24 to 72 hours; fever is absent or moderate.

- Signs of severity:
 - Significant deterioration in general condition, toxic appearance (pallor, greyish colouration)
 - Apnoea, cyanosis (check lips, buccal mucosa, fingernails)
 - Respiratory distress (nasal flaring, sternal and chest wall indrawing)
 - Anxiety and agitation (hypoxia), altered level of consciousness
 - Respiratory rate > 60/minute
 - Decreased respiratory distress and slow respirations (< 30/minute below the age of 1 year and < 20/minute below the age of 3 years, exhaustion). Exercise caution in interpreting these signs as indicators of clinical improvement.
 - Sweats, tachycardia at rest and in the absence of fever
 - Silence on auscultation (severe bronchospasm)
 - Difficulty drinking or sucking (reduced tolerance for exertion)

Treatment

Treatment is symptomatic. Obstructive signs and symptoms last for about 10 days; cough may persist for 2 weeks longer.

Hospitalise children with one of the following criteria:

- Presence of any sign of severity
 -
 -
 -
 -
 -
 -
 -
 -

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Small, frequent feedings to reduce vomiting triggered by bouts of coughing.
- Increased fluids if fever and/or significant secretions are present.
- Treat [fever](#) (Chapter 1).
- Handle the patient the patient as little as possible and avoid unnecessary procedures.

Hospitalisation

- In all cases:
 - Place the infant in a semi-reclining position ($\pm 30^\circ$).
 - Nasal irrigation, small, frequent feeds, treatment of fever: as for outpatient treatment.
 - Gentle oro-pharyngeal suction if needed.
 - Monitor fluid intake: normal requirements are 80 to 100 ml/kg/day + 20 to 25 ml/kg/day with high fever or very profuse secretions.
- According to symptoms:
 - Humidified nasal oxygen (1 to 2 litres/minute).
 - When there is vomiting or significant fatigue when sucking, fluid requirements may be administered by nasogastric tube (small volumes on a frequent basis) or the IV route, for the shortest possible time. Avoid breastfeeding or oral feeds in children with severe tachypnoea, but do not prolong NG feeds (respiratory compromise) or IV infusions any longer than necessary.
 - Bronchodilator therapy: this therapy may be considered after a trial treatment has been given (**salbutamol** inhaler, 100 micrograms/puff: 2 to 3 puffs with spacer, repeated twice at an interval of 30 minutes). If inhaled salbutamol appears effective in relieving symptoms, the treatment is continued (2 to 3 puffs every 6 hours in the acute phase, then gradual reduction as recovery takes place). If the trial is ineffective, the treatment is discontinued.
 - Antibiotics are not indicated unless there is concern about complications such as secondary bacterial pneumonia.

Prevention and control

The risk of transmission of the virus is increased in hospital settings:

- Children with bronchiolitis should be grouped together, away from other children (cohorting).
- As infection is most commonly transmitted by the hands, the most important prevention measure is hand-washing after any contact with patients, and objects or surfaces in contact with patients on which the virus may survive for several hours.
- In addition, staff should wear gowns, gloves and surgical masks when in contact with patients.

Footnotes

(a) Lie the child on his back, head turned to the side and instill 0.005% NaCl into the nose one nostril at a time.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute pneumonia

- [Pneumonia in children under 5 years of age](#)
- [Pneumonia in children over 5 years and adults](#)
- [Persistent pneumonia](#)

Acute pneumonia is a viral, bacterial (pneumococcus, *Haemophilus influenzae*, staphylococcus, atypical bacteria) or parasitic (pneumocystosis) infection of the pulmonary alveoli.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Pneumonia in children under 5 years of age

The most common causes are viruses, pneumococcus and *Haemophilus influenzae*.

Clinical features

- Cough or difficulty breathing
- Fever often high ($> 39^{\circ}\text{C}$), but the child may present with low-grade fever or may have no fever (often a sign of serious illness)

Clinical examination must be done on a calm child in order to correctly count the respiratory rate and look for signs of serious illness.

- A child has tachypnoea (increased respiratory rate) if:
 - RR ≥ 60 breaths/minute in children under 1 months
 - RR ≥ 50 breaths/minute in children from 1 to 11 months
 - RR ≥ 40 breaths/minute in children from 12 months to 5 years
- On pulmonary auscultation: dullness with diminished vesicular breath sounds, crepitations and sometimes bronchial breathing or normal pulmonary auscultation.
- Signs of serious illness (severe pneumonia):
 - Chest indrawing: the inferior thoracic wall depresses on inspiration as the superior abdomen expands
 - Cyanosis (lips, oral mucosa, fingernails) or $\text{SpO}_2 < 90\%$
 - Nasal flaring
 - Altered consciousness (child is abnormally sleepy or difficult to wake)
 - Stridor (hoarse noise on inspiration)
 - Grunting (a short repetitive noise produced by a partial closure of the vocal cords) on expiration
 - Refusal to drink or feed
 - Children under 2 months
 - Severe malnutrition

Notes:

- In malnourished children, the RR thresholds should be decreased by 5 breaths/minute from those listed above.
- Chest indrawing is significant if it is clearly visible and present at all times. If it is observed when a child is upset or feeding and is not visible when the child is resting, there is no chest indrawing.
- In children under 2 months of age, moderate chest indrawing is normal as the thoracic wall is flexible.
- If only the soft tissues between the ribs or above the clavicles depress, there is no chest indrawing.

Consider also:

- Malaria in endemic areas, as it may also cause cough and tachypnoea.
- [Staphylococcal pneumonia](#) in patients with empyema or painful abdominal swelling and diarrhoea.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Children under 2 months

The first line treatment is the combination **ampicillin** slow IV (3 minutes) for 10 days + **gentamicin** slow IV (3 minutes) or IM for 5 days:

Children 0 - 7 days	< 2 kg	ampicillin 50 mg/kg every 12 hours + gentamicin 3 mg/kg once daily
	≥ 2 kg	ampicillin 50 mg/kg every 8 hours + gentamicin 5 mg/kg once daily
Children 8 days - < 1 month		ampicillin 50 mg/kg every 8 hours + gentamicin 5 mg/kg once daily
Children 1 month - < 2 months		ampicillin 50 mg/kg every 6 hours + gentamicin 6 mg/kg once daily

For ampicillin, IV route is preferred but IM route may be an alternative.

If ampicillin is not available, alternatives may be **cefotaxime** slow IV (3 minutes) or infusion (20 minutes) or IM for 10 days (for doses, see [Meningitis](#), Chapter 7), or, as a last resort: **ceftriaxone** slow IV^b (3 minutes) or infusion (30 minutes; 60 minutes in neonates) or IM: 50 mg/kg once daily for 10 days.

If the child's condition does not improve^c after 48 hours of well administered treatment, add **cloxacillin** IV for 10 to 14 days:

Children 0 - 7 days	< 2 kg	cloxacillin 50 mg/kg every 12 hours
	≥ 2 kg	cloxacillin 50 mg/kg every 8 hours
Children > 7 days	< 2 kg	cloxacillin 50 mg/kg every 8 hours
	≥ 2 kg	cloxacillin 50 mg/kg every 6 hours

Children from 2 months to 5 years

The first line treatment is:

ceftriaxone IM or slow IV^b (3 minutes): 50 mg/kg once daily

or

ampicillin slow IV (3 minutes) or IM: 50 mg/kg every 6 hours

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

If the child's condition does not improve after 48 hours with ceftriaxone + cloxacillin, consider tuberculosis. For the diagnosis, refer to the guide [Tuberculosis](#), MSF.

If tuberculosis is unlikely, continue with ceftriaxone + cloxacillin and add azithromycin (see [Atypical pneumonia](#)).

Notes:

- For malnourished children, refer to specific protocol.
- In the event of moderate-large empyema, assess if drainage is required. Administer antibiotics active against pneumococci and staphylococci (see [Staphylococcal pneumonia](#)).

Adjvant therapy

- [Fever](#): paracetamol PO (Chapter 1).
- Infants: keep warm.
- Install on an incline (head elevated) or in semi-sitting position.
- Clear the airway (nasal irrigation with 0.9% sodium chloride if needed).
- Oxygen at the flow rate required to maintain $\text{SpO}_2 \geq 90\%$ or, if pulse oximeter is not available, minimum 1 litre/minute.
- Maintain adequate hydration and nutrition:
 - In children with severe respiratory difficulty: place an IV line and give 70% of normal maintenance fluids. Resume oral feeding as soon as possible (no severe respiratory difficulty, ability to eat normally). Use a nasogastric tube only if an IV line cannot be established: children under 12 months: 5 ml/kg/hour; children over 12 months: 3 to 4 ml/kg/hour; alternate milk and water. Resume normal oral feeding as soon as possible.
 - In the absence of severe respiratory difficulty: breastfeed on demand; milk/food and water by spoon on demand.
 - ORS when required ([Dehydration](#), Chapter 1).

Pneumonia with no signs of serious illness

Children under 2 months

Admit the child for inpatient care and treat for [severe pneumonia](#).

Children from 2 months to 5 years

Treat as outpatient, except infants.

amoxicillin PO: 30 mg/kg 3 times daily for 5 days

Follow-up in 48 to 72 hours or sooner if the child's condition deteriorates:

- If the condition is improving^c: continue with the same antibiotic to complete treatment.
- If there is no improvement after 3 days of correct administration: add azithromycin (see [Atypical pneumonia](#)).
- If the condition is deteriorating: hospitalise and treat as severe pneumonia.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Pneumonia in children over 5 years and adults

The most common causes are viruses, pneumococcus, and *Mycoplasma pneumoniae*.

Clinical features

- Cough, with or without purulent sputum, fever, thoracic pain, tachypnoea
- On pulmonary auscultation: decreased vesicular breath sounds, dullness, localised foci of crepitations, sometimes bronchial wheeze.

Sudden onset with high fever (higher than 39 °C), thoracic pain and oral herpes are suggestive of pneumococcal infection. Symptoms may be confusing, particularly in children with abdominal pain, meningeal syndrome, etc.

Signs of serious illness (severe pneumonia) include:

- Cyanosis (lips, oral mucosa, fingernails)
- Nasal flaring
- Intercostal or subclavial indrawing
- RR > 30 breaths/minute
- Heart rate > 125 beats/minute
- Altered level of consciousness (drowsiness, confusion)

Patients at risk include the elderly, patients suffering from heart failure, sickle cell disease or severe chronic bronchitis; immunocompromised patients (severe malnutrition, HIV infection with CD4 < 200).

Treatment

Severe pneumonia (inpatient treatment)

ceftriaxone IM or slow IV^a (3 minutes)

Children: 50 mg/kg once daily

Adults: 1 g once daily

The treatment is given by parenteral route for at least 3 days then, if the clinical condition has improved^b and oral treatment can be tolerated, switch to **amoxicillin** PO to complete 7 to 10 days of treatment:

Children: 30 mg/kg 3 times daily (max. 3 g daily)

Adults: 1 g 3 times daily

or

ampicillin slow IV (3 minutes) or IM

Children: 50 mg/kg every 6 hours

Adults: 1 g every 6 to 8 hours

Ampicillin is preferably administered in 4 divided doses. If the context does not permit it, the daily dose must be divided

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

If the clinical condition deteriorates or does not improve after 48 hours of correct administration, administer ceftriaxone as above + **cloxacillin** IV infusion:

Children: 25 to 50 mg/kg every 6 hours

Adults: 2 g every 6 hours

After clinical improvement and 3 days with no fever, switch to **amoxicillin/clavulanic acid (co-amoxiclav)** PO to complete 10 to 14 days of treatment. Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin:

Children < 40 kg: 50 mg/kg 2 times daily

Children ≥ 40 kg and adults:

Ratio 8:1: 3000 mg daily (2 tablets of 500/62.5 mg 3 times daily)

Ratio 7:1: 2625 mg daily (1 tablet of 875/125 mg 3 times daily)

If the clinical condition does not improve after 48 hours with ceftriaxone + cloxacillin, consider tuberculosis. For the diagnosis, refer to the guide [Tuberculosis](#), MSF.

If tuberculosis is unlikely, continue with ceftriaxone + cloxacillin and add azithromycin (see [Atypical pneumonia](#)).

Adjuvant therapy

- **Fever:** paracetamol PO (Chapter 1).
- Clear the airway (nasal irrigation with 0.9% sodium chloride if needed).
- Oxygen at the flow rate required to maintain $\text{SpO}_2 \geq 90\%$ or, if pulse oximeter is not available, minimum 1 litre/minute.
- Maintain adequate hydration and nutrition.

Pneumonia without signs of serious illness (outpatient treatment)

amoxicillin PO

Children: 30 mg/kg 3 times daily (max. 3 g daily) for 5 days

Adults: 1 g 3 times daily for 5 days

Follow-up in 48 to 72 hours or sooner if the child's condition deteriorates:

- If the condition is improving^b: continue with the same antibiotic to complete treatment.
- If there is no improvement after 3 days of correct administration: add azithromycin (see [Atypical pneumonia](#)).
- If the condition is deteriorating: hospitalise and treat as severe pneumonia.

Footnotes

(a) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must never be administered by IV route. For IV administration, water for injection must always be used.

(b) Improvement criteria include: fever reduction, diminished respiratory distress, improved SpO_2 , improved appetite and/or

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Persistent pneumonia

Last update: November 2022

In patients not responding to therapy, consider atypical pneumonia, tuberculosis, pneumocystosis ([HIV infection and AIDS](#), Chapter 8).

Bacteria responsible for atypical pneumonia are mainly *Mycoplasma pneumoniae* and *Chlamydophila pneumoniae*. If suspected, one of the following antibiotics may be used:

First choice, **azithromycin** PO

Children: 10 mg/kg (max. 500 mg) once daily for 5 days

Adults: 500 mg on D1 then, 250 mg once daily from D2 to D5

If not available,

erythromycin PO

Children: 10 mg/kg (max. 500 mg) 4 times daily for 10 to 14 days

Adults: 500 mg 4 times daily for 10 to 14 days

or

doxycycline PO (except in pregnant or breastfeeding women)

Children under 45 kg: 2 to 2.2 mg/kg (max. 100 mg) 2 times daily for 10 to 14 days

Children 45 kg and over and adults: 100 mg 2 times daily for 10 to 14 days

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Staphylococcal pneumonia

Pneumonia due to *Staphylococcus aureus* affecting young children, often those in a poor general condition (malnutrition, skin lesions, etc.). Staphylococcal pneumonia is a classic complication of measles.

Clinical features

- General signs: change in overall condition, pallor, high fever or hypothermia, frequently signs of shock; presence of skin lesions (point of bacterial entry), however, skin lesions may be absent.
- Gastrointestinal signs: nausea, vomiting, diarrhoea, painful abdominal distension.
- Respiratory signs: dry cough, tachypnoea, signs of distress (nasal flaring, chest indrawing). Pulmonary auscultation is often normal; sometimes dullness indicating pleural effusion.

Paraclinical investigations

- Chest x-ray (if available): may show multilobar consolidation, cavitation, pneumatoceles, spontaneous pneumothorax.

Treatment

Treatment is urgent as patients deteriorate quickly: hospitalise.

- Antibiotic treatment: if staphylococcal aetiology cannot be confirmed or while waiting for confirmation, a broad spectrum antibiotic therapy is recommended:

ceftriaxone IM or slow IV^a (at least 3 minutes): 50 mg/kg once daily

+ **cloxacillin** IV infusion (60 minutes)^b

Neonates 0 to 7 days (< 2 kg): 50 mg/kg every 12 hours

Neonates 0 to 7 days (\geq 2 kg): 50 mg/kg every 8 hours

Neonates 8 days to < 1 month (< 2 kg): 50 mg/kg every 8 hours

Neonates 8 days to < 1 month (\geq 2 kg): 50 mg/kg every 6 hours

Children 1 month and over: 25 to 50 mg/kg every 6 hours (max. 8 g daily)

After clinical improvement^c, 3 days with no fever, and drain removal if any, switch to **amoxicillin/clavulanic acid** PO to complete 10 to 14 days. Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin: 50 mg/kg 2 times daily

In the event of large empyema: same treatment but switch to the oral route after 7 days with no fever and treat for 3 weeks.

Clindamycin IV may be an alternative to cloxacillin: 10 mg/kg every 8 hours then switch to clindamycin PO at the same dose according to the criteria above

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- If there is significant pleural effusion: pleural tap with drainage (for pyopneumothorax; insert 2 drains, one anterior and one posterior) or without drainage (for suppurative pleurisy, make repetitive taps with an IV catheter).

Clinical evolution

- There is a serious risk of decompensation from pneumothorax or suppurative pleurisy or pyopneumothorax.
- On a paediatric ward, adequate equipment for urgent pleural drainage should always be available.

Footnotes

- (a) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must never be administered by IV route. For IV administration, water for injection must always be used.
- (b) Cloxacillin powder for injection should be reconstituted in 4 ml of water for injection. Then dilute each dose of cloxacillin in 5 ml/kg of 0.9% sodium chloride or 5 % glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (c) Improvement criteria include: fever reduction, diminished respiratory distress, improved SpO₂, improved appetite and/or activity.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Asthma

- [Asthma attack \(acute asthma\)](#)
- [Chronic asthma](#)

Asthma is a chronic inflammatory disorder of the airways associated with airway hyperresponsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness and coughing. These episodes are usually associated with airflow obstruction within the lung, often reversible, either spontaneously or with treatment.

Factors that precipitate/aggravate asthma include: allergens, infection, exercise, drugs (aspirin), tobacco, etc.

In young children, most initial episodes of asthma-like symptoms are associated with a respiratory tract infection, with no symptoms between infections. Wheezing episodes usually become less frequent with time; most of these children do not develop asthma.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Asthma attack (acute asthma)

Asthma attack is a substantial worsening of asthma symptoms. The severity and duration of attacks are variable and unpredictable.

Assessment of the severity of asthma attack

The severity of the asthma attack must be rapidly evaluated by the following clinical criteria. Not all signs are necessarily present.

Assessment of severity in children over 2 years and adults

Mild to moderate attack	Severe attack	Life threatening attack
Able to talk in sentences Respiratory rate (RR) Children 2-5 years \leq 40/min Children > 5 years \leq 30/min Heart rate Children 2-5 years \leq 140/min Children > 5 years \leq 125/min and No criteria of severity	Cannot complete sentences in one breath or Too breathless to talk or feed RR Children 2-5 years $>$ 40/min Children > 5 years $>$ 30/min Adults \geq 25/min Heart rate Children 2-5 years $>$ 140/min Children > 5 years $>$ 125/min Adults \geq 110/min SpO₂ \geq 92%	Altered level of consciousness (drowsiness, confusion, coma) Exhaustion Silent chest Paradoxical thoracoabdominal movement Cyanosis Collapse Bradycardia in children or arrhythmia/hypotension in adults SpO₂ < 92%

Treatment

Treatment and follow-up depend on the severity of the attack and the patient's response:

Mild to moderate attack

- Reassure the patient; place him in a 1/2 sitting position.
- Administer:
 - **salbutamol (aerosol):** 2 to 4 puffs every 20 to 30 minutes, up to 10 puffs if necessary during the first hour. In children under 4 years, administer 1/2 dose (one puff). Single puffs should be

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Severe attack

- Hospitalise the patient; place him in a 1/2 sitting position.
- Administer:
 - **oxygen** continuously, at least 5 litres/minute or maintain the SpO₂ between 94 and 98%.
 - **salbutamol** (aerosol): 2 to 4 puffs every 20 to 30 minutes, up to 10 puffs if necessary in children under 5 years, up to 20 puffs in children over 5 years and adults. Use a spacer to increase effectiveness, irrespective of age. or **salbutamol** (solution for nebulisation), see [Life-threatening attack](#).
 - **prednisolone** PO: one dose of 1 to 2 mg/kg
In the case of vomiting, until the patient can tolerate oral prednisolone, use **hydrocortisone** IV:
Children 1 month to < 5 years: 4 mg/kg every 6 hours (max. 100 mg per dose)
Children 5 years and over and adults: 100 mg every 6 hours
- If the attack is completely resolved, observe the patient for at least 4 hours. Continue the treatment with **salbutamol** for 24 to 48 hours (2 to 4 puffs every 4 hours) and **prednisolone** PO (1 to 2 mg/kg once daily) to complete 3 days of treatment.
Reassess after 10 days: consider long-term treatment if the asthma attacks have been occurring for several months. If the patient is already receiving long-term treatment, reassess the severity of the asthma (see [table](#)) and review compliance and correct use of medication and adjust treatment if necessary.
- If symptoms worsen or do not improve, see [Life-threatening attack](#).

Life-threatening attack (intensive care)

- Insert an IV line.
- Administer:
 - **oxygen** continuously, at least 5 litres/minute or maintain the SpO₂ between 94 and 98%.
 - **salbutamol + ipratropium** nebuliser solutions using a nebuliser:

Children 1 month to < 5 years	salbutamol 2.5 mg + ipratropium 0.25 mg every 20 to 30 minutes
Children 5 to < 12 years	salbutamol 2.5 to 5 mg + ipratropium 0.25 mg every 20 to 30 minutes
Children 12 years and over and adults	salbutamol 5 mg + ipratropium 0.5 mg every 20 to 30 minutes

The two solutions can be mixed in the nebuliser reservoir.

- corticosteroids (prednisolone PO or hydrocortisone IV) as for [severe attack](#).
- If the attack is resolved after one hour: switch to salbutamol aerosol and continue prednisolone PO as for [severe attack](#).
- If symptoms do not improve after one hour:
 - administer a single dose of **magnesium sulfate** by IV infusion in 0.9% sodium chloride over 20 minutes,

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Footnotes

- (a) If a conventional spacer is not available, use a 500 ml plastic bottle: insert the mouthpiece of the inhaler into a hole made in the bottom of the bottle (the seal should be as tight as possible). The child breathes from the mouth of the bottle in the same way as he would with a spacer. The use of a plastic cup instead of a spacer is not recommended (ineffective).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Chronic asthma

Clinical features

- Asthma should be suspected in patients with episodic respiratory symptoms (wheezing, chest tightness, shortness of breath and/or cough) of variable frequency, severity and duration, disturbing sleep, and causing the patient to sit up to breathe. These symptoms may appear during or after exercise.
- Chest auscultation may be normal or demonstrate diffuse sibilant wheezes.
- Atopic disorders or a personal or family history of atopy (eczema, allergic rhinitis/conjunctivitis) or a family history of asthma increases probability of asthma but their absence does not exclude asthma.

Patients with typical symptoms of asthma and a history of disease that is characteristic of asthma should be considered as having asthma after exclusion of other diagnoses.

The assessment of the frequency of daytime and nighttime symptoms and limitations of physical activity determines whether asthma is intermittent or persistent.

Treatment

Only patients with persistent asthma need long-term treatment. The mainstay of treatment is inhaled corticosteroids. Treatment is started at the step most appropriate to initial severity then, re-evaluated and adjusted according to clinical response. It aims to abolish symptoms with the lowest possible dose of inhaled corticosteroids. An intervening severe exacerbation or loss of control necessitates reassessment to re-evaluate treatment.

Long-term treatment does not mean treatment for life. Asthma attacks may occur over months or years, with intervening asymptomatic intervals when long-term treatment is not required.

Long-term treatment of asthma according to severity

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Categories	Treatment
Intermittent asthma <ul style="list-style-type: none"> • Day time symptoms < once weekly • Night time symptoms < 2 times monthly • Normal physical activity 	No long term treatment Inhaled salbutamol when symptomatic
Mild persistent asthma <ul style="list-style-type: none"> • Day time symptoms > once weekly, but < once daily • Night time symptoms > 2 times monthly • Symptoms may affect activity 	Continuous treatment with inhaled beclometasone + Inhaled salbutamol when symptomatic
Moderate persistent asthma <ul style="list-style-type: none"> • Daily symptoms • Symptoms affect activity • Night time symptoms > once weekly • Daily use of salbutamol 	Continuous treatment with inhaled beclometasone + Inhaled salbutamol (1 puff 4 times daily)
Severe persistent asthma <ul style="list-style-type: none"> • Daily symptoms • Frequent night time symptoms • Physical activity limited by symptoms 	Continuous treatment with inhaled beclometasone + Inhaled salbutamol (1-2 puff/s 4 to 6 times daily)

Inhaled corticosteroid treatment: **beclometasone** dose varies according to the severity of asthma. Find the minimum dose necessary to both control the symptoms and avoid local and systemic adverse effects:

Children: 50 to 100 micrograms 2 times daily depending on the severity; increase to 200 micrograms 2 times daily if necessary (max. 800 micrograms daily)

Adults: 100 to 250 micrograms 2 times daily depending on the severity; increase to 500 micrograms 2 times daily if necessary (max. 1500 micrograms daily)

 The number of puffs of beclometasone depends on its concentration in the inhaled aerosol: 50, 100 or 250 micrograms per puff.

Do not restrict exercise. If exercise is a trigger for asthma attacks, administer 1 or 2 puffs of salbutamol 10 minutes beforehand.

In pregnant women, poorly controlled asthma increases the risk of pre-eclampsia, eclampsia, haemorrhage, in utero growth retardation, premature delivery, neonatal hypoxia and perinatal mortality. Long-term treatment remains inhaled salbutamol and beclometasone at the usual dosage for adults. Whenever possible, avoid oral corticosteroids.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Pulmonary tuberculosis

Pulmonary tuberculosis is a bacterial infection due to *Mycobacterium tuberculosis*, spread from person to person through inhalation of infected respiratory droplets.

After infection, *M. tuberculosis* multiplies slowly in the lungs and is usually eliminated spontaneously or lies dormant.

Only 10% of cases develop active tuberculosis. The risk of progressing to active tuberculosis is higher in immunocompromised patients. In certain countries, half of newly diagnosed tuberculosis patients are co-infected with HIV^[1].

For more information on tuberculosis, refer to the guide [Tuberculosis](#), MSF.

Clinical features

- Prolonged cough (> 2 weeks) with or without sputum production and/or haemoptysis, prolonged fever, night sweats, anorexia, weight loss, chest pain and fatigue.
- Differential diagnosis includes pneumonia, chronic obstructive pulmonary disease (COPD), lung cancer, pulmonary distomatosis ([Flukes](#), Chapter 6) and melioidosis (Southeast Asia).

In an endemic area, the diagnosis of tuberculosis is to be considered, in any patient consulting for respiratory symptoms for over 2 weeks who does not respond to non-specific antibacterial treatment.

Laboratory

- In the general population: Xpert® MTB/RIF test which simultaneously detects *M. tuberculosis* (MTB) in sputum and resistance to rifampicin (RIF). If not available perform sputum smear microscopy^[2].
- If HIV co-infection suspected or diagnosed: Xpert® MTB/RIF test and point-of-care, urine LF-LAM (lateral flow urine lipoarabinomannan assay)^[2].

Treatment

For pulmonary tuberculosis, the standard treatment is a combination of four antituberculosis drugs (isoniazid, rifampicin, pyrazinamide, ethambutol). The regimen is organised into 2 phases (initial phase and continuation phase) and lasts 6 months.

If the strain is drug-resistant, the treatment is longer and different drug combinations are used.

It takes significant investment to cure tuberculosis, both from the patient and the medical team. Only uninterrupted treatment will lead to cure and prevent the development of resistance. It is essential that the patient understands the importance of treatment adherence and has access to correct case management until treatment is completed.

Prevention

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

1. World Health Organization. Global tuberculosis report 2018.
<https://apps.who.int/iris/handle/10665/274453> [Accessed 21 October 2019]
2. Global Laboratory Initiative. GLI model TB diagnostic algorithms. 2018.
http://www.stoptb.org/wg/gli/assets/documents/GLI_algorithms.pdf [Accessed 21 October 2019]
3. World Health Organization. Weekly epidemiological record/Relevé épidémiologique hebdomadaire 23rd February 2018, 93rd year/23 Février 2018, 93e année. No 8, 2018, 93, 73–96.
https://www.who.int/immunization/policy/position_papers/bcg/en/ [Accessed 21 October 2019]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 3: Gastrointestinal disorders

[Acute diarrhoea](#)

[Shigellosis](#)

[Amoebiasis](#)

[Disorders of the stomach and duodenum](#)

[Gastro-oesophageal reflux](#)

[Gastric and duodenal ulcers in adults](#)

[Dyspepsia](#)

[Stomatitis](#)

[Oral and oropharyngeal candidiasis](#)

[Oral herpes](#)

[Other infectious causes](#)

[Stomatitis from scurvy \(vitamin C deficiency\)](#)

[Other lesions resulting from a nutritional deficiency](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Acute diarrhoea

Acute diarrhoea is defined as at least 3 liquid stools per day for less than 2 weeks.

- There are 2 clinical types of acute diarrhoea:
 - **Diarrhoea without blood**, caused by viruses in 60% of cases (rotavirus, enterovirus), bacteria (*Vibrio cholerae*, enterotoxigenic *Escherichia coli*, non Typhi *Salmonella*, *Yersinia enterocolitica*) or parasites (giardiasis). Diseases, such as malaria, acute otitis media, respiratory tract infections, etc. can be accompanied by this type of diarrhoea.
 - **Diarrhoea with blood**, caused by bacteria (*Shigella* in 50% of cases, *Campylobacter jejuni*, enteroinvasive or enterohaemorrhagic *Escherichia coli*, *Salmonella*) or parasites (intestinal amoebiasis).
- Infectious diarrhoeas are transmitted by direct (dirty hands) or indirect (ingestion of contaminated water or food) contact.
- The high mortality rate from diarrhoeal diseases, even benign, is due to acute dehydration and malnutrition. This can be prevented by adequate rehydration and nutrition.

Clinical features

- First assess for signs of dehydration (see [Dehydration](#), Chapter 1).
- Then look for other signs:
 - profuse watery diarrhoea (cholera, enterotoxigenic *E. coli*),
 - repeated vomiting (cholera),
 - fever (salmonellosis, viral diarrhoea),
 - presence of red blood in stools: see also [Shigellosis](#) and [Amoebiasis](#) (Chapter 3).
- In a patient over 5 years with severe and rapid onset of dehydration, suspect cholera.

Treatment

General principles:

- Prevent or treat dehydration: rehydration consists of prompt replacement of fluid and electrolyte losses as required, until the diarrhoea stops.
- Administer zinc sulfate to children under 5 years.
- Prevent malnutrition.
- Do not systematically administer antimicrobials: only certain diarrhoeas require antibiotics (see [Antimicrobial treatment](#)).
- Do not administer anti-diarrhoeal drugs or antiemetics.
- Treat the underlying condition if any (malaria, otitis, respiratory infection, etc.).

Prevention and treatment of dehydration

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Zinc sulfate is given in combination with oral rehydration solution in order to reduce the duration and severity of diarrhoea, as well as to prevent further occurrences in the 2 to 3 months after treatment:

zinc sulfate PO

Children under 6 months: 10 mg (½ tablet) once daily for 10 days

Children from 6 months to 5 years: 20 mg (1 tablet) once daily for 10 days

Place the half-tablet or full tablet in a teaspoon, add a bit of water to dissolve it, and give the entire spoonful to the child.

Antimicrobial treatment

Diarrhoea without blood

Most acute diarrhoeas are caused by viruses unresponsive to antimicrobials. Antimicrobials can be beneficial in the event of cholera or giardiasis.

- **Cholera:** the most important part of treatment is rehydration. In the absence of resistance (perform antibiotic-sensitivity testing at the beginning of an outbreak), antibiotic treatment shortens the duration of diarrhoea. See the guide [Management of a cholera epidemic](#), MSF.
- **Giardiasis:** see [Intestinal protozoan infections](#), Chapter 6.

Diarrhoea with blood

- **Shigellosis** is the most frequent cause of bloody diarrhoea (amoebiasis is much less common). If there is no laboratory diagnosis to confirm the presence of amoebae, first line treatment is for [shigellosis](#) (Chapter 3).
- **Amoebiasis:** antiparasitic treatment only if motile *Entamoeba histolytica* amoebae are found in stools or if a correct shigellosis treatment has been ineffective (see [Amoebiasis](#), Chapter 3).

Prevention

- Breastfeeding reduces infant morbidity and mortality from diarrhoea and the severity of diarrhoea episodes.
- When the child is weaned preparation and storage of food are associated with the risk of contamination by faecal micro-organisms: discourage bottle-feeding; food must be cooked well; milk or porridge must never be stored at room temperature.
- Access to sufficient amounts of clean water and personal hygiene (washing hands with soap and water before food preparation and before eating, after defecation etc.) are effective methods of reducing the spread of diarrhoea.
- In countries with a high rotavirus diarrhoea fatality rate, the WHO recommends routine rotavirus vaccination in children between 6 weeks and 24 months of age^[1].

References

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Shigellosis

Shigellosis is a highly contagious bacterial infection resulting in bloody diarrhoea. There are 4 serogroups of shigella: *S. dysenteriae*, *S. sonnei*, *S. flexneri*, *S. boydii*.

S. dysenteriae type 1 (Sd1) is the only strain that causes large scale outbreaks. It has the highest case fatality rate (up to 10%).

Patients at risk of death are children under 5 years, malnourished patients, children after measles, adults over 50 years.

Clinical features

- Diarrhoea with bright red blood visible in stool^a, with or without fever
- Abdominal and rectal pain frequent
- Signs of serious illness: fever above 39 °C; severe dehydration; seizures, altered mental status
- Complications (more frequent with Sd1): febrile seizures (5 to 30% of children), rectal prolapse (3%), septicaemia, intestinal obstruction or perforation, moderate to severe haemolytic uraemic syndrome

Laboratory

Shigellosis in an epidemic context:

- Confirm the causal agent (stool culture) and perform antibiotic sensitivity tests.
- Perform monthly culture and sensitivity tests (antibiotic resistance can develop rapidly, sometimes during the course of an outbreak).

Treatment

- Patients with signs of serious illness or with life-threatening risk factors must be admitted as inpatients.
- Treat patients with neither signs of serious illness nor risk factors as outpatients.
- Antibiotherapy:

First-line treatment	
ciprofloxacin PO for 3 days Children: 15 mg/kg 2 times daily (max. 1 g daily) Adults: 500 mg 2 times daily	<ul style="list-style-type: none">• if the strain is sensitive• if there is no antibiotic sensitivity test• if oral administration is possible
ceftriaxone IM for 3 days Children: 50 to 100 mg/kg once daily	<ul style="list-style-type: none">• in patients with severe infection and/or oral administration is not possible

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

azithromycin PO for 5 days

Children: one dose of 12 mg/kg on D1 then 6 mg/kg once daily from D2 to D5

Adults: one dose of 500 mg on D1 then 250 mg once daily from D2 to D5

or

cefixime PO for 5 days

Children: 8 mg/kg once daily (max. 400 mg daily)

Adults: 400 mg once daily

If there is no improvement 48 hours after starting second-line treatment, treat for [amoebiasis^{\[1\]\[2\]}](#).

- For pain and/or fever:

paracetamol PO (see [Pain](#), Chapter 1). All opioid analgesics are contra-indicated as they slow peristalsis.

- Supportive therapy:

▫ nutrition: nutritional supplement with frequent meals

+ 2500 kcal daily during hospitalisation

+ 1000 kcal daily as outpatients

▫ rehydration: administration of ORS according to WHO protocols (see [Dehydration](#), Chapter 1).

▫ zinc supplement in children under 5 years (see [Acute diarrhoea](#), Chapitre 3).

- Never give loperamide or any other antidiarrhoeal.

- Management of complications: rectal prolapse reduction, septicaemia (see [Septic shock](#), Chapter 1), etc.

Shigellosis in an epidemic context

- Isolation of hospitalised patients; school exclusion of children treated as outpatients.
- Hygiene (handwashing, hygienic preparation and storage of food, home hygiene, etc.).
- Management if signs worsen or bloody diarrhoea in entourage (seek medical attention).

Footnotes

(a) This definition excludes: blood detected on microscope examination; stool containing digested blood (melaena); streaks of blood on the surface of normal stool (haemorrhoids, anal or rectal lesion, etc.).

(b) Ciprofloxacin should be avoided in pregnant women. Nevertheless, if ceftriaxone is not available, the other antibiotics can be used, including ciprofloxacin if necessary.

References

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Amoebiasis

Amoebiasis is a parasitic infection due to the intestinal protozoa *Entamoeba histolytica*. Transmission is faecal-oral, by ingestion of amoebic cysts from food or water contaminated with faeces. Usually, ingested cysts release non-pathogenic amoebae and 90% of carriers are asymptomatic.

In 10% of infected patients, pathogenic amoebae penetrate the mucous of the colon: this is the intestinal amoebiasis (amoebic dysentery). The clinical picture is similar to that of shigellosis, which is the principal cause of dysentery. Occasionally, the pathogenic amoebae migrate via the blood stream and form peripheral abscesses. Amoebic liver abscess is the most common form of extra-intestinal amoebiasis.

Clinical features

- Amoebic dysentery
 - diarrhoea containing red blood and mucus
 - abdominal pain, tenesmus
 - no fever or moderate fever
 - possibly signs of dehydration
- Amoebic liver abscess
 - painful hepatomegaly; mild jaundice may be present
 - anorexia, weight loss, nausea, vomiting
 - intermittent fever, sweating, chills; change in overall condition

Investigations

- Amoebic dysentery: identification of mobile trophozoites (*E. histolytica histolytica*) in fresh stool samples
- Amoebic liver abscess: indirect haemoagglutination and ELISA
- POCUS^a: perform an EFAST (extended focused assessment with sonography for trauma) examination, with additional views of the liver and spleen to evaluate for signs of amoebic lesions. Contact an expert (local or via telemedicine services) to help interpret the images and differentiate amoebic abscesses from other pathologies with similar characteristics.

Treatment

- Amoebic dysentery
 - The presence of cysts alone should not lead to the treatment of amoebiasis.
 - Amoebiasis confirmed with a parasitological stool examination:
tinidazole PO
Children: 50 mg/kg once daily for 3 days (max. 2 g daily)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Footnotes

(a) POCUS should only be performed and interpreted by trained clinicians.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Disorders of the stomach and duodenum

- [Gastro-oesophageal reflux](#)
- [Gastric and duodenal ulcers in adults](#)
- [Dyspepsia](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Gastro-oesophageal reflux

Clinical features

Burning stomachache or heartburn, generally relieved by antacids; acid regurgitation (often postural: while sitting forward or lying down). In the absence of dysphagia (oesophageal stenosis), these signs are benign.

Treatment

- First instance: encourage the patient to avoid alcohol and tobacco use.
Give **aluminium hydroxide/magnesium hydroxide** PO (400 mg/400 mg tablet)^a : 1 to 2 tablets 3 times daily 20 minutes to one hour after meals, or 1 tablet during painful attacks.
- If antacids are insufficient:
omeprazole PO: 20 mg once daily in the morning for 3 days
- In young children: no drug treatment, rest and sleep on an incline (30° to 45°).

Footnotes

- (a) Aluminium hydroxide/magnesium hydroxide may decrease intestinal absorption of drugs taken at the same time:
 - atazanavir, chloroquine, digoxin, doxycycline, iron salts, gabapentin, itraconazole, levothyroxine (take at least 2 hours apart).
 - ciprofloxacin (take ciprofloxacin 2 hours before or 4 hours after antacids), dolutegravir (take dolutegravir 2 hours before or 6 hours after antacids), velpatasvir (take 4 hours apart).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Gastric and duodenal ulcers in adults

Clinical features

Burning epigastric pain or epigastric cramps between meals, that wake the patient at night. Recurrent episodes characteristically last a few days and are often accompanied by nausea and even vomiting.

The most common complications are perforation and bleeding.

Treatment of non-complicated ulcers

- For an isolated episode:
 - identify patients taking NSAID or acetylsalicylic acid; stop treatment;
 - encourage patients to avoid alcohol and tobacco use;
 - **omeprazole** PO: 20 mg once daily in the morning for 7 to 10 days. In severe or recurrent cases, dose can be increased to 40 mg once daily and the treatment can be prolonged for up to 8 weeks.
- If the patient has frequent recurrences unrelated to NSAID use, that require repeated treatment with antiulcer drugs: see [eradication of *Helicobacter pylori*](#).

Treatment of complicated ulcers

Perforation

Perforation should be considered in patients presenting with sudden onset intense epigastric pain, particularly if there is rigidity of the abdominal wall. The risk of peritonitis is increased if the perforation occurs on a full stomach.

- To start:
 - place the patient on a strict fast (NPO); insert a nasogastric tube and aspirate if possible;
 - insert an intravenous line and hydrate (Ringer lactate);
 - treat acute pain (see [Pain](#), Chapter 1);
 - **omeprazole** IV infusion: 40 mg once daily over 20 to 30 minutes
- Refer to a surgeon.
- If referral not possible, risk of mortality is high:
 - Continue conservative management including maintenance fluid (alternate 5% glucose and Ringer lactate).
 - Start IV antibiotics (see [Shock](#), Chapter 1).
 - If after 3 days, the patient's clinical condition has improved, cautiously restart oral feeding, remove the nasogastric tube and start PO treatment to eradicate *Helicobacter pylori* (see [eradication of *Helicobacter pylori*](#)).

Gastrointestinal bleeding

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Intensive care and transfusion according to the severity of the bleeding (see [haemorrhagic shock](#), Chapter 1).
- Emergency surgical intervention.

Eradication of *Helicobacter pylori*

Most peptic ulcers are caused by *Helicobacter pylori* infection. If a diagnosis of ulcer is probable, treatment to eradicate *H. pylori* should be considered if the patient has frequent attacks requiring repeated and/or prolonged treatments with antiulcer drugs over 8 weeks or in cases of complicated ulcers (perforation or gastrointestinal bleeding). Infection should be confirmed with a test where possible.

H. pylori resistance to antibiotics varies globally, follow national recommendations where available. If not, for information, administer a triple therapy for 7 days:

omeprazole PO 20 mg 2 times daily + **clarithromycin** PO 500 mg 2 times daily + **amoxicillin** PO 1 g 2 times daily^a.

In immunocompromised patients, consider mycobacterium avium complex (MAC) infection or other nontuberculous mycobacterium (NTM) infection prior to starting a clarithromycin-containing triple therapy.

If symptoms continue despite treatment, consider the differential diagnosis of gastric cancer. Refer for investigations if possible.

Notes:

- Acetylsalicylic acid (aspirin) and NSAID (ibuprofen, diclofenac, etc.) are contra-indicated in patients suffering from or with a history of ulcers.
- Omeprazole is as effective PO as IV.

Footnotes

- (a) In penicillin-allergic patients, amoxicillin PO can be substituted with **metronidazole** PO 500 mg 2 times daily.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Dyspepsia

Last updated: December 2020

Clinical features

Epigastric pain or discomfort following meals, often accompanied by bloating, sensation of fullness and nausea.

Dyspepsia is most commonly functional. The diagnosis of functional dyspepsia is based on clinical assessment after ruling out organic causes ([Gastro-oesophageal reflux](#), [Gastric and duodenal ulcers](#), drug-induced symptoms, gastric cancer). If possible, test for *Helicobacter pylori*.

Treatment

In adults:

- In case of patients who test positive for *H. pylori*, see [Eradication of *Helicobacter pylori*](#)^[1].
- **Omeprazole** PO (10 mg once daily) for 4 weeks may help even in *H. pylori*-negative patients^{[2][3]}.

Note: consider and treat possible intestinal parasites (see [Intestinal protozoan infections](#), [Cestodes](#), [Nematode infections](#), Chapter 6; [Amoebiasis](#), Chapter 3).

References

1. Ford AC, Mahadeva S, Carbone MF, Lacy BE, Talley NJ. Functional dyspepsia. Lancet. 2020 Nov 21;396(10263):1689-1702.
2. Moayyedi PM, Lacy BE, Andrews CN, et al. ACG and CAG clinical guideline: management of dyspepsia. Am J Gastroenterol. 2017 Jul;112(7):988-1013.
http://www.cag-acg.org/images/publications/CAG_CPG_Dyspepsia_AJG_Aug2017.pdf [Accessed 24 November 2020]
3. National Institute for Health and Care Excellence. Gastro-oesophageal reflux disease and dyspepsia in adults: investigation and management. Sept 2014.
<https://www.nice.org.uk/guidance/CG184/chapter/1-Recommendations#interventions-for-functional-dyspepsia> [Accessed 24 November 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Stomatitis

- [Oral and oropharyngeal candidiasis](#)
- [Oral herpes](#)
- [Other infectious causes](#)
- [Stomatitis from scurvy \(vitamin C deficiency\)](#)
- [Other lesions resulting from a nutritional deficiency](#)

Stomatitis is an inflammation of the mucous membranes of the mouth caused by a fungal, viral or bacterial infection, a vitamin deficiency, an injury, etc.

Prolonged or painful stomatitis may contribute to dehydration or may cause loss of appetite with denutrition, particularly in children.

In infants, examine routinely the mouth in the event of breast refusal or difficulties in sucking.

In all cases:

- Maintain adequate hydration and feeding; offer foods that will not irritate the mucosa (soft, non-acidic). Use a nasogastric tube for a few days if pain is preventing the patient from eating.
- Keep the mouth clean to prevent complications and recurrence.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Oral and oropharyngeal candidiasis

Infection due to *Candida albicans*, common in infants, immunocompromised or diabetic patients. Other risk factors include treatment with oral antibiotics or high-dose inhaled corticosteroids.

Clinical features

- White patches on the tongue, inside the cheeks, that may spread to the pharynx.
- In patients with frequent recurrences or extensive forms invading the esophagus (swallowing difficulty and pain), consider HIV infection.

Treatment

nystatin oral suspension for 7 days

Children and adults: 400 000 IU daily, i.e. 1 ml of the oral suspension (100 000 IU) 4 times daily
or

miconazole oral gel for 7 days

Children 6 months to 2 years: 1.25 ml 4 times daily

Children over 2 years and adults: 2.5 ml 4 times daily

Apply the oral suspension of nystatin or the oral gel of miconazole between meals; keep in the mouth for 2 to 3 minutes, then swallow. In young children, apply to the tongue and inside of each cheek.

Show the mother how to treat since, in most cases, candidiasis will be treated at home.

In immunocompromised patients: see [HIV infection and AIDS](#), Chapter 8.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Oral herpes

Infection due to the herpes simplex virus. Primary infection typically occurs in children aged 6 months to 5 years and may cause acute gingivostomatitis, sometimes severe. After primary infection, the virus remains in the body and causes in some individuals periodic recurrences which are usually benign (herpes labialis).

Clinical features

- Primary herpetic gingivostomatitis
Multiple vesicles on the oral mucosa and lips which rupture to form painful, yellowish, at times extensive ulcers.
Local lesions are usually associated with general malaise, regional lymphadenopathy and fever.
- Recurrent herpes labialis
Clusters of vesicles at the junction between the lip and the skin.

In patients with frequent recurrences or extensive forms, consider HIV infection (see [HIV infection and AIDS](#), Chapter 8).

Treatment

Primary herpetic gingivostomatitis

- Treat [pain](#): paracetamol or ibuprofen PO (Chapter 1)
- In the event of severe lesions, inability to drink and significant pain:
 - Admit the child to hospital (high risk of dehydration).
 - If the child presents within the first 96 hours of symptoms onset, **aciclovir** PO for 5 to 7 days:
Children under 2 years: 200 mg 5 times daily
Children 2 years and over and adults: 400 mg 5 times daily
- In the event of secondary bacterial infection: amoxicillin PO 7 days.

In immunocompromised patients: see [HIV infection and AIDS](#), Chapter 8.

Recurrent herpes labialis

Spontaneous resolution within 7 to 10 days. An antiseptic (chlorhexidine or povidone iodine) may be applied; paracetamol PO if necessary.

Both forms of herpes are contagious: do not touch lesions (or wash hands afterwards); avoid oral contact.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Other infectious causes

See [Pharyngitis](#) (Chapter 2), [Diphtheria](#) (Chapter 2), [Measles](#) (Chapter 8).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Stomatitis from scurvy (vitamin C deficiency)

Clinical features

Bleeding gums, associated in infants with lower limb pain caused by subperiosteal haemorrhage. It is common in contexts of poor food quality or in populations completely dependent on food aid (refugee camps).

Treatment

ascorbic acid (vitamin C) PO

The optimal dose has not been established. For information:

Children 1 month to 11 years: 100 mg 3 times daily

Children 12 years and over and adults: 250 mg 3 times daily

or

Children 1 month to 3 years: 100 mg 2 times daily

Children 4 to 11 years: 250 mg 2 times daily

Children 12 years and over and adults: 500 mg 2 times daily

Treatment is administered at least 2 weeks or longer (until symptoms resolve), then preventive treatment is given (children and adults: 50 mg daily as long as the situation requires).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Other lesions resulting from a nutritional deficiency

Other vitamin deficiencies may provoke mouth lesions: angular stomatitis of the lips and glossitis from vitamin B₂ (riboflavin), niacin (see [Pellagra](#), Chapter 4) or vitamin B₆ (pyridoxine) deficiencies.

Iron deficiency may also provoke angular stomatitis (see [Anaemia](#), Chapter 1).

Give the corresponding vitamins at curative doses. Multivitamins are insufficient to treat true vitamin deficiencies.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Chapter 4: Skin diseases

[Dermatology](#)

[Scabies](#)

[Lice \(pediculosis\)](#)

[Superficial fungal infections](#)

[Bacterial skin infections](#)

[Impetigo](#)

[Furuncles and carbuncles](#)

[Erysipelas and cellulitis](#)

[Cutaneous anthrax](#)

[Endemic treponematoses](#)

[Leprosy](#)

[Herpes simplex and herpes zoster](#)

[Herpes simplex](#)

[Herpes zoster \(shingles\)](#)

[Other skin disorders](#)

[Eczema](#)

[Seborrheic dermatitis](#)

[Urticaria](#)

[Pellagra](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Dermatology

Skin diseases, particularly infectious skin diseases, are very common. They must be treated individually or collectively, but must also be considered as indicators of the sanitary condition of a population. A high prevalence of infectious skin diseases may reflect a problem of insufficient water quantity and lack of hygiene in a population.

Dermatological examination

- Observe the type of lesion:
 - **Macule:** flat, non palpable lesion that is different in colour than the surrounding skin
 - **Papule:** small (< 1 cm) slightly elevated, circumscribed, solid lesion
 - **Vesicle** (< 1 cm), **bulla** (> 1 cm): clear fluid-filled blisters
 - **Pustule:** vesicle containing pus
 - **Nodule:** firm, elevated palpable lesion (> 1 cm) that extend into the dermis or subcutaneous tissue
 - **Erosion:** loss of the epidermis that heals without leaving a scar
 - **Excoriation:** erosion caused by scratching
 - **Ulcer:** loss of the epidermis and at least part of the dermis that leaves a scar
 - **Scale:** flake of epidermis that detaches from the skin surface
 - **Crust:** dried serum, blood, or pus on the skin surface
 - **Atrophy:** thinning of the skin
 - **Lichenification:** thickening of the skin with accentuation of normal skin markings
- Look at the distribution of the lesions over the body; observe their arrangement: isolated, clustered, linear, annular (in a ring). Ask if the lesions are itchy.
- Look for a possible cause: insect bites; scabies, lice, other parasitic skin infections; contact with plants, animals, jewellery, detergents, etc.
- Ask about any past or ongoing treatment: topical, oral or parenteral.
- Look for local or regional signs (secondary infection, lymphangitis, adenopathy, erysipelas) and/or systemic signs (fever, septicaemia, secondary focus).
- Consider the sanitary conditions of the family, particularly for contagious skin diseases (scabies, scalp ringworm, lice).
- Check tetanus vaccination status.

Patients with skin disease often present late. At this stage, primary lesions and specific signs may be masked by secondary infection. In these cases, it is necessary to re-examine the patient, after treating the secondary infection, in order to identify and treat the underlying skin disease.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Scabies

Scabies is a cutaneous parasitosis due to the presence of the mite *Sarcoptes scabiei hominis* within the epidermis. It exists in two forms: ordinary scabies, relatively benign and moderately contagious; and crusted scabies, favoured by immune deficiency, extremely contagious and refractory to conventional treatment.

Person to person transmission takes place chiefly through direct skin contact, and sometimes by indirect contact (sharing clothing, bedding). The challenge in management is that it must include simultaneous treatment of both the patient and close contacts, and at the same time, decontamination of clothing and bedding of all persons undergoing treatment, in order to break the transmission cycle.

Clinical features

Ordinary scabies

In older children and adults

- Itching, worse at night, very suggestive of scabies if close contacts have the same symptom and
 - Typical skin lesions:
 - Scabies burrows (common): fine wavy lines of 5 to 15 mm, corresponding to the tunnels made by the parasite within the skin. Burrows are most often seen in the interdigital spaces of the hand and flexor aspect of the wrist, but may be present on the areolae, buttocks, elbows, axillae. The back and the face are spared. Burrows may be associated with vesicles, corresponding to the entry point of the parasite in the skin.
 - Scabies nodules (less common): reddish-brown nodules, measuring 2 to 20 mm, on the genitals in men, persisting after effective treatment (they are not necessarily indicative of active infection).
- and/or
- Secondary skin lesions: resulting from scratching (excoriations, crusts) or super-infection (impetigo).

Typical lesions and secondary lesions may co-exist, or specific lesions may be entirely masked by secondary lesions.

In infants and young children

- Vesicular eruption; often involving palms and soles, back, face, and limbs. Secondary infection or eczematisation is frequent. Isolated scabies nodules in the axillae may be the only manifestation.
- Examination of the mother's hands may support the diagnosis.

Crusted (Norwegian) scabies

Thick, scaly, erythematous plaques, generalised or localised, resembling psoriasis, with or without itching (50% of cases). Delay in diagnosis may lead to a scabies epidemic.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Topical treatment

Topical scabicides are applied over the entire body (including the scalp, post-auricular areas, umbilicus, palms and soles), avoiding mucous membranes and face, and the breasts in breastfeeding women. Particular attention should be paid to common infestation sites. The recommended contact time should not be shortened or exceeded; the patient must not wash his hands while the product is in use (or the product should be reapplied if the hands are washed). In children under 2 years, the hands must be wrapped to prevent accidental ingestion of the product and contact with eyes. Topical scabicides should not be applied to broken or inflamed skin. Treatment of secondary bacterial infection, if present, should be initiated 24 to 48 hours before use of topical scabicides (see [Impetigo](#)).

The preferred treatment is **5% permethrin** cream:

Children 2 months and over and adults: one application, with a contact time of 8 hours, then rinse thoroughly. Repeat the application after 7 days.

or, if not available, **25% benzyl benzoate** lotion:

See the table below for dilution (depending on age), contact time and number of applications.

	Children < 2 years	Children 2 to 12 years	Children > 12 years and adults	Pregnant women
Dilution	Lotion must be diluted before use: 1 part 25% lotion + 3 parts water	Lotion must be diluted before use: 1 part 25% lotion + 1 part water	Use undiluted 25% lotion	Use undiluted 25% lotion
Contact time	12 hours (6 hours for infants < 6 months) then rinse thoroughly	24 hours then rinse thoroughly	24 hours then rinse thoroughly	12 hours then rinse thoroughly
Number of applications	One application	Two applications (e.g. 24 hours apart, with a rinse between the 2 applications; or 2 successive applications, 10 minutes apart, when the first application has dried with a rinse after 24 hours)	One application	

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Weight	15 to 24 kg	25 to 35 kg	36 to 50 kg	51 to 65 kg
Ivermectin 3 mg tab	1 tab	2 tab	3 tab	4 tab

Treatment effectiveness is judged on clinical grounds. Itching may persist for 1 to 3 weeks after elimination of the parasite.

Persistence of typical burrows beyond 4 weeks should lead to suspicion of treatment failure (insufficient treatment, e.g. the scalp was not included in topical treatment or the patient washed his hands during the treatment period), or early re-infestation (contacts and environment not treated). In these cases, patient and contacts should be retreated.

Persistent itching may be due to another condition, initially masked by scabies.

Crusted scabies

Treatment combines simultaneous administration of oral ivermectin and topical scabicide at regular intervals, e.g. every week for 2 to 3 weeks or more, according to severity and clinical response.

Crusts should be softened (salicylic acid ointment) and removed before applying local treatment (otherwise, local treatment is ineffective).

As exfoliated skin scales may spread the parasite, the patient should be isolated during the treatment, staff should use protection (gloves, gowns and hand washing after contact), and environment (bedding, floors and surfaces) should be decontaminated.

Footnotes

- (a) Treatment with ivermectin in these patients is reserved for severe cases for which no alternative exists (see [Crusted scabies](#)).
- (b) In areas where loiasis is endemic, certain precautions are recommended before administering ivermectin: e.g. measure the Loa loa microfilaraemia, if possible, or ensure that the patient has no history of loiasis (migration of an adult worm under the conjunctiva or transient « Calabar » swellings), nor history of severe adverse reactions following a previous treatment with ivermectin, or if in doubt, use topical treatment in preference to oral.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Lice (pediculosis)

Pediculosis is a benign contagious parasitic infection due to 3 species of lice specific to humans: head lice, body lice and pubic lice. Transmission from person to person occurs through direct or indirect contact.

Body lice are potential vectors of [relapsing fever](#) (Chapter 7), typhus ([Eruptive rickettsioses](#), Chapter 7) and trench fever.

Clinical features

- Head lice mainly affect children: itching and scratch marks (nape of neck and around the ears), which may become secondarily infected (impetigo) in prolonged infestation; presence of live lice and/or live (shiny, grey) nits attached to the hair shaft within 5 mm of the scalp.
- Body lice mainly affect populations living under poor conditions (refugees, prisoners, the homeless): itching and scratch marks (back, belt line and armpits), often inflamed and infected; presence of lice and nits in the clothing (parasites are not found on the body).
- Pubic lice are considered to be a sexually transmitted infection (STI): itching and scratch marks (pubic and perianal area), but other hairy areas may also be affected (armpits, thighs, eyelashes); lice and nits at the base of the hair shaft, rarely visible.
- Examine contacts; check for associated systemic infection (body lice) or STI (pubic lice).

Treatment

Head lice

- Apply lotion to scalp and dry hair, paying particular attention to the areas behind the ears and around the nape of the neck. Do not reduce or exceed the recommended duration of application.

4% dimeticone lotion

Children 6 months and over and adults: leave on hair for 8 hours, then rinse thoroughly.

Keep away from flames and/or intense heat sources (including cigarettes) during application and until rinsing (risk of ignition).

or, if dimeticone is not available or in children 2 to 6 months:

1% permethrin lotion.

Children 2 months and over and adults: leave on hair for 10 minutes, then rinse thoroughly.

- Repeat application of either treatment after 7 days.
- Decontaminate combs, headwear and bedding (wash $\geq 60^{\circ}\text{C}$ /30 minutes, iron or dry in the sun or, if not feasible, seal in a plastic bag for 2 weeks).
- Treat as above contacts with live lice and/or live nits. Do not treat those with dead nits alone (dull, white, $> 1\text{ cm}$ from scalp).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Individual treatment

Disinfection of clothing and bedding as above or as for head lice.

Pubic lice

Shave and/or apply **1% permethrin** lotion to hairy areas (as for head lice). Treat the partner at the same time.

Decontaminate clothing and bedding (as for head lice). Repeat the application after 7 days.

Treatment of secondary bacterial infection, if present, should begin 24 to 48 hours before local antiparasitic treatment (see [Impetigo](#)); local treatment is applied later when tolerated.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Superficial fungal infections

Superficial fungal infections are benign infections of the skin, scalp and nails caused by *Candida albicans* or dermatophytes.

Clinical features and treatment

Candidiasis

Candidal diaper dermatitis

Erythema of the perianal area with peripheral desquamation and sometimes pustules. Secondary infection may develop.

- Buttocks must be kept clean (ordinary soap and water) and dry.
- Avoid humidity: according to the context, expose the buttocks to air or change diapers more frequently; remove plastic pants.
- Protect the skin with **zinc oxide ointment** if diarrhoea is present.
- If diaper dermatitis is severe and persistent despite these measures, consider an intestinal infection (**nystatin PO: 100 000 IU 4 times daily for 20 days**).

Other candidiasis

- Candidiasis of skin folds: **miconazole 2%** cream, one application 2 times daily for 2 to 4 weeks
- Oral candidiasis: see [Stomatitis](#), Chapter 3.
- Vulvovaginal candidiasis: see [Abnormal vaginal discharge](#), Chapter 9.

Dermatophytoses

Dermatophytes cause various clinical lesions, depending on the anatomic site involved: scalp, glabrous (hairless) skin, folds or nails.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Anatomic site ^(a)	Clinical features	Treatment
Scalp Scalp ringworm <i>Tinea capitis</i>	<p>Common in children. Depending on the species:</p> <ul style="list-style-type: none"> • One or more round, scaly, erythematous plaques with the ends of broken hairs. • Inflammation, suppuration, crusting and peripheral lymphadenopathy (kerion). • Permanent hair loss (favus). <p>Some scalp ringworms are contagious: simultaneously examine (and treat) symptomatic contacts.</p>	<ul style="list-style-type: none"> • Shave or cut hair short on and around the lesions. • Local treatment: 2 times daily, clean with soap and water, dry and apply miconazole 2% cream or Whitfield's ointment for 2 weeks or longer if necessary. • Administer systemic treatment as local treatment alone does not cure scalp ringworm: griseofulvin PO for 6 weeks minimum (up to 8 to 12 weeks) Children 1 to 12 years: 10 to 20 mg/kg once daily (max. 500 mg daily) Children ≥ 12 years and adults: 500 mg to 1 g once daily, depending on severity or itraconazole PO Children: 3 to 5 mg/kg once daily for 4 to 6 weeks (max. 200 mg daily) Adults: 200 mg once daily for 2 to 4 weeks • Suppurative lesions: treat superinfection (see <u>Impetigo</u>) before applying local antifungal treatment. • For painful kerion: paracetamol PO. <i>In pregnant/lactating/breastfeeding women:</i> oral antifungals are contraindicated. Apply a topical treatment (miconazole 2% cream or Whitfield's ointment) to limit the spread of infection until it is possible to treat orally.
Glabrous skin Ringworm of the	Erythematous, scaly, pruritic macule with a well-demarcated, raised, vesicular border and central	<ul style="list-style-type: none"> • For non widespread, localised tinea: Local treatment: 2 times daily, clean

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<i>Tinea pedis</i> (athlete's foot)	Pruritus, fissure and whitish scales in the 3 rd and/or 4 th interdigital spaces ^(b) .	
<i>Tinea cruris</i>	<ul style="list-style-type: none"> • Groin (<i>Tinea cruris</i>): Circumscribed, pruritic, erythematous plaque, with a pale centre surrounded by vesiculo- pustules, extending outward from the groin. 	lesions, use miconazole 2% cream only (do not use Whitfield's ointment).

- (a) Dermatophytosis may affect the nails (*Tinea unguium*, onychomycosis). Treatment is prolonged (12 to 18 months with griseofulvin) thus, in practice, difficult. Failures and relapses are frequent.
- (b) In candidal intertrigo, lesions are usually located in the 1st and 2nd interdigital spaces.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Bacterial skin infections

- [Impetigo](#)
- [Furuncles and carbuncles](#)
- [Erysipelas and cellulitis](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Impetigo

Impetigo is a benign, contagious infection of the epidermis due to group A β -haemolytic streptococcus and *Staphylococcus aureus*. Co-infection is common. Transmission is by direct contact. Lack of water, and poor hygiene, increase spread.

Primary infections are most common in children. Secondary infections complicating preexisting pruritic dermatoses (lice, scabies, eczema, herpes, chickenpox, etc.) are more common in adults.

Clinical features

- Non bullous impetigo (classic form): flaccid vesicle on erythematous skin which becomes pustular and forms a yellowish crust. Different stages of the infection may be present simultaneously. The lesion does not leave a scar. The most common sites of infection are around the nose and mouth, on the limbs or on the scalp.
- Bullous impetigo: large flaccid bullae and erosions of the skin in the ano-genital region in newborns and infants.
- Ecthyma: an ulcerative form of impetigo that leaves scars. This form is most common in the immunocompromised (e.g. HIV infection, malnutrition), diabetics and alcoholics.
- Regardless of the type of impetigo: absence of fever or systemic signs.
- Possible complications:
 - abscess, pyodermitis, cellulitis, lymphangitis, osteomyelitis, septicaemia;
 - acute glomerulonephritis (routinely look for signs of glomerulonephritis).

Treatment

- **Localised non bullous impetigo** (max. 5 lesions in a single skin area):
 - Clean with soap and water and dry before applying mupirocin.
 - **2% mupirocin** ointment: one application 3 times daily for 7 days. Reassess after 3 days. If there is no response, switch to oral antibiotic therapy (see below).
 - Keep fingernails short. Avoid touching the lesions, keep them covered with gauze if possible.
- **Extensive non bullous impetigo** (more than 5 lesions or impetigo involving more than one skin area), bullous impetigo, ecthyma, impetigo with abscess; immunocompromised patient; topical treatment failure:
 - Clean with soap and water and dry 2 to 3 times daily.
 - Keep fingernails short. Avoid touching the lesions, keep them covered with gauze if possible.
 - Incise abscesses if present.
 - Administer oral antibiotic therapy^a:
cefalexin PO for 7 days
Neonates under 7 days: 25 mg/kg 2 times daily
Neonates 7 to 28 days: 25 mg/kg 3 times daily
Children 1 month to 12 years: 25 mg/kg 2 times daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Look for and treat any underlying dermatosis: [lice](#), [scabies](#), [eczema](#), [herpes](#), [scalp ringworm](#), or an ENT infection.
- Trace and treat contacts.
- Check for proteinuria (use urine dipstick) 3 weeks after the infection.

Footnotes

- (a) In penicillin-allergic patients only (resistance to macrolides is common), **azithromycin** PO for 3 days (children: 10 mg/kg once daily; adults: 500 mg once daily).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Furuncles and carbuncles

Necrotising perifollicular infection, usually due to *Staphylococcus aureus*. Risk factors include: nasal carriage of *S. aureus*, maceration, breaks in the skin, poor hygiene; diabetes mellitus, malnutrition, iron deficiency or immunodeficiency.

Clinical features

- Furuncle: red, warm, painful nodule with a central pustule, usually around a hair follicle. It becomes fluctuant, discharges a core of purulent exudate, and leaves a depressed scar. It occurs most frequently on the thighs, groin, buttocks, armpits, neck and back. There is no fever.
- Carbuncle: a cluster of interconnected furuncles, sometimes with fever and peripheral lymphadenopathy. It leaves a depressed scar.

Treatment

- Single furuncle:
 - Clean with soap and water 2 times daily and cover with a dry dressing.
 - Apply warm moist compresses to the furuncle in order to encourage it to drain.
 - After drainage, clean and apply a dry dressing until the lesion has healed.
- Furuncle on the face, multiple furuncles, carbuncles or in immunocompromised patients:
 - Same local care.
 - Add systematically an antibiotic for 7 days^a:
cefalexin PO
Neonates under 7 days: 25 mg/kg 2 times daily
Neonates 7 to 28 days: 25 mg/kg 3 times daily
Children 1 month to 12 years: 25 mg/kg 2 times daily
Children 12 years and over and adults: 1 g 2 times daily
or
amoxicillin/clavulanic acid (co-amoxiclav) PO. Use formulations in a ratio of 8:1 or 7:1. The dose is expressed in amoxicillin:
Children < 40 kg: 25 mg/kg 2 times daily
Children ≥ 40 kg and adults:
Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)
Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)
- In all cases: wash hand frequently, wash bedding.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Erysipelas and cellulitis

Last updated: October 2020

Acute skin infections, due to bacteria (usually Group A beta-haemolytic streptococcus and sometimes *Staphylococcus aureus*, including methicillin resistant *S. aureus*–MRSA) that enter through a break in the skin. The main risk factors are: venous insufficiency, obesity, oedema or lymphoedema, history of erysipelas or cellulitis, immunosuppression and cutaneous inflammation (e.g. dermatosis, wound). Erysipelas is a superficial infection (affecting the dermis and superficial lymph vessels), while cellulitis affects the deeper tissues (deep dermis layers and subcutaneous fat). Generally, these infections affect the lower extremities and sometimes the face. If the orbital and periorbital tissues are infected, see [Periorbital and orbital cellulitis](#), Chapter 5. If the infection is perifollicular, see [Furuncles and carbuncles](#), Chapter 4.

Clinical signs

- Warm, tender, swollen well-demarcated erythematous plaque.
- Fever, lymphadenopathy and lymphangitis.
- Look for a portal of entry (bite, ulcer, wound, intertrigo, eczema, fungal infection, etc.).
- In case of intense pain disproportionate to the skin lesion, hypoesthesia, rapidly progressing local signs, crepitation, skin necrosis or critically ill appearing patient, consider necrotising fasciitis that is a surgical emergency (see [Necrotising infections of the skin and soft tissues](#), Chapter 10).
- Other complications: septicaemia (see [Septic shock](#), Chapter 1), acute glomerulonephritis, osteomyelitis, septic arthritis.
- The main differential diagnoses include: contact dermatitis, stasis dermatitis due to venous insufficiency, venous thrombosis and erythema migrans characteristic of Lyme disease.

Paraclinical investigations

- Ultrasound: can detect signs of cellulitis and rule out an underlying abscess, deep vein thrombosis or a foreign body.
- Radiography: can detect a foreign body, underlying osteomyelitis (or gas in the subcutaneous tissue in case of a necrotising infection, nevertheless the absence of gas does not rule out this diagnosis).
- Test for proteinuria with urine dipstick 3 weeks after infection to look for glomerulonephritis.

Treatment

- In all cases:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Hospitalize for the following: children younger than 3 months, critically ill appearing patient ^b, local complications, debilitated patient (chronic conditions, the elderly) or if there is a risk of non-compliance with or failure of outpatient treatment. Treat other patients as outpatients.

- Outpatient antibiotic therapy ^c :

cefalexin PO for 7 to 10 days

Children 1 month to under 12 years: 25 mg/kg 2 times daily

Children 12 years and over and adults: 1 g 2 times daily

or

amoxicillin/clavulanic acid (co-amoxiclav) PO for 7 to 10 days.

Use formulations in a ratio of 8:1 or 7:1. The dose is expressed in amoxicillin:

Children < 40 kg: 25 mg/kg 2 times daily

Children ≥ 40 kg and adults:

Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)

Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)

In the event of worsening clinical signs after 48 hours of antibiotic treatment, consider IV route.

- Inpatient antibiotic therapy ^d :

▫ First line therapy:

cloxacillin IV infusion over 60 minutes ^e

Children 1 month to under 12 years: 12.5 to 25 mg/kg every 6 hours

Children 12 years and over and adults: 1 g every 6 hours

or

amoxicillin/clavulanic acid (co-amoxiclav) by slow IV injection (3 minutes) or IV infusion (30 minutes). The dose is expressed in amoxicillin:

Children under 3 months: 30 mg/kg every 12 hours

Children 3 months and over: 20 to 30 mg/kg every 8 hours (max. 3 g daily)

Adults: 1 g every 8 hours

If there is clinical improvement after 48 hours (afebrile and erythema and oedema have improved) switch to cefalexin or amoxicillin/clavulanic acid PO at the doses indicated above to complete 7 to 10 days of treatment.

▫ If there is no clinical improvement after 48 hours, consider MRSA:

clindamycin IV infusion over 30 minutes ^f

Children 1 month and over: 10 mg/kg every 8 hours

Adults: 600 mg every 8 hours

After 48 hours, change to clindamycin PO at the doses indicated above to complete 7 to 10 days of treatment.

Footnotes

(a) The erythema will regress if the treatment is effective. If the erythema spreads consider a treatment failure (MRSA or a

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

- (f) Dilute each dose of clindamycin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Cutaneous anthrax

Last updated: September 2022

Anthrax is caused by the bacterium *Bacillus anthracis* that primarily affects herbivores (sheep, goats, cows, camels, horses, etc.). Humans may become infected through contact of broken skin with a dead or sick animal. People at risk include livestock farmers and those that manipulate skins, wool or carcasses of infected animals.

The disease is found in Eastern Europe, Central Asia, the Mediterranean Basin, Africa and South America. Pulmonary (acquired by inhalation) and intestinal (acquired by eating infected meat) forms also exist.

Clinical features

- Papule, then pruritic vesicle on uncovered skin surfaces (face, neck, arms, legs). The vesicle ulcerates and becomes a painless black eschar surrounded by oedema, often associated with lymphangitis and regional lymphadenopathy.
- The following are criteria of severity:
 - lesion located on the head or neck, or
 - presence of systemic symptoms (fever, malaise, headache, tachycardia, tachypnoea, hypotension, hyper/hypothermia), or
 - presence of extensive oedema, or
 - multiple, extensive or bullous lesions.

Laboratory

- From vesicular fluid^a: culture and drug susceptibility testing (rarely available) or Gram stain for microscopic examination.
- PCR testing (reference laboratory).

Treatment

Uncomplicated cutaneous anthrax

- Do not excise the eschar; daily dry dressings.
- Antibiotic treatment for 7 to 10 days:
 - First-line antibiotics:
ciprofloxacin PO (including in pregnant or breastfeeding women and children)
Children: 15 mg/kg (max. 500 mg) 2 times daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

amoxicillin PO, if penicillins are effective (documented susceptibility)

Children: 30 mg/kg (max. 1 g) 3 times daily

Adults: 1 g 3 times daily

Severe cutaneous anthrax

- Combined antibiotic treatment for 14 days:



Do not mix the two antibiotics in the same infusion bag (incompatibility).

- First-line:

ciprofloxacin IV infusion over 60 minutes^b

Children: 10 mg/kg (max. 400 mg) every 8 hours

Adults: 400 mg every 8 hours

+ **clindamycin** IV infusion over 30 minutes^b

Children 1 month and over: 10 to 13 mg/kg (max. 900 mg) every 8 hours

Adults: 900 mg every 8 hours

- Alternative, if penicillins are effective (documented susceptibility):

ampicillin IV infusion over 30 minutes^b

Children 1 month and over: 50 mg/kg (max. 3 g) every 6 hours or 65 mg/kg (max. 4 g) every 8 hours

Adults: 3 g every 6 hours or 4 g every 8 hours

+ **clindamycin** IV infusion as above.

Change to oral treatment as soon as possible to complete 14 days of treatment with ciprofloxacin + clindamycin or amoxicillin + clindamycin as for uncomplicated cutaneous anthrax.

- Intensive care: symptomatic treatment of shock (see [Shock](#), Chapter 1); tracheostomy and ventilatory support may be necessary.

Prevention

- Antibiotic prophylaxis in case of known skin exposure: treat for 10 days PO as for uncomplicated cutaneous anthrax.
- Livestock vaccination; burial or burning of animal carcasses.

Footnotes

(a) Samples can be stored (including transport time) for 7 days max. in cold chain (if not available, at a temperature < 30 °C).

(b) Dilute each dose of ciprofloxacin, clindamycin or ampicillin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and above and in adults. Administer ciprofloxacin more slowly than clindamycin or ampicillin.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Endemic treponematoses

Endemic treponematoses are bacterial infections caused by 3 different types of treponema (other than *Treponema pallidum*). Human-to-human transmission may be direct or indirect.

The 3 endemic treponematoses result in positive syphilis serology (TPHA-VDRL), but these tests are not necessary as diagnosis is clinical. There is no laboratory test that can distinguish between the different treponematoses.

For the diagnosis and treatment of syphilis, see [Genital infections](#), Chapter 9.

Clinical features

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

	Yaws	Pinta	Bejel
Pathogen	<i>Treponema pertenue</i>	<i>Treponema carateum</i>	<i>Treponema pallidum</i> type M
Geographic distribution	Tropical and humid forests	Tropical zones of Latin America	Arid areas, semi-desert of the Middle East and Africa
Population	Children between 4 and 14 years	Children and adults	Nomadic populations, particularly children
First stage	<p>Yaws chancre: skin coloured lesion, non-indurated, itchy, on the lower limbs in 95% of cases, with peripheral adenopathy.</p> <p>Spontaneous healing or development of a large yaw surrounded by smaller yaws.</p>	<p>Annular, erythematous, scaly plaques, usually on uncovered body parts (face, extremities), resemble dermatophytes.</p> <p>Lesions heal spontaneously leaving scars.</p>	<p>Discrete chancre: moist papule, most commonly on the mucous membranes or in dermal folds, with peripheral adenopathy.</p>
Second stage	<p>Lesions appear 3 weeks after the initial chancre, occur in crops and heal spontaneously:</p> <ul style="list-style-type: none"> • Frambesioma (papillomatous lesion, vegetal, very contagious) • Isolated or associated with yaws (round, squamous papules, not very contagious) • Osteoperiostitis of the long bones (phalanges, nasal process of the maxilla, tibia) 	<p>Pintids: plaques of various colours (bluish, reddish, whitish). May occur anywhere on the body.</p>	<ul style="list-style-type: none"> • Mucous patches of the mouth common: very contagious ulcerated, round in form, indurated, with white coating, bleed easily, usually occur on the inside of the lips, cheek and tongue or labial folds • Condyloma in the anogenital region (rare) • Cutaneous lesions are rare: vegetal aspect, in dermal folds • Bone destruction identical to that of yaws, in the legs and forearms
Late stage	After some years of	Symmetrical white patches on	After several years of latency:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment

Yaws

azithromycin PO^[1]

Children and adults: 30 mg/kg single dose (max. 2 g)

or, if not available,

benzathine benzylpenicillin IM^{[2][3]}

Children under 10 years: 1.2 MIU single dose

Children 10 years and over and adults: 2.4 MIU single dose

Pinta and bejel

benzathine benzylpenicillin IM.

As for yaws.

For patients allergic to penicillin:

doxycycline PO (except in children under 8 years and pregnant or lactating women)

Children 8 years and over: 50 mg 2 times daily for 14 days

Adults: 100 mg 2 times daily for 14 days

Notes:

- Antibiotic treatment will cure early stage cases and may relieve the pain of osteitis. It may be ineffective for late stage infections.
- Syphilis serology will remain positive despite clinical cure.

Treatment of contacts and latent cases

The same treatment should be administered to all symptomatic and asymptomatic contacts and to all latent cases (asymptomatic individuals with positive serologic test for syphilis) in endemic zones.

References

1. World Health Organization (2012). Yaws: recognition booklet for communities. Reprinted with changes, 2014. <http://www.who.int/iris/handle/10665/75360> [Accessed 15 May 2018]
2. Oriol Mitjà, David Mabey. Yaws, bejel, and pinta (last updated. May 07, 2018). UpToDate [Accessed 15 May 2018].
3. Michael Marks, Anthony W Solomon, David C Mabey. Endemic treponemal diseases. Transactions of The Royal Society of Tropical Medicine and Hygiene, Volume 108, Issue 10, 1 October 2014, Pages 601–607. <https://doi.org/10.1093/trstmh/tru128> [Accessed 15 May 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Leprosy

Leprosy is a chronic bacterial infection due to *Mycobacterium leprae*.

It is transmitted by frequent close contact, mainly between household members.

It mainly affects young adults. 94% of reported cases globally were in Bangladesh, Brazil, Democratic Republic of Congo, Ethiopia, India, Indonesia, Madagascar, Myanmar, Nepal, Nigeria, the Philippines, Sri Lanka and the United Republic of Tanzania.^[1]

Clinical features

Leprosy should be considered in any patient presenting with:

- Hypopigmented or erythematous skin lesion(s) with partial or complete loss of sensation to touch, pain, heat;
- Infiltrated pigmented nodules, initially with no sensory loss, on the face, ear lobes and the upper and lower limbs;
- Tender, infiltrated and hypertrophied peripheral nerve (ulnar, radial, median, popliteal, tibial etc.) with possible paraesthesia of the extremities, trophic changes (perforating ulcer of the foot) or paralysis (steppage gait, deformities of hands and feet, facial nerve paralysis).

There are different clinical forms and classification systems of leprosy.

Ridley-Jopling classification

This classification differentiates 5 forms based on the bacteriological index. These forms correlate with the immunological response to *M. leprae*. Patients with tuberculoid leprosy (TT) are resistant to the bacillus and infection is localised. Patients with lepromatous leprosy (LL) are extremely sensitive to the bacillus and the infection is disseminated. Borderline forms (BT, BB, BL) are between the two ends of the spectrum (TT and LL).

Paucibacillary forms (least contagious forms)		Multibacillary forms (most contagious forms)		
Tuberculoid	Borderline Tuberculoid	Borderline	Borderline Lepromatous	Lepromatous
T.T.	B.T.	B.B.	B.L.	L.L.

WHO classification

In order to simplify diagnosis and to promote rapid implementation of treatment, the WHO has simplified clinical classification of leprosy and differentiates only 2 forms:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment

Countries where leprosy is endemic have a control programme. Check national recommendations.

First-line treatment regimens recommended by the WHO

Age	Multibacillary leprosy (more than 5 skin lesions)	Paucibacillary leprosy (1 to 5 skin lesions)
Children 10 to 14 years	rifampicin PO: 450 mg once monthly + clofazimine PO: 150 mg once monthly and 50 mg on alternate days + dapsone PO: 50 mg once daily	rifampicin PO: 450 mg once monthly + clofazimine PO: 150 mg once monthly and 50 mg on alternate days + dapsone PO: 50 mg once daily
Children 15 years and over and adults	rifampicin PO: 600 mg once monthly + clofazimine PO: 300 mg once monthly and 50 mg once daily + dapsone PO: 100 mg once daily	rifampicin PO: 600 mg once monthly + clofazimine PO: 300 mg once monthly and 50 mg once daily + dapsone PO: 100 mg once daily
Duration	12 months	6 months

Note: the monthly doses of rifampicin and clofazimine are administered under direct observation by medical staff whereas the daily doses of clofazimine and dapsone are taken by the patient at home. Rifampicin should be taken on an empty stomach to improve absorption.

Teach the patient to recognise and quickly report a lepra reaction or relapse in order to modify or restart treatment.

Leprosy reactions

These reactions usually occur during the course of treatment in patients with multibacillary leprosy (BL and LL). They are associated with the immunological response to *M. leprae* antigens. Urgent treatment is required to avoid irreversible disability. Do not interrupt ongoing leprosy treatment.

Clinical features

- Reversal reactions:
 - Exacerbation of the skin lesions that become erythematous and oedematous and risk or ulceration. Onset or worsening of numbness of skin lesions;
 - Onset of acute painful hypertrophic neuritis.
- Erythema nodosum leprosum:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Week 3 and 4: 30 mg once daily
Week 5 and 6: 20 mg once daily
Week 7 and 8: 15 mg once daily
Week 9 and 10: 10 mg once daily
Week 11 and 12: 5 mg once daily

- Erythema nodosum leprosum:
 - **prednisolone (or prednisone)** PO as for reversal reactions, for 3 months.^[2]
 - Fever: paracetamol PO (see [Fever](#), Chapter 1)

References

1. World Health Organization. Global Leprosy Programme. Global leprosy strategy 2016-2020. Accelerating towards a leprosy-free world, 2016.
http://apps.who.int/iris/bitstream/handle/10665/208824/9789290225096_en.pdf?sequence=14&isAllowed=y [Accessed 17 October 2018]
2. World Health Organization. WHO Expert Committee on Leprosy. Eighth report. WHO technical report series, n° 968. Geneva, 2012.
http://www.searo.who.int/entity/global_leprosy_programme/publications/8th_expert_comm_2012.pdf [Accessed 17 October 2018]
3. World Health Organization. A guide to eliminating leprosy as a public health problem. Leprosy Elimination Group, 2000.
http://apps.who.int/iris/bitstream/handle/10665/66612/WHO_CDS_CPE_2000.14.pdf?sequence=1 [Accessed 17 October 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Herpes simplex and herpes zoster

- [Herpes simplex](#)
- [Herpes zoster \(shingles\)](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Herpes simplex

Recurrent viral infection of the skin and mucous membranes due to the *Herpes simplex* virus. Recurrent lesions have a different presentation than primary infection.

Clinical features

- Recurrent herpes labialis: tingling feeling followed by an eruption of vesicles on an erythematous base, located on the lips ('fever blisters') and around the mouth, they may extend onto the face. Recurrence corresponds to a reactivation of the latent virus after a primary infection. No associated malaise, adenopathy or fever.
- Carefully consider other sites: buccal ([Stomatitis](#), Chapter 3), genital ([Genital ulcers](#), Chapter 9), ophthalmic, and secondary bacterial infections.

Treatment

- Clean with soap and water 2 times daily until the lesions have healed.
- For patients with secondary bacterial infections: antibiotic treatment as for [impetigo](#).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Herpes zoster (shingles)

Acute viral infection due to the varicella-zoster virus. Chickenpox is the primary infection and herpes zoster the reactivation of the latent virus.

Clinical features

- Unilateral neuralgic pain followed by an eruption of vesicles on a erythematous base, that follow the distribution of a nerve pathway.
- Lesions most commonly occur on the thorax, but herpes zoster may also develop on the face with a risk of ophthalmic complications.
- Herpes zoster is more common in adults than in children.

Treatment

- Similar to that of herpes simplex, with the addition of systematic analgesics: paracetamol PO (see [Pain](#), Chapter 1).
- Aciclovir PO given within the first 48 hours after the eruption of lesions is only indicated for severe forms: necrotic or extensive lesions or lesion on the face which may spread to the eyes (see [HIV infection and AIDS](#), Chapter 8).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Other skin disorders

- [Eczema](#)
- [Seborrheic dermatitis](#)
- [Urticaria](#)
- [Pellagra](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Eczema

Acute eczema: erythematous plaque, pruritic, vesicular, oozing, with poorly demarcated and crumbly borders.

Chronic eczema: erythematous plaque, scaly, dry, poorly demarcated and pruritic.

Look for a cause (contact allergic dermatitis, fungal or bacterial infection with a distant focus, malnutrition) and ask about family history.

Treatment

- Clean with soap and water 2 times daily.
- Then:
 - for acute eczema: **calamine** lotion, one application 2 times daily
 - for chronic eczema: **zinc oxide** ointment, one application 2 times daily
- Look for and treat any pre-existing condition (scabies, lice etc.).
- For patients with secondary infections: treat as [impetigo](#).
- For patients with intense pruritus, antihistamines for a few days (see [Urticaria](#)).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Seborrheic dermatitis

Seborrheic dermatitis is an inflammatory chronic dermatosis that can be localized on areas rich with sebaceous glands. This dermatosis is more common in infected patients with HIV.

Clinical features

- Erythematous plaques covered by greasy yellow scales that can be localized on the scalp, the face (nose wings, eyebrows, edge of the eyelids), sternum, spine, perineum, and skin folds.

Treatment

- Clean with soap and water 2 times daily; shampooing the scalp.
- **Hydrocortisone 1%** cream: one application once daily or 2 times daily to the affected area only, in thin layer, for 7 days maximum
- Do not apply if pre-existing bacterial infection; treat first the infection (see [Impetigo](#)).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Urticaria

Last updated: July 2022

Papules: transient, erythematous, oedematous, pruritic, resembling nettle stings.

Look for a cause: food or drug (particularly antibiotic) allergy, insect bites; the invasive stage of a bacterial or parasitic infection (ascariasis, strongyloidiasis, ancylostomiasis, schistosomiasis, loiasis), viral infection (hepatitis B or C); generalised disease (cancer, lupus, dysthyroidism, vasculitis).

Treatment

- If the pruritus is intense, antihistamines for a few days:

loratadine PO

Children over 2 years and under 30 kg: 5 mg (5 ml) once daily

Children over 30 kg and adults: 10 mg (1 tab) once daily

- In the event of anaphylactic reaction, see [Shock](#) (Chapter 1).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Pellagra

Pellagra is a dermatitis resulting from niacin and/or tryptophane deficiency (in persons whose staple food is sorghum; patients with malabsorption, or during famine).

Clinical features

Classically, disease of the 'three Ds': dermatitis, diarrhoea and dementia.

- Dark red plaques, well demarcated, symmetric, located on exposed areas of the body (forehead, neck, forearms, legs). The skin becomes very scaly, pigmented, sometimes with haemorrhagic bullae.
- Gastrointestinal (glossitis, stomatitis and diarrhoea) and neuropsychiatric symptoms are seen in more serious forms.

Treatment

- **nicotinamide (vitamin PP) PO^[1]**

Children and adults: 100 mg 3 times daily, give with a diet rich in protein until the patient is fully cured.

- In the event of an epidemic of pellagra, for example in a refugee camp, it is vital that the food ration be modified (add groundnuts or dry vegetables) in order to meet the daily requirements (approximately 15 mg daily for adults).

References

1. World Health Organization, United Nations High Commissions for Refugees. Pellagra and its prevention and control in major emergencies. World Health Organization, 2000.
http://www.who.int/nutrition/publications/en/pellagra_prevention_control.pdf [Accessed 23 May 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 5: Eye diseases

[Xerophthalmia \(vitamin A deficiency\)](#)

[Conjunctivitis](#)

[Neonatal conjunctivitis](#)

[Viral epidemic keratoconjunctivitis](#)

[Trachoma](#)

[Periorbital and orbital cellulitis](#)

[Other pathologies](#)

[Onchocerciasis \(river blindness\)](#)

[Loiasis](#)

[Pterygium](#)

[Cataract](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Xerophthalmia (vitamin A deficiency)

The term xerophthalmia covers all the ocular manifestations of vitamin A deficiency. Xerophthalmia can progress to irreversible blindness if left untreated.

In endemic areas, vitamin A deficiency and xerophthalmia affect mainly children (particularly those suffering from malnutrition or measles) and pregnant women.

Disorders due to vitamin A deficiency can be prevented by the routine administration of retinol.

Clinical features

- The first sign is hemeralopia (crepuscular blindness): the child cannot see in dim light, may bump into objects and/or show decreased mobility.
- Then, other signs appear gradually:
 - Conjunctival xerosis: bulbar conjunctiva appears dry, dull, thick, wrinkled and insensitive
 - Bitot's spots: greyish foamy patches on the bulbar conjunctiva, usually in both eyes (specific sign, however not always present)
 - Corneal xerosis: cornea appears dry and dull
 - Corneal ulcerations
 - Keratomalacia (the last and most severe sign of xerophthalmia): softening of the cornea, followed by perforation of the eyeball and blindness (extreme care must be taken during ophthalmic examination due to risk of rupturing cornea)

Treatment

Treat early symptoms to avoid the development of severe complications. Vision can be saved provided that ulcerations affect less than a third of the cornea and the pupil is spared. Even if deficiency has already led to keratomalacia and irreversible loss of sight, it is imperative to administer treatment, in order to save the other eye and the life of the patient.

retinol (vitamin A) PO:

- Treatment is the same regardless of the clinical stage, except in pregnant women.

Age	200 000 IU capsule ^(a)
Children < 6 months ^(b)	50 000 IU (2 drops) once daily on D1, D2 and D8
Children 6 months to < 1 year	100 000 IU (4 drops) once daily on D1, D2 and D8

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- If the cornea is affected, risk of blindness outweighs teratogenic risk. Administer 200 000 IU once daily on D1, D2 and D8.

Corneal lesions are a medical emergency. In addition to the immediate administration of retinol, treat or prevent secondary bacterial infections with **1% tetracycline eye ointment**, one application 2 times daily (do not apply eye drops containing corticosteroids) and protect the eye with an eye-pad after each application.

Prevention

- Systematically administer retinol PO to children suffering from measles (one dose on D1 and D2).
- In areas where vitamin A deficiency is endemic^a, routine supplementation of **retinol** PO:

Age	200 000 IU capsule^(c)
Children < 6 months	50 000 IU (2 drops) single dose
Children 6 months to < 1 year	100 000 IU (4 drops) every 4 to 6 months
Children 1 to < 5 years	200 000 IU (one capsule) every 4 to 6 months
Women after delivery	200 000 IU (one capsule) single dose

(c) Capsules must not be swallowed whole. Cut the end of the capsule and deliver the dose directly into the mouth.

To avoid excessive dosage, record any doses administered on the health/immunisation card and do not exceed indicated doses. Vitamin A overdose may cause raised intracranial pressure (bulging fontanelle in infants; headache, nausea, vomiting) and, in severe cases, impaired consciousness and convulsions. These adverse effects are transient; they require medical surveillance and symptomatic treatment if needed.

Footnotes

- (a) For more information country-specific prevalence of vitamin A deficiency, see:
<https://www.thelancet.com/action/showPdf?pii=S2214-109X%2815%2900039-X>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Conjunctivitis

- [Neonatal conjunctivitis](#)
- [Viral epidemic keratoconjunctivitis](#)

Conjunctivitis is an acute inflammation of the conjunctiva due to a bacterial or viral infection, allergy, or irritation.

Conjunctivitis may be associated with measles or rhinopharyngitis in children.

In the absence of hygiene and effective treatment, secondary bacterial infections may develop, affecting the cornea (keratitis).

Clinical features

- Clinical signs of all conjunctivitis include: redness of the eye and irritation. Visual acuity is not affected.
- Depending on the cause:
 - abundant and purulent secretions, eyelids stuck together on waking, unilateral infection at onset: bacterial conjunctivitis;
 - watery (serous) secretions, no itching: viral conjunctivitis;
 - excessive lacrimation, eyelid oedema, intense itching: allergic conjunctivitis.
- In endemic areas, turn both upper eyelids up to check for signs of trachoma (see [Trachoma](#)).
- Suspect keratitis if patient reports intense pain (more than is usually associated with conjunctivitis) and photophobia. Instill one drop of **0.5% fluorescein** to check for possible ulcerations.
- Always check for foreign bodies (subconjunctival or corneal) and remove after administering **0.4% oxybuprocaine** anaesthetic eye drops. Never give bottle of eye drops to the patient.

Treatment

Bacterial conjunctivitis

- Clean eyes 4 times daily with boiled water or 0.9% sodium chloride.
- Apply into both eyes **1% tetracycline eye ointment**: one application 2 times daily for 7 days
- Never use corticosteroid drops or ointment.

Viral conjunctivitis

- Clean eyes 4 times daily with boiled water or 0.9% sodium chloride.
- Apply local antibiotics if there is a (risk of) secondary bacterial infection (see above).

Allergic conjunctivitis

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Neonatal conjunctivitis

Conjunctivitis due to *Neisseria gonorrhoeae* and/or *Chlamydia trachomatis* in neonates born to mothers with genital gonococcal and/or chlamydial infections at the time of delivery.

Neonatal conjunctivitis is a medical emergency. Without prompt treatment, risk of corneal lesions and visual impairment.

Clinical features

- Unilateral or bilateral purulent conjunctivitis in the first 28 days of life.

Treatment

- Clean eyes with isotonic sterile solution (0.9% sodium chloride or Ringer lactate) 4 times daily to remove secretions.
- Antibiotic treatment:
 - for all neonates with conjunctivitis in the first 28 days of life
 - for all neonates born to mothers with a genital infection (purulent cervical discharge) at the time of delivery

	0 to 7 days	8 to 28 days
First line	ceftriaxone IM: 50 mg/kg single dose (max. 125 mg)	ceftriaxone IM: 50 mg/kg single dose (max. 125 mg) + azithromycin PO: 20 mg/kg once daily for 3 days
Alternatives	If ceftriaxone contra-indicated: cefotaxime IM: 100 mg/kg single dose	If azithromycin unavailable: erythromycin PO: 12.5 mg/kg 4 times daily for 14 days

If symptoms persist 48 hours after parenteral treatment alone, administer **azithromycin** PO (or **erythromycin** PO as above).

Notes:

- When systemic treatment is not immediately available, clean both eyes and apply 1% tetracycline eye ointment every hour, until systemic treatment is available.
- In all cases, treat the genital infection of the mother and partner (see [Genital infections](#), Chapter 9).
- Azithromycin and erythromycin are associated with an increased risk of pyloric stenosis in neonates. The risk is [increased](#).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

1. Lund M et al. Use of macrolides in mother and child and risk of infantile hypertrophic pyloric stenosis: nationwide cohort study. *BMJ*. 2014; 348: g1908.
<https://www.bmjjournals.org/content/348/bmj.g1908> [Accessed 16 April 2021]
2. Murchison L et al. Post-natal erythromycin exposure and risk of infantile hypertrophic pyloric stenosis: a systematic review and meta-analysis. *Pediatr Surg Int*. 2016 Dec; 32(12): 1147-1152.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5106491/> [Accessed 16 April 2021]
3. Almaramhy HH et al. The association of prenatal and postnatal macrolide exposure with subsequent development of infantile hypertrophic pyloric stenosis: a systematic review and meta-analysis. *Ital J Pediatr*. 2019 Feb 4; 45(1)20.
<https://ijponline.biomedcentral.com/articles/10.1186/s13052-019-0613-2> [Accessed 16 April 2021]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Viral epidemic keratoconjunctivitis

Corneal and conjunctival lesions

- Treat as viral conjunctivitis. If possible, refer to an ophthalmologist.
- Protect the eye with a compress as long as photophobia lasts. Remove as soon as possible.
- If necessary, administer a preventive dose of [vitamin A](#).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Trachoma

Trachoma is a highly contagious keratoconjunctivitis due to *Chlamydia trachomatis*. The disease is endemic in the poorest rural areas of Africa, Asia, Central and South America and the Middle East.

Infection is usually first contracted early in childhood by direct or indirect contact (dirty hands, contaminated towels, flies). In the absence of hygiene and effective treatment, the inflammation intensifies with successive infections, causing scars and deformities on the upper tarsal conjunctiva. The resulting ingrowing eyelashes (trichiasis) cause corneal lesions followed by permanent blindness, usually in adulthood.

The WHO classifies trachoma into 5 stages. Early diagnosis and treatment of first stages is essential to avoid the development of trichiasis and associated complications.

Clinical features

Several stages can occur simultaneously^{[1][2]} :

- Stage 1: trachomatous inflammation - follicular (TF)
Presence of five or more follicles in the upper tarsal conjunctiva. Follicles are whitish, grey or yellow elevations, paler than the surrounding conjunctiva.
- Stage 2: trachomatous inflammation - intense (TI)
The upper tarsal conjunctiva is red, rough and thickened. The blood vessels, normally visible, are masked by a diffuse inflammatory infiltration or follicles.
- Stage 3: trachomatous scarring (TS)
Follicles disappear, leaving scars: scars are white lines, bands or patches in the tarsal conjunctiva.
- Stage 4: trachomatous trichiasis (TT)
Due to multiple scars the margin of the eyelid, usually the upper lid, turns inwards (entropion); the eyelashes rub against the cornea and cause ulcerations and chronic inflammation.
- Stage 5: corneal opacity (CO)
Cornea gradually loses its transparency, leading to visual impairment and blindness.

Treatment

- Stages 1 and 2:
 - Clean eyes and face several times per day.
 - Antibiotic treatment^[3] :
The treatment of choice is **azithromycin PO**:
Children: 20 mg/kg single dose
Adults: 1 g single dose

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Note: epilation of ingrowing eyelashes is not recommended since it offers only temporary relief and regrowing eyelashes are more abrasive to the cornea.

- Stage 5: no treatment

Prevention

Cleaning of the eyes, face and hands with clean water reduces direct transmission and the development of secondary bacterial infections.

References

1. Solomon AW et al. The simplified trachoma grading system, amended. Bull World Health Organ. 2020;98(10):698-705.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7652564/> [Accessed 20 April 2021]
2. Thylefors B et al. A simple system for the assessment of trachoma and its complications. Bull World Health Organ. 1987;65(4):477-83.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2491032/> [Accessed 20 April 2021]
3. Evans JR et al. Antibiotics for trachoma. Cochrane Database Syst Rev. 2019 Sep 26;9:CD001860.
<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD001860.pub4/full> [Accessed 20 April 2021]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Periorbital and orbital cellulitis

Periorbital cellulitis is a common, usually benign, bacterial infection of the eyelids. It arises principally following trauma to the eyelids (insect bite or abrasion).

Orbital cellulitis is a serious infection involving the contents of the orbit (fat and ocular muscles) that may lead to loss of vision or a brain abscess. It usually arises secondary to spread from sinusitis (e.g. as a complication of ethmoid sinusitis).

Periorbital and orbital cellulitis are more common in children than in adults.

The most common organisms causing periorbital and orbital cellulitis are *Staphylococcus aureus*, *Streptococcus pneumoniae* and other streptococci, as well as *Haemophilus influenzae type b* (Hib) in children living in countries where rates of immunisation with Hib remain low.

Clinical features

- Signs common to both periorbital and orbital cellulitis: acute eyelid erythema and oedema; the oedema has a violaceous hue if secondary to *H. influenzae*.
- In case of orbital cellulitis only:
 - Pain with eye movements;
 - Ophthalmoplegia (paralysis of eye movements) often with diplopia (double vision);
 - Protrusion of the eye (eye bulges out of the socket);
 - High fever, systemic signs.

Treatment

- Hospitalize for the following: orbital cellulitis, children younger than 3 months, critically ill appearing patient^a, local complications, debilitated patient (chronic conditions, the elderly), if there is a risk of non-compliance with or failure of outpatient treatment. Treat the other patients as outpatients.

- Outpatient antibiotic therapy^b:

cefalexin PO for 7 to 10 days

Neonates 0 to 7 days: 25 mg/kg 2 times daily

Neonates 8 days to 1 month: 25 mg/kg 3 times daily

Children over 1 month: 25 mg/kg 2 times daily (max. 2 g daily)

Children ≥ 40 kg and adults: 1 g 2 times daily

or

amoxicillin/clavulanic acid (co-amoxiclav) PO for 7 to 10 days

Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin:

Children < 40 kg: 50 mg/kg 2 times daily

Children ≥ 40 kg and adults:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Neonates 0 to 7 days (≥ 2 kg): 50 mg/kg every 8 hours

Neonates 8 days to < 1 month (< 2 kg): 50 mg/kg every 8 hours

Neonates 8 days to < 1 month (≥ 2 kg): 50 mg/kg every 6 hours

Children 1 month and over: 25 to 50 mg/kg every 6 hours (max. 8 g daily)

Children ≥ 40 kg and adults: 2 g every 6 hours

If there is clinical improvement (patient afebrile and erythema and oedema have improved) after 5 days, change to amoxicillin/clavulanic acid PO at the doses indicated above to complete 7 to 10 days of treatment.

If there is no improvement in the first 48 hours (suspicion of methicillin resistant *S. aureus*), replace cloxacillin with:

clindamycin IV infusion (30 minutes)^f

Neonates 0 to 7 days (< 2 kg): 5 mg/kg every 12 hours

Neonates 0 to 7 days (≥ 2 kg): 5 mg/kg every 8 hours

Neonates 8 days to < 1 month (< 2 kg): 5 mg/kg every 8 hours

Neonates 8 days to < 1 month (≥ 2 kg): 10 mg/kg every 8 hours

Children 1 month and over: 10 mg/kg every 8 hours (max. 1800 mg daily)

Adults: 600 mg every 8 hours

After 5 days, change to clindamycin PO at the same doses to complete 7 to 10 days of treatment.

- If orbital cellulitis is unresponsive to IV antibiotics, consider an abscess. Transfer patient to a surgical centre for drainage.

Footnotes

- (a) Critically ill appearing child: weak grunting or crying, drowsy and difficult to arouse, does not smile, disconjugate or anxious gaze, pallor or cyanosis, general hypotonia.
- (b) For penicillin-allergic patients, **clindamycin** PO for 7 to 10 days:
Children: 10 mg/kg 3 times daily; adults: 600 mg 3 times daily
- (c) For penicillin-allergic patients, **clindamycin** IV infusion (doses as above).
- (d) For administration by IV route, ceftriaxone powder should be reconstituted in water for injection only. For administration by IV infusion, dilute each dose of ceftriaxone in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children over 20 kg and in adults.
- (e) Cloxacillin powder for injection should be reconstituted in 4 ml of water for injection. Then dilute each dose of cloxacillin in 5 ml/kg of 0.9% sodium chloride or 5 % glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children over 20 kg and in adults.
- (f) Dilute each dose of clindamycin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children over 20 kg and in adults.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Other pathologies

- [Onchocerciasis](#)
- [Loiasis](#)
- [Pterygium](#)
- [Cataract](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Onchocerciasis (river blindness)

Ocular lesions result from the invasion of the eye by microfilariae. They generally develop in adults and progress to blindness in the absence of early treatment.

Clinical features and treatment

Ocular lesions are always associated with onchocercal skin lesions (see [Onchocerciasis](#), Chapter 6).

- Pruritus, hemeralopia (crepuscular blindness), decrease in visual acuity, narrowing of the visual field, awareness of microfilariae in the visual field (the patient sees “little wiggling worms before his eyes”).
- Lesions of the cornea (punctuate, then sclerosing, keratitis), iris (iridocyclitis) or posterior segment (chorioretinopathy and optic atrophy); microfilariae within the anterior chamber or vitreous humor (slit lamp).

For treatment, see [Onchocerciasis](#), Chapter 6. Ivermectin treatment may improve anterior segment lesions (sclerosing keratitis, iridocyclitis) and visual acuity. Severe lesions (chorioretinal lesions, optic atrophy) continue to progress despite treatment.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Loiasis

Clinical features and treatment

Migration of an adult worm under the palpebral or bulbar conjunctiva (white, filiform worm, measuring 4 to 7 cm in length, mobile) and ocular pruritus, lacrimation, photophobia or eyelid oedema.

For treatment, see [Loiasis](#), Chapter 6. The migration of the worm is often of very brief duration. Do not attempt to extract it, or administer anaesthetic drops; simply reassure the patient, the event is harmless. Surgical removal is likewise futile if the worm is dead/calcified.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Pterygium

A whitish, triangular growth of fibrovascular tissue extending slowly from the conjunctiva to the cornea. It occurs most frequently in patients who are exposed to wind, dust, or arid climates and never disappears spontaneously.

Clinical features and treatment

Two stages:

- Benign pterygium develops slowly, does not reach the pupil: no treatment.
- Progressive vascularized pterygium: red and inflamed growth covers the pupil and may impair vision:
 - Clean eye with sterile water or 0.9% sodium chloride.
 - Surgical removal if facilities are available.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Cataract

Opacity of the lens that causes a progressive loss of visual acuity. Cataract is common in the tropics and can occur at a younger age than in Europe. The presence of cataract in both eyes leads to blindness. Surgery is the only treatment.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Chapter 6: Parasitic diseases

[Malaria](#)

[Human African trypanosomiasis \(sleeping sickness\)](#)

[American trypanosomiasis \(Chagas disease\)](#)

[Leishmaniases](#)

[Intestinal protozoan infections \(parasitic diarrhoea\)](#)

[Flukes](#)

[Schistosomiases](#)

[Cestodes](#)

[Nematode infections](#)

[Filariasis](#)

[Onchocerciasis \(river blindness\)](#)

[Loiasis](#)

[Lymphatic filariasis \(LF\)](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Malaria

Malaria is a parasitic infection due to protozoa of the genus *Plasmodium*, transmitted to humans by the bite of *Anopheles* mosquitoes. Transmission by transfusion of parasite infected blood and transplacental transmission are also possible.

5 species of *Plasmodium* cause malaria in humans: *P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae* and *P. knowlesi*. All species may cause uncomplicated malaria. Severe malaria (defined by the presence of complications) is almost always due to *P. falciparum*: and, less frequently, *P. vivax* and *P. knowlesi*.

Uncomplicated malaria can rapidly progress to severe malaria, and severe malaria may cause death within a few hours if left untreated.

Clinical features

Malaria should always be considered in patients living in or coming from, an endemic area, who presents with fever (or history of fever in the previous 48 hours).

Uncomplicated malaria

Fever is frequently associated with chills, sweating, headache, muscular ache, malaise, anorexia or nausea. In children, fever may be associated with abdominal pain, diarrhoea and vomiting. Mild to moderate anaemia is frequent in children and pregnant women.

Severe malaria

In addition to the above, patients presenting with one or more of the following complications^[1] should be hospitalised immediately:

- Impaired consciousness, including coma.
- Seizures: more than 2 episodes of generalised or focal (e.g. abnormal eye movements) seizures within 24 hours.
- Prostration: extreme weakness; in children: inability to sit or drink/suck.
- Respiratory distress: rapid, laboured breathing or slow, deep breathing.
- Shock: cold extremities, weak or absent pulse, capillary refill time ≥ 3 seconds, cyanosis.
- Jaundice: yellow discolouration of mucosal surfaces of the mouth, conjunctivae and palms.
- Haemoglobinuria: dark red urine.
- Abnormal bleeding: skin (petechiae), conjunctivae, nose, gums; blood in stools.
- Acute renal failure: oliguria (urine output < 12 ml/kg/day in children and < 400 ml/day in adults) despite adequate hydration.

Laboratory

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Microscopy

Thin and thick blood films enable parasite detection, species identification, quantification and monitoring of parasitaemia.

Blood films may be negative due to sequestration of the parasitized erythrocytes in peripheral capillaries in severe malaria, as well as in placental vessels in pregnant women.

Note: even with positive diagnostic results, rule out other causes of fever.

Additional examinations

Haemoglobin (Hb) level

To be measured routinely in all patients with clinical anaemia, and in all patients with severe malaria.

Blood glucose level

To be measured routinely to detect hypoglycaemia in patients with severe malaria and those with malnutrition (see [Hypoglycaemia](#), Chapter 1).

Treatment of malaria due to *P. vivax*, *P. ovale*, *P. malariae*, *P. knowlesi*

chloroquine (CQ) PO^b

Children and adults:

Day 1: 10 mg base/kg

Day 2: 10 mg base/kg

Day 3: 5 mg base/kg

In general *P. vivax* remains sensitive to CQ but resistance is found in several countries. Where such resistance is high (>10%), or in countries which have de-registered CQ due to *P. falciparum* resistance, an artemisinin-based combination therapy (ACT)^c should be used instead^[1]. For dosing information, see [Treatment of uncomplicated falciparum malaria](#).

Relapses can occur with *P. vivax* and *P. ovale* due to activation of dormant parasites in the liver. Primaquine PO for 14 days (0.25 to 0.5 mg/kg once daily in children ≥ 15 kg; 15 mg once daily in adults) can be given to eliminate these parasites, after the initial treatment with CQ or an ACT. However, this treatment is only recommended for patients living in areas where reinfection is unlikely, i.e. non-endemic, low transmission areas or in countries aiming for elimination of malaria. This treatment is contra-indicated in individuals with G6PD deficiency. If G6PD deficiency cannot be tested individually, the decision to prescribe primaquine must take into account the prevalence of deficiency in the population.

Treatment of uncomplicated falciparum malaria

Antimalarial treatment

During pregnancy, see [Antimalarial treatment in pregnant women](#).

Treatment is an artemisinin-based combination therapy (ACT)^c given by the oral route for 3 days^[1]. The first-line ACT is chosen according to therapeutic efficacy in the area where the patient is living. If the first line ACT is unavailable

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Treatment of uncomplicated falciparum malaria^b

ACT	Presentation	Dosage
artemether/lumefantrine (AL)	Coformulated tablets of 20 mg artemether/120 mg lumefantrine Blister child 5 to < 15 kg, 6 tab/blister Blister child 15 to < 25 kg, 12 tab/blister Blister child 25 to < 35 kg, 18 tab/blister Blister child ≥ 35 kg and adult, 24 tab/blister Coformulated tablets of 80 mg artemether/ 480 mg lumefantrine Blister child ≥ 35 kg and adult, 6 tab/blister	On D1, the first dose is given at 0 hour and the 2 nd dose at 8-12 hours. Subsequent doses on D2 and D3 are given 2 times daily (morning and evening). ==> 1 tab 2 times daily on D1, D2, D3 ==> 2 tab 2 times daily on D1, D2, D3 ==> 3 tab 2 times daily on D1, D2, D3 ==> 4 tab 2 times daily on D1, D2, D3 ==> 1 tab 2 times daily on D1, D2, D3
artesunate/amodiaquine (AS/AQ)	Coformulated tablets Blister child 4.5 to < 9 kg, tab of AS 25 mg/AQ base 67.5 mg, 3 tab/blister Blister child 9 to < 18 kg, tab of AS 50 mg/AQ base 135 mg, 3 tab/blister Blister child 18 to < 36 kg, tab of AS 100 mg/AQ base 270 mg, 3 tab/blister Blister child ≥ 36 kg and adult, tab of AS 100 mg/AQ base 270 mg, 6 tab/blister	==> 1 tab once daily on D1, D2, D3 ==> 1 tab once daily on D1, D2, D3 ==> 1 tab once daily on D1, D2, D3 ==> 2 tab once daily on D1, D2, D3
dihydroartemisinin/piperaquine (DHA/PPQ)	Coformulated tablets Blister child, tab of DHA 20 mg/PPQ 160 mg, 3 tab/blister Blister child, tab of DHA 40 mg/PPQ 320 mg, 3 tab/blister Blister child, tab of DHA 40 mg/PPQ 320 mg, 6 tab/blister Blister adolescent-adult, tab of DHA 40 mg/PPQ 320 mg, 9 tab/blister Blister adolescent-adult, tab of DHA 40 mg/PPQ 320 mg, 12 tab/blister	5 to < 8 kg: 1 tab 20/160 mg once daily on D1, D2, D3 8 to < 11 kg: 1½ tab 20/160 mg once daily on D1, D2, D3 11 to < 17 kg: 1 tab 40/320 mg once daily on D1, D2, D3 17 to < 25 kg: 1½ tab 40/320 mg once daily on D1, D2, D3 25 to < 36 kg: 2 tab 40/320 mg once daily on D1, D2, D3 36 to < 60 kg: 3 tab 40/320 mg once daily on D1, D2, D3 60 to < 80 kg: 4 tab 40/320 mg once daily on D1, D2, D3 ≥ 80 kg: 5 tab 40/320 mg once daily on D1, D2, D3

In low malaria endemic areas, in addition to ACT, all individuals (except in children < 30 kg, pregnant women or breastfeeding women of infants aged < 6 months) diagnosed with *P. falciparum* malaria, should be given a single dose of 0.25 mg/kg **primaquine** to reduce the risk of transmission^[3].

Notes:

- In infants below the age/weight mentioned in the table above, there is little data on efficacy and safety of ACTs.
- The combinations AL, AS/AQ and DHA/PPQ can be used. The dose should be calculated so as to correspond to 10-16 mg/kg/dose of lumefantrine; 10 mg/kg daily of amodiaquine; 20 mg/kg daily of piperaquine.
- Clinical condition of young children can deteriorate rapidly; it may be preferable to start parenteral treatment straight away (see below).

Quinine PO is not recommended as standard treatment, however still remains in some national protocols:

quinine PO for 7 days^b

Children and adults under 50 kg: 10 mg/kg 3 times daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

During pregnancy, see [Antimalarial treatment in pregnant women](#).

Pre-referral treatment

If the patient needs to be transferred, administer before transfer:

- At community level, for children under 6 years: one dose of rectal **artesunate^d** (10 mg/kg)
 - Children 2 months to < 3 years (≤ 10 kg): 1 rectal capsule (100 mg)
 - Children 3 to < 6 years (≤ 20 kg): 2 rectal capsules (200 mg)
- or
- At dispensary level, for children and adults: the first dose of artesunate or, if unavailable, the first dose of artemether. For dosing information, see below.

In either case, provide patients, especially children, with some sugar prior to or during transfer.

Inpatient treatment

The drug of choice is artesunate, preferably IV, or if not possible IM.

For patients in shock: IM route is not appropriate, use artesunate IV only.

artesunate slow IV injection (3 to 5 minutes) or, if not possible, slow IM injection, into the anterior thigh:

Children under 20 kg: 3 mg/kg/dose

Children 20 kg and over and adults: 2.4 mg/kg/dose

- One dose on admission (H0)
- One dose 12 hours after admission (H12)
- One dose 24 hours after admission (H24)
- Then one dose once daily

Treat parenterally for at least 24 hours (3 doses), then, if the patient can tolerate the oral route, change to a complete 3-day course of an ACT. If not, continue parenteral treatment once daily until the patient can change to oral route (without exceeding 7 days of parenteral treatment).

If artesunate is not unavailable, artemether may be an alternative:

artemether IM into the anterior thigh (never administer by IV route)

Children and adults: 3.2 mg/kg on admission (D1) then 1.6 mg/kg once daily

Treat parenterally for at least 24 hours (2 doses), then, if the patient can tolerate the oral route, change to a complete 3-day course of an ACT. If not, continue parenteral treatment once daily until the patient can change to oral route (without exceeding 7 days of parenteral treatment).

Note: if patient is still on parenteral treatment on Day 5, continue on the same treatment until Day 7. In this case it is not necessary to start an ACT.

Quinine IV is still recommended in some national protocols. It may be used in treatment of malaria with shock if artesunate IV is not available. The dose is expressed in quinine salt:

- Loading dose: 20 mg/kg to be administered over 4 hours, then, keep the vein open with an infusion of 5% glucose over 4 hours, then

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Symptomatic treatment and management of complications

Hydration

Maintain adequate hydration. As a guide, for volume to be administered per 24 hours by oral or IV route, see [Appendix 1](#).

Adjust the volume according to clinical condition in order to avoid dehydration or fluid overload (risk of pulmonary oedema).

Fever

Paracetamol in the event of high fever only ([Fever](#), Chapter 1).

Severe anaemia

For treatment, see [Anaemia](#), Chapter 1.

Hypoglycaemia

For treatment, see [Hypoglycaemia](#), Chapter 1.

Notes:

- In an unconscious or prostrated patient, in case of emergency or when venous access is unavailable or awaited, use granulated sugar by the sublingual route to correct hypoglycaemia^e.
- The risk of hypoglycaemia is higher in patients receiving IV quinine.

Coma

Check/ensure the airway is clear, measure blood glucose level and assess level of consciousness.

In the event of hypoglycaemia or if blood glucose level cannot be measured, administer glucose.

If the patient does not respond to administration of glucose, or if hypoglycaemia is not detected:

- Insert a urinary catheter; place the patient in the recovery position.
- Monitor vital signs, blood glucose level, level of consciousness, fluid balance (urine output and fluid input) hourly until stable, then every 4 hours.
- Rule out meningitis (lumbar puncture) or proceed directly to administration of an antibiotic (see [Meningitis](#), Chapter 7).
- Reposition the patient every 2 hours; ensure eyes and mouth are kept clean and moist, etc.

Seizures

See [Seizures](#), Chapter 1. Address possible causes (e.g. hypoglycaemia; fever in children).

Respiratory distress

- Rapid laboured breathing:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Look first for dehydration ([Dehydration](#), Chapter 1), especially due to inadequate fluid intake or excessive fluid losses (high fever, vomiting, diarrhoea). Treat dehydration if present. Be aware of the risk of fluid overload and acute pulmonary oedema. Monitor for the return of urine output.

Acute renal failure (ARF) is found almost exclusively in adults and is more common in Asia than Africa. Insert a urinary catheter, measure output. Restrict fluids to 1 litre/day (30 ml/kg/day in children), plus additional volume equal to urine output. Renal dialysis is often necessary.

Antimalarial treatment in pregnant women

Uncomplicated *P. vivax*, *P. ovale*, *P. malariae*, *P. knowlesi* malaria

As other patients.

Primaquine should not be given in pregnancy.

Uncomplicated falciparum malaria

All ACT included in the table [Treatment of uncomplicated falciparum malaria](#) can be used in all trimesters.

If ACTs are not available, quinine PO (for dosing, see [Treatment of uncomplicated falciparum malaria](#)) combined with **clindamycin** PO if possible (10 mg/kg 2 times daily for 7 days) may be an alternative to ACT. Primaquine should not be given in pregnancy.

Severe malaria

Artesunate, or if unavailable artemether, is recommended in all trimesters.

Quinine IV is not recommended as standard treatment, however it still remains in some national protocols.

Prevention

- For pregnant women in areas with high risk of infection with *P. falciparum*, refer to the guide [Essential obstetric and newborn care](#), MSF.
- In areas with seasonal malaria transmission (in particular across the Sahel sub-region), seasonal malaria chemoprevention in children < 5 years reduces mortality: administer amodiaquine + SP at monthly intervals for 4 months during the transmission period^[4].
- In malaria endemic areas and in epidemic-prone contexts, all in-patient facilities (including HIV treatment centres and feeding centres), should be furnished with long-lasting insecticidal nets (LLINs). For more information, refer to the guide [Public health engineering](#), MSF.
- See specialised literature for information regarding anti-vector measures and prevention in travellers.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- (c) ACT: a combination of artemisinin or one of its derivatives (e.g. artesunate, artemether) with another antimalarial of a different class.
- (d) If it is impossible to refer a patient to a center capable of providing parenteral treatment, rectal artesunate should be given according to the same schedule as artesunate slow IV injection (H0, H12, H24, then once daily).
- (e) Place a level teaspoon of sugar, moistened with a few drops of water, under the tongue, then place the patient in the recovery position. Repeat after 15 minutes if the patient has not regained consciousness. As with other methods for treating hypoglycaemia, maintain regular sugar intake, and monitor.

References

1. World Health Organization. Guidelines for the treatment of malaria, 3rd ed. World Health Organization. 2015.
<https://apps.who.int/iris/handle/10665/162441>
2. World Health Organization. Compendium of WHO malaria guidance: prevention, diagnosis, treatment, surveillance and elimination. 2019.
<https://apps.who.int/iris/handle/10665/312082>
3. World Health Organization. WHO policy brief on single-dose primaquine as gametocytocide in Plasmodium falciparum malaria. 2015.
https://www.who.int/malaria/publications/atoz/who_htm_gmp_2015.1.pdf?ua=1
4. World Health Organization. WHO Policy Recommendation: Seasonal Malaria Chemoprevention (SMC) for *Plasmodium falciparum* malaria control in highly seasonal transmission areas of the Sahel sub-region in Africa. 2012.
https://www.who.int/malaria/publications/atoz/smc_policy_recommendation_en_032012.pdf?ua=1

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Human African trypanosomiasis (sleeping sickness)

Human African trypanosomiasis (HAT) is a zoonosis caused by protozoa (trypanosomes), transmitted to humans through the bite of a tsetse fly (*Glossina*). Transmission by contaminated blood transfusion and transplacental transmission are also possible.

The disease is found only in sub-Saharan Africa. There are two forms: *Trypanosoma brucei gambiense* HAT in western and central Africa and *Trypanosoma brucei rhodesiense* HAT in eastern and southern Africa.

Clinical features

Inoculation may be followed by an immediate local reaction (trypanosomal chancre). This chancre arises in about 50% of all rhodesiense but rarely in gambiense.

Gambiense HAT

- Incubation lasts from a few days to several years.
- The first stage (haemolymphatic stage) corresponds to the haematogenous and lymphatic dissemination of the parasite. Signs include intermittent fever, joint pain, lymphadenopathy (firm, mobile, painless lymph nodes, mainly cervical), hepatosplenomegaly and skin signs (facial oedema, pruritus).
- The second stage (meningoencephalitic stage) corresponds to the invasion of the central nervous system. Signs of the haemolymphatic stage recede or disappear and varying neurological signs progressively develop: sensory disturbances (deep hyperesthesia), psychiatric disorders (apathy or agitation), disturbance of the sleep cycle (with daytime somnolence alternating with insomnia at night), impaired motor functions (paralysis, seizures, tics) and neuroendocrine disorders (amenorrhoea, impotence).
- In the absence of treatment: cachexia, lethargy, coma and death.

Rhodesiense HAT

The first stage is the same as above, but the incubation period is shorter (< 3 weeks), the disease evolves more rapidly and symptoms are more severe. Patients often die of myocarditis in 3 to 6 months without having developed signs of the meningo-encephalitic stage.

In practice, gambiense and rhodesiense HAT can be difficult to differentiate: e.g., there exist cases of acute gambiense infection and others of chronic rhodesiense infection.

Laboratory

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Meningoencephalitic stage: evidence of trypanosomes OR > 5 white cells/mm³

Treatment (except in pregnant women)

- Due to the toxicity of trypanocides, detection of the parasite is essential before initiating treatment. In the absence of parasitological confirmation, treatment may nevertheless be justified in certain cases: very strong clinical suspicion, patients in life-threatening condition, strong serological suspicion (CATT 1:16 positive) in a population where the disease is highly prevalent ($> 2\%$).
- Several treatment regimens exist. Check national recommendations and local resistance levels.
- Treatment must be administered under close medical supervision. Patients receiving pentamidine can be treated as outpatients but those receiving suramin, eflornithine (with or without nifurtimox) or melarsoprol should be hospitalised.
- After treatment, patients should be checked every 6 months (clinical examination, lumbar puncture and examination for trypanosomes) over 24 months, to look for relapse.

Haemolymphatic stage (Stage I)

Gambiense HAT

pentamidine isetionate deep IM

Children and adults: 4 mg/kg once daily for 7 to 10 days

Patients should receive a source of glucose (meal, sweet tea) one hour before injection (risk of hypoglycaemia); they should remain supine during administration and one hour after injection (risk of hypotension).

Rhodesiense HAT

suramin slow IV

Children and adults:

D1: test dose of 4 to 5 mg/kg

D3, D10, D17, D24, D31: 20 mg/kg (max. 1 g per injection)

Suramin may cause anaphylactic reactions, a test dose is recommended prior to starting treatment. In the event of an anaphylactic reaction after the test dose, the patients must not be given suramin again.

Meningoencephalitic stage (Stage II)

Before administrating trypanocides, the priority is to improve the patient's general condition (rehydration, treatment of malaria, intestinal worms, malnutrition, bacterial infections). It is nonetheless recommended not to postpone the trypanocidal treatment for more than 10 days.

Gambiense HAT

- First choice: nifurtimox-eflornithine combination therapy (NECT)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Children under 12 years: 150 mg/kg every 6 hours for 14 days

Children 12 years and over and adults: 100 mg/kg every 6 hours for 14 days

- In the event of a relapse after NECT or eflornithine:

melarsoprol slow IV

Children and adults: 2.2 mg/kg once daily for 10 days

Melarsoprol is highly toxic: reactive encephalopathy (coma, or recurrent or prolonged seizures) in 5 to 10% of treated patients, fatal in around 50% of cases; peripheral neuropathy, invasive diarrhoea, severe skin rash, phlebitis, etc.

Prednisolone PO (1 mg/kg once daily) is frequently combined throughout the duration of treatment.

Rhodesiense HAT

melarsoprol slow IV

Children and adults: 2.2 mg/kg once daily for 10 days

Prednisolone PO (1 mg/kg once daily) is frequently combined throughout the duration of treatment.

Treatment in pregnant women

All trypanocides are potentially toxic for the mother and the foetus (risk of miscarriage, malformation, etc.). However, due to the life-threatening risk for the mother and the risk of mother-to-child transmission, treatment must be initiated as follows:

- Haemolymphatic stage:
pentamidine for gambiense HAT as of the second trimester and suramin for rhodesiense HAT.
- Meningoencephalitic stage: treatment depends on the mother's condition:
 - If in immediately life-threatening condition: treatment with NECT or eflornithine cannot be deferred until after delivery.
 - If not immediately life-threatening condition: pentamidine for gambiense HAT and suramin for rhodesiense HAT.
Treatment with NECT or eflornithine is to be administered after delivery.

Prevention and control

- Individual protection against tsetse fly bites: long sleeves and trousers, repellents, keeping away from risk areas (e.g. near rivers).
- Disease control: mass screening and treatment of patients (*T.b. gambiense*), trypanocide treatment of cattle (*T.b. rhodesiense*), vector control using tsetse fly traps or insecticides.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

American trypanosomiasis (Chagas disease)

Chagas disease is a zoonosis caused by the protozoa *Trypanosoma cruzi*. It is transmitted to humans by contact of triatomine bug faeces with a break in the skin (often caused by a bite from the triatomine bug), or with mucous membranes. Transmission by contaminated blood transfusion, accidental exposure to blood, mother-to-child (during pregnancy or childbirth) or consumption of contaminated food and water is also possible.

Chagas disease has two phases: an acute phase, which lasts approximately 4 to 6 weeks, and a chronic phase, which is lifelong if left untreated.

The disease is primarily found on the American continent^a. It is significantly underdiagnosed.^[1]

Clinical features

Acute phase

- Most cases are asymptomatic.
- If transmitted through a break in the skin: a red swelling on the skin (chagoma) or unilateral painless purplish periorbital oedema (Romaña's sign) with local lymphadenopathy, headache and fever.
- Rarely: multiple lymphadenopathies, hepatosplenomegaly, myocarditis (chest pain, dyspnoea), meningoencephalitis (seizures, paralysis).

Chronic phase

- Many cases remain asymptomatic (indeterminate phase).
- Up to 30% of cases develop organ damage:^[2]
 - cardiac lesions (conduction disorders, dilated cardiomyopathy): arrhythmia, dyspnoea, chest pain, heart failure;
 - gastrointestinal lesions (dilation of the oesophagus or colon i.e. megaoesophagus, megacolon): difficulty swallowing, severe constipation.
 - Individuals with immunosuppression have a higher risk of developing organ damage than the general population.

Diagnosis

Laboratory^[1]

- Acute phase:
 - Identification of *Trypanosoma cruzi* by direct microscopy of fresh blood or blood concentrated by microhematocrit method.
 - In case of strong clinical suspicion despite no definitive diagnosis from direct microscopy, perform serologic tests after a delay of approximately 1 month (see "Chronic phase").
- Chronic phase:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Aetiological treatment

- Acute or chronic Chagas disease can be treated with either benznidazole or nifurtimox. However, treatment is not recommended if patient has already developed cardiac or digestive complications.
- Close clinical monitoring should be provided due to the frequent occurrence of adverse effects. Where available, blood tests (complete blood count, liver and renal function tests) should be performed before, during and after treatment.
- Protocols vary according to the country, follow national recommendations.

For information:

	Age	Dose and duration
benznidazole PO ^(a)	2 to 12 years ^[3]	5 to 8 mg/kg daily in 2 divided doses for 60 days
	> 12 years and adults ^[4]	5 to 7 mg/kg daily in 2 divided doses for 60 days
nifurtimox PO ^{(b)[3]}	≤ 10 years	15 to 20 mg/kg daily in 3 to 4 divided doses for 90 days
	11 to 16 years	12.5 to 15 mg/kg daily in 3 to 4 divided doses for 90 days
	≥ 17 years and adults	8 to 10 mg/kg daily in 3 to 4 divided doses for 90 days

(a) Benznidazole is contra-indicated in pregnancy, breastfeeding and in patient with severe hepatic/renal impairment.

(b) Nifurtimox is contra-indicated in pregnancy, breastfeeding, patients with severe hepatic/renal impairment or history of severe mental disorders or seizures. Adverse effects (gastrointestinal disturbances, agitation, sleeping disorders, seizure) are frequent and reversible and should not necessarily result in discontinuation of treatment. Avoid alcohol and fatty meals during treatment.

Symptomatic treatment

See [Seizures](#) (Chapter 1), [Pain](#) (Chapter 1) and [Heart failure](#) (Chapter 12).

Prevention

- Individual protection against bite from triatomine bugs: use of long-lasting insecticidal net.
- In healthcare settings: standard precautions to avoid contamination with soiled materials or potentially infected body fluids.
- Blood transfusions: advise patients with Chagas disease not to donate blood. In endemic areas, screen donor blood for *Trypanosoma cruzi* antibodies.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

1. Pan American Health Organization. Guidelines for diagnosis and treatment of Chagas disease. Washington, D.C. 2019.
http://iris.paho.org/xmlui/bitstream/handle/123456789/49653/9789275120439_eng.pdf?sequence=6&isAllowed=y
2. Rassi A, Marin-Neto J. Seminar: Chagas disease. The Lancet, Volume 375, ISSUE 9723, P1388-1402, April 17, 2010.
3. Centers for Disease Control and Prevention. Parasites - American Trypanosomiasis.
<https://www.cdc.gov/parasites/chagas/> [Accessed 17 February 2020]
4. World Health Organization. WHO Model Prescribing Information: Drugs Used in Parasitic Diseases - Second Edition. Geneva. 1995.
<https://apps.who.int/iris/handle/10665/41765> [Accessed 6 May 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Leishmaniases

The leishmaniases are a group of parasitic diseases caused by protozoa of the genus *Leishmania*, transmitted by the bite of a sandfly. Over 20 species cause disease in man.

- **Cutaneous** leishmaniasis is endemic in more than 70 countries in South and Central America, Middle East, Central Asia, and Africa.
- **Mucocutaneous** leishmaniasis occurs in Latin America and, more rarely, in Africa (Ethiopia, Sudan).
- **Visceral** leishmaniasis occurs in more than 60 countries in East and North Africa, South and Central Asia, Southern Europe, and South and Central America.

Clinical features

Cutaneous and mucocutaneous leishmaniasis

- Single or multiple lesions on the uncovered parts of the body: an erythematous papule begins at the sandfly bite, enlarges to a nodule and extends in surface and depth to form a scabbed ulcer. Ulcers are painless, unless there is secondary bacterial or fungal infection.
Usually, lesions heal spontaneously, leaving a scar, and result in lifelong protection from disease.
- Lesions may also spread to the mucosa (mouth, nose, conjunctiva) giving rise to the mucocutaneous form, which may cause severe disfigurement.

Visceral leishmaniasis

Visceral leishmaniasis (kala azar) is a systemic disease, resulting in pancytopenia, immunosuppression, and death if left untreated.

- Prolonged (> 2 weeks) irregular fever, splenomegaly, and weight loss are the main signs.
- Other signs include: anaemia, diarrhoea, epistaxis, lymphadenopathy, moderate hepatomegaly.
- Bacterial diarrhoea, pneumonia, and tuberculosis may develop due to immunosuppression.

Post-kala azar dermal leishmaniasis

Macular, nodular or papular skin rash of unknown aetiology, particularly on the face, and typically occurring after apparent cure of visceral leishmaniasis.

Laboratory

Cutaneous and mucocutaneous leishmaniasis

- Parasitological diagnosis: identification of Giemsa-stained parasites in smears of tissue biopsy from the edge of

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment

The various species of *Leishmania* respond differently to drugs. Follow national recommendations.

For information:

Cutaneous and mucocutaneous leishmaniasis

- Cutaneous lesions generally heal spontaneously in 3 to 6 months. Treatment is only indicated if lesions are persistent (> 6 months), disfiguring, ulcerating, or disseminated.
- Forms with a single lesion or few lesions: start with local treatment with a pentavalent antimonial: **sodium stibogluconate** or **meglumine antimoniate**, 1 to 2 ml infiltrated into the lesion if it is a nodule and into the edges and base around the crust if it is an ulcer.
It should be repeated every 3 to 7 days for 2 to 4 weeks. Once healing begins, the treatment can be stopped and healing will continue.
- IM treatment with a pentavalent antimonial (20 mg/kg daily for 10 to 20 days) is restricted to severe cases and must be administered under close medical supervision.
- Miltefosine PO (as for visceral leishmaniasis) for 28 days is effective in many forms of cutaneous leishmaniasis.
- Ulcers are often secondarily infected with streptococci and staphylococci: administer suitable antibiotics.
- Mucocutaneous forms: as for visceral leishmaniasis.

Visceral leishmaniasis

Visceral leishmaniasis in East Africa

- First-line treatment:
a pentavalent antimonial IM or slow IV: 20 mg/kg daily for 17 days
+ **paromomycin** IM: 15 mg (11 mg base)/kg daily for 17 days
- Second-line treatment for relapse and for specific vulnerable groups: severe disease, pregnant women, patients over 45 years:
liposomal amphotericin B IV infusion: 3 to 5 mg/kg once daily for 6 to 10 days up to a total dose of 30 mg/kg
- Treatment in HIV co-infected patients:
liposomal amphotericin B IV infusion: 3 to 5 mg/kg once daily for 6 to 10 days up to a total dose of 30 mg/kg
+ **miltefosine** PO for 28 days:
Children 2 to 11 years: 2.5 mg/kg once daily
Children ≥ 12 years and < 25 kg: 50 mg once daily
Children ≥ 12 years and adults 25 to 50 kg: 50 mg 2 times daily
Adults > 50 kg: 50 mg 3 times daily

Visceral leishmaniasis in South Asia

- First-line treatment:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Post-kala azar dermal leishmaniasis (PKDL)

Only patients with severe or disfiguring disease or with lesions remaining for > 6 months, and young children with oral lesions that interfere with feeding, are treated.

PKDL in East Africa

a **pentavalent antimonial** IM or slow IV: 20 mg/kg daily for 17 to 60 days

+ **paromomycin** IM: 15 mg (11 mg base)/kg daily for 17 days

or

liposomal amphotericin B IV infusion: 2.5 mg/kg once daily for 20 days

or

miltefosine PO for 28 days (as for visceral leishmaniasis) may be beneficial in HIV co-infected patients

PKDL in South Asia

liposomal amphotericin B IV infusion: 5 mg/kg 2 times weekly up to a total dose of 30 mg/kg

Prevention

- Insecticide-treated mosquito nets.
- Vector control and elimination of animal reservoir hosts.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Intestinal protozoan infections (parasitic diarrhoea)

The most important intestinal protozoan infections are amoebiasis (*Entamoeba histolytica*), giardiasis (*Giardia lamblia*), cryptosporidiosis (*Cryptosporidium sp*), cyclosporiasis (*Cyclospora cayetanensis*) and isosporiasis (*Isospora belli*).

Intestinal protozoa are transmitted by the faecal-oral route (soiled hands, ingestion of food or water contaminated with faeces) and may cause both individual cases of diarrhoea and epidemic diarrhoea outbreaks.

Clinical features

- Amoebiasis gives rise to bloody diarrhoea (see [Amoebiasis](#), Chapter 3).
- Clinical presentation of giardiasis, cryptosporidiosis, cyclosporiasis and isosporiasis is very similar:
 - Diarrhoea is usually mild and self-limiting, except in children and patients with advanced HIV disease (CD4 < 200). These patients are likely to develop severe, intermittent or chronic diarrhoea that may be complicated by malabsorption with significant wasting (or failure to gain weight in children) or severe dehydration.
 - Stools are usually watery, but steatorrhoea (pale, bulky, fatty stools) may be found in the event of secondary fat malabsorption; stools may contain mucus.
 - Diarrhoea is usually associated with non-specific gastrointestinal symptoms (abdominal distension and cramps, flatulence, nausea, anorexia), but patients have low-grade fever or no fever.

Laboratory

Definitive diagnosis relies on parasite identification in stool specimens (trophozoites and cysts for giardia; oocysts for cryptosporidium, cyclospora, isospora). Two to three samples, collected 2 to 3 days apart are necessary, as pathogens are shed intermittently.

Treatment

- Correct dehydration if present (for clinical features and management, see [Dehydration](#), Chapter 1).
- If the causal agent has been identified in the stool:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Giardiasis	tinidazole PO single dose Children: 50 mg/kg (max. 2 g) Adults: 2 g or metronidazole PO for 3 days Children: 30 mg/kg once daily Adults: 2 g once daily
Cryptosporidiosis	In immunocompetent patients, no aetiological treatment; spontaneous resolution in 1 to 2 weeks.
Cyclosporiasis	co-trimoxazole PO for 7 days Children: 25 mg SMX + 5 mg TMP/kg 2 times daily Adults: 800 mg SMX + 160 mg TMP 2 times daily In immunocompetent patients, symptoms usually resolve spontaneously in 1 to 3 weeks. Treatment is given in case of severe or prolonged symptoms.
Isosporiasis	co-trimoxazole PO for 7 to 10 days Adults: 800 mg SMX + 160 mg TMP 2 times daily In immunocompetent patients, symptoms usually resolve spontaneously in 2 to 3 weeks. Treatment is given in case of severe or prolonged symptoms.

- If reliable stool examination cannot be carried out: parasitic diarrhoeas cannot be differentiated on clinical grounds, nor is it possible to distinguish these from non-parasitic diarrhoeas. An empirical treatment (using tinidazole or metronidazole and co-trimoxazole as above, together or in succession) may be tried in the case of prolonged diarrhoea or steatorrhoea. In patients with HIV infection, see empirical treatment ([HIV infections and AIDS](#), Chapter 8).
- In patients with advanced HIV disease, cryptosporidiosis, cyclosporiasis and isosporiasis are opportunistic infections; the most effective intervention is the treatment of the underlying HIV infection with antiretrovirals. Patients remain at high risk for dehydration/death until immunity is restored.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Flukes

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Infection/Epidemiology	Clinical features/Diagnosis	Treatment
Lung flukes <i>Paragonimus</i> sp <i>Distribution:</i> South-East Asia, China, parts of Cameroon, Nigeria, Gabon, Congo, Colombia, Peru <i>Transmission:</i> eating raw freshwater crustaceans	The two most prominent symptoms are prolonged (> 2 weeks) productive cough and intermittent haemoptysis (rusty-brown sputum). In endemic areas, paragonimosis should be considered whenever pulmonary tuberculosis is suspected as the clinical and radiological features overlap. Paragonimosis is confirmed when eggs are detected in sputum (or possibly in stools).	praziquantel PO Children 4 years and over and adults: 25 mg/kg 3 times daily for 2 days
Hepatobiliary flukes <i>Fasciola hepatica</i> and <i>gigantica</i> <i>Distribution:</i> worldwide, in areas where sheep and cattle are raised <i>Transmission:</i> eating uncooked aquatic plants	<p><i>During migration phase:</i> asthenia, prolonged fever, myalgia, right upper quadrant pain, mild hepatomegaly; sometimes, allergic signs (e.g. pruritus). At this stage, the diagnosis is rarely considered and can only be confirmed through serology; parasitological examination of stools is always negative.</p> <p><i>Once adult flukes are present in the biliary tract:</i> presentation resembles cholelithiasis: right upper quadrant pain, recurrent episodes of obstructive jaundice/ febrile cholangitis. The diagnosis is confirmed when parasite eggs are detected in stools (or flukes are seen in the biliary tract with sonography).</p>	triclabendazole PO Children and adults: 10 mg/kg single dose May repeat in 24 hours in the event of severe infection
<i>Opisthorchis felineus</i> (Asia, Eastern Europe) <i>Opisthorchis viverrini</i> (Cambodia, Laos, Vietnam, Thailand) <i>Clonorchis sinensis</i> (China, Korea, Vietnam) <i>Transmission:</i> eating raw/undercooked freshwater fish	Abdominal pain and diarrhoea. With heavy infection, hepatobiliary symptoms: hepatomegaly, right upper quadrant pain, jaundice or episodes of febrile cholangitis. The diagnosis is confirmed when parasite eggs are detected in stools.	praziquantel PO Children 4 years and over and adults: 25 mg/kg 3 times daily for 2 days
Intestinal flukes <i>Fasciolopsis buski</i> (India, Bangladesh, South-East Asia)	Symptoms are limited to diarrhoea and epigastric or abdominal pain. With massive infection, <i>F. buski</i> can cause oedematous allergic reactions (including ascites, anasarca). The diagnosis is confirmed when parasite eggs are detected in	praziquantel PO Children 4 years and over and

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

raw/undercooked fish
(other species)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Schistosomiases

Schistosomiases are acute or chronic visceral parasitic diseases due to 5 species of trematodes (schistosomes). The three main species infecting humans are *Schistosoma haematobium*, *Schistosoma mansoni* and *Schistosoma japonicum*. *Schistosoma mekongi* and *Schistosoma intercalatum* have a more limited distribution.

Humans are infected while wading/bathing in fresh water infested with schistosome larvae. Symptoms occurring during the phases of parasite invasion (transient localized itching as larvae penetrate the skin) and migration (allergic manifestations and gastrointestinal symptoms during migration of schistosomules) are frequently overlooked. In general, schistosomiasis is suspected when symptoms of established infection become evident. Each species gives rise to a specific clinical form: genito-urinary schistosomiasis due to *S. haematobium*, intestinal schistosomiasis due *S. mansoni*, *S. japonicum*, *S. mekongi* and *S. intercalatum*.

The severity of the disease depends on the parasite load. Heavily infected patients are prone to visceral lesions with potentially irreversible sequelae. Children aged 5 to 15 years are particularly at risk: prevalence and parasite load are highest in this age group.

An antiparasitic treatment should be administered to reduce the risk of severe lesions, even if there is a likelihood of re-infection.

Clinical features

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

	Parasite/Epidemiology^a	Clinical features/Diagnosis (established infection)
Genito-urinary schistosomiasis	<i>S. haematobium</i> <i>Distribution:</i> Africa, Madagascar and the Arabian peninsula	<ul style="list-style-type: none"> • Urinary manifestations: <ul style="list-style-type: none"> ▫ In endemic areas, urinary schistosomiasis should be suspected in any patients who complain of macroscopic haematuria (red coloured urine throughout, or at the end of, micturition). Haematuria is frequently associated with polyuria/dysuria (frequent and painful micturition). ▫ In patients, especially children and adolescents, with urinary symptoms, visual inspection of the urine (and dipstick test for microscopic haematuria if the urine appears grossly normal) is indispensable. ▫ Presumptive treatment is recommended in the presence of macro- or microscopic haematuria, when parasitological confirmation (parasite eggs detected in urine) cannot be obtained. • Genital manifestations: <p>In women, symptoms of genital infection (white-yellow or bloody vaginal discharge, itching, lower abdominal pain, dyspareunia) or vaginal lesions resembling genital warts or ulcerative lesions on the cervix; in men, haematospermia (blood in the semen).</p> • If left untreated: risk of recurrent urinary tract infections, fibrosis/calcification of the bladder and ureters, bladder cancer; increased susceptibility to sexually transmitted infections and risk of infertility. • In endemic areas, genito-urinary schistosomiasis may be a differential diagnosis to the genito-urinary tuberculosis, and in women, to the sexually transmitted infections (especially in women with a history of haematuria).
Intestinal schistosomiasis	<i>S. mansoni</i> <i>Distribution:</i> tropical Africa, Madagascar, the Arabian peninsula, South America (especially Brazil) <i>S. japonicum</i> <i>Distribution:</i> China,	<ul style="list-style-type: none"> • Non-specific digestive symptoms (abdominal pain; diarrhoea, intermittent or chronic, with or without blood) and hepatomegaly. • For <i>S. intercalatum</i>: digestive symptoms only (rectal pain, tenesmus, rectal prolapse, bloody diarrhoea). • If left untreated: risk of hepatic fibrosis, portal hypertension, cirrhosis, gastrointestinal haemorrhage (hematemesis, melena,

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment

praziquantel PO^{[1][2]}

Children 4 years and over and adults^b :

- *S. haematobium, S. mansoni, S. intercalatum*: 40 mg/kg single dose or 2 doses of 20 mg/kg administered 4 hours apart
- *S. japonicum, S. mekongi*: 2 doses of 30 mg/kg or 3 doses of 20 mg/kg administered 4 hours apart

Footnotes

(a) For more information on geographic distribution of schistosomiasis:
https://www.who.int/schistosomiasis/Schistosomiasis_2012-01.png?ua=1

(b) For the treatment of schistosomiasis, praziquantel may be administered to pregnant women.

References

1. Treatment Guidelines from The Medical Letter. Vol. 11 (Suppl). Drugs for Parasitic Infections. 2013.
https://www.uab.edu/medicine/gorgas/images/docs/syllabus/2015/03_Parasites/RxParasitesMedicalLetter2013.pdf [Accessed 25 May 2020]
2. Centers for Disease Control and Prevention (CDC). Schistosomiasis. Resources for Health Professionals. 2018.
https://www.cdc.gov/parasites/schistosomiasis/health_professionals/index.html#tx [Accessed 25 May 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Cestodes

Cestodes (adult forms)

Parasites	Clinical features/Laboratory	Treatment	Transmission/Prevention
Taeniasis <i>Taenia saginata</i> <i>Taenia solium</i> (worldwide)	Often asymptomatic Segments expelled in the stools, sometimes gastrointestinal disturbances (epigastric or abdominal pain, nausea, diarrhoea) Laboratory: eggs in stools or collected from perianal skin (scotch tape method), segments in stools	praziquantel PO ^(a) Children 4 years and over and adults: 5 to 10 mg/kg single dose	Transmission by eating raw or under-cooked meat: <ul style="list-style-type: none">• beef for <i>T. saginata</i>• pork for <i>T. solium</i> Prevention: <ul style="list-style-type: none">• individual: cook meat thoroughly• collective: slaughterhouse monitoring
Diphyllobothriasis <i>Diphyllobothrium latum</i> (temperate or cold lake areas)	Often asymptomatic In the event of heavy infection: mild gastrointestinal disturbances, anaemia due to vitamin B ₁₂ deficiency associated with (rare) neurological sequelae Laboratory: eggs in stools	praziquantel PO ^(a) Children 4 years and over and adults: 5 to 10 mg/kg single dose If anaemia: vitamin B₁₂ + folic acid	Transmission by eating raw or under-cooked freshwater fish Prevention: <ul style="list-style-type: none">• individual: cook fish thoroughly
Hymenolepiasis <i>Hymenolepis nana</i> (worldwide)	Often asymptomatic In the event of heavy infection: gastrointestinal disturbances (epigastric pain) Laboratory: eggs in stools	praziquantel PO ^(a) Children 4 years and over and adults: 15 to 25 mg/kg single dose	Transmission by faecal-oral route or auto-infection Prevention: <ul style="list-style-type: none">• individual: hand washing, nail cutting• collective: hygiene and sanitation (water, latrines, etc.)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Parasites	Clinical features/Laboratory	Treatment	Transmission/Prevention
Cysticercosis <i>Taenia solium</i> (worldwide)	<ul style="list-style-type: none"> Muscular: asymptomatic or myalgia Subcutaneous: nodules Neurological (neurocysticercosis): headache, convulsions, coma, etc. Ocular: exophthalmia, strabismus, iritis, etc. <p>Laboratory: hypereosinophilia in blood and cerebrospinal fluid</p>	<p>Neurological and ocular cysticercosis should be managed in specialized facilities.</p> <p>Antiparasitic treatment without diagnosis of location by computerised tomography and/or magnetic resonance imaging can worsen the symptoms even threaten the life. Neurosurgical treatment can be required.</p>	<p>Transmission: by eating food contaminated with <i>T. solium</i> eggs or auto-infection</p> <p>Prevention:</p> <ul style="list-style-type: none"> individual: treat <i>T. solium</i> carriers, hygiene, cook meat thoroughly
Hydatid cyst <i>Echinococcus granulosus</i> (South America, North, East and South Africa, Western Europe)	<p>Cysts located in the liver (60% of cases); lungs (30% of cases), and, less frequently, in other sites including the brain.</p> <p>Long asymptomatic period. The cyst becomes symptomatic when complications develop (biliary obstruction; anaphylactic shock in the event of rupture into peritoneal cavity, vessels or an organ; febrile painful jaundice in the event of rupture into the biliary tree, etc.).</p>	<p>First-line treatment: surgical excision</p> <p>albendazole PO^(b) is useful in addition to, or instead of, surgery:</p> <p>Children over 2 years and adults under 60 kg: 7.5 mg/kg 2 times daily</p> <p>Adults over 60 kg: 400 mg 2 times daily</p> <p>Treatment duration: In addition to surgery (pre-operatively or post-operatively): continuous course of minimum 2 months or at least two 28-day courses with a drug-free interval of 14 days.</p> <p>Inoperable cases: 28-day courses with drug-</p>	<p>Transmission:</p> <ul style="list-style-type: none"> direct: contact with dogs indirect: water and food contaminated by dog faeces <p>Prevention:</p> <ul style="list-style-type: none"> individual: avoid contact with dogs collective: eliminate stray dogs, monitor slaughterhouses

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Nematode infections

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Infection/Epidemiology	Clinical features/Diagnosis	Treatment
Ascariasis (roundworms) ^(a) <i>Ascaris lumbricoides</i> <i>Distribution:</i> worldwide, mainly in tropical and subtropical <i>Transmission:</i> ingestion of ascaris eggs	<ul style="list-style-type: none"> • <i>During larval migration</i> Loeffler's syndrome: transient pulmonary symptoms (dry cough, dyspnoea, wheezing) and mild fever. • <i>Once adult worms are present in the intestine</i> Abdominal pain and distension. In general, the diagnosis is made when adult worms are expelled from the anus (or occasionally from the mouth). Ascaris are large (15–30 cm), cylindrical worms, pinkish-white, with slightly tapered ends. • <i>Complications</i> Ascariasis is usually benign, but massive infestation may cause intestinal obstruction (abdominal pain, vomiting, constipation), especially in children < 5 years. Worms may accidentally migrate to gall bladder, liver or peritoneum, causing jaundice, liver abscess, or peritonitis. • Ascaris eggs may be detected through parasitological examination of stools. 	albendazole PO single dose Children > 6 months and adults: 400 mg (200 mg in children > 6 months but < 10 kg) or mebendazole PO for 3 days Children > 6 months and adults: 100 mg 2 times daily (50 mg 2 times daily in children > 6 months but < 10 kg)
Trichuriasis (whipworms) ^(a) <i>Trichuris trichiura</i> <i>Distribution and transmission:</i> as for <i>A. lumbricoides</i>	<ul style="list-style-type: none"> • In heavy infection: abdominal pain and diarrhoea. • In massive infection: chronic bloody diarrhea, tenesmus, rectal prolapse due to frequent attempts to defecate, especially in children. Worms may sometimes be seen on the rectal mucosa when prolapsed: these are grayish-white, 3–5 cm in length, in the shape of a whip, with a thickened body and a long, threadlike extremity. • Trichuris eggs may be detected through parasitological examination of stools. 	albendazole PO for 3 days Children > 6 months and adults: 400 mg once daily (200 mg once daily in children > 6 months but < 10 kg) or mebendazole PO for 3 days, as for ascariasis. A single dose of albendazole or mebendazole is often insufficient.
Ankylostomiasis ^(a) <i>Ancylostoma duodenale</i> <i>Necator americanus</i> <i>Distribution:</i> tropical and	<ul style="list-style-type: none"> • <i>During larval penetration/migration</i> Cutaneous signs (pruritic papulo-vesicular rash at the site of penetration, usually the feet) and pulmonary symptoms (similar to ascariasis). 	albendazole single dose (as for ascariasis) is much more effective than mebendazole single dose.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<p><i>Strongyloides stercoralis</i></p> <p><i>Distribution:</i> humid tropical regions</p> <p><i>Transmission:</i> larval skin penetration and auto-infection</p>	<ul style="list-style-type: none"> ▫ During larval penetration/migration: cutaneous signs (erythema and pruritus at the site of penetration, which may persist several weeks) and pulmonary symptoms (similar to ascariasis). ▫ Once larvae are present in the intestine: gastrointestinal symptoms (bloating, abdominal and epigastric pain, vomiting, diarrhoea). • <i>Chronic strongyloidiasis</i> Intestinal larvae may re-infect their host (auto-infection) by penetrating through the intestinal wall or by migrating transcutaneously from perianal skin. Chronic infections result in prolonged or recurrent pulmonary and gastrointestinal symptoms. Transcutaneous migration of intestinal larvae gives rise to a typical rash (<i>larva currens</i>), mainly in the anal region and on the trunk: sinuous, raised, linear, migrating lesion, intensely pruritic, moving rapidly (5 to 10 cm/hour) and lasting several hours or days. • <i>Complications</i> Hyperinfection (massive infestation) results in exacerbation of pulmonary and gastrointestinal symptoms, and possible dissemination of larvae to atypical locations, (CNS, heart, etc.). This form occurs mainly in patients receiving immunosuppressive therapy (e.g. corticosteroids). • Strongyloides larvae may be detected through parasitological examination of stools. 	<p>ivermectin PO^(b) single dose Children > 15 kg and adults: 200 micrograms/kg, on an empty stomach</p> <p>While less effective, a 3-day treatment with albendazole PO (as for trichuriasis) may be an alternative.</p> <p>Hyperinfections are refractory to conventional therapy. Prolonged or intermittent multiple-dose regimens are required.</p>
<p><i>Enterobiasis</i> (pinworms)</p> <p><i>Enterobius vermicularis</i></p> <p><i>Distribution:</i> worldwide</p> <p><i>Transmission:</i> faecal-oral route or auto-infection</p>	<ul style="list-style-type: none"> • Anal pruritus, more intense at night, vulvovaginitis in girls (rare). In practice, the diagnosis is most often made when worms are seen on the perianal skin (or in the stool in heavy infestation). Pinworms are small (1 cm), mobile, white, cylindrical worms with slightly tapered ends. • Pinworm eggs may be collected from the anal 	<p>albendazole PO single dose, as for ascariasis</p> <p>or</p> <p>mebendazole PO single dose Children > 6 months and adults: 100 mg /50 mg in children < 6 months</p>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

<p><i>Transmission:</i> consumption of raw or undercooked meat containing trichinella larvae (pork, wart-hog, bear, dog, etc.)</p>	<ul style="list-style-type: none"> • Muscular phase (about 1 week after ingestion) High fever; muscular pain [ocular [pain on eye movement], masseters [limitation of mouth opening], throat and neck [pain with swallowing and speech], trunk and limbs]; facial or bilateral peri-orbital oedema; conjunctival haemorrhage, subungual haemorrhage; headache. Typical features are not always present and the patient may present with a non-specific flu-like syndrome. Other features, such as dietary habits (consuming pork/raw meat), suggestive symptoms (fever > 39 °C and myalgia and facial oedema) in several individuals who have shared the same meal (e.g. ceremony) or hypereosinophilia > 1000/mm³, reinforce the clinical suspicion. • Definitive diagnosis: muscle biopsy; serology (ELISA, Western Blot). 	<p>400 mg 2 times daily or mebendazole PO for 10 to 15 days Children > 2 years: 2.5 mg/kg 2 times daily Adults: 200 mg 2 times daily <i>plus, regardless of which antihelminthic is chosen:</i> prednisolone PO 0.5 to 1 mg/kg once daily for the duration of treatment</p>
--	---	--

- (a) Roundworms, whipworms and hookworms frequently co-infect the same host. This should be taken into account when prescribing antihelminthic treatment.
- (b) The migrating larvae of *Ancylostoma braziliense* and *caninum* (hookworms of cats and dogs) also present as a pruritic, inflammatory, creeping eruption in humans (cutaneous larva migrans) but with a slower rate of progression and a longer duration (several weeks or months). Treatment is with **albendazole** (400 mg single dose or once daily for 3 days in children > 6 months and adults; 200 mg in children > 6 months but < 10 kg) or **ivermectin** (200 micrograms/kg single dose).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Filariasis

- [Onchocerciasis \(river blindness\)](#)
- [Loiasis.](#)
- [Lymphatic filariasis \(LF\)](#)

Filiariases are helminthiases due to tissue-dwelling nematode worms (filariae). Human to human transmission takes place through the bite of an insect vector.

The most important pathogens are outlined in the table below. Mixed infections are common in co-endemic regions.

Each filarial species is found in 2 principal developmental stages: macrofilariae (adult worms) and microfilariae (larval offspring). The treatment depends on the pathogenic stage of the species considered and targets microfilariae for *O. volvulus* and macrofilariae for the other species.

Species/Infections	Location of macrofilariae	Location of microfilariae	Pathogenic stage	Presence of <i>Wolbachia</i>
<i>Onchocerca volvulus</i> (onchocerciasis)	Subcutaneous nodules	Skin and eye	Microfilariae	Yes
<i>Loa loa</i> (loiasis)	Subcutaneous tissue	Blood	Macrofilariae	No
<i>Wuchereria bancrofti, Brugia malayi</i> and <i>Brugia timori</i> (lymphatic filariasis)	Lymph vessels	Blood	Macrofilariae	Yes

Classical antifilarial agents include diethylcarbamazine (DEC), ivermectin and albendazole. Doxycycline is used solely in the treatment of *O. volvulus* and lymphatic filarial worms, which harbour an endosymbiotic bacterium (*Wolbachia*) sensitive to doxycycline.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Onchocerciasis (river blindness)

The distribution of onchocerciasis is linked to that of its vector (*Simulium*), which reproduces near rapidly flowing rivers in intertropical Africa (99% of cases), Latin America (Guatemala, Mexico, Ecuador, Colombia, Venezuela, Brazil) and Yemen.

Clinical features

In endemic areas, the following signs, alone or in combination, are suggestive of onchocerciasis:

- Onchocercomas: painless subcutaneous nodules containing adult worms, usually found over a bony prominence (iliac crest, trochanters, sacrum, rib cage, skull, etc.), measuring several mm or cm in size, firm, smooth, round or oval, mobile or adherent to underlying tissue; single, or multiple and clustered.
- Acute papular onchodermititis: papular rash, sometimes diffuse but often confined to the buttocks or lower extremities, intensely itchy, associated with scratch marks, often superinfected ("filarial scabies")^a. This arises from dermal invasion by microfilariae.
- Late chronic skin lesions: patchy depigmentation on the shins ("leopard skin"), skin atrophy or areas of dry, thickened, peeling skin (lichenification; "lizard skin").
- Visual disturbances and ocular lesions: see [Onchocerciasis](#), Chapter 5.

Laboratory

- Detection of the microfilariae in the skin (skin snip biopsy, iliac crest).
- If the skin biopsy is positive, look for loiasis in regions where loiasis is co-endemic (mainly in Central Africa).

Treatment

Antiparasitic treatment

- Diethylcarbamazine is contra-indicated (risk of severe ocular lesions).
- **Doxycycline** PO (200 mg once daily for 4 weeks; if possible 6 weeks) kills a significant percentage of adult worms and progressively reduces the number of *O. volvulus* microfilariae^b. It is contraindicated in children < 8 years and pregnant or breast-feeding women.
- **Ivermectin** PO is the drug of choice: 150 micrograms/kg single dose; a 2nd dose should be administered after 3 months if clinical signs persist. Repeat the treatment every 6 or 12 months to maintain the parasite load below the threshold at which clinical signs appear^c. Ivermectin is not recommended in children < 5 years or < 15 kg and pregnant women.
- In case of co-infection with *Loa loa* or in regions where loiasis is co-endemic, ivermectin should be administered with caution (risk of severe adverse reactions in patients with high *L. loa* microfilarial load):

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- In the case of concomitant lymphatic filariasis: administer ivermectin first then start treatment for lymphatic filariasis with doxycycline PO (see [Lymphatic filariasis](#)) one week later.

Nodulectomy (surgical removal of onchocercomas)

Nodules are benign, often deep, and their ablation does not treat onchocerciasis. Thus, nodulectomy is reserved for cranial nodules (their proximity to the eye is a risk factor for visual compromise) or nodules which are cosmetically unacceptable. In other cases, refrain from nodulectomy. Nodulectomy is performed under local anaesthesia, in an appropriately equipped facility.

Footnotes

- (a) Differential diagnosis is sarcoptic scabies ([Scabies](#), Chapter 4).
- (b) Elimination of *Wolbachia* reduces the longevity and fertility of the macrofilariae, and thus the production of new microfilariae within the organism.
- (c) Ivermectin kills microfilariae and disrupts production of microfilariae by adult worms. However the treatment must be administered at regular intervals since it does not kill adult worms.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Loiasis

The distribution of loiasis is linked to that of its vector (*Chrysops*) in forests or savannah with gallery forests in West or Central Africa (limits West: Benin; East: Uganda; North: Sudan and South: Angola).

Clinical features

- The subconjunctival migration of an adult worm is pathognomonic of *Loa loa* infection.
- Localised subcutaneous swellings, allergic in origin, transient (several hours or days), painless, non-pitting, appearing anywhere on the body, frequently the upper extremities and face, often associated with localised or generalised pruritus (« Calabar swellings »).
- Onset of pruritus, in the absence of other signs.
- Subcutaneous migration of an adult worm: pruritic, palpable red cord-like linear lesion, sinuous, advancing (1 cm/hour), disappearing rapidly with no trace^a. Such migration generally arises following treatment with diethylcarbamazine, rarely spontaneously.

Laboratory

- Detection of microfilariae in the peripheral blood (thick film, stained with Giemsa). Blood specimens should be collected between 10 am and 5 pm. Quantify microfilaraemia even if the diagnosis is certain, since treatment is determined by the intensity of the parasite load.
- If the thick film is positive, look for onchocerciasis in regions where onchocerciasis is coendemic (mainly in Central Africa).

Treatment

Antiparasitic treatment

- Diethylcarbamazine (DEC) is the only macrofilaricide available but is contra-indicated in:
 - Patients with microfilaraemia > 2000 mf/ml (risk of severe encephalopathy, with poor prognosis).
 - Patients co-infected with *O. volvulus* (risk of severe eye lesions).
 - Pregnant women, infants, and patients in poor general condition.
- Ivermectin (and possibly albendazole) is used to reduce microfilaraemia before administration of DEC; however, ivermectin administration may trigger encephalopathy in patients with very high *Loa loa* microfilaraemia (> 30 000 mf/ml).
- Doxycycline is not indicated since *Loa loa* does not harbour *Wolbachia*.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

2) *L. loa* microfilaraemia is between 2,000 and 8,000 mf/ml

Reduce microfilaraemia with **ivermectin** (150 micrograms/kg single dose); repeat the treatment every month if necessary; administer DEC when the microfilaraemia is < 2000 mf/ml.

3) *L. loa* microfilaraemia is between 8,000 and 30,000 mf/ml

Treatment with **ivermectin** (150 micrograms/kg single dose) may cause marked functional impairment for several days. Close supervision and support from family member(s) are necessary^b. Prescribe paracetamol as well for 7 days.

4) *L. loa* microfilaraemia is > 30,000 mf/ml

- ▷ If the loiasis is well tolerated, it is preferable to refrain from treatment: the disease is benign and treatment with ivermectin may cause very severe adverse reactions (encephalopathy), albeit rarely.
- ▷ If loiasis has a significant clinical impact and/or the patient presents with symptomatic onchocerciasis requiring treatment, **ivermectin** (150 micrograms/kg single dose) is administered for 5 days under supervision in hospital^c. An attempt to first reduce *L. loa* microfilaraemia using **albendazole** (200 mg 2 times daily for 3 weeks) is an option. When *L. loa* microfilaraemia is < 30 000 mf/ml, treat with ivermectin under close supervision and support, then DEC when the microfilaraemia is < 2000 mf/ml.

Extraction of macrofilariae

Subcutaneous migration of a microfilaria usually results from treatment with DEC; the worm will die beneath the skin and extracting it serves no purpose.

Removal of an adult worm from the conjunctiva: see [Loasis](#), Chapter 5.

Footnotes

- (a) For differential diagnosis, see [cutaneous larva migrans](#).
- (b) Patients may present with various pain syndromes, be unable to move without help or unable to move at all. Monitoring is necessary to determine whether the patient can manage activities of daily living, and provide assistance if necessary. If the patient remains bedridden for several days, ensure pressure sores do not develop (mobilisation, repositioning).
- (c) A severe reaction may occur on D2-D3. It is usually preceded by haemorrhages of the palpebral conjunctiva on D1-D2. Routinely check for this sign by turning back the eyelids. Symptoms of post ivermectin encephalopathy are reversible and the prognosis favourable, if the patient is correctly managed; the treatment is symptomatic until symptoms resolve. Avoid the use of steroids due to adverse effects.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Lymphatic filariasis (LF)

The distribution of LF is linked to that of its mosquito vectors (Anopheles, Culex, Aedes, etc.):

- *W. bancrofti*: sub-Saharan Africa, Madagascar, Egypt, India, South East Asia, Pacific region, South America, The Caribbean
- *B. malayi*: South East Asia, China, India, Sri Lanka
- *B. timori*: Timor

90% of LF is due to *W. bancrofti* and 10% to *Brugia* spp.

Clinical features

- Acute recurrent inflammatory manifestations
 - Adenolymphangitis: lymph node(s) and red, warm, tender oedema along the length of a lymphatic channel, with or without systemic signs (e.g. fever, nausea, vomiting). The inflammation may involve the lower limbs, external genitalia and breast.
 - In men: acute inflammation of the spermatic cord (funiculitis), epididymis and testicle (epididymo-orchitis).
 - Attacks resolve spontaneously within a week and recur regularly in patients with chronic disease.
- Chronic manifestations
 - Lymphoedema: oedema of the lower extremity or external genitalia or breast, secondary to obstruction of the lymphatics by macrofilariae. The oedema is reversible initially but then becomes chronic and increasingly severe: hypertrophy of the area affected, progressive thickening of the skin (fibrous thickening with formation of creases, initially superficial, but then deep, and verrucous lesions). The final stage of lymphoedema is elephantiasis.
 - In men: increase in volume of fluid due to accumulation within the tunica vaginalis (hydrocoele, lymphocoele, chylocoele); chronic epididymo-orchitis.
 - Chyluria: milky or rice-water urine (disruption of a lymphatic vessel in the urinary tract).
 - In patients parasitized by *Brugia* spp, genital lesions and chyluria are rare: lymphoedema is usually confined to below the knee.

Laboratory

- Detection of microfilariae in the peripheral blood (thick film)^a; blood specimens should be collected between 9 pm and 3 am.
- In regions where loiasis and/or onchocerciasis are co-endemic, check for co-infection if the LF diagnosis is positive.

Treatment

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Ivermectin (weak or absent macrofilaricidal effect) and albendazole should not be used for the treatment of individual cases (no effect on symptoms).
- In the case of confirmed or probable co-infection with *O. volvulus*: treat [onchocerciasis](#) first, then administer doxycycline.

Control/prevention of inflammatory manifestations and infectious complications

- Acute attacks: bed rest, elevation of the affected limb without bandaging, cooling of the affected limb (wet cloth, cold bath) and analgesics; antibacterial or antifungal cream if necessary; antipyretics if fever (paracetamol) and hydration.
- Prevention of episodes of lymphangitis and lymphoedema: hygiene of the affected extremity^b, comfortable footwear, immediate attention to secondary bacterial/fungal infections and wounds.
- Established lymphoedema: bandaging of the affected limb by day, elevation of the affected extremity (after removal of the bandage) when at rest, simple exercises (flexion-extension of the feet when recumbent or upright, rotation of the ankles); skin hygiene, as above.

Surgery

May be indicated in the treatment of chronic manifestations: advanced lymphoedema (diversion-reconstruction), hydrocoele and its complications, chyluria.

Footnotes

- (a) When test results are negative in a clinically suspect case, consider detection of antigens (ICT rapid test) and/or ultrasound of the inguinal area in search of the « filarial dance sign ».
- (b) Wash at least once daily (soap and water at room temperature), paying special attention to folds and interdigital areas; rinse thoroughly and dry with a clean cloth; nail care.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 7: Bacterial diseases

[Bacterial meningitis](#)

[Tetanus](#)

[Enteric \(typhoid and paratyphoid\) fevers](#)

[Brucellosis](#)

[Plague](#)

[Leptospirosis](#)

[Relapsing fever \(borreliosis\)](#)

[Louse-borne relapsing fever \(LBRF\)](#)

[Tick-borne relapsing fever \(TBRF\)](#)

[Eruptive rickettsioses](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Bacterial meningitis

Meningitis is an acute bacterial infection of the meninges, which may affect the brain and lead to irreversible neurological damage and auditory impairment.

Bacterial meningitis is a medical emergency. The treatment is based on early parenteral administration of antibiotics that penetrates well into the cerebrospinal fluid (CSF). Empiric antibiotic therapy is administered if the pathogen cannot be identified or while waiting for laboratory results.

The main bacteria responsible vary depending on age and/or context:

Meningitis in a non-epidemic context

- Children 0 to 3 months:
 - Children ≤ 7 days: Gram-negative bacilli (*Klebsiella spp*, *E. coli*, *S. marcescens*, *Pseudomonas spp*, *Salmonella spp*) and group B streptococcus
 - Children > 7 days: *S. pneumoniae* accounts for 50% of all bacterial meningitis. *L. monocytogenes* is occasionally responsible for meningitis during this period.
- Children 3 months-5 years: *S. pneumoniae*, *H. influenza B* and *N. meningitidis*
- Children > 5 years and adults: *S. pneumoniae* and *N. meningitidis*

Special conditions:

- Immunodepressed patients (HIV, malnourished): high percentage of Gram- negative bacilli (specially *Salmonella spp*) and also *M. tuberculosis*.
- Sickle cell anaemia: *Salmonella spp* and *Staphylococcus aureus* are frequent causes.
- Meningitis may be related to *S. aureus* when associated with skin infection or skull fracture.

Meningitis in an epidemic context

In the Sahelian region (but not exclusively, e.g. Rwanda, Angola, Brazil), during the dry season, epidemics of meningococcal meningitis (*Neisseria meningitidis* A or C or W135) affect children from 6 months of age, adolescents and adults. In these regions, whether during epidemics or not, all the above pathogens can be found, especially in young children.

Clinical features

The clinical presentation depends on the patient's age.

Children over 1 year and adults

- Fever, severe headache, photophobia, neck stiffness
- Brudzinski's sign (neck flexion in a supine patient results in involuntary flexion of the knees) and Kernig's sign
(attempt to extend the knee from the flexed thigh position and with the hip flexed)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Laboratory

- Lumbar puncture (LP):
 - Macroscopic examination of CSF: antibiotic therapy should be initiated immediately if the LP yields a turbid CSF.
 - Microscopic examination: Gram stain (but a negative examination does not exclude the diagnosis) and white blood cell count (WBC).
 - In an epidemic context, once the meningococcal aetiology has been confirmed, there is no need for routine LP for new cases.

	Pressure	Aspect	WBC (leucocytes/mm ³)	Protein	Other tests
Normal CSF		Clear	< 5	Pandy- < 40 mg/dl	-
Bacterial meningitis	++++	Cloudy, turbid	100-20 000 mainly neutrophiles In neonates: > 20 In immunocompromised, the WBC may be < 100	Pandy+ 100-500 mg/dl	Gram stain +
Viral meningitis	Normal to +	Clear	10-700 mainly lymphocytes	Pandy-	-
TB meningitis	+++	Clear or yellowish	< 500 mainly lymphocytes	Pandy+	AFB
Cryptococcal meningitis	++++	Clear	< 800 mainly lymphocytes	Pandy-	India ink

- Rapid test for detection of bacterial antigens.

Note: in an endemic area, it is essential to test for severe malaria (rapid test or thin/thick films).

Treatment in a non-epidemic context

Antibiotic therapy

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	No associated skin infection		Associated skin infection (including umbilical cord infection)	
	First line	Alternative	First line	Alternative
0 to 7 days < 2 kg	ampicillin IV 100 mg/kg every 12 hours + cefotaxime IV 50 mg/kg every 12 hours	ampicillin IV 100 mg/kg every 12 hours + gentamicin IV 3 mg/kg once daily	cloxacillin IV 50 mg/kg every 12 hours + cefotaxime IV 50 mg/kg every 12 hours	cloxacillin IV 50 mg/kg every 12 hours + gentamicin IV 3 mg/kg once daily
0 to 7 days ≥ 2 kg	ampicillin IV 100 mg/kg every 8 hours + cefotaxime IV 50 mg/kg every 8 hours	ampicillin IV 100 mg/kg every 8 hours + gentamicin IV 5 mg/kg once daily	cloxacillin IV 50 mg/kg every 8 hours + cefotaxime IV 50 mg/kg every 8 hours	cloxacillin IV 50 mg/kg every 8 hours + gentamicin IV 5 mg/kg once daily
8 days to < 1 month ≥ 2 kg	ampicillin IV 100 mg/kg every 8 hours + cefotaxime IV 50 mg/kg every 8 hours	ampicillin IV 100 mg/kg every 8 hours + gentamicin IV 5 mg/kg once daily	cloxacillin IV 50 mg/kg every 6 hours + cefotaxime IV 50 mg/kg every 8 hours	cloxacillin IV 50 mg/kg every 6 hours + gentamicin IV 5 mg/kg once daily
1 to 3	ampicillin IV	ampicillin IV	cloxacillin IV	cloxacillin

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

> 3 months	ceftriaxone IV Children: 100 mg/kg on D1 then starting on D2: 100 mg/kg once daily or 50 mg/kg every 12 hours (max. 4 g daily)	cloxacillin IV Children < 40 kg: 50 mg/kg every 6 hours Children ≥ 40 kg: 2 g every 6 hours + ceftriaxone IV Children: 100 mg/kg on D1 then starting on D2: 100 mg/kg once daily or 50 mg/kg every 12 hours (max. 4 g daily)
Adults	ceftriaxone IV: 4 g once daily or 2 g every 12 hours	cloxacillin IV: 2 g every 6 hours + ceftriaxone IV: 4 g once daily or 2 g every 12 hours

Duration of antibiotic therapy:

1) According to the pathogen:

- *Haemophilus influenzae*: 7 days
- *Streptococcus pneumoniae*: 10-14 days
- Group B streptococcus and *Listeria*: 14-21 days
- Gram-negative bacilli: 21 days
- *Neisseria meningitidis*: see antibiotic therapy in an epidemic context

2) If the pathogen is unknown:

- Children < 3 months: 2 weeks beyond the first sterile CSF culture or 21 days
- Children > 3 months and adults: 10 days. Consider extending treatment or alternative diagnoses if fever persists beyond 10 days. On the other hand, a 7-day course of ceftriaxone is sufficient in patients who are making an uncomplicated recovery.

Additional treatment

- Dexamethasone reduces the risk of hearing loss in patients with *H. influenzae* or *S. pneumoniae*. Early administration is indicated in meningitis caused by these pathogens or when the pathogen is unknown, except in neonates (and in presumed meningococcal meningitis in an epidemic context). **dexamethasone IV**^{[1][2]}
Children > 1 month: 0.15 mg/kg (max. 10 mg) every 6 hours for 2 to 4 days
Adults: 10 mg every 6 hours for 2 to 4 days
The treatment should be started before or with the first dose of antibiotic, otherwise, the treatment offers no benefit.
- Ensure that the patient is well fed and well hydrated (infusions or nasogastric tube if necessary).
- [Seizures](#) (Chapter 1).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Age	Treatment ^[3]
Children < 2 months	ceftriaxone IV ^(a) or IM ^(b) for 7 days 100 mg/kg once daily
Children ≥ 2 months and adults	ceftriaxone IV ^(a) or IM ^(b) or 5 days Children 2 months to < 5 years: 100 mg/kg once daily (max. 2 g daily) Children ≥ 5 years and adults: 2 g once daily

(a) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must never be administered by IV route. For IV administration, water for injection must always be used.

(b) For IM administration, divide the dose into 2 injections if needed, half-dose in each buttock.

Note:

A short treatment with a single dose of ceftriaxone IM can be used in children 2 years and older and in adults during a meningococcal meningitis epidemic if 1) confirmed by a reliable laboratory 2) the number of cases exceeds management capacities with the 5-day treatment. Check national recommendations. Nevertheless, it is essential to ensure a monitoring of cases after 24 hours.

ceftriaxone IM^a

Children 2 to < 12 years: 100 mg/kg single dose

Children ≥ 12 years and adults: 4 g single dose

If there is no clinical improvement (fever > 38.5 °C, repeated seizures, appearance or aggravation of a reduced level of consciousness or of neurological signs) 24 hours after the injection, continue the treatment with ceftriaxone for 5 days.

Additional treatment

- Ensure that the patient is well fed and well hydrated (infusions or nasogastric tube if necessary).
- [Seizures](#) (Chapter 1).
- Coma: prevention of bed sores, care of the mouth and eyes, etc.
- Dexamethasone is not indicated.

Footnotes

(a) For IM administration, divide the dose into 2 injections if needed, half-dose in each buttock.

References

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Tetanus

Last updated: August 2022

Tetanus is a severe infection due to the bacillus *Clostridium tetani*, found in soil, and human and animal waste. The infection is noncontagious.

Clostridium tetani is introduced into the body through a wound and produces a toxin whose action on the central nervous system is responsible for the symptoms of tetanus.

Tetanus is entirely preventable by vaccination. It occurs in people who have not been fully vaccinated before exposure or have not received adequate post-exposure prophylaxis. In these individuals, most breaks in the skin or mucous membranes carry a risk of tetanus, but the wounds with the greatest risk are: the stump of the umbilical cord in neonates, puncture wounds, wounds with tissue loss or contamination with foreign material or soil, avulsion and crush injuries, sites of non-sterile injections, chronic wounds (e.g. lower extremity ulcers), burns and bites. Surgical or obstetrical procedures performed under non-sterile conditions also carry a risk of tetanus.

Clinical features

Generalised tetanus is the most frequent and severe form of the infection. It presents as muscular rigidity, which progresses rapidly to involve the entire body, and muscle spasms, which are very painful. Level of consciousness is not altered.

Children and adults

- Average time from exposure to onset of symptoms is 7 days (3 to 21 days).
- Muscular rigidity begins in the jaw muscles (difficulty with then inability to open mouth [trismus] preventing the patient from speaking, eating), spreading to the face (fixed smile), neck (difficulty with swallowing), to the trunk (restriction of respiratory muscles; hyperextension of spine [opisthotonus]), to the abdomen (guarding) and to the limbs (flexion of the upper limbs and extension of the lower limbs).
- Muscle spasms, which are very painful, appear at the onset or when muscular rigidity becomes generalised. They are triggered by stimuli (noise, light, touch) or arise spontaneously. Spasms of the thoracic and laryngeal muscles may cause respiratory distress or aspiration.

Neonates

- In 90% of cases, initial symptoms appear within 3 to 14 days of birth.
- The first signs are significant irritability and difficulty sucking (rigidity of the lips, trismus) then rigidity becomes generalised, as in adults. Any neonate, who initially sucked and cried normally, presenting with irritability and difficulty sucking 3 to 28 days after birth and demonstrating rigidity and muscle spasms should be assumed to have neonatal tetanus.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Handle the patient carefully, while sedated and as little as possible; change position every 3 to 4 hours to avoid bedsores.
- Teach family the danger signs and instruct them to call the nurse for the slightest respiratory symptom (cough, difficulty breathing, apnoea, excessive secretions, cyanosis, etc.).
- Establish IV access for hydration, IV injections.
- Gentle suction of secretions (mouth, oropharynx).
- Insert a nasogastric tube for hydration, feeding and administration of oral medications.
- Provide hydration and nutrition in feeds divided over 24 hours. In neonates, give expressed breast milk every 3 hours (risk of hypoglycaemia).

Neutralisation of toxin

human tetanus immunoglobulin IM

Neonates, children and adults: 500 IU single dose, injected into 2 separate sites

Inhibition of toxin production

metronidazole^a IV infusion (30 minutes; 60 minutes in neonates) for 7 days

- Neonates:
 - 0 to 7 days: 15 mg/kg on D1 then, after 24 hours, 7.5 mg/kg every 12 hours
 - 8 days to < 1 month (< 2 kg): same doses
 - 8 days to < 1 month (≥ 2 kg): 15 mg/kg every 12 hours
- Children 1 month and over: 10 mg/kg every 8 hours (max. 1500 mg daily)
- Adults: 500 mg every 8 hours

Control of rigidity and spasms, and sedation of the patient

Diazepam should decrease the frequency and intensity of spasms without causing respiratory depression. The dose and frequency of administration depend on the patient's clinical response and tolerance.



- There is a high risk of respiratory depression and hypotension when using diazepam, especially in children and elderly patients. Constant and close monitoring of the patient's respiratory rate (RR) and oxygen saturation (SpO_2) is essential, with immediate availability of equipment for manual ventilation (Ambu bag, face mask) and intubation, suction (electric if possible) and Ringer lactate.
- A continuous IV infusion of diazepam requires the use of a dedicated vein (no other infusion/injection in this vein); avoid the antecubital fossa if possible.
- Do not stop treatment abruptly; an abrupt stop can cause spasms.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<p>Neonates</p>	<p>diazepam emulsion for injection (10 mg ampoule, 5 mg/ml, 2 ml)</p> <ul style="list-style-type: none"> • 0.1 to 0.3 mg/kg by slow IV injection (3 to 5 minutes) every 1 to 4 hours depending on the severity and the persistence of the spasms as long as the RR is ≥ 30. • If despite hourly diazepam the spasms persist, start a continuous infusion of diazepam with an electric syringe: 0.1 to 0.5 mg/kg/hour (2.4 to 12 mg/kg every 24 hours). Start with 0.1 mg/kg/hour and if symptoms persist, increase by 0.1 mg/kg/hour as long as RR is ≥ 30. • If in spite of 0.5 mg/kg/hour symptoms persist, the dose can be increased up to 0.8 mg/kg/hour as long as the RR ≥ 30. • Diluted diazepam emulsion does not keep for more than 6 hours. <p>Example:</p> <p>Neonate weighing 3 kg (administration by electric syringe)</p> $0.1 \text{ mg/kg/hour} \times 3 \text{ kg} = 0.3 \text{ mg/hour}$ <p>Dilute one 10 mg ampoule of diazepam emulsion for injection in 50 ml of 10% glucose to obtain a solution containing 0.2 mg of diazepam per ml.</p> <p>Administer 1.5 ml/hour [dose (in mg/hour) \div dilution (in mg/ml) = dose in ml/hour i.e. 0.3 (mg/hour) \div 0.2 (mg/ml) = 1.5 ml/hour].</p> <p>If an electric syringe is not available, diluting the diazepam emulsion in an infusion bag for continuous infusion may be considered. Weigh the risks associated with this mode of administration (accidental bolus or insufficient dose). The infusion should be monitored closely to avoid any change, however small, of the prescribed rate.</p>
<p>Children > 1 month and adults</p>	<p>Same doses and protocol as in neonates but:</p> <ul style="list-style-type: none"> • Use diazepam solution for injection 5 mg/ml: (10 mg ampoule, 5 mg/ml, 2 ml). ^(a) • These doses can be administered as long as the RR is: <ul style="list-style-type: none"> ≥ 30 in children under 1 year ≥ 25 in children 1 to 4 years ≥ 20 in children 5 to 12 years ≥ 14 in children over 12 years ≥ 12 in adults <p>Examples:</p> <ul style="list-style-type: none"> • Child weighing 6 kg (continuous IV infusion using a pediatric infusion set; 1 ml = 60 drops) $0.1 \text{ mg/kg/hour} \times 6 \text{ kg} = 0.6 \text{ mg/hour}$ <p>Dilute one 10 mg ampoule of diazepam solution for injection in 50 ml of 5% glucose (10% glucose if child < 3 months) to obtain a solution containing 0.2 mg of diazepam per ml.</p> <p>Administer 3 ml/hour [dose (in mg/hour) \div dilution (in mg/ml) = dose in ml/hour i.e. 0.6 (mg/hour) \div 0.2 (mg/ml) = 3 ml/hour] or 3 drops/minute (in a paediatric infusion set ml/hour = drops/minute).</p> • Adult weighing 60 kg (standard adult infusion set, 1 ml = 20 drops) $0.1 \text{ mg/kg/hour} \times 60 \text{ kg} = 6 \text{ mg/hour}$

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

When the frequency and severity of the spasms have decreased, start weaning the diazepam (gradually decrease the rate of infusion):

- Calculate the total daily dose of IV diazepam and administer it orally in 4 divided doses, 6 hours apart, via nasogastric (NG)^b tube.
- Give first NG dose and decrease rate of IV infusion by 50%.
- Give second NG dose and stop IV diazepam infusion.
- If withdrawal signs^c appear, wean more slowly.
- Once on diazepam PO, wean by 10 to 20% of the original dose daily, until at a dose of 0.05 mg/kg every 6 hours.
- Then increase the interval from every 6 hours to every 8 hours for 24 hours as tolerated (wean more slowly if withdrawal signs appear).
- Continue to increase the interval between the doses from every 8 hours to every 12 hours and then to every 24 hours before stopping the diazepam.
- Each step should be for 24 hours or more if withdrawal signs appear.

Notes:

- It is often at these smaller doses that it is difficult to wean diazepam. If this is the case, slow the wean further: dropping the % wean (e.g. 5% wean every 24 hours instead of 10% wean) or increasing the interval between weans (e.g. going from every 24 hours to every 48 hours).
- If the patient is also receiving morphine, wean diazepam first then, wean morphine.
- Non-pharmacological measures to reduce withdrawal: reduce environmental stimuli; swaddle infants, frequent feedings.
- Infants who have had tetanus remain hypertonic, even when they are no longer having spasms.

Treatment of pain

morphine PO (via nasogastric tube) if necessary (see [Pain](#), Chapter 1).

When morphine is administered with diazepam the risk of respiratory depression is increased, thus closer monitoring is required. When morphine is no longer required, wean the same way as diazepam.

Treatment of the entry point and associated infections

- Search systematically the entry wound. Provide local treatment under sedation: cleansing and for deep wounds, irrigation and debridement.
- Cord infection: do not excise or debride; treat bacterial omphalitis and sepsis, add to metronidazole IV: cloxacillin IV + cefotaxime IV or cloxacillin IV + gentamicin IV (for doses, see [Bacterial meningitis](#)).

Tetanus vaccination

As tetanus does not confer immunity, vaccination against tetanus must be administered once the patient has recovered.

In case of neonatal tetanus, initiate the vaccination of the mother.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Depending on pre-exposure vaccination status:
Tetanus vaccine (TV)^d and immunoglobulin: see indications below.

Type of wound	Complete vaccination (3 or more doses) Time since administration of last dose:			Incomplete vaccination (less than 3 doses) or no vaccination or unknown status
	< 5 years	5-10 years	> 10 years	
Minor, clean	None	None	TV 1 booster dose	Initiate or complete TV
Other	None	TV 1 booster dose	TV 1 booster dose	Initiate or complete TV and administer tetanus immunoglobulin

tetanus vaccine IM

Children and adults: 0.5 ml per dose

If no vaccination or unknown vaccination status: administer at least 2 doses at an interval of 4 weeks.

If incomplete vaccination: administer one dose.

Then, to ensure long-lasting protection, administer additional doses to complete a total of 5 doses, as indicated in the table below.

human anti-tetanus immunoglobulin IM

Children and adults: 250 IU single dose; 500 IU for wounds more than 24 hours old.

Inject the vaccine and the immunoglobulin in 2 different sites, using a separate syringe for each.

2) Routine vaccination (pre-exposure prophylaxis)

- Children: 6 doses in total: a first series of 3 doses of DTP or DTP + HepB or DTP + HepB + Hib before the age of 1 year, administered at an interval of 1 month (e.g. at the age of 6, 10 and 14 weeks), then a dose of a vaccine containing tetanus toxoid between the age of 12 and 23 months, a dose between the age of 4 to 7 years, then a dose between the age of 12 and 15 years.
- Women of childbearing age: 5 doses during the reproductive years: a series of 3 doses of Td with an interval of at least one month between the first and second dose and an interval of at least 6 months between the second and third dose, then two other doses, each at minimum interval of one year, e.g. during pregnancies (see table below).
- Pregnant women: if a woman has never been vaccinated or if her vaccination status is unknown: 2 doses of Td during the pregnancy to reduce the risk of tetanus in mother and neonate: the first as soon as possible during the pregnancy and the second at least 4 weeks later and at least 2 weeks before delivery. This vaccination schedule protects more than 80% of neonates from tetanus. A single dose offers no protection

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Dose	Vaccination schedule in adults	Degree and duration of protection
TV1	On first contact with the health care system or as soon as possible during pregnancy	No protection
TV2	At least 4 weeks after TV1	80% 1 to 3 years
TV3	6 months to 1 year after TV2 or during the following pregnancy	95% 5 years
TV4	1 to 5 years after TV3 or during the following pregnancy	99% 10 years
TV5	1 to 10 years after TV4 or during the following pregnancy	99% Throughout the reproductive years

Footnotes

- (a) Clindamycin IV for 7 days is an alternative (for doses, see [Periorbital and orbital cellulitis](#), Chapter 5).
- (b) Administration of oral diazepam tablets to infants: calculate the exact dose of diazepam, e.g. to obtain 0.5 mg of diazepam, cut a scored diazepam 2 mg tablet in half along scoring then split in half again. Crush quarter tablet and dissolve in expressed breast milk or infant formula.
- (c) Withdrawal signs: excessive irritability, tremors, increased muscle tone, frequent yawning, poor feeding, watery stools and sweating.
- (d) Tetanus-containing vaccine, such as Td or DTP or DTP + HepB or DTP + HepB + Hib according to availability and patient's age.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Enteric (typhoid and paratyphoid) fevers

Last updated: September 2022

Enteric fevers include typhoid fever, due to *Salmonella enterica* serotype Typhi (*S. Typhi*) and paratyphoid fever, due to *Salmonella enterica* serotype Paratyphi A, B or C (*S. Paratyphi*).

Enteric fevers are acquired by the ingestion of water or food contaminated with excreta of symptomatic or asymptomatic carriers or by direct contact (dirty hands).

Enteric fevers are endemic in South, Central and Southeast Asia, sub-Saharan Africa, Oceania and, to a lesser extent, in Latin America.

Effective treatment significantly reduces the risk of complications and death.

Clinical features

Clinical manifestations of typhoid and paratyphoid fevers are the same. Enteric fevers have insidious onset and vary from mild to severe.

- The characteristic sign is prolonged fever, which gradually increases during the first week, plateaus the second week then decreases between the third and fourth week.
- Non-specific signs and symptoms are frequently associated: gastrointestinal disturbances (abdominal pain, constipation or diarrhoea, vomiting), headache, malaise, chills, fatigue, non productive cough and/or hepatosplenomegaly.
- Erythematous maculopapular rash on the trunk extreme fatigue and/or relative bradycardia (heart rate-temperature dissociation) may be present.
- Serious complications affect about 27% of hospitalised patients^[1] and usually occur during the second or third week of illness. These may include decreased level of consciousness, intestinal haemorrhage or perforation or peritonitis, shock, or nephritis. In pregnant women, severe infection may lead to foetal complications (miscarriage, preterm delivery, intrauterine death).
- Relapse may occur 2 to 3 weeks after recovery. It is usually not due to antibiotic resistance, and re-treatment is required.

Clinical diagnosis is difficult as enteric fevers resembles other infections present in regions where they are endemic. The main differential diagnoses are: [malaria](#), [brucellosis](#), [leptospirosis](#), [typhus](#), [rickettsiosis](#), [sepsis](#) and [dengue](#).

Laboratory

- Culture of *S. Typhi* or Paratyphi and drug susceptibility test (blood and stool specimens).
- In all cases, rapid test for malaria in endemic regions (and antimalarial treatment if needed. see [Malaria](#). Chapter 6).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Strains resistant to chloramphenicol, ampicillin/amoxicillin and co-trimoxazole (multidrug-resistant, MDR strains) are present in most parts of the world.
- Ciprofloxacin is used as first-line treatment in some countries, however fluoroquinolone resistance is endemic in Asia and is increasing in several parts of the world^[2].
- Ceftriaxone resistance has been identified in several regions^[2].
- MDR strains also resistant to fluoroquinolones and third-generation cephalosporins (extensively drug-resistant, XDR strains) have developed^[3].

Uncomplicated cases (outpatients)

Uncomplicated cases (the vast majority of cases) can be treated with an oral antibiotic treatment.

- First-line antibiotics:

azithromycin PO for 7 days (including for MDR and XDR cases, and pregnant women)

Children: 10 to 20 mg/kg (max. 1 g) once daily

Adults: 500 mg to 1 g once daily or 1 g on D1 then 500 mg once daily

or

cefixime PO for 10 to 14 days (except for third-generation cephalosporin resistance and XDR cases)

Children: 10 mg/kg (max. 200 mg) 2 times daily

Adults: 200 mg 2 times daily
- Alternatives include, only if recent data show susceptibility of strains to these antibiotics in the region:

amoxicillin PO for 14 days

Children: 30 mg/kg (max. 1 g) 3 times daily

Adults: 1 g 3 times daily

or

co-trimoxazole PO for 14 days

Children: 20 mg SMX + 4 mg TMP/kg (max. 800 mg SMX + 160 mg TMP) 2 times daily

Adults: 800 mg SMX + 160 mg TMP 2 times daily

Severe cases (inpatients)

- Severe cases include:
 - toxic appearance or decreased level of consciousness or medical or surgical complication;
 - oral administration not possible due to persistent vomiting.

These cases should be treated under close monitoring. Antibiotic treatment is initially parenteral, then oral when there is decreasing fever, clinical improvement and the patient can tolerate oral treatment.
- Start with **ceftriaxone IV^a** (including for pregnant women)
- Children: 50 to 100 mg/kg (max. 4 g) once daily
- Adults: 2 g once daily or 2 times daily
- Then change to azithromycin PO (as above) to complete at least 7 days of treatment.
- For suspected or confirmed ceftriaxone resistance or XDR stains, use meropenem IV, including for pregnant women, then change to azithromycin PO to complete at least 7 days of treatment.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Hygiene measures common to all diarrhoeas: handwashing; consumption of treated water (chlorinated, boiled, bottled, etc.); washing/cooking of food, etc.
- In hospitals: for patients with watery diarrhoea, consider disinfection of excreta with chlorinated solution, if stools are collected in buckets.
- Vaccination with the **typhoid conjugate vaccine** in endemic regions^c. This vaccine can be used to control typhoid outbreaks. It does not protect against paratyphoid fever.

Footnotes

- (a) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must NEVER be administered by IV route. For IV administration, water for injection must always be used.
- (b) Do not add metronidazole if the patient receives meropenem (meropenem already covers anaerobic bacteria).
- (c) For more information, see Typhoid vaccines: WHO position paper:
<http://apps.who.int/iris/bitstream/handle/10665/272272/WER9313.pdf?ua=1>

References

1. Cruz Espinoza LM, McCready E, Holm M, et al. Occurrence of typhoid fever complications and their relation to duration of illness preceding hospitalization: a systematic literature review and meta-analysis. Clin Infect Dis 2019;69(Suppl 6):S435-48. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6821330/> [Accessed 28 June 2022]
2. Browne AJ, Hamadani BHK, Kumaran EAP, Rao P, et al. Drug-resistant enteric fever worldwide, 1990 to 2018: a systematic review and meta-analysis. BMC Medicine 2020;18:1+22. <https://doi.org/10.1186/s12916-019-1443-1> [Accessed 23 February 2022]
3. Klemm EJ, Shakoor S, Page AJ, Qamar FN, et al. Emergence of an Extensively Drug-Resistant *Salmonella enterica* Serovar Typhi Clone Harboring a Promiscuous Plasmid Encoding Resistance to Fluoroquinolones and Third-Generation Cephalosporins. mBio. 2018 Jan-Feb; 9(1): e00105-18. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5821095/> [Accessed 26 June 2022]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Brucellosis

Last updated: September 2022

Brucellosis is a zoonosis that mainly affects livestock animals.

The main routes of transmission to humans are:

- digestive, by ingestion of unpasteurized milk (or unpasteurized milk products) from an infected animal;
- cutaneous, by direct contact with infected animals or carcasses of infected animals.

Brucellosis is caused by bacteria of the genus *Brucella*, particularly *B. melitensis* (sheep and goats), *B. abortus* (cattle), *B. suis* (pigs).

The disease is found worldwide and mainly in rural areas.

After primary infection relapses may occur (5 to 15% of cases, even months after end of initial treatment) or the infection may become chronic.

Clinical features

Acute form (primary infection)

- Remittent or intermittent fever (39-40 °C), associated with several signs or symptoms: chills, night sweats, joint and muscle pain, weight loss, fatigue, malaise, headache; adenopathies (particularly in children).
- May be associated with: non-specific gastrointestinal disorders, cough, hepato and/or splenomegaly, arthritis (knee), orchitis.

Diagnosis is difficult because of the broad spectrum of fluctuating and non-specific clinical manifestations. In patients with unexplained fever, brucellosis should be considered when risk factors are present: consumption of unpasteurized milk products; exposure to livestock (e.g. livestock farmers, veterinarians, butchers, slaughterhouse workers).

Localised form

Primary infection may progress to localised infection (even several months or years later), mainly:

- osteoarticular: sacroiliac joint and often particularly lower limbs joints; spine (intervertebral disk infection, vertebral osteomyelitis)
- genito-urinary: orchitis, epididymitis
- pulmonary: bronchitis, pneumonia, pleurisy
- neurological : meningitis, encephalitis, polyneuritis

Paraclinical investigations

Laboratory

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Radiography

- Joint pain (hips, knees, ankles, vertebrae, sacroiliac joint): small erosions or destruction or joint space narrowing. Often involves the spine, particularly the lumbar spine, causing spondylodiskitis.
- Pulmonary signs: chest x-ray often normal. There may be consolidation, nodules, lymphadenopathy, or pleural effusion.

Treatment

Check national recommendations on antibiotic therapy. For information:

Children under 8 years	co-trimoxazole + rifampicin or co-trimoxazole + gentamicin
Children 8 years and over	doxycycline + rifampicin or doxycycline + gentamicin
Adults	doxycycline + rifampicin or doxycycline + streptomycin or gentamicin
Pregnant/breast-feeding women	rifampicin

co-trimoxazole PO for 6 weeks

Children < 8 years: 20 mg SMX + 4 mg TMP/kg (max. 800 mg SMX + 160 mg TMP) 2 times daily

doxycycline PO for 6 weeks

Children ≥ 8 years and < 45 kg: 2 to 2.2 mg/kg (max. 100 mg) 2 times daily

Children ≥ 45 kg and adults: 100 mg 2 times daily

rifampicin PO for 6 weeks

Children: 15 to 20 mg/kg (max. 600 mg) once daily

Adults: 600 to 900 mg once daily

gentamicin IM for 2 weeks

Children and adults: 5 mg/kg once daily

streptomycin IM for 2 weeks

Adults: 1 g once daily

For localised forms of the infection, same treatment but for a period of 6 weeks to 4 months depending on the focus.

Prevention

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Plague

Last updated: September 2022

Plague is a zoonosis caused by the Gram-negative bacillus *Yersinia pestis* that affects many wild and domestic mammals, particularly rodents.

Plague is transmitted to man by infected animals (direct contact or inhalation of their respiratory secretions), the bite of a flea of infected animals, or inhalation of respiratory secretions of individuals with pneumonic plague.

Natural foci of infection include Africa, Asia, North and South America, and parts of Europe.

Bubonic plague is the most common form, usually resulting from the bite of an infected flea. Without prompt treatment, the bacteria may be disseminated by haematogenous route, producing a more severe form (see below) with a high mortality rate.

The following forms of plague may be primary or secondary to bubonic plague:

- Pneumonic plague can rapidly progress to respiratory distress, shock, and death without prompt treatment.
- Septicaemic plague is a fulminant illness that can progress to disseminated intravascular coagulation, respiratory distress, shock, and death.
- Plague meningitis is a rare but very severe form of plague.

Clinical features

See [table](#) below.

Main differential diagnoses include:

- Other causes of lymphadenitis (e.g. some [bacterial skin infections](#), tularemia, [lymphogranuloma venereum, chancroid](#))
- [Acute pneumonia](#) (Chapter 2)
- Other causes of septicaemia (see [Shock](#), Chapter 1) or meningitis (see [Bacterial meningitis](#), Chapter 7)

Laboratory

- Collect pre-treatment specimens: lymph node aspirate (bubonic plague), sputum (pneumonic plague), blood (septicaemic plague), or cerebrospinal fluid (plague meningitis).
- Send specimens^a to reference laboratory for:
 - Rapid diagnostic test for detection of F1 capsular antigen of *Y. pestis*
 - PCR
 - Culture of *Y. pestis* and drug susceptibility test
- In all cases, rapid test for malaria in endemic regions (and antimalarial treatment if needed. see [Malaria](#). Chapter 6).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Treatment of suspected cases

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Forms of plague	Clinical features	Antibiotic treatment ^[1]
Bubonic	<ul style="list-style-type: none"> • Fever, chills, malaise, headache <p>AND</p> <ul style="list-style-type: none"> • Lymph node (bubo), painful, usually inguinal, (one or more) 	<p>Children (including < 8 years) and adults:</p> <p>doxycycline PO:</p> <ul style="list-style-type: none"> • Under 45 kg: 4.4 mg/kg (max. 200 mg) on D1, then 2.2 mg/kg (max. 100 mg) 2 times daily • 45 kg and over: 200 mg on D1, then 100 mg 2 times daily <p>or</p> <p>gentamicin IM or IV:^(a)</p> <ul style="list-style-type: none"> • Children: 4.5 to 7.5 mg/kg once daily • Adults: 5 mg/kg once daily <p>or</p> <p>ciprofloxacin PO:</p> <ul style="list-style-type: none"> • Children: 15 mg/kg 2 to 3 times daily (max. 750 mg 2 times daily or 500 mg 3 times daily) • Adults: 750 mg 2 times daily
Pneumonic	<ul style="list-style-type: none"> • Fever, chills, malaise, headache <p>AND</p> <ul style="list-style-type: none"> • Dyspnea, chest pain, productive cough with purulent or blood-stained sputum • Respiratory distress or failure and sepsis in severe or advanced, untreated disease 	<p>Children and adults:</p> <p>If mild disease:</p> <p>gentamicin IM or IV^(a) (as above)</p> <p>or</p> <p>ciprofloxacin PO (as above) or IV^(b)</p> <p>If severe disease:</p> <p>gentamicin + ciprofloxacin (as above)</p> <p>or, if not available, gentamicin + doxycycline (as above)</p> <p>After clinical improvement, change to ciprofloxacin or doxycycline PO (as above).</p>
Septicaemic	<ul style="list-style-type: none"> • Frequently no localizing symptoms or signs • Gastrointestinal disturbances (abdominal pain, vomiting, diarrhoea, etc.) often present 	As severe pneumonic plague
Meninaitis^(c)	Signs of meninaitis.	Children and adults:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- (a) **Streptomycin** IM may be an alternative to gentamicin (except in pregnant women):
 - Children: 15 mg/kg (max. 1 g) every 12 hours
 - Adults: 1 g every 12 hours
- (b) Use **ciprofloxacin** IV when oral route is not possible:
 - Children: 10 mg/kg (max. 400 mg) every 8 or 12 hours
 - Adults: 400 mg every 8 hours
- (c) If plague meningitis develops, add chloramphenicol to the existing regimen, and continue the combined regimen for an additional 10 days.

Treatment in pregnant women

- Bubonic, pneumonic, and septicaemic plague: **gentamicin** IM or IV (as above) + **ciprofloxacin** PO (500 mg 3 times daily) or IV (as above)
- Plague meningitis: **chloramphenicol** IV + **ciprofloxacin** PO (500 mg 3 times daily) or IV (as above)

Infection prevention and control (in hospitals)

- Bubonic plague: no isolation, standard precautions (handwashing, gowns, gloves, eye protection, etc.) with respect to lymph node aspiration or discharge and other body fluids.
- Pneumonic plague: isolation (in single room if possible), standard precautions, plus, for 48 hours after the start of antibiotic treatment, droplet precautions (medical mask for healthcare workers and for patients during contact). Only for aerosol-generating procedures, airborne precautions (FFP2 or N95 respirators) for health workers exposed to aerosols.
- Elimination of fleas (e.g. bedding, clothing, corpse): refer to the guide [Public health engineering](#), MSF.

Post-exposure prophylaxis of contacts

In the event of contact (distance less than 2 meters without appropriate personal protective equipment) with a pneumonic plague patient or direct contact with infected body fluids or tissues of any plague patient and within one week after the end of exposure:

doxycycline PO for 7 days

Children: 2.2 mg/kg (max. 100 mg) 2 times daily

Adults (including pregnant women): 100 mg 2 times daily

or

ciprofloxacin PO for 7 days

Children: 20 mg/kg (max. 750 mg) 2 times daily

Adults: 500 to 750 mg 2 times daily

Pregnant women: 500 mg 3 times daily

Prevention

- Flea vector control, sanitation and rodent reservoir control, refer to the guide [Public health engineering](#), MSF.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

References

1. Nelson CA, Meaney-Delman D, Fleck-Derderian S, Cooley KM, et al. Antimicrobial treatment and prophylaxis of plague: recommendations for naturally acquired infection and bioterrorism response. MMWR Recomm Rep 2021;70(No. RR-3):1-27. https://www.cdc.gov/mmwr/volumes/70/rr/rr7003a1.htm?s_cid=rr7003a1_w [Accessed 25 January 2022]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Leptospirosis

Last update: October 2022

Leptospirosis is a zoonosis that affects many domestic and wild animals, mainly rodents (particularly rats) but also dogs and cattle, etc.

It is transmitted to humans by contact through skin lesions or mucous membranes (e.g. eyes, mouth) with:

- freshwater or moist soil contaminated with urine of an infected animal (indirect contact);
- urine, blood and other body fluids or tissues of an infected animal (direct contact).

It is caused by bacteria (spirochetes) of the genus *Leptospira*.

Leptospirosis occurs worldwide, particularly in tropical and subtropical regions. There are often outbreaks after heavy rainfall or flooding.

Clinical features

Approximately 90% of cases are asymptomatic or mild with a favourable outcome. 5 to 15% of cases present a severe form with multiple organ dysfunction and a high mortality rate without prompt treatment.

Mild form

- Acute phase (septicaemic):
 - Sudden onset of high fever with chills, headache, myalgia (especially calf and lumbar pain), photophobia, ocular pain. Bilateral conjunctival suffusion affecting the bulbar conjunctiva (redness without discharge) is a characteristic sign, but not always present.
 - May be associated with: gastrointestinal symptoms (anorexia, abdominal pain, nausea, vomiting), non-productive cough, lymphadenopathy, hepatomegaly, and sometimes, skin rash.
- Immune phase:
 - The signs of the acute phase regress after 5 to 7 days then reappear for a few days usually in a milder form (milder fever, less severe myalgia) then disappear.
 - Signs of meningitis (thought to be of immune origin) are however very common during this phase.

Severe or ictero-haemorrhagic form

The onset is the same but a few days later the symptoms worsen: renal disorders (oliguria or polyuria), hepatic disorder (jaundice), widespread haemorrhages (purpura, ecchymoses, epistaxis, haemoptysis, etc.), pulmonary signs (chest pain) or cardiac signs (myocarditis, pericarditis).

Diseases with different forms of the bacterial culture of Leptospiral isolates. Patients that present the following

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- [Malaria](#)
- [Typhoid fever, brucellosis, rickettsioses](#)

Laboratory

Diagnosis

- Collect pre-treatment specimens and send them to reference laboratory:
 - Acute phase (first week of illness): blood and/or serum for IgM screening, PCR, and acute specimen for microscopic agglutination test (MAT);
 - Immune phase (second week of illness): serum for IgM screening and convalescent specimen for MAT, and urine for PCR.
- In all cases, rapid test for malaria in endemic regions (and antimalarial treatment if needed, see [Malaria](#), Chapter 6).

Other investigations

(if available)

- Serum creatinine: elevated if renal dysfunction.
- Full blood count: possible neutrophilia and thrombocytopenia (acute phase), or anaemia secondary to haemorrhage (immune phase).
- Cerebrospinal fluid (immune phase): features of aseptic meningitis in CSF (see [viral meningitis](#), Chapter 7).
- Urine: mild proteinuria, leukocyturia, possible microscopic haematuria (acute phase).

Treatment

Start empiric antibiotic treatment as soon as leptospirosis suspected, before results of diagnosis tests are available.

Mild form (outpatients)

Symptomatic treatment

- Rest and treatment of pain and [fever](#): paracetamol PO (Chapter 1).
- Acetylsalicylic acid (aspirin) is contra-indicated (risk of haemorrhage).

Antibiotic treatment

- **doxycycline** PO for 7 days

Children under 45 kg: 2 to 2.2 mg/kg (max. 100 mg) 2 times daily

Children 45 kg and over and adults: 100 mg 2 times daily

or, particularly in pregnant women:

- **azithromycin** PO for 3 days

Children: 10 mg/kg (max. 500 mg) on D1 then 5 mg/kg (max. 250 mg) once daily on D2 and D3

Adults : 1 g on D1 then 500 mg once daily on D2 and D3

or, if not available,

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Specific management according to organs affected. Oliguria generally responds to correction of hypovolaemia.
- Rest and treatment of pain and **fever**: paracetamol PO (Chapter 1). Avoid or use paracetamol with caution in patients with hepatic involvement.

Antibiotic treatment

- **ceftriaxone** IV for 7 days^a

Children: 80 to 100 mg/kg (max. 2 g) once daily

Adults: 2 g once daily

or

- **benzylpenicillin** IV for 7 days

Children: 50 000 IU (30 mg)/kg (max. 2 MIU or 1200 mg) every 6 hours

Adults: 1 to 2 MIU (600 to 1200 mg) every 6 hours

Prevention

- Avoid bathing in freshwater in endemic areas.
- Disinfect laundry and objects soiled by urine of infected animal or patient.
- Vaccination and protective clothing (only for professionals at risk of exposure).

Footnotes

(a) For IV administration of ceftriaxone, dilute with water for injection only.

References

1. World Health Organization. Human leptospirosis: guidance for diagnosis, surveillance and control. World Health Organization, 2003.
https://apps.who.int/iris/bitstream/handle/10665/42667/WHO_CDS_CSR_EPH_2002.23.pdf?%20sequence=1&isAllowed=y
[Accessed 5 September 2022]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Relapsing fever (borreliosis)

- [Louse-borne relapsing fever \(LBRF\)](#)
- [Tick-borne relapsing fever \(TBRF\)](#)

Relapsing fever (FR) is caused by spirochetes of the genus *Borrelia*, transmitted to humans by arthropod vectors.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Louse-borne relapsing fever (LBRF)

Last updated: October 2022

LBRF is caused by *Borrelia recurrentis*. It occurs in epidemic waves when conditions favourable to the transmission of body lice are met: cold climate/season, overcrowding and very poor sanitation (e.g. refugee camps, prisons). Endemic foci of LBRF are mainly the Sudan and the Horn of Africa (especially Ethiopia). LBRF can be associated with louse-borne typhus (see [Eruptive rickettsioses](#)). The mortality rate for untreated LBRF ranges from 15 to 40%.

Clinical features

- Relapsing fever is characterized by febrile episodes separated by afebrile periods of approximately 7 days (4 to 14 days).
- The initial febrile episode lasts up to 6 days:
 - Sudden onset of high fever (axillary temperature > 39 °C), severe headache and asthenia, diffuse pain (muscle, joint, back pain), often associated with gastrointestinal disturbances (anorexia, abdominal pain, vomiting, diarrhoea).
 - Splenomegaly is common; bleeding signs (e.g. petechiae, subconjunctival haemorrhage, epistaxis, bleeding gums), jaundice or neurological symptoms may be observed.
 - The febrile episode terminates in a crisis with an elevation in temperature, heart rate and blood pressure, followed by a fall in temperature and blood pressure, which may last for several hours.
- Following the initial febrile episode, the cycle usually recurs; each episode is less severe than the previous one and the patient develops temporary immunity.
- Complications:
 - collapse during defervescence, myocarditis, cerebral haemorrhage;
 - during pregnancy: abortion, preterm delivery, in utero foetal death, neonatal death.

In practice, in an applicable epidemiological setting (see above), a suspect case of LBRF is, according to WHO, a patient with high fever and two of the following symptoms: severe joint pain, chills, jaundice or signs of bleeding (nose or other bleeding) or a patient with high fever who is responding poorly to antimalarial drugs. Clothing should be checked for the presence of body lice and nits.

Laboratory

The diagnosis is confirmed by detection of *Borrelia* in thick or thin blood films (Giemsa stain). Blood samples must be collected during febrile periods. Spirochetes are not found in the peripheral blood during afebrile periods. In addition, the number of circulating spirochetes tends to decrease with each febrile episode.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

or

azithromycin PO

Children: 10 mg/kg (max. 500 mg) single dose

Adults: 500 mg single dose

- Treatment of pain and [fever](#) (paracetamol PO) and prevention or treatment of dehydration in the event of associated diarrhoea.
- Elimination of body lice is essential in control of epidemics (see [Pediculosis](#), Chapter 4).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Tick-borne relapsing fever (TBRF)

Last update: October 2022

TBRFs are caused by different *Borrelia* species. They are endemic in temperate and warm regions of the world, especially in Africa (Tanzania, DRC, Senegal, Mauritania, Mali, the Horn of Africa) and mainly in rural areas. TBRF is a major cause of morbidity and mortality in children and pregnant women. The mortality rate for untreated TBRF ranges from 2 to 15%.

Clinical features

The clinical manifestations and complications of TBRF are similar to those of LBRF but central nervous system (CNS) involvement (particularly lymphocytic meningitis) is more frequent than in LBRF and the number of relapses is higher.

The clinical diagnosis is difficult, especially during the first episode: cases occur sporadically rather than in outbreaks; the tick bite is painless and usually unnoticed by the patient; symptoms are very similar to those of [malaria](#), [typhoid fever](#), [leptospirosis](#), certain arbovirosis (yellow fever, [dengue](#)) or [rickettsiosis](#), and [meningitis](#).

Laboratory

- As for LBRF, the diagnosis is confirmed by detection of *Borrelia* in the patient's blood.
- Repeat the examination if the first smear is negative despite strong clinical suspicion.
- In all cases, rapid test for malaria in endemic regions (and antimalarial treatment if needed, see [Malaria](#), Chapter 6).

Treatment

- Antibiotic treatment:

doxycycline PO for 7 to 10 days

Children under 45 kg: 2.2 mg/kg (max. 100 mg) 2 times daily

Children 45 kg and over and adults: 100 mg 2 times daily

or

azithromycin PO for 7 to 10 days (if doxycycline is contra-indicated or not available)

Children: 10 mg/kg (max. 500 mg) once daily

Adults: 500 mg once daily

or

ceftriaxone IV^a for 10 to 14 days (for pregnant women or in case of CNS involvement)

Children: 50 to 75 mg/kg (max. 2 g) once daily

Adults: 2 g once daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Footnotes

(a) For IV administration of ceftriaxone, dilute with water for injection only.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Eruptive rickettsioses

Last update: October 2022

Rickettsioses are eruptive fevers caused by bacteria of the genus *Rickettsia* and transmitted to man by an arthropod vector. Three main groups are distinguished: typhus group, spotted fever group and scrub typhus group.

Clinical features

- Common to all forms:
 - Sudden onset of fever (temperature of over 39 °C) with severe headache and myalgias.
 - 3 to 5 days later; onset of generalised cutaneous eruption (see below).
 - Hypotension; non-dissociated rapid heart rate (variable).
 - Typhoid state: prostration, obnubilation, confusion and extreme asthenia, particularly marked in typhus forms.
 - Inoculation eschar: painless, black crusted lesion surrounded by a erythematous halo at the site of the bite.
Always check for this significant sign.
 - Non-cutaneous signs vary from one form to another, and are atypical and variable (see below).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Group	Typhus		Spotted fever			Scrub typhus
Form	Epidemic typhus	Murine typhus	Mediterranean spotted fever	Rocky Mountain spotted fever	Other Old-World tick-borne fevers	Scrub typhus
Pathogen	<i>R. prowasekii</i>	<i>R. typhi</i>	<i>R. conorii</i>	<i>R. rickettsii</i>	<i>R. sibirica</i> , <i>R. australis</i>	<i>O. tsutsugamushi</i>
Vector	body lice	rat fleas	ticks	ticks	ticks	mites
Reservoir	man	rats	dogs	rodents	rodents, dogs, etc.	rodents
Occurrence	epidemic	endemic	endemic	endemic	endemic	sporadic
Geographical distribution	worldwide, conflicts; main sites: Burundi/Rwanda, Ethiopia	worldwide	around the mediterranean, Sub-Saharan Africa	North America, Central America, Columbia, Brazil	Southern Africa, Australia, Siberia	Far East, India, South Pacific Area
Rash	maculopapular	maculopapular	maculopapular	purpural	maculopapular	macular
Eschar	0	0	black necrotic area	rare	black necrotic area	black necrotic area
Typhoid state	+++	+++	+/-	+/-	+/-	+++
Extra-cutaneous signs	cough, myalgia, meningeal signs	gastrointestinal signs	meningeal signs	gastrointestinal and neurological signs, hypotension	variables	meningeal signs
Case fatality (%)	30 (without treatment)	5	2	5	1	0-30

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Treatment

- Symptomatic treatment:
 - Hydration (PO or IV if the patient is unable to drink).
 - Fever: paracetamol PO (Chapter 1). Acetylsalicylic acid (aspirin) is contra-indicated due to the risk of haemorrhage.
- Antibiotic^a for 5 to 7 days or until 3 days after the fever has disappeared:
doxycycline PO

Children under 45 kg: 2.2 mg/kg (max. 100 mg) 2 times daily
Children 45 kg and over and adults: 100 mg 2 times daily
In severe infections, a loading dose of doxycycline is recommended:
Children under 45 kg: 4.4 mg/kg (max. 200 mg) on D1 then 2.2 mg /kg (max. 100 mg) 2 times daily
Children 45 kg and over and adults: 200 mg on D1 then 100 mg 2 times daily
- In a context of epidemic typhus, **doxycycline** PO is the choice treatment, but there is a risk of recurrence:

Children: 4 mg/kg (max. 100 mg) single dose
Adults: 200 mg single dose

Prevention

- Epidemic typhus: control of body lice (see [Pediculosis](#), Chapter 4).
- Murine typhus: control of fleas and then rats.
- Spotted fevers: avoid tick bites by wearing clothing and using repellents.
- Scrub typhus: use of repellents, chemoprophylaxis with **doxycycline** PO (200 mg once weekly in adults).

Footnotes

- (a) Unlike borrelioses, antibiotic treatment of rickettsioses does not provoke a Jarisch-Herxheimer reaction. However, the geographical distribution of borrelioses and rickettsioses may overlap, and thus a reaction may occur due to a possible co-infection (see [Borreliosis](#)).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 8: Viral diseases

[Measles](#)

[Poliomyelitis](#)

[Rabies](#)

[Viral hepatitis](#)

[Dengue](#)

[Viral haemorrhagic fevers](#)

[HIV infection and AIDS](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Measles

Measles is a highly contagious acute viral infection, transmitted by the airborne route (inhalation of respiratory droplets spread by infected individuals). The disease mainly affects children under 5 years of age and can be prevented by immunization.

For more information, refer to the guide [Management of a measles epidemic](#), MSF.

Clinical features

The average incubation period is 10 days.

Prodromal or catarrhal phase (2 to 4 days)

- High fever (39-40 °C) with cough, coryza (nasal discharge) and/or conjunctivitis (red and watery eyes).
- Koplik's spots: tiny bluish-white spots on an erythematous base, found on the inside of the cheek. This sign is specific of measles infection, but may be absent at the time of examination. Observation of Koplik's spots is not required for diagnosing measles.

Eruptive phase (4 to 6 days)

- On average 3 days after the onset of symptoms: eruption of erythematous, non-pruritic maculopapules, which blanch with pressure. The rash begins on the forehead then spreads downward to the face, neck, trunk (second day), abdomen and lower limbs (third and fourth day).
- As the rash progresses, prodromal symptoms subside. In the absence of complications, the fever disappears once the rash reaches the feet.
- The rash fades around the fifth day in the same order that it appeared (from the head to the feet).

The eruptive phase is followed by skin desquamation during 1 to 2 weeks, very pronounced on pigmented skin (the skin develops a striped appearance).

In practice, a patient presenting with fever and erythematous maculopapular rash and at least one of the following signs: cough or runny nose or conjunctivitis, is a clinical case of measles.

Complications

Most measles cases experience at least one complication:

- Respiratory and ENT: pneumonia, otitis media, laryngotracheobronchitis
- Ocular: purulent conjunctivitis, keratitis, xerophthalmia (risk of blindness)
- Gastrointestinal: diarrhoea with or without dehydration, benign or severe stomatitis
- Neurological: febrile seizures; rarely, encephalitis
- Acute malnutrition, provoked or aggravated by measles (post-measles period)

Pneumonia and dehydration are the most common immediate causes of death.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Severe oral lesions that prevent eating
- Acute malnutrition
- Treat as outpatient children with no major complications, no complications or minor complications:
 - Pneumonia without severe signs
 - Acute otitis media
 - Purulent conjunctivitis (no corneal lesions)
 - Diarrhoea without dehydration
 - Oral candidiasis that does not interfere with eating

If in doubt, keep the child under observation for a few hours.
- Isolation
 - Isolation of hospitalized patients
 - Measles cases treated as outpatients should be kept at home during this period.

Treatment

Supportive and preventive treatment

- Treat fever: paracetamol ([Fever](#), Chapter 1).
- Make the child drink (high risk of dehydration).
- Give smaller, more frequent meals or breastfeed more frequently (every 2 to 3 hours).
- Clear the nasopharynx (nose-blown or nasal lavages) to prevent secondary respiratory infection and improve the child's comfort.
- Clean the eyes with clean water 2 times daily and administer retinol on D1 and D2 (see [Xerophthalmia](#), Chapter 5) to prevent ocular complications.
- In children under 5 years: amoxicillin PO for 5 days as a preventive measure (reduction of respiratory and ocular infections).
- In the event of watery diarrhoea without dehydration: oral rehydration according to WHO *Plan A* (see [Dehydration](#), Chapter 1).
- Insert a nasogastric tube for a few days if oral lesions prevent the child from drinking.

Treatment of complications

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

	Treatment of complications
Severe pneumonia	ceftriaxone IV or IM + cloxacillin IV then change to amoxicillin/clavulanic acid PO (see Chapter 2) + oxygen if cyanosis or SpO ₂ < 90% + salbutamol if expiratory wheezing and sibilant rales on auscultation In all cases, close monitoring.
Pneumonia without severe signs	amoxicillin PO for 5 days
Croup	Inpatient monitoring (risk of worsening). Keep the child calm. Agitation and crying exacerbate the symptoms. For severe croup: dexamethasone IM : 0.6 mg/kg single dose + nebulized epinephrine (adrenaline) , 1 mg/ml ampoule): 0.5 ml/kg (max. 5 ml) + oxygen if cyanosis or SpO ₂ < 90% Intensive monitoring until symptoms resolve.
Acute otitis media	See Otitis , Chapter 2.
Dehydration	Per oral route or IV depending on the degree of dehydration.
Oral candidiasis	See Stomatitis , Chapter 3.
Purulent conjunctivitis	See Conjunctivitis , Chapter 5.
Keratitis/keratoconjunctivitis	tetracycline 1% eye ointment 2 times daily for 7 days + retinol PO one dose on D1, D2 and D8 (see Xerophthalmia , Chapter 5) + eye protection and pain management (see Pain , Chapter 1). No topical corticosteroids.
Xerophthalmia	See Xerophthalmia , Chapter 5.
Febrile seizures	See Seizures , Chapter 1.

Prevention

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Footnotes

- (a) Symptoms (hoarse crying or voice, difficulty breathing, a high-pitched inspiratory wheeze [inspiratory stridor], characteristic "barking" cough) are caused by inflammation and narrowing of the larynx. Croup is considered benign or "moderate" if the stridor occurs when the child is agitated or crying, but disappears when the child is calm. The child should be monitored during this period, however, because his general and respiratory status can deteriorate rapidly. Croup is severe when the stridor persists at rest or is associated with signs of respiratory distress.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Poliomyelitis

Poliomyelitis is an acute viral infection due to a poliovirus (serotypes 1, 2 or 3). Human-to-human transmission is direct (faecal-oral) or indirect (ingestion of food and water contaminated by stool). Humans are the only reservoir of the virus. In principle the disease can be eradicated by mass vaccination.

In endemic areas, poliomyelitis mainly affect children under 5 years not (or not fully) vaccinated, but the infection can affect persons of any age, especially in areas where population immunity is low.

Clinical features

- Up to 90% of cases are asymptomatic or present mild symptoms^[1].
- **Non-paralytic form:** a non-specific febrile illness with muscle pain, headache, vomiting, backache; no neurological involvement. As spontaneous recovery usually occurs within 10 days, diagnosis is rarely made outside epidemic contexts.
- **Paralytic form:** in less than 1% of cases, after the non-specific signs, the patient develops rapid onset (from the morning to the evening) asymmetrical acute flaccid paralysis, predominantly of the lower limbs, with ascending progression. The muscles become soft with diminished reflexes. Sensation is maintained. The disease is life threatening if paralysis involves the respiratory muscles or muscles of swallowing. Initial urinary retention is common. Gastrointestinal disturbances (nausea, vomiting, diarrhoea), muscle pain and meningeal symptoms may also occur.

Laboratory

Look for the polio virus in stool samples. The virus is excreted for one month after infection, but only intermittently; therefore, 2 samples must be collected with an interval of 24-48 hours, and within 14 days of onset of symptoms^[2]. Send the stool samples to a reference laboratory, with a clinical description of the patient. The stool samples must be stored and transported between 0 °C and 8 °C.

Treatment

- Hospitalise patients with the paralytic form: rest, prevent bed sores in bedridden patients, give analgesics (do not give IM injections to patients in the febrile phase), ventilate patients with respiratory paralysis.
- Physiotherapy once the lesions are stable to prevent muscle atrophy and contractures.
- Care for sequelae: physiotherapy, surgery and prosthetics.

Outbreak control in case of acute flaccid paralysis (AFP)

- Consider any patient with AFP as a suspected case of poliomyelitis.
- Send stool samples to a reference laboratory to confirm the diagnosis.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Prevention

- 3 types of vaccines exist:
 - a trivalent injectable inactivated poliovirus vaccine (IPV),
 - a bivalent oral live attenuated poliovirus vaccine (bOPV), containing serotypes 1 and 3,
 - a monovalent oral type 2 vaccine (mOPV or nOPV) exclusively used for responding to epidemics.
- Vaccination schedule: depends on the epidemiology of the virus.

Protocols vary according to the country, follow national recommendations. For information, the WHO recommends:

Schedule	Primary vaccination	
	Endemic or at risk zones ^(a)	Other zones
Birth	1 dose bOPV ^(b)	-
6 weeks	1 dose bOPV	1 dose bOPV
10 weeks	1 dose bOPV	1 dose bOPV
14 weeks	1 dose bOPV + 1 dose IPV	1 dose bOPV + 1 dose IPV

(a) Countries where poliomyelitis is endemic or zones at high risk of importation and subsequent spread of the virus.

(b) The first dose of bOPV is administered at birth, or as soon as possible, to optimise seroconversion rates after subsequent doses and induce mucosal protection.

In children who start routine vaccination late (after the age of 3 months), the dose of IPV is administered together with the first dose of bOPV, followed by 2 doses of bOPV alone administered 4 weeks apart.

There is also an 'IPV only' schedule: 3 doses administered at least 4 weeks apart (e.g. at 6, 10 and 14 weeks) and a booster dose at least 6 months later.

IPV should eventually completely replace bOPV.

References

1. World Health Organization. Poliomyelitis (polio). https://www.who.int/health-topics/poliomyelitis#tab=tab_1 [Accessed 08 June 2021]
2. Centers for Disease Control and Prevention. Poliomyelitis. 2020.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Rabies

Rabies is a viral infection of wild and domestic mammals, transmitted to humans by the saliva of infected animals through bites, scratches or licks on broken skin or mucous membranes.

In endemic areas (Africa and Asia), 99% of cases are due to dog bites and 40% of cases are children under 15 years of age^[1].

Before symptoms develop, rabies can effectively be prevented by post-exposure prophylaxis.

Once symptoms develop, rabies is fatal. There is no curative treatment; care is palliative.

Clinical features

- The incubation period averages 20 to 90 days from exposure (75% of patients), but can be shorter (in severe exposure, e.g. bites to face, head and hands; multiple bites), or longer (20% of patients develop symptoms between 90 days and 1 year, and 5% more than 1 year after exposure).
- Prodromal phase: itching or paraesthesiae or neuropathic pain around the site of exposure, and non-specific symptoms (fever, malaise, etc.).
- Neurologic phase:
 - Encephalitic form (furious form): psychomotor agitation or hydrophobia (throat spasms and panic, triggered by attempting to drink or sight/sound/touch of water) and aerophobia (similar response to a draft of air); sometimes seizures. The patient is calm and lucid between episodes. Infection evolves to paralysis and coma.
 - Paralytic form (less common, 20% of cases): progressive ascending paralysis resembling Guillain-Barré syndrome; evolves to coma.

Diagnosis is often difficult: there may be no history of scratch or bite (exposure through licking) or wounds may have healed; a reliable history may be difficult to obtain.

Post-exposure prophylaxis

Definitions of exposure categories (WHO)

Category I	Contact with animal, or licks on intact skin	No exposure
Category II	Nibbles on exposed skin Minor bite(s) or scratch(es) without bleeding	Minor exposure
Category III	Transdermal bite(s) or scratch(es) Licks on broken skin <small>Contamination of mucous membranes by animal's saliva (licks)</small>	Severe exposure

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Prolonged cleansing of the wound or contact site for 15 minutes to eliminate the virus, as soon as possible after exposure, is of critical importance. For skin: use soap, rinse copiously with running water, remove all foreign material; application of a disinfectant (povidone iodine 10% or other) is an additional precaution which does not take the place of thorough wound washing. For mucous membranes (eye, mouth, etc.): rinse thoroughly with water or 0.9% sodium chloride. Local cleansing is indicated even if the patient presents late.

According to condition/type of wound

In order to avoid inoculating virus deeper into the tissues, wounds are either not sutured at all (e.g. superficial, non-mutilating or puncture wounds), or are left open and re-evaluated in 48-72 hours, with a view to possible closure. Highly contaminated wounds, or wounds that may compromise function, require surgical management (exploration, removal of foreign material, excision of necrotic tissue, copious irrigation with sterile 0.9% sodium chloride or Ringer lactate, with local or general anaesthesia). When suturing is indicated (face), rabies immunoglobulin should be administered several hours before wound closure (see below). Infected wounds are not sutured and reassessed daily.

Passive and active immunisation

Given the duration of incubation, administration of vaccine/immunoglobulin is always a priority, even for patients exposed several months previously.

Anti-rabies serotherapy

Rabies immunoglobulin is indicated after:

- Category III exposures (except in patients who have received a full course of pre-exposure prophylaxis against rabies, see [Prevention](#));
- Category II and III exposures in immunocompromised patients^b (even in patients who have received a full course of pre-exposure prophylaxis against rabies).

It is intended to neutralize virus in the exposure site. It is given as a single dose on D0, with the first dose of rabies vaccine.

human rabies immunoglobulin:

Children and adults: 20 IU/kg

or

highly purified rabies immunoglobulin F(ab')2 fragments:

Children and adults: 40 IU/kg

Infiltrate rabies immunoglobulin into and around the previously washed wound(s). Ensure it is not injected into a blood vessel (risk of shock).

For finger wounds, infiltrate very cautiously to avoid increased pressure in the tissue compartment (compartment syndrome).

In the event of multiple wounds, dilute the dose 2- to 3-fold with sterile 0.9% sodium chloride to obtain a sufficient quantity to infiltrate all the sites exposed.

Infiltrate rabies immunoglobulin into the wound even if it has already healed.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

A complete prophylaxis series is indicated for Category II and III exposures. It should be started on D0 and continued to completion if the risk of rabies has not been excluded^c. Several different types of rabies vaccines prepared from cell cultures (CCEEV) exist. These vaccines must replace nerve tissue vaccines (NTV).

Prophylaxis schedules may vary from country to country, check national recommendations. The patient must be administered the full course of doses indicated.

Main post-exposure prophylaxis regimens^[1]

Date	No pre-exposure prophylaxis or unknown prophylaxis status or incomplete pre-exposure prophylaxis or complete pre-exposure prophylaxis with an NTV		
	IM route ^(a) 1 dose = 0.5 or 1 ml depending on the manufacturer	ID route ^(b) 1 dose = 0.1 ml	
D0	2 doses ^(c) (1 dose in each arm or thigh)	1 dose ^(c)	2 doses ^(c) (1 dose in each arm)
D3		1 dose	2 doses (1 dose in each arm)
D7	1 dose	1 dose	2 doses (1 dose in each arm)
D14		1 dose ^(d)	
D21	1 dose		

(a) IM route: there are two possible schedules, the Zagreb regimen (2-0-1-0-1) over 21 days or the 4-dose Essen regimen (1-1-1-0) over 14 to 28 days. The IM injection is administered into the anterolateral part of the thigh in children < 2 years; into the deltoid muscle (arm) in children ≥ 2 years and adults; do not administer into the gluteal muscle.

(b) ID route: inject into the deltoid muscle (or the suprascapular region or the anterolateral part of the thigh). Incorrect ID technique results in failure of post-exposure prophylaxis. If correct ID technique cannot be assured, use IM route.

(c) As well as a single dose of rabies immunoglobulin on D0 if indicated.

(d) The last injection can be administered between D14 and D28.

Notes:

- In immunocompromised patients: 1 dose on D0, 1 dose on D7 and 1 dose between D21 and D28.^[1]
- In patients that have received a full course of pre-exposure prophylaxis (see [Prevention](#)), the post-exposure

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Infection present	No infection and	No infection and
<ul style="list-style-type: none"> local: redness, oedema, serosanguinous or purulent drainage locoregional or general: lymphangitis, lymphadenopathy, localised cellulitis, bone or joint infection, fever 	<ul style="list-style-type: none"> wounds on the face or hands or genital region wounds involving joint, tendon, ligament or fracture deep puncture wounds wounds with crush injury wounds very contaminated or requiring debridement wounds where correct debridement is not possible immunocompromised patients 	<ul style="list-style-type: none"> no criteria requiring antibiotic prophylaxis wounds more than 24-48 hours old
Antibiotherapy PO 7 days in the event of local non severe infection; 14 days in the event of severe local infection, or widespread generalised infection.	Antibiotic prophylaxis PO 5 to 7 days	No antibiotic prophylaxis

The same dosage is used for both treatment and prophylaxis:

The treatment of choice is **amoxicillin/clavulanic acid (co-amoxiclav) PO^d**

Use formulations in a ratio of 8:1 or 7:1. The dose is expressed in amoxicillin:

Children < 40 kg: 25 mg/kg 2 times daily

Children ≥ 40 kg and adults:

Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)

Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)

Tetanus vaccination and serotherapy

Check prophylaxis status. If unknown or not up-to-date, see [Tetanus](#), Chapter 7.

Prevention

Pre-exposure prophylaxis with a CCEEV for people at risk (prolonged stay in rabies endemic areas, professionals in contact with animals susceptible of carrying the virus, etc): 1 dose by IM route or 2 doses by ID route on D0 and D7.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

(d) In penicillin-allergic patients:

- Children: **co-trimoxazole** (30 mg SMX + 6 mg TMP/kg 2 times daily) + **clindamycin** (10 mg/kg 3 times daily)
- Adults: **co-trimoxazole** (800 mg SMX + 160 mg TMP 2 times daily) or **doxycycline** (100 mg 2 times daily or 200 mg once daily, except in pregnant and lactating women) + **metronidazole** (500 mg 3 times daily).

References

1. Weekly epidemiological record/Relevé épidémiologique hebdomadaire, 20 April 2018, 93th year/20 avril 2018, 93e année. No 16, 2018, 93, 201–220.
<http://apps.who.int/iris/bitstream/handle/10665/272371/WER9316.pdf?ua=1> [Accessed 25 october 2018]
2. Spencer O, Banerjee S. Animal bites. BMJ Best practice 2018 [Accessed 25 october 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Viral hepatitis

Last updated: October 2021

Several viral infections of the liver are grouped under the heading of viral hepatitis: hepatitis A, B, C, D (delta) and E. The different hepatitis viruses are present throughout the world, but their prevalence varies by country. Hepatitis A and B are common in developing countries where the vast majority of infections occur during childhood.

The clinical characteristics of all five diseases are similar enough to make differential diagnosis difficult; however, there are epidemiological, immunological and pathological differences. Patients with hepatitis B, C and D may later develop chronic liver disease.

The main characteristics of each type of viral hepatitis are summarized in the table below.

Clinical features

- **Asymptomatic forms**

Mild or anicteric forms are the most common, irrespective of the causal virus.

- **Icteric forms**

Insidious or sudden onset with symptoms of varying intensity: fever, fatigue, nausea, gastrointestinal disturbance, followed by jaundice, dark coloured urine and more or less claycoloured stool.

- **Fulminant forms**

Hepatocellular failure with severe cytolysis that can be fatal. This form is most frequent in hepatitis B patients with secondary infection with the D virus, and in the event of pregnant women infected with hepatitis E during their third trimester.

- **Chronic hepatitis**

Hepatitis B, C and D may lead to cirrhosis and/or hepatocellular carcinoma (HCC).

The various forms of viral hepatitis

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

	Hepatitis A	Hepatitis B	Hepatitis C	Hepatitis D	Hepatitis E
Age group most at risk	Children	Children	Young adults	Young adults	Young adults
Transmission	Faecal-oral Contaminated food and water Transfusion (rare)	Vertical ^(a) Close contact with infected person (especially intra-familial). Exposure to blood (transfusion; material contaminated with blood) Sexual	Exposure to blood (transfusion; material contaminated with blood) Sexual (low) Intranasal (implements shared by intranasal drug users) Vertical ^(a)	Exposure to blood (transfusion; material contaminated with blood) Sexual Possibly vertical ^(a)	Faecal-oral Contaminated food and water
Incubation period	2 to 6 weeks	4 to 30 weeks (average 10 weeks)	2 to 25 weeks	Co-infection B/D: as for hepatitis B Secondary infection of hepatitis B: approximately 5 weeks	2 to 8 weeks
Fulminant forms	0.2 to 0.4%	1 to 3%	More rare than in hepatitis B	Much more common in patients with secondary infection of hepatitis B than in patients with B/D co-infection	20% mortality in pregnant women
Prognosis	No chronic	Chronicity: 0.2 to 10% (risk)	Chronicity: up	Chronicity: <	No chronic

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Individual prevention	Polyvalent immunoglobulin	Specific anti-HBs immunoglobulin Safe sex (condoms)	Specific anti-HBs immunoglobulin may be effective	As for hepatitis B (the D virus can only develop with B)	Cook meat (pork)
Vaccination	Anti-hepatitis A	Anti-hepatitis B	Does not exist	Anti-hepatitis B	Does not exist
Collective prevention	Hygiene, sanitation	Limit transfusion, screen blood prior to transfusion Single use of disposable material			Hygiene, sanitation

- (a) Vertical transmission: transmission of the virus from the mother to the child during pregnancy, at the time of delivery, or during the first 28 days after birth.

Laboratory

Diagnosis

- HAV, HDV and HEV infection: detection of IgM anti-HAV, anti-HDV and anti-HEV antibodies, respectively.
- HBV infection: detection of HBsAg; chronic hepatitis B: presence of HBsAG for longer 6 months; chronic active hepatitis B: detection of HBeAg and/or HBV DNA.
- HCV infection: detection of anti-HCV antibodies and HCV RNA; chronic hepatitis C: viraemia persists for longer than 6 months.

Other tests

- ALT (or AST) level, platelet count, creatinine, HCV diagnosis and HBV viral load to decide treatment of chronic active hepatitis B.
- APRI score (evaluation of liver fibrosis in chronic hepatitis): $[(\text{patient's AST level}/\text{normal AST level}) \times 100]/\text{platelet count}$ (10^9 platelets/litre). An APRI score > 1 indicates probable severe fibrosis.
- HIV test.

Other investigations

Elastography (Fibroscan®): measures the elasticity of the liver to determine stage of liver fibrosis, scored from F0 (absence of fibrosis) to F4 (cirrhosis).

Treatment

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

dual therapy with lamivudine or emtricitabine (risk of developing HIV drug resistance).

- Patients without HIV co-infection

Treatment is indicated in the event of cirrhosis or advanced hepatic fibrosis (APRI score > 1.5 or Fibroscan F3-F4 > 10 kPa); HBsAg positive with persistently elevated ALT or AST > 2 times the normal values in 2 samples taken 3 or 6 months apart; or persistently elevated ALT or AST with a high viral load (> 20 000 IU/ml).

tenofovir PO (300 mg tab, equivalent to 245 mg of tenofovir disoproxil), lifelong therapy:

Children ≥ 12 years and adults, including pregnant women: one tablet once daily taken with a meal

Treatment of chronic hepatitis C^[1]

Genotypes 1, 2, 3, 4, 5, 6 without cirrhosis or with compensated cirrhosis	sofosbuvir/velpatasvir PO (400 mg SOF/100 mg VEL tablet) 1 tablet once daily for 12 weeks
Genotypes 1, 2, 4, 5, 6 without cirrhosis or with compensated cirrhosis Genotype 3 without cirrhosis	sofosbuvir/daclatasvir PO (400 mg SOF/60 mg DCV tablet) 1 tablet once daily for 12 weeks
Genotype 3 with compensated cirrhosis	sofosbuvir/daclatasvir PO (400 mg SOF/60 mg DCV tablet) 1 tablet once daily for 24 weeks

In case of decompensated cirrhosis (presence of ascites or jaundice or mental confusion or signs of gastrointestinal haemorrhage): same treatment but for 24 weeks.

Treatment is contra-indicated during pregnancy and breastfeeding.

For women of childbearing age: provide a contraceptive; do not start treatment in women who do not want contraception.

Vaccination

- Routine vaccination of neonates and infants^[2] (according to national vaccination schedule):
 - 3 dose schedule: one dose as soon as possible after birth, preferably within the first 24 hours of life, then one dose at 6 weeks and one dose at 14 weeks^a
 - 4 dose schedule: one dose as soon as possible after birth, preferably within the first 24 hours of life, then one dose at 6 weeks, one dose at 10 weeks and one dose at 14 weeks^a
- Catch-up vaccination (unvaccinated individuals):
3 dose schedule (0-1-6): 2 doses 4 weeks apart, then a third dose 6 months after the first dose
- Post-exposure prophylaxis:
One dose on D0, one dose on D7 and one dose between D21 and D30 then a booster dose 12 months after the first dose

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

References

1. World Health Organization. Guidelines for the care and treatment of persons diagnosed with chronic hepatitis C virus infection. July 2018.
<http://apps.who.int/iris/bitstream/handle/10665/273174/9789241550345-eng.pdf?ua=1> [Accessed 21 December 2018]
2. Weekly epidemiological record/Relevé épidémiologique hebdomadaire 7 JULY 2017, 92th YEAR / 7 JUILLET 2017, 92e ANNÉE No 27, 2017, 92, 369–392
<http://apps.who.int/iris/bitstream/handle/10665/255841/WER9227.pdf?sequence=1> [Accessed 22 November 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Dengue

Last update: October 2022

Dengue fever is an arbovirus transmitted to humans by the bite of a mosquito (*Aedes*). Transmission by transfusion of contaminated blood and transplacental transmission to the foetus have also been reported.

Four different serotypes of dengue have been described. Infection with one serotype provides a lifelong immunity to that specific serotype, but only partial, short-term immunity to other serotypes. There is no specific antiviral treatment.

Dengue is a mainly urban disease, present in tropical and subtropical regions^a, in particular in Asia, Central and South America and the Caribbean. Outbreaks have been described in Eastern Africa.

Primary infection may be asymptomatic or present as mild or occasionally severe dengue fever. Subsequent infections increase the risk of severe dengue.

Clinical features

After the incubation period (4 to 10 days), the illness occurs in 3 phases:

- **Febrile phase:** high fever (39 to 40 °C) lasting 2 to 7 days, often accompanied by generalized aches, a maculopapular rash and mild haemorrhagic manifestations.
- **Critical phase** (between the third and seventh day): at the end of the febrile phase, temperature decreases. The majority of patients will have dengue without warning signs and proceed to the recovery phase. Certain patients will develop dengue with warning sign(s) at this stage. These patients are at higher risk for developing severe dengue.
- **Recovery phase:** patient improves, vital signs normalise, gastrointestinal symptoms subside and appetite returns. At times, bradycardia and generalized pruritus.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Symptoms according to severity (adapted from PAHO^[1])

Dengue without warning signs	Fever + 2 of the following symptoms: <ul style="list-style-type: none"> • Nausea, vomiting • Rash resembling measles • Generalized aches (headache, retro-orbital pain, myalgias, arthralgias) • Benign mucocutaneous bleeding (petechiae, positive tourniquet test^(a), epistaxis, gingival bleeding) • Leucopenia
Dengue with warning signs	Presence of at least one of these symptoms: <ul style="list-style-type: none"> • Intense abdominal pain • Persistent vomiting • Fluid accumulation (ascites, pleural effusion) • Mucosal bleeding • Hepatomegaly (> 2 cm) • Postural hypotension • Agitation or lethargy • Increasing haematocrit
Severe dengue	<ul style="list-style-type: none"> • Severe fluid accumulation (ascites, pleural effusion) with respiratory distress and/or shock • Severe mucocutaneous bleeding • Severe organ involvement (e.g.: transaminases > 1000 IU/litre, myocarditis, altered mental status)

(a) Tourniquet test: inflate a blood pressure cuff on the upper arm to a point midway between the systolic and diastolic pressure for 5 min. The test is positive when 20 or more petechiae per 2.5 cm square are observed.

Major differential diagnoses

Other conditions to consider include a wide range of acute febrile illnesses, e.g.:

- Chikungunya, Zika, influenza, mononucleosis, [measles](#), rubella, [viral haemorrhagic fevers](#)
- [Malaria](#)
- Meningococcemia, [typhoid fever](#), [leptospirosis](#), [rickettsioses](#), other causes of sepsis
- Leukaemia

Laboratory

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Monitoring disease course

- Haematocrit (Hct) or if available full blood count (FBC) at baseline, then daily if possible.
 - A progressive increase in Hct is a warning sign. It indicates haemoconcentration due to increased vascular permeability (plasma leakage). Hct should be monitored frequently (before and after fluid administration) in patients with warning signs up to the end of the fluid treatment.
 - Leukopenia and thrombocytopenia are common and improve as the recovery phase begins. Leukocytosis may occur with severe bleeding.
- Liver function tests if possible at baseline, then according to results.

Treatment

Patients in Group A (outpatients)

Patients with no warning signs, able to drink sufficiently and with a normal urine output.

- Bed rest and good hydration.
- Fever and pain: **paracetamol** PO at the usual doses (see [Fever](#), Chapter 1), maintaining a 6 to 8 hour interval between doses. Do not prescribe acetylsalicylic acid, ibuprofen or other NSAIDs.
- Seek medical attention if: no clinical improvement, persistent vomiting, cold extremities, agitation or lethargy, breathing difficulties or absence of urine output.
- If follow-up is impossible or symptoms cannot be monitored at home (patients living far from the health care facility/living alone), hospitalise for observation.

Patients in Group B (inpatients)

Patients with any of the following:

- Warning sign(s)
- Acute (e.g. severe dehydration or malaria) or chronic (e.g. diabetes, cardiovascular, renal or haemolytic disease, obesity) co-morbidities
- Risk factors for bleeding (e.g. anticoagulation, coagulopathy, peptic ulcer or gastritis, treatment with NSAIDs)
- Pregnant women, patients under 1 year or 65 years and over or patients with difficulty drinking

In all cases:

- Place the patient under a mosquito net; encourage oral fluid intake (including oral rehydration solution (ORS) if needed).
- Avoid invasive procedures (nasogastric tube, IM injections) to minimize the risk of bleeding.
- Fever and pain: **paracetamol** PO^[2] with caution and without exceeding:
 - children: 10 mg/kg every 6 to 8 hours
 - adults: 500 mg every 6 to 8 hours
- In case of elevated transaminases ≥ 10 times the upper limit of normal, do not administer paracetamol. Use tepid sponging for reducing fever.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Monitor clinical status (warning signs, general symptoms, vital signs, capillary refill time), IV and oral fluid intake, urine output, hourly for at least 4 hours, then every 4 hours while the patient is on IV fluid treatment.
- Place an intravenous line and administer a bolus of Ringer lactate:
 - children and adults: 10 ml/kg over one hour
 - patients 65 years and over or with co-morbidities : 5 ml/kg over one hour
- Re-assess the patient:
 - If no improvement after first bolus: administer a second bolus as above. If necessary, a total of 3 boluses can be administered. If still no improvement after 3 bolus, consider as severe dengue (patients in Group C) and transfer to intensive care unit.
 - If improvement after the first, second, or third bolus, reduce Ringer lactate:
 - ▷ children and adults: 5 to 7 ml/kg/hour over 2 to 4 hours
 - ▷ patients 65 years and over or with co-morbidities: 5 ml/kg/hour over 2 to 4 hours
 - If continuing improvement, reduce Ringer lactate (then stop as soon as possible to reduce the risk of fluid overload):
 - ▷ children and adults: 3 to 5 ml/kg/hour over 2 to 4 hours, then 2 to 4 ml/kg/hour over 24 to 48 hours
 - ▷ patients 65 years and over or with co-morbidities: 3 ml/kg/hour over 2 to 4 hours, then 2 ml/kg/hour over 24 to 48 hours
 - If the patient deteriorates after initial improvement, resume the bolus therapy with Ringer lactate (up to 3 bolus) as above.

Patients in Group C (intensive care unit)

Patients with severe dengue requiring emergency treatment for managing shock and other complications (e.g. severe bleeding, acidosis, coagulopathy).

Prevention

- Individual protection: long sleeves and trousers, repellents, mosquito net (*Aedes* bites during the day).
- Elimination of mosquito breeding sites (small collections of water in discarded tires, flower pots, and other containers).

Footnotes

- (a) For more information:
http://gamapserver.who.int/mapLibrary/Files/Maps/Global_DengueTransmission_ITHRiskMap.png?ua=1
- (b) Adequate urine output: at least 1 ml/kg/hour in children and 0.5 ml/kg/hour in adults. If unavailable, ensure that the patient is urinating at least every 4 hours.
- (c) Remove 50 ml of Ringer lactate (RL) from a 500 ml RL bottle or bag, then add 50 ml of 50% glucose to the remaining 450 ml of RL to obtain 500 ml of 5% glucose-RL solution

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Viral haemorrhagic fevers

Several diseases with different aetiologies and different modes of transmission are grouped under this term as they present with common clinical signs.

Dengue haemorrhagic fever is a viral haemorrhagic fever that is described in a specific chapter (see [Dengue](#), Chapter 8).

Clinical features

- Common syndrome (CS):
 - Fever higher than 38.5 °C;
 - Haemorrhagic symptoms (purpura, epistaxis, haematemesis, melaena, etc.).
- The clinical signs are often nonspecific; the severity varies depending on the aetiology.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	Reservoir/ Vector <i>Geographical distribution</i>	Isolation of patients	Clinical features	Estimated case fatality rate
Ebola^(a) Marburg	Bats (?) <i>Africa</i>	Strict isolation	CS + sudden onset general malaise, vomiting and diarrhoea	60-80%
Lassa^(a)	Rodents <i>West Africa^(b)</i>	Strict isolation	CS + general malaise, headache, muscle pain, facial oedema, pharyngitis, proteinuria on reagent strip	15-20%
Junin and Machupo^(a)	Rodents <i>South America</i>	Isolation	CS + vomiting, erythema of the face and, depending on the aetiology: <ul style="list-style-type: none"> • periorbital oedema, cervical adenopathy, pharyngitis • pharyngitis, reddened conjunctivae • oedema of the soft palate, generalised petechial rash • proteinuria on reagent strip 	15-30%
Omsk	Ticks <i>Europe, Asia</i>	None		2-5%
Crimean Congo^(a)	Livestock/Ticks <i>Africa, Asia</i>	Strict isolation		5-20%
FHSR (hantavirus)^(a)	Rodents <i>Asia and Europe</i>	None		< 1%
Kyasanur	Small mammals/Ticks <i>India</i>	None	CS + headache, muscle pain, prostration	2-10%
Rift Valley^(a)	Livestock/Mosquitoes <i>Africa</i>	Mosquito nets	Clinical signs: <ul style="list-style-type: none"> • isolated fever • SC • encephalitis • retinitis and blindness 	30-50%
Yellow fever^(a)	Primates/Mosquitoes <i>Africa, South America</i>	Mosquito nets	CS + jaundice, proteinuria on reagent strip, oliguria, headache	10-30%

(a) Viral haemorrhagic fever with epidemic potential.

(b) For more information on geographic distribution of Lassa fever:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Management

Suspicion of haemorrhagic fever

Isolated case of fever with haemorrhagic symptoms in an endemic area

- Isolation: isolation room (or if not available, use screens/partitions); restrict visitors (if a carer is strictly necessary, s/he must be protected with gown, gloves, mask).
- Standard precautions:
The majority of hospital-acquired infections have occurred due to a lack of respect for these precautions:
 - Hand washing;
 - Gloves for patient examination and when touching blood, body fluids, secretions, excretions, mucous membranes, non-intact skin;
 - Gowns to protect skin and prevent soiling of clothing during consultations and activities that are likely to generate splashes or sprays of blood, body fluids, secretions, or excretions;
 - Surgical mask and goggles, or face shield, to protect mucous membranes of the eyes, nose, and mouth during activities that may generate splashes of blood, body fluids, secretions, and excretions;
 - Adequate procedures for the routine cleaning and disinfection of objects and surfaces;
 - Rubber gloves to handle soiled laundry;
 - Safe waste management;
 - Safe injection practices.

Confirmed cases of Ebola, Marburg, Lassa, Crimean-Congo fevers or epidemics of unknown origin

- Strict isolation in a reserved area separate from other patient areas, with a defined circuit for entrance/exit and changing room at the entrance/exit; dedicated staff and equipment/supplies; use of disposable material if possible.
- Standard precautions (as above)

PLUS

- Droplet precautions AND contact precautions including personal protective equipment (PPE).
The PPE is to be worn systematically prior to entry into isolation area, regardless the tasks to be performed (care, cleaning, distribution of meals, etc.) and to be removed before leaving the isolation area:
 - two pairs of gloves,
 - double gown or coverall suit,
 - surgical cap or hood, mask, protective glasses,
 - impermeable apron,
 - rubber boots.
- Disinfection of surfaces, objects, clothing and bedding with chlorine solution; safe handling and on site disposal of waste and excreta, etc.
- In the event of a death, do not wash the body. Prompt and safe burial of the dead as quickly as possible, using a body bag.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Aetiological treatment: ribavirine for Lassa fever and Crimean-Congo fever.
- Symptomatic treatment:
 - Fever: paracetamol (Chapter 1). Acetylsalicylic acid (aspirin) is contra-indicated.
 - Pain: mild (paracetamol), moderate (tramadol), severe (sublingual morphine): see Pain, Chapter 1.
 - Dehydration: oral rehydration salts and/or IV rehydration with Ringer lactate, see Dehydration, Chapter 1.
 - Seizures (Chapter 1).
 - Vomiting: **ondansetron** PO^[1]
 Children 6 months to < 2 years: 2 mg once daily
 Children 2 to < 4 years: 2 mg 2 times daily
 Children 4 to < 12 years: 4 mg 2 times daily
 Children ≥ 12 years and adults: 4 to 8 mg 2 times daily
- For Ebola and Marburg haemorrhagic fevers: invasive procedures must be strictly limited. Health care staff is at risk of contamination when inserting and maintaining IV lines. An IV line must be well secured so that the patient, often confused, cannot pull it out.

Prevention

- Vaccination against yellow fever^[2]:

Children and adults: 0.5 ml single dose

 - Routine vaccination : children from 9 months of age, along with the measles vaccine.
 - Mass vaccination campaign during an epidemic: children from 6 months and adults ; for pregnant women, only administer during an epidemic.
- Vaccination against Rift Valley fever: only during an epidemic.
- Vector control programmes for known vectors.
- Infection control measures are essential in all cases.

References

1. World Health Organization. Clinical management of patients with viral haemorrhagic fever. A pocket guide for front-line health workers. Interim emergency guidance for country adaptation, February 2016.
http://apps.who.int/iris/bitstream/handle/10665/205570/9789241549608_eng.pdf;jsessionid=15E17DE39631519C2051413DDCBBC8A7?sequence=1 [Accessed 11 January 2019]
2. Weekly epidemiological record-Relevé épidémiologique hebdomadaire 5 july 2013, 88th year / 5 juillet 2013, 88e année No. 27, 2013, 88, 269–284.
<https://www.who.int/wer/2013/wer8827.pdf?ua=1> [Accessed 10 december 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

HIV infection and AIDS

Acquired immune deficiency syndrome (AIDS) is the most advanced stage of infection with human immunodeficiency virus (HIV).

Two subtypes of HIV have been identified. HIV-1 is more widespread than HIV-2, the latter mainly being found in West Africa. HIV-2 is less virulent and less transmissible than HIV-1.

HIV weakens the immune system by causing a deficit in CD4 T lymphocytes.

Evolution of the disease

- **Primary infection or acute retroviral syndrome:** 50 to 70% of newly infected individuals develop during seroconversion (from 15 days to 3 months post exposure), a viral syndrome with fever, malaise, and lymphadenopathy.
- **Asymptomatic HIV infection** (after seroconversion): a period of clinical latency, but not viral latency. The time period for progression from HIV infection to the development of severe immune deficiency in western countries is approximately 10 years. This period appears to be shorter in developing countries.
- **Symptomatic HIV infection:** with progressive destruction of the immune system, common and more severe diseases occur more frequently, and with higher mortality, in seropositive individuals.
- **AIDS:** this stage corresponds to the development of severe opportunistic infections and neoplasms. From a biological point of view, AIDS is defined as a CD4 count < 200 cells/mm³. Without treatment the disease progresses rapidly towards death.

The World Health Organization (WHO) has proposed a clinical classification of HIV infection in 4 stages of severity for adults and adolescents and for children.^[1]

Laboratory

Diagnosis of HIV infection

- The diagnosis is made with serological (detection of antibodies against the virus) or virological (especially in infants) testing.
- Testing should always be done **voluntarily with informed consent**.
- All HIV test results must be strictly **confidential** in order to avoid discrimination.
- The individual should have access to services offering pre-test and post-test counselling, treatment and support.
- A diagnosis of HIV infection can be made only after at least 2 different test results (2 different brands) are clearly positive: the positive result of an initial (highly sensitive) test must be confirmed through use of a second (highly specific) test. In areas where HIV prevalence is low, diagnosis is confirmed after 3 positive test results.

CD4 lymphocyte counts

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment of HIV infection

Antiretroviral (ARV) treatment^a

A multi-drug (at least 3) antiretroviral therapy (ART) is the reference treatment. It does not eradicate the virus, but slows the progression of the disease and improves the patient's clinical state by reducing viral replication and consequently increasing the CD4 cell count to levels beyond the threshold of opportunistic infections.

Therapeutic classes

Four major classes ARV are used:

- NRTI (nucleoside/nucleotide reverse transcriptase inhibitors): zidovudine (AZT), lamivudine (3TC), abacavir (ABC), tenofovir (TDF), emtricitabine (FTC).
- NNRTI (non-nucleoside reverse transcriptase inhibitors): efavirenz (EFV), nevirapine (NVP), etravirine (ETR). HIV-2 is naturally resistant to NNRTIs.
- PI (protease inhibitors): atazanavir (ATV), lopinavir (LPV), ritonavir (RTV), darunavir (DRV).
- INI (integrase inhibitors): dolutegravir, raltegravir.

Principles of ARV treatment

- Daily triple therapy must be taken for life to prevent the rapid development of resistance. It is important that the patient understands this and that adherence to treatment is optimal.
- Follow the ART protocols recommended by national HIV program.
- The most widely used and easiest regimens to administer are 2 NRTI + 1 NNRTI: e.g. TDF/3TC/EFV.
- In the event of treatment failure, all 3 drugs should be replaced with a second-line regimen: 2 other NRTIs + 1 PI.

Other possible combinations exist which are less commonly used or more difficult to manage.

Criteria for ARV treatment

As a priority ART should be initiated in all patients with WHO clinical stage 3 or 4 and patients with CD4 < 350 /mm³. However, those with higher CD4 counts can initiate ART.

Monitoring of ARV treatment

HIV viral load is an essential tool for monitoring the effectiveness of ARV. CD4 count is useful for identifying severely immunosuppressed. Other tests such as blood count, tests for liver (ALAT) and renal function (creatinine clearance) are not essential, but can be useful in detecting adverse effects.

Treatment of opportunistic and other infections

With progressive immunosuppression, HIV-infected patients who are not receiving triple therapy (or patients on ART but with poor adherence) become increasingly susceptible to infections. For conditions of clinical stages 2 and 3, standard treatments are usually effective. Patients may benefit from primary prophylaxis against opportunistic

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

The most reliable method of prevention is the use of male or female condoms.

Male circumcision decreases significantly the risk of HIV transmission.

Early diagnosis and treatment of sexually transmitted infections is essential as they increase the transmission of HIV (see [Chapter 9](#)).

ART to HIV positive and adherent partner does protect the negative partner from HIV infection.

Occupational transmission

(accidental needle stick injuries or injuries with contaminated objects, contact between a patient's blood and unprotected broken skin or mucous membranes)

Prevention is based on use of standard precautions to avoid contamination with soiled material or potentially infected body fluids.

Post-exposure prophylaxis (PEP): e.g. in the event of rape or occupational accidental exposure to blood, ARV treatment initiated as soon as possible within 72 hours of exposure for a duration of 1 month may reduce the risk of infection.

Nosocomial transmission

Prevention of nosocomial HIV infection is based on the rational use of injections and strict respect for hygiene and sterilization and disinfection procedures for medical material.

For transfusion: strict respect of indications for transfusion and systematic serological screening of the donor's blood are the two indispensable precautions in the prevention of HIV transmission through transfusions.

Transmission in injection drug users

Needle and syringe exchange programs with disposable needles and syringes for users can reduce the risk.

Mother-to-child transmission (MTCT)

The global rate of vertical transmission varies from 20 to 40%. The risk of transmission through breast-feeding is evaluated at approximately 12% and persists for the duration of breast-feeding.

- **In pregnant women:** HIV transmission from mother-to-child may be reduced by ART. The protocol called Option B+ is the internationally preferred protocol. All HIV-infected pregnant women receive lifelong triple-drug therapy, regardless of the CD4 count or clinical stage, both for their own health and to prevent transmission to the child. The most commonly recommended ART is TDF/3TC/EFV or TDF/FTC/EFV. Check national recommendations. In addition, ARVs are administered to the newborn.
Programs targeting pregnant women also include other preventive measures such as avoiding artificial rupture of the membranes and systematic episiotomy.
- **In breast-feeding women:** exclusive breast-feeding for the first 6 months of life, introduction of complementary (solid) foods at 6 months, gradual cessation of breast-feeding to the age of 12 months.

Prevention of opportunistic infections

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Infections	Primary prophylaxis
Pneumocystosis	co-trimoxazole PO
Cerebral toxoplasmosis	Children: 50 mg SMX + 10 mg TMP/kg once daily
Isosporiasis	Adults: 800 mg SMX + 160 mg TMP once daily
Various bacterial infections	
Malaria	

Secondary prophylaxis

For patients who develop a specific opportunistic infection, in order to prevent recurrence once treatment for the infection is completed.

Infections	Secondary prophylaxis	Comments
Pneumocystosis		Alternative dapsone PO Children: 2 mg/kg once daily (max. 100 mg daily) Adults: 100 mg once daily
Toxoplasmosis	co-trimoxazole PO Children: 50 mg SMX + 10 mg TMP/kg once daily Adults: 800 mg SMX + 160 mg TMP once daily	Alternative Adults: dapsone PO: 200 mg once weekly or 50 mg once daily + pyrimethamine PO: 75 mg once weekly + folinic acid PO: 25 to 30 mg once weekly
Isosporiasis		-
Penicilliosis Histoplasmosis	itraconazole PO Adults: 200 mg once daily	-
Cryptococcal meningitis	fluconazole PO Children: 6 mg/kg once daily Adults: 200 mg once daily	-

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Symptoms	Definitions and aetiologies	Diagnosis	Treatment
Diarrhoea with or without blood (also see <u>Chapter 3</u>)	<p>Diarrhoea is defined as at least 3 liquid stools per day.</p> <p>Aetiologies:</p> <p>Parasitic infections</p> <ul style="list-style-type: none"> • <i>Isospora belli</i> • <i>Cryptosporidium</i> • <i>Microsporidium</i> • <i>Giardia lamblia</i> • <i>Entamoeba histolytica</i> <p>Bacterial infections</p> <ul style="list-style-type: none"> • <i>Shigella</i> • <i>Salmonella enteritis</i> • <i>Campylobacter enteritis</i> <p>Mycobacterial infections</p> <ul style="list-style-type: none"> • <i>Mycobacterium tuberculosis</i> (gastrointestinal TB) • <i>Mycobacterium avium</i> complex <p>Helminthiasis</p> <ul style="list-style-type: none"> • <i>Strongyloides stercoralis</i> <p>Viral infections</p> <ul style="list-style-type: none"> • Cytomegalovirus (CMV) <p>Other causes</p> <ul style="list-style-type: none"> • Kaposi sarcoma • Lymphoma • Idiopathic (HIV infection) • Antiretrovirals (especially lopinavir and ritonavir) 	<p>1. History and clinical examination</p> <p>2. Microscopic examination of stool for ova and parasites (2 to 3 samples)</p> <p><i>Note:</i> <i>I. belli, Cryptosporidium, Microsporidium, MAC and CMV are unlikely if CD4 count > 200 cells.</i></p>	<ul style="list-style-type: none"> • Persistent (> 2 weeks) or chronic (> 4 weeks) diarrhoea is often associated with weight loss and dehydration. • Prevention or treatment of dehydration is critical (<u>Dehydration</u>, Chapter 1). • Depending on the results of the stool examinations: give appropriate treatment. • If there is no laboratory support: Acute bloody diarrhoea • First-line treatment: Children: azithromycin PO: 20 mg/kg once daily for 5 days or ciprofloxacin PO: 15 mg/kg 2 times daily for 7 days Adults: ciprofloxacin PO: 500 mg 2 times daily for 7 days • If amoebiasis suspected: tinidazole or metronidazole PO (<u>Amoebiasis</u>, Chapter 3).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Non-bloody persistent or chronic diarrhea

Persistent or chronic diarrhoea suggests advanced immunocompromised state. For patients who qualify for ARVs by CD4 count (or unknown CD4 count), ARV initiation is urgent and will usually resolve symptoms in 14 to 28 days.

- *Isospora belli*: **co-trimoxazole** PO

Children: 40 mg SMX + 8 mg TMP/kg 2 times daily for 10 days then 25 mg SMX + 5 mg TMP/kg 2 times daily for 3 weeks

Adults: 800 mg SMX + 160 mg TMP 2 times daily for 7 to 10 days then 400 mg SMX + 80 mg TMP 2 times daily for 3 weeks

- *Cryptosporidium*: no specific treatment in HIV-infected patients

- *Microsporidium*: **albendazole** PO (limited efficacy)

Children: 10 mg/kg 2 times daily (max. 800 mg daily) for 7 days

Adults: 400 mg 2 times daily for 2 to 4 weeks

- *Helminthiasis*: **albendazole** PO for 3 days

Children > 6 months but ≤ 10 kg: 200 mg once daily

Children > 6 months and adults: 400 mg once daily

- *Giardiasis*: **tinidazole** or **metronidazole** (Intestinal protozoan infections, Chapter 6).

- If no improvement (and no contraindications such as bloody diarrhoea) symptomatic treatment

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

			<p>Nutrition ++++</p> <p>Children: continue to breastfeed; increase daily calorie intake:</p> <p>6-11 months: add 150 kcal daily</p> <p>12-23 months: add 200 kcal daily</p> <p>2-5 years: add 250 kcal daily</p> <p>6-9 years: add 350 kcal daily</p> <p>10-14 years: add 400 kcal daily</p> <p>Eliminate fresh milk, give porridge prepared with rice water or soup or yoghurts. Give 2.5 ml of oil per meal.</p> <p>Any child 0-5 years should receive zinc sulfate (<u>Acute diarrhoea</u>, Chapter 3).</p> <p>Adults: increase the calorie and protein intake (at least 2 g protein/kg daily). No food is excluded but avoid raw food, fresh milk and foods high in fibre. Encourage small, frequent meals.</p>
Oral and oesophageal lesions	<p>Fungal infections</p> <ul style="list-style-type: none"> • Oral candidiasis: see <u>Stomatitis</u>, Chapter 3. • Oesophageal candidiasis: pain on swallowing, dysphagia. May result in weight loss. <p>Viral infections</p> <ul style="list-style-type: none"> • Oral hairy leukoplakia (keratosis on the lateral sides of the tongue due to the Epstein-Barr virus) • Oral and oesophageal herpes <p>Aphthous ulcers</p>	<p>Clinical examination is enough to make a diagnosis.</p> <p>Consider all severe oral candidiasis (if the pharynx is involved) as oesophageal candidiasis even in the absence of dysphagia.</p>	<ul style="list-style-type: none"> • Mild oral candidiasis nystatin PO Children and adults: 100 000 IU (= 1 ml) 4 times daily or miconazole oral gel Children 6 months-2 years: 1.25 ml 4 times daily Children over 2 years and adults: 2.5 ml 4 times daily The treatment lasts 7 to 14 days. • Moderate to severe oral candidiasis and oesophageal candidiasis fluconazole PO Children: 3 to 6 mg/kg once daily Adults: 50 to 200 mg once daily up to 400 mg daily if necessary The treatment lasts 7 to 14 days for oral candidiasis and 14 to 21 days for oesophageal candidiasis.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

			<ul style="list-style-type: none"> • Oral herpes: Analgesics (paracetamol, ibuprofen). For recurrent or extensive forms affecting the oesophagus, add: aciclovir PO for 7 days Children under 2 years: 200 mg 5 times daily Children 2 years and over and adults: 400 mg 5 times daily <i>Secondary prophylaxis only for patients with frequent recurrences.</i>
Respiratory problems (also see <u>Chapter 2</u>)	<p>Cough and/or thoracic pain and/or dyspnoea in a symptomatic HIV infected patient.</p> <p>Aetiologies:</p> <p>Bacterial infections</p> <ul style="list-style-type: none"> • <i>Streptococcus pneumoniae</i> • <i>Haemophilus influenzae</i> • <i>Staphylococcus aureus</i> <p>Mycobacterial infections</p> <ul style="list-style-type: none"> • <i>M. tuberculosis</i>, MAC <p>Protozoal infections</p> <ul style="list-style-type: none"> • <i>Pneumocystis jiroveci</i> (PCP) <p>Fungal infections</p> <ul style="list-style-type: none"> • <i>Cryptococcus neoformans</i> • <i>Histoplasma capsulatum</i> • <i>Coccidioides immitis</i> • <i>Aspergillus</i> spp • <i>Penicillium marneffei</i> <p>Viral infections</p> <ul style="list-style-type: none"> • CMV <p>Neoplasms</p>	<p>1. History and clinical examination: Blood in the sputum? If fever < 7 days, dyspnoea: unlikely TB. If cough > 21 days, weight loss, thoracic pain > 15 days, no dyspnoea: likely TB. Pulmonary auscultation: bilateral lobar pneumonia?</p> <p>2. If possible:</p> <ol style="list-style-type: none"> a) Look for AFB in sputum b) Chest x-ray <ul style="list-style-type: none"> • PCP: bilateral interstitial infiltrates • TB: miliary shadowing, large heart, pleural effusion, enlarged lymph nodes inside the chest. <p><i>Notes</i></p> <ul style="list-style-type: none"> • MAC, PCP, CMV and fungal infections are unlikely in patients with a CD4 count > 200 cells/mm³ 	<ul style="list-style-type: none"> • For the diagnosis and treatment of upper respiratory tract infections, particularly pneumonia: see <u>Chapter 2</u>. • If the chest x-ray is consistent with staphylococcal pneumonia: Children: see <u>Staphylococcal pneumonia</u>, Chapter 2. • Adults: ceftriaxone IM or slow IV 1 g once daily + cloxacillin IV 2 g every 6 hours • If the sputum examination is AFB+, treat for TB.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<p>Others</p> <ul style="list-style-type: none"> • Lymphoid interstitial pneumonia • Pleural effusion (often TB) • Pericardial effusion (often TB) • Pneumothorax (may be due to PCP) 	<ul style="list-style-type: none"> • If the sputum examination is negative and the chest x-ray is consistent with PCP: <p>co-trimoxazole PO for 21 days Children: 50 mg SMX + 10 mg TMP/kg 2 times daily Adults: 1600 SMX + 320 TMP 3 times daily</p> <p>Note: the symptoms may become worse during the first phase of treatment, effectiveness can only be evaluated after one week of treatment.</p> <p>Add prednisolone PO for patients with severe PCP with hypoxia: Children: start with 2 mg/kg daily then decrease the dose following the adult example Adults: 40 mg 2 times daily for 5 days, then 40 mg once daily for 5 days then 20 mg once daily for 10 days</p> <p><i>Secondary prophylaxis is recommended.</i></p> <ul style="list-style-type: none"> • Fungal infections (cryptococcosis, penicilliosis, histoplasmosis): Adults: amphotericin B IV: 0.7 to 1 mg/kg once daily for 2 weeks (cryptococcosis, penicilliosis) or 1 to 2 weeks (histoplasmosis), then: fluconazole PO: 400 mg daily for 8 weeks (cryptococcosis) itraconazole PO: 200 mg 2 times daily for 10 weeks (penicilliosis) itraconazole PO: 200 mg 3 times daily for 3 days then 200 to 400 mg daily for 12 weeks (histoplasmosis) <p><i>Secondary prophylaxis is recommended.</i></p>
--	--

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	<p>Persistent generalised lymphadenopathy (PGL):</p> <ul style="list-style-type: none"> • 2 or more extra-inguinal sites • lymph nodes > 1.5 cm • enlarged for 3 or more months PGL is usually due to HIV infection. <p>Aetiologies:</p> <p>HIV infection</p> <p>Infections</p> <ul style="list-style-type: none"> • TB • Syphilis • Histoplasmosis • Toxoplasmosis • CMV <p>Neoplasms</p> <ul style="list-style-type: none"> • Kaposi sarcoma • Lymphoma 	<p>2. Suspected TB: lymph node aspiration, look for AFB, chest x-ray</p> <p><i>Note:</i> in HIV infected patients, TB is often extrapulmonary.</p> <p>3. Suspected syphilis: serology</p> <p>4. If all examinations are negative: biopsy is useful to exclude lymphoma, Kaposi's sarcoma and fungal or mycobacterial infections (see notes for patients in stage 1).</p>	<ul style="list-style-type: none"> • Early syphilis: <p>benzathine benzylpenicillin IM</p> <p>Adults: 2.4 MIU single dose (1.2 MIU in each buttock)</p> <p>or, if not available:</p> <p>azithromycin PO</p> <p>Adults: 2 g single dose</p> <p><i>Note:</i> in patients in stage 1, no further investigation (other than 1, 2 and 3 in this table) or treatment are required.</p>
<p>Skin lesions</p> <p>(also see Chapter 4)</p>	<p>Bacterial infections</p> <ul style="list-style-type: none"> • Furunculosis • Impetigo and pyoderma • Axillary hidradenitis • Pyomyositis • Syphilis <p>Viral infections</p> <ul style="list-style-type: none"> • Herpes zoster • Herpes simplex • Genital warts • <i>Molluscum contagiosum</i> <p>Fungal infections</p> <ul style="list-style-type: none"> • Candidiasis, dermatophytoses and deep mycoses (penicilliosis, cryptococcosis, histoplasmosis, etc.) <p>Neoplasms</p>		<p>Bacterial infections</p> <ul style="list-style-type: none"> • Furunculosis, impetigo, chronic folliculitis: see Bacterial skin infections, Chapter 4. • Suppurative axillary hidradenitis: local treatment + doxycycline PO: 200 mg once daily for 6 weeks (in adults) • Pyomyositis: antibiotics and surgical drainage, see Pyomyositis, Chapter 10. • Primary and secondary syphilis: see Genital ulcers, Chapter 9.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Other skin infections

- Chronic prurigo or urticaria
- Severe seborrhoeic dermatitis
- Psoriasis
- Scabies
- Diffuse cutaneous xerosis

Rash caused by medication**Bed sores****Viral infections**

- Herpes zoster: see Herpes simplex and herpes zoster, Chapter 4.

For necrotic, extensive forms, eruption on the face, ophthalmic zoster, add **aciclovir** within 48 hours of the onset of lesions:

Children (IV route): 5 to 10 mg/kg every 8 hours for 7 days

Adults (oral route): 800 mg 5 times daily for 7 days

- Herpes simplex: see

Herpes simplex and herpes zoster, Chapter 4.

- Genital warts: see Venereal warts, Chapter 9.

Fungal infections

- Candidiasis: **2% miconazole** cream, one application 2 times daily
- Dermatophytoses: see Superficial fungal infections, Chapter 4.

Treatment of Kaposi sarcoma (KS)

- Start promptly ART.
- KS tumours with oedema or ulceration or presence of extensive oral or gastrointestinal or pulmonary KS +/- systemic illness: chemotherapy

Other skin infections

- Prurigo, urticaria: see Other skin disorders, Chapter 4.
- Seborrhoeic dermatitis: **Whitfield's** ointment or **2% miconazole**, one application 2 times daily. For severe inflammation, use a topical corticosteroid in combination with miconazole.

• Xerosis: **zinc oxide** ointment or

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

 Strictly necessary Functional Performance Cookies

disorders in adults	<p>Infections</p> <ul style="list-style-type: none"> • TB meningitis • Cryptococcal meningitis • Cerebral toxoplasmosis • Neurosyphilis • CMV encephalitis • HIV encephalopathy • Progressive multifocal leuko-encephalopathy • Cerebral malaria <p>Neoplasms</p> <ul style="list-style-type: none"> • Primary CNS lymphoma <p>Common causes of headache unrelated to HIV infection: sometimes more frequent in HIV infected patients (sinusitis, problems with accommodation etc.)</p> <p>Adverse effects of ARVs</p>	<p>examination:</p> <ul style="list-style-type: none"> • Change in mental state • Focal deficits • Seizures • Signs of meningeal irritation • Raised intracranial pressure • Motor problems, ataxia <p>In settings where cryptococcal infection is common, screen all adults with CD4 < 100 prior to initiation of ART, using a rapid CrAg test on serum or plasma.</p> <p>In endemic areas: check for malaria (if febrile). Lumbar puncture (LP) if not contra-indicated.</p> <p>Elements in favour of neurosyphilis:</p> <ul style="list-style-type: none"> • VDRL positive in blood and/or CSF • cells in the CSF • high protein in the CSF 	<p>Chapter 6.</p> <p>If focal signs, treat for toxoplasmosis:</p> <p>co-trimoxazole PO: 25 mg SMX + 5 mg TMP/kg 2 times daily for 4 to 6 weeks</p> <p>or</p> <p>pyrimethamine PO: 100 mg morning and evening on D1, then 75 to 100 mg daily + sulfadiazine PO: 2 g 2 to 3 times daily + folic acid PO: 15 mg once daily, for 6 weeks</p> <p><i>A secondary prophylaxis is recommended.</i></p>
----------------------------	---	--	--

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

If the LP is positive:

- Bacterial meningitis: see Chapter 7.
- TB meningitis: see the guide Tuberculosis, MSF.

• Cryptococcal meningitis^[2] :
amphotericin B IV: 1 mg/kg once daily + **flucytosine** PO: 25 mg/kg 4 times daily for 1 week
then **fluconazole** PO: 1200 mg once daily for 1 week then 800 mg once daily for 8 weeks
or, if not available
amphotericin B IV: 1 mg/kg once daily + **fluconazole** PO: 1200 mg once daily for 2 weeks
then **fluconazole** PO alone: 800 mg once daily for 8 weeks
or
fluconazole PO: 1200 mg once daily + **flucytosine** PO: 25 mg/kg 4 times daily for 2 weeks
then **fluconazole** PO alone: 800 mg once daily for 8 weeks
During the induction phase: give fluconazole IV (same doses) if the patient cannot take oral treatment; liposomal amphotericin B (3 mg/kg daily 2 weeks) may be used instead of conventional amphotericin B in case of renal impairment.
A secondary prophylaxis is recommended.
Note: intracranial pressure (ICP) is often raised in cryptococcal meningitis. To lower ICP, repeated 'therapeutic' punctures to drain CSF may be necessary at the beginning of treatment.
Neurosypilis:
benzylpenicillin IV: 2 to 4 MII 1/1 2

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

disorders in children	<ul style="list-style-type: none"> • Bacterial meningitis • TB meningitis • Cryptococcal meningitis • Cerebral toxoplasmosis • CMV meningo-encephalitis • Cerebral malaria 	<p>only patients with acute episodes benefit from specific aetiological treatment (seizures, meningeal syndrome, focal signs).</p> <p>In endemic areas, check for malaria (if febrile). Lumbar puncture (LP) if not contra-indicated.</p>	<p>Chapter 6.</p> <p>If LP is not possible:</p> <ul style="list-style-type: none"> • Treat for bacterial meningitis if patient febrile and/or meningeal syndrome (see Chapter 7). • If focal signs, treat for toxoplasmosis: <p>co-trimoxazole PO: 25 mg SMX + 5 mg TMP/kg 2 times daily for 4 to 6 weeks</p> <p>or</p> <p>pyrimethamine PO: 1 mg/kg 2 times daily for 2 days then 1 mg/kg once daily + sulfadiazine PO: 40 mg/kg 2 times daily + folinic acid PO: 10 mg once daily, for 8 weeks</p> <p><i>A secondary prophylaxis is recommended.</i></p>
------------------------------	--	---	---

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

			<p>If the LP is positive:</p> <ul style="list-style-type: none"> • Bacterial meningitis: see Chapter 7. • TB meningitis: see the guide Tuberculosis, MSF. • Cryptococcal meningitis (in order of preference)^[2]: <p>amphotericin B IV: 1 mg/kg once daily + flucytosine PO: 25 mg/kg 4 times daily for 1 week then fluconazole PO: 12 mg/kg once daily (max. 800 mg daily) for 1 week then 6-12 mg/kg once daily (max. 800 mg daily) for 8 weeks or, if not available amphotericin B IV: 1 mg/kg once daily + fluconazole PO: 12 mg/kg once daily (max. 800 mg daily) for 2 weeks then fluconazole PO alone: 6-12 mg/kg once daily for 8 weeks (max. 800 mg daily) or fluconazole PO: 12 mg/kg once daily (max. 800 mg daily) + flucytosine PO: 25 mg/kg 4 times daily for 2 weeks then fluconazole PO alone: 6-12 mg/kg once daily (max. 800 mg daily) for 8 weeks During the induction phase: give fluconazole IV (same doses) if the child cannot take oral treatment; liposomal amphotericin B (3 mg/kg daily, 2 weeks) may be used instead of conventional amphotericin B in case of renal impairment.</p> <p><i>A secondary prophylaxis is recommended.</i></p>
Persistent or recurrent fever	Temperature > 38 °C, chronic (lasting more than 5 days) or recurrent	1. History and clinical examination: look for a ENT or urinary infection	<p>Positive malaria test: see Malaria, Chapter 6.</p> <p>If testing is not available in endemic</p>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<p>Infections</p> <ul style="list-style-type: none"> • Common childhood diseases • Severe bacterial infections (TB, pneumonia, typhoid fever, septicaemia, meningitis, endocarditis, etc.) • Occult bacterial infections (sinusitis, otitis, urinary tract infections) • Opportunistic infections (TB, mycosis, toxoplasmosis) • Malaria <p>Neoplasms</p> <ul style="list-style-type: none"> • Non-Hodgkin's lymphoma <p>HIV infection</p> <p>Fever caused by medication</p>	<p>4. Chest x-ray, CBC, blood cultures, urinalysis, stool culture, serology, lumbar puncture (LP). If the child is under treatment, consider adverse effects of medication.</p>	<p>Identified or suspected focus of infection:</p> <ul style="list-style-type: none"> • ENT: see Chapter 2; urinary: see Chapter 9, etc. • TB: see the guide Tuberculosis, MSF.
--	---	---

Footnotes

- (a) For more information: The use of antiretroviral drugs for treating and preventing HIV infection. Recommendations for a public health approach. World Health Organization, second edition, 2016.
http://apps.who.int/iris/bitstream/10665/208825/1/9789241549684_eng.pdf?ua=1

References

1. World Health Organization. WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification de HIV-related disease in adults and children, 2007.
<http://www.who.int/hiv/pub/guidelines/HIVstaging150307.pdf> [Accessed 17 May 2018]
2. Word Health Organization. Guidelines for the diagnosis, prevention, and management of cryptococcal disease in HIV-infected adults, adolescents and children, Geneva, 2018.
<http://apps.who.int/iris/bitstream/handle/10665/260399/9789241550277-eng.pdf?sequence=1> [Accessed 17 May 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 9: Genito-urinary diseases

[Nephrotic syndrome in children](#)

[Urolithiasis](#)

[Acute cystitis](#)

[Acute pyelonephritis](#)

[Acute prostatitis](#)

[Genital infections](#)

[Urethral discharge](#)

[Abnormal vaginal discharge](#)

[Genital ulcers](#)

[Lower abdominal pain in women](#)

[Upper genital tract infections \(UGTI\)](#)

[Venereal warts](#)

[Major genital infections \(summary\)](#)

[Abnormal uterine bleeding \(in the absence of pregnancy\)](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

[Submit](#)

[Accept All](#)

Nephrotic syndrome in children

Nephrotic syndrome (NS) is characterized by the presence of oedema, heavy proteinuria, hypoalbuminemia, and hyperlipidaemia.

Primary or idiopathic NS is the most common cause of NS in children between 1 and 10 years. It usually responds to corticosteroids.

Secondary NS is associated with infectious diseases (e.g. post-infectious glomerulonephritis, endocarditis, hepatitis B and C, HIV infection, malaria, and schistosomiasis) and may respond to treatment of the underlying cause.

Children with NS are at increased risk of thromboembolism, severe bacterial infections (in particular, due to *S. pneumoniae*) and malnutrition. Untreated NS may progress to renal failure.

Clinical features

- Typically, the child presents with soft, pitting and painless oedema, which varies in location based on position and activity. Upon awaking, the child has periorbital or facial oedema, which over the day decreases as oedema of the legs increases.
As oedema worsens, it may localize to the back or genitals, or become generalized with ascites and pleural effusions.
- This oedema should be differentiated from the oedema of severe acute malnutrition (SAM): in SAM, the child presents with bilateral pitting oedema of the feet and lower legs that does not vary with position. Oedema extends upwards to hands and face in severe cases. It is usually associated with typical skin and hair changes (see Kwashiorkor: [Severe acute malnutrition](#), Chapter 1).
- Once SAM is excluded, the following two criteria must be met to make a clinical diagnosis of primary NS:
 - Presence of heavy proteinuria,
 - and
 - Absence of associated infections: see [Hepatitis B and C](#) and [HIV infection](#) (Chapter 8), [Malaria](#) and [Schistosomiases](#) (Chapter 6).

Laboratory

- Urine
 - Measure protein with urinary dipstick on three separate voided urine samples (first voided urine if possible). In NS, proteinuria is equal or greater than +++ or equal or greater than 300 mg/dl or 30 g/litre^a. NS is excluded if heavy proteinuria is not consistently present.
 - In case of macroscopic haematuria, or microscopic haematuria $\geq +$, consider glomerulonephritis.
- Blood tests (if available)
 - Serum albumin concentration less than 30 g/litre and hyperlipidaemia.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies


Submit

Accept All

- Exclude active tuberculosis and/or start antituberculous treatment.

- Corticosteroid treatment

See algorithm below. Total length of initial treatment is 2 to 4 months.



a Prednisone may be used interchangeably with prednisolone in this algorithm.

b If child has relapsed more than once, treat until proteinuria disappears but then taper prednisolone down to 0.5 mg/kg every other day rather than discontinuing entirely and treat for 12 months. Continue as long as proteinuria remains negative. If proteinuria recurs, treat as relapse. Child has steroid dependent NS.

c Frequent relapses: 2 or more in the first 6 months or 4 or more in a 12 month period.

- Nutrition, fluid intake, nursing and follow-up

- No salt-added diet.
- Do not restrict fluids (risk of thrombosis due to hypercoagulability). If oedema is very severe, fluids may initially be restricted (e.g. 75% of usual intake) while monitoring urine output.
- Encourage child to walk and play to prevent thromboembolism.
- Discharge child when stable, follow-up at least monthly, more frequently if indicated, weight and urine dipstick at each visit.
- Instruct the parent to continue no salt-added diet and to seek medical advice in case of fever, abdominal pain, respiratory distress or signs of thromboembolism.

- Management of infections

Treat infections as soon as they appear but do not routinely give prophylactic antibiotics.

- Immunization

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Signs include decreased urine output with any one of the following: capillary refill \geq 3 seconds, poor skin perfusion/mottling, cold extremities, low blood pressure.

If signs are present, administer **human albumin 5% IV**: 1 g/kg. If albumin is not available, administer **Ringer lactate or 0.9% sodium chloride**: 10 ml/kg over 30 minutes.

If signs of shock are present, see [Shock](#), Chapter 1.

- Respiratory distress due to severe oedema (rare)

This is the only situation in which diuretics should be used and only if there are no signs of intravascular volume depletion or after hypovolaemia has been corrected:

furosemide PO: 0.5 mg/kg 2 times daily

If not effective, discontinue furosemide. If creatinine is normal, administer **spironolactone** PO: 1 mg/kg 2 times daily. The dose can be increased to 9 mg/kg daily in resistant cases of ascites.

While on diuretics, monitor for dehydration, thromboembolism and hypokalaemia.

Specialized advice and management (including further investigations such as renal biopsy) are required:

- In children less than 1 year or more than 10 years,
- In case of steroid resistant NS,
- In case of mixed nephrotic and nephritic clinical picture.

In case of steroid-resistant NS, when referral is impossible and as a last resort, the following palliative measure may reduce proteinuria and delay renal failure:

enalapril PO: 0.1 to 0.3 mg/kg 2 times daily (start with the lowest dose and increase gradually if necessary until reduction of proteinuria). If available, monitor for hyperkalaemia.

This is a palliative measure and the prognosis for steroid-resistant NS is poor in the absence of specialized treatment.

Footnotes

- (a) Nephrotic range proteinuria in children is defined as urinary protein excretion greater than 50 mg/kg daily. Quantitative measurement of protein excretion is normally based on a timed 24-hour urine collection. However, if this test cannot be performed, urine dipstick measurements can be substituted.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Urolithiasis

Last updated: December 2020

Urolithiasis is the formation and passage of calculi (stones) in the urinary tract.

Clinical features

- Many calculi do not cause symptoms; they may be found incidentally through radiology exams.
- Symptoms arise when calculi cause partial or complete obstruction and/or infection:
 - Intermittent, acute flank to pelvic pain (renal colic). Pain can be severe and typically causes nausea and vomiting. Abdomen/flank may be tender to palpation. Patients are typically restless, finding no comfortable position.
 - Haematuria and/or gravel (calculi) passed in urine.
 - Fever and signs of pyelonephritis if secondary infection develops (see [Acute pyelonephritis](#), Chapter 9).

Note: if available, ultrasound may demonstrate calculi and hydronephrosis.

Treatment

- Encourage the patient to drink fluids.
- Administer analgesics according to the intensity of pain (see [Pain](#), Chapter 1).
- In case of secondary infection: antibiotic treatment as for [pyelonephritis](#). The effectiveness will depend on the passage of calculi.

Note: the majority of calculi pass spontaneously. If there are signs of significant renal dysfunction or secondary infection that does not improve with antibiotic treatment, consider surgical referral.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute cystitis

Last updated: July 2021

Cystitis is an infection of the bladder and urethra that affects mainly women and girls from 2 years of age. *Escherichia coli* is the causative pathogen in at least 70% of cases. Other pathogens include *Proteus mirabilis*, *Enterococcus* sp, *Klebsiella* sp and in young women, *Staphylococcus saprophyticus*.

Clinical features

- Burning pain on urination and urinary urgency and frequency; in children: crying when passing urine; involuntary loss of urine.
AND
- No fever (or mild fever), no flank pain; no systemic signs and symptoms in children.

It is essential to rule out [pyelonephritis](#).

The symptom 'burning pain on urination' alone is insufficient to make the diagnosis. See [Abnormal vaginal discharge](#).

Investigations

- Urine dipstick test:
Perform dipstick analysis for nitrites (which indicate the presence of enterobacteria) and leukocytes (which indicate an inflammation) in the urine.
 - If dipstick analysis is positive for nitrites and/or leukocytes, a urinary infection is likely.
 - In women, if dipstick analysis is negative for both nitrites and leukocytes, a urinary infection is excluded.
- Microscopy/culture: when a dipstick analysis is positive, it is recommended to carry out urine microscopy/culture in order to confirm the infection and identify the causative pathogen, particularly in children and pregnant women.
When urine microscopy is not feasible, an empirical antibiotic therapy should be administered to patients with typical signs of cystitis and positive dipstick urinalysis (leukocytes and/or nitrites).

Note: aside of these results, in areas where urinary schistosomiasis is endemic, consider schistosomiasis in patients with macroscopic haematuria or microscopic haematuria detected by dipstick test, especially in children from 5 to 15 years, even if the patient may suffer from concomitant bacterial cystitis.

- POCUS^a: in cases of recurrent cystitis, perform FAST views to evaluate for signs of urinary tract pathologies.

Treatment

Cystitis in girls ≥ 2 years

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- If dipstick analysis is negative for nitrites but positive for leukocytes, the infection may be due to *S. saprophyticus*. Fosfomycin is not active against this pathogen. Use nitrofurantoin as above.
- Whatever the antibiotic used, symptoms may persist for 2 to 3 days despite adequate treatment.
- In the event of treatment failure (or recurrent cystitis i.e. > 3-4 episodes per year), **ciprofloxacin** PO: 500 mg 2 times daily for 3 days
- For patients with recurrent cystitis, consider bladder stones, urinary schistosomiasis, urinary tuberculosis or gonorrhoea (examine the partner).

Cystitis in pregnant or lactating women

fosfomycin-trometamol PO: 3 g single dose

or

nitrofurantoin PO (contraindicated in the last month of pregnancy): 100 mg 3 times daily for 7 days

or

cefixime PO: 200 mg 2 times daily for 5 days

Footnotes

(a) POCUS should only be performed and interpreted by trained clinicians.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute pyelonephritis

Pyelonephritis is an infection of the renal parenchyma, more common in women than in men.

The pathogens causing pyelonephritis are the same as those causing cystitis (see [Acute cystitis](#), Chapter 9).

Pyelonephritis is potentially severe, especially in pregnant women, neonates and infants.

Management depends on the presence of signs of severity or complications or risk of complications.

Clinical features

Neonates and infant

- Symptoms are not specific: fever, irritability, vomiting, poor oral intake. Palpation of the lower abdomen may show abdominal tenderness. The absence of fever does not rule out the diagnosis. On the other hand, fever—with no obvious cause—may be the only manifestation.
- Neonates may present with fever or hypothermia, altered general condition, altered conscious state, pale/grey colour, shock.

In practice, a urinary tract infection should be suspected in children with unexplained fever or septic syndrome with no obvious focus of infection.

Older children and adults

- Signs of cystitis (burning pain on urination and urinary urgency and frequency, etc.)
AND
- Fever $> 38^{\circ}\text{C}$ and unilateral flank pain or abdominal tenderness
- Nausea and/or vomiting are common.

Laboratory

See [Acute cystitis](#), Chapter 9.

Treatment

- Criteria for hospital admission:
 - Patients at risk of complications: children, pregnant women, men^a, functional or structural abnormality of the urinary tract (lithiasis, malformation, etc.), severe immunodeficiency;
 - Patients with complicated pyelonephritis: urinary tract obstruction, renal abscess, emphysematous pyelonephritis in diabetic patients;
 - Patients with signs of severe infection: sepsis (infection with signs of organ dysfunction) and septic shock, dehydration or nausea/vomiting preventing hydration and oral treatment;

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Children 0 to 7 days (≥ 2 kg): 5 mg/kg once daily

Children 8 days to < 1 month: 5 mg/kg once daily

or

cefotaxime slow IV (3 minutes) for 7 to 10 days

Children 0 to 7 days (< 2 kg): 50 mg/kg every 12 hours

Children 0 to 7 days (≥ 2 kg): 50 mg/kg every 8 hours

Children 8 days to < 1 month: 50 mg/kg every 8 hours

▫ **Children one month and over**

ceftriaxone IM or slow IV^b (3 minutes): 50 mg/kg once daily until the child's condition improves (at least 3 days) then change to oral route to complete 10 days of treatment with:

amoxicillin/clavulanic acid PO (dose expressed in amoxicillin)

Children < 40 kg: 25 mg/kg 2 times daily

Children ≥ 40 kg:

Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)

Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)

• Antibiotherapy in adults^[1]

▫ **Uncomplicated pyelonephritis**

ceftriaxone IM: 1 g single dose or **gentamicin** IM: 5 mg/kg single dose

+

ciprofloxacin PO: 500 mg 2 times daily for 7 days

or

amoxicillin/clavulanic acid PO (dose expressed in amoxicillin) for 10 to 14 days

Ratio 8:1: 2000 mg daily (2 tablets of 500/62.5 mg 2 times daily)

Ratio 7:1: 1750 mg daily (1 tablet of 875/125 mg 2 times daily)

or

cefixime PO: 200 mg 2 times daily or 400 mg once daily for 10 to 14 days

▫ **Pyelonephritis with criteria for hospital admission**

ampicillin slow IV (3 minutes): 2 g every 6 hours for at least 3 days + **gentamicin** IM: 5 mg/kg once daily for 3 days

then change to amoxicillin/clavulanic acid PO (or another antibiotic depending on the antibiotic susceptibility test) to complete 10 to 14 days of treatment

or

ceftriaxone IV^b : 1 g once daily for at least 3 days + **gentamicin** IM: 5 mg/kg once daily for 3 days in the event of sepsis then change to amoxicillin/clavulanic acid PO (or another antibiotic depending on the antibiotic susceptibility test) to complete 10 to 14 days of treatment

Preferably use the combination ampicillin + gentamicin to cover enterococci.

Pyelonephritis with abscess formation or emphysematous pyelonephritis may require longer antibiotherapy.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

- (b) The solvent of ceftriaxone for IM injection contains lidocaine. Ceftriaxone reconstituted using this solvent must never be administered by IV route. For IV administration, water for injection must always be used.

References

1. Gupta K, Hooton TM, Naber KG, Wullt B, Colgan R, Miller LG, Moran GJ, Nicolle LE, Raz R, Schaeffer AJ, Soper DE, Infectious Diseases Society of America, European Society for Microbiology and Infectious Diseases. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: A 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. Clin Infect Dis. 2011;52(5):e103.
<https://academic.oup.com/cid/article/52/5/e103/388285> [Accessed 17 December 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Acute prostatitis

Prostatitis is an acute bacterial infection of the prostate.

The most common causative pathogen is *Escherichia coli*. Other pathogens include *Proteus mirabilis*, *Klebsiella* sp, *Pseudomonas aeruginosa* and *Enterococcus* sp.

Progression to chronic prostatitis is possible.

Clinical features

- Fever (often high) and chills.
- Signs of cystitis (burning on urination and urinary frequency).
- Perineal, urethral, penile or rectal pain.
- Urinary retention.

On examination:

- Very painful digital rectal examination. Fluctuant mass in case of prostatic abscess.
- Leukocyturia, pyuria, possible macroscopic haematuria.

Treatment

- Antibiotic therapy:
ciprofloxacin PO: 500 mg 2 times daily for 14 days then review the patient. Stop treatment if signs and symptoms have completely resolved. If signs and symptoms are ongoing continue the same treatment for a further 14 days.^[1]
- Symptomatic treatment:
 - Ensure adequate hydration (1.5 litres daily).
 - Treat fever (Chapter 1) and pain (Chapter 1).
- Refer to a surgeon in case of suspected prostatic abscess.

References

1. National Institute for Health and Care Excellence. NICE guideline [NG110] Prostatitis (acute): antimicrobial prescribing, 2018. <https://www.nice.org.uk/guidance/ng110/resources/visual-summary-pdf-6544018477> [Accessed 4 March 2020]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Genital infections

Last updated: August 2021

The diagnosis and treatment of genital infections (GI) present several difficulties: clinical features are not specific; many infections are asymptomatic; laboratory tests available in the field are not always reliable; mixed infections are common; sexual partners need to be treated simultaneously in case of sexually transmitted infections^a and the risk of recurrence or treatment failure is increased in HIV-infected patients.

Thus, the WHO has introduced the syndromic management of GI and developed standardised case management flowcharts: based on the identification of consistent groups of signs and symptoms (syndromes), patients are treated for the pathogens/infections^b that may cause each syndrome.

Look for a GI if a patient complains of:	See
Urethral discharge Painful or difficult urination (dysuria)	<u>Urethral discharge</u>
Abnormal vaginal discharge Vulvar itching/burning Pain with intercourse (dyspareunia) Painful or difficult urination (dysuria)	<u>Abnormal vaginal discharge</u>
Genital blisters or sores Burning sensation in the vulva or perineum	<u>Genital ulcers</u>
Skin growths in the genital (or anal) area	<u>Venereal warts</u>
Lower abdominal pain (in women)	<u>Lower abdominal pain</u> <u>Upper genital tract infections</u>

Basic principles of GI management

- The patient can be effectively treated without laboratory testing. Some tests may help in diagnosing vaginal and urethral discharge, but they should never delay treatment (results should be available within one hour).
- The patient should be treated at his/her first encounter with the health care provider (no patient should be sent home without treatment, e.g. while waiting for laboratory results).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Taking into consideration the physical, psychological, legal and social consequences of sexual violence, *medical care* is not limited to the diagnosis and treatment of genital lesions or infections.

Care includes listening to the victim's story, a complete physical examination, laboratory tests if available, and completion of a medical certificate.

During the consultation, prophylactic or curative treatments must be proposed to the patient.

- **Prophylactic treatment:**

- priority is given to:
 - ▷ a) the risk of HIV transmission. Start antiretroviral therapy as early as possible if the patient is seen within 48-72 hours after exposure (see [HIV infection and AIDS](#), Chapter 8);
 - ▷ b) the risk of pregnancy resulting from rape. Administer emergency contraception as soon as possible, ideally within 72 hours after the rape^c :
levonorgestrel PO, one 1.5 mg tablet single dose (including in women receiving HIV post-exposure prophylaxis); double the dose (3 mg) only if the patient was already taking an enzyme-inducing drug (e.g. rifampicin, carbamazepine, certain antiretrovirals) before the rape;
or **ulipristal** PO, one 30 mg tablet single dose;
or a **copper intrauterine device** (except in case of active genital infection);
- prevention of sexually transmitted infections: a single dose of **azithromycin** PO 2 g + **ceftriaxone** IM 500 mg (or, if ceftriaxone is not available, **cefixime** PO 400 mg). If necessary, treatment of trichomoniasis may be started later than the other treatments (**tinidazole** or **metronidazole** PO, 2 g single dose);
- tetanus prophylaxis and/or vaccination (see [Tetanus](#), Chapter 7) if there are any wounds;
- vaccination against hepatitis B (accelerated vaccination schedule, see [Viral hepatitis](#), Chapter 8).

- **Curative treatment:**

- of any related pathologies/infections if the assault is not recent.
- of wounds,

Mental health care is necessary irrespective of any delay between the event and the patient arriving for a consultation. Care is based on immediate attention (one-on-one reception and listening) and if necessary, follow-up care with a view to detecting and treating any psychological and/or psychiatric sequelae (anxiety, depression, post-traumatic stress disorder, etc.). See [Chapter 11](#).

Footnotes

- (a) GI may be sexually transmitted (e.g. gonorrhoea, chlamydia) or not (e.g. most cases of candidiasis).
- (b) Keep in mind that in *Schistosoma haematobium* endemic areas, genital symptoms may also be due to, or associated with, genitourinary schistosomiasis (see [Schistosomiasis](#), Chapter 6).
- (c) Nevertheless, between 72 and 120 hours (5 days) after the rape, emergency contraception is still sufficiently effective to be administered.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)


Urethral discharge

Last updated: August 2022

Urethral discharge is seen almost exclusively in men. The principal causative organisms are *Neisseria gonorrhoeae* (gonorrhoea) and *Chlamydia trachomatis* (chlamydia).

Abnormal discharge should be confirmed by performing a clinical examination^a. In males, the urethra should be milked gently if no discharge is visible. Furthermore, specifically check for urethral discharge in patients complaining of painful or difficult urination (dysuria).

Case management



Laboratory

- *C. trachomatis* cannot easily be identified in a field laboratory. In the absence of validated rapid diagnostic tests, the treatment is empiric.
- In men, a methylene blue or Gram stained smear from a urethral swab may be used to detect gonococci (Gram negative intracellular diplococci).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment for chlamydia		Treatment for gonorrhoea
azithromycin PO: 1 g single dose or doxycycline PO: 100 mg 2 times daily for 7 days	PLUS	ceftriaxone IM: 500 mg single dose or, if ceftriaxone is not available, cefixime PO: 400 mg single dose

If urethral discharge persists or reappears after 7 days:

- Verify that the patient has received an effective treatment (i.e. one of the combinations above).
- Gonococcal resistance is a possibility if another treatment (e.g. co-trimoxazole or kanamycin) has been administered: re-treat for gonorrhoea as above (chlamydia is rarely resistant).
- If an effective antibiotic therapy has been given, consider trichomoniasis (**tinidazole** or **metronidazole** PO, 2 g single dose); also consider reinfection.

Treatment of the partner

The sexual partner receives the same treatment as the patient, whether or not symptoms are present.

Footnotes

- (a) In areas where lymphatic filariasis is endemic, be careful not to confuse purulent urethral discharge with milky or rice-water urine (chyluria) suggestive of lymphatic filariasis.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Abnormal vaginal discharge

Last updated: August 2022

Abnormal vaginal discharge is defined as discharge that is different from usual with respect to colour/odour/consistency (e.g. discoloured or purulent or malodorous).

Abnormal discharge is often associated with vulvar pruritus or pain with intercourse (dyspareunia), or painful or difficult urination (dysuria) or lower abdominal pain. Routinely check for abnormal vaginal discharge in women presenting with these symptoms.

Abnormal vaginal discharge may be a sign of infection of the vagina (vaginitis) and/or the cervix (cervicitis) or upper genital tract infection.

Abnormal discharge must be clinically confirmed: inspection of the vulva, speculum exam checking for cervical/vaginal inflammation or discharge.

Abdominal and bimanual pelvic examinations should be performed routinely in all women presenting with vaginal discharge to rule out upper genital tract infection (lower abdominal pain and cervical motion tenderness).

The principal causative organisms are:

- In vaginitis: *Gardnerella vaginalis* and other bacteria (bacterial vaginosis), *Trichomonas vaginalis* (trichomoniasis) and *Candida albicans* (candidiasis).
- In cervicitis: *Neisseria gonorrhoeae* (gonorrhoea) and *Chlamydia trachomatis* (chlamydia).
- In upper genital tract infections: see [Upper genital tract infections](#).


Case management

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All



Cervicitis may be difficult to diagnose. When in doubt, administer treatment for cervicitis to women with abnormal vaginal discharge and any of the following risk factors:

- Urethral discharge in the partner
- Context of sexual violence or prostitution
- New partner or more than one partner in the preceding 3 months

Laboratory

- Xpert molecular (PCR) tests are recommended for the detection of *C. trachomatis* and *N. gonorrhoea*.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Treatment for chlamydia		Treatment for gonorrhoea
Non-pregnant women		
azithromycin PO: 1 g single dose or doxycycline PO: 100 mg 2 times daily for 7 days	PLUS	ceftriaxone IM: 500 mg single dose or, if not available, cefixime PO: 400 mg single dose
Pregnant women		
azithromycin PO: 1 g single dose or erythromycin PO: 1 g 2 times daily or 500 mg 4 times daily for 7 days	PLUS	ceftriaxone IM: 500 mg single dose or, if not available, cefixime PO: 400 mg single dose

Bacterial vaginosis and trichomoniasis

tinidazole PO: 2 g single dose
or **metronidazole** PO: 2 g single dose

In the case of treatment failure:

tinidazole PO: 500 mg 2 times daily for 5 days
or **metronidazole** PO: 400 to 500 mg 2 times daily for 7 days

Vulvovaginal candidiasis

clotrimazole (500 mg vaginal tab): 1 tablet inserted deep into the vagina at bedtime, single dose

If the patient has extensive vulvar involvement, **miconazole 2%** cream (one application to the vulva 2 times daily for 7 days) may be used in combination with the intravaginal treatment above. Miconazole cream may complement, but does not replace, treatment with clotrimazole.

Treatment of the partner

When the patient is treated for vaginitis or cervicitis, the partner receives the same treatment as the patient, whether or not symptoms are present.

In the case of vulvovaginal candidiasis, the partner is treated only if symptomatic (itching and redness of the glans/prepuce): **miconazole 2%** cream, one application 2 times daily for 7 days.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit


Accept All

Genital ulcers

Genital ulcers, defined as single or multiple vesicular, ulcerative or erosive lesions of the genital tract, with or without inguinal lymphadenopathy, should lead to consideration of sexually transmitted infection.

The principal causative organisms are *Treponema pallidum* (syphilis), *Haemophilus ducreyi* (chancroid) and *Herpes simplex* (genital herpes). *Chlamydia trachomatis* (lymphogranuloma venereum) and *Calymmatobacterium granulomatis* (donovanosis)^a are less frequent.

Case management



Laboratory

Laboratory testing available in the field is of little value: e.g., in syphilis, a negative RPR or VDRL result does not exclude primary syphilis in early stage, and a positive test may reflect previous infection in a successfully treated patient.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

In patients with a first episode, treatment may reduce the duration of symptoms when given within 5 days after the onset of symptoms: 400 mg 3 times daily for 7 days.

In patients with recurrence, give the same dose for 5 days, but treatment is only effective if initiated during the prodromal phase or within 24 hours after the onset of symptoms.

In patients with frequent recurrences (more than 6 episodes per year), see [HIV infection and AIDS](#), Chapter 8.

- Treatment of [pain](#): paracetamol PO (Chapter 1).

Syphilis

benzathine benzylpenicillin IM: 2.4 MUI per injection (half the dose in each buttock)^[1].

Early syphilis (primary, secondary, or early latent infection of less than 12 months duration): single dose

Late latent syphilis (infection of more than 12 months duration or of unknown duration): one injection weekly for 3 weeks

or, for penicillin-allergic patients or if penicillin is not available:

erythromycin PO: 1 g 2 times daily or 500 mg 4 times daily for 14 days (early syphilis) or 30 days (late latent syphilis)

or

doxycycline PO: 100 mg 2 times daily for 14 days (early syphilis) or 30 days (late latent syphilis)^b

or

azithromycin PO: 2 g single dose (only in cases of early syphilis and only if the strain is sensitive)^[2]

Chancroid

azithromycin PO: 1 g single dose

or

ceftriaxone IM: 250 mg single dose

or

erythromycin PO: 1 g 2 times daily or 500 mg 4 times daily for 7 days

Fluctuant lymph nodes may be aspirated through healthy skin as required. Do not incise and drain lymph nodes.

Note: treat simultaneously for syphilis AND chancroid as both are frequent, and cannot be correctly distinguished on clinical grounds.

Lymphogranuloma venereum

erythromycin PO: 1 g 2 times daily or 500 mg 4 times daily for 14 days

or

doxycycline PO: 100 mg 2 times daily for 14 days^b

Fluctuant lymph nodes may be aspirated through healthy skin as required. Do not incise and drain lymph nodes.

Donovanosis

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

The sexual partner receives the same treatment as the patient, whether or not symptoms are present, except in the case of genital herpes (the partner is treated only if symptomatic).

Footnotes

- (a) Lymphogranuloma venereum is endemic in East and West Africa, India, Southeast Asia, South America and the Caribbean. Donovanosis is endemic in South Africa, Papua New Guinea, India, Brazil and the Caribbean.
- (b) Doxycycline is contra-indicated in pregnant and breast-feeding women.

References

1. Centers for Disease Control and Prevention. Syphilis Pocket Guide for Providers. 2017.
<https://www.cdc.gov/std/syphilis/Syphilis-Pocket-Guide-FINAL-508.pdf>
2. World Health Organization. WHO guidelines for the treatment of *Treponema pallidum* (syphilis), Geneva, 2016.
<http://apps.who.int/iris/bitstream/handle/10665/249572/9789241549806-eng.pdf?sequence=1>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Lower abdominal pain in women


Upper genital tract infection should be suspected in women with lower abdominal pain (see [Upper genital tract infections](#)).

Gynaecological examination should be routinely performed:

- Inspection of the vulva, speculum examination: check for purulent discharge or inflammation.
- Abdominal exam and bimanual pelvic exam: check for pain on mobilising the cervix.

If available, POCUS^a: perform FAST views to evaluate for free fluid and urological abnormalities. Perform pelvic views to evaluate for uterine and adnexal pathologies. Consult a gynaecologist (local or via telemedicine services).

Case management



This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Upper genital tract infections (UGTI)

Upper genital tract infections are bacterial infections of the uterus (endometritis) and/or the fallopian tubes (salpingitis), which may be complicated by peritonitis, pelvic abscess or septicaemia.

UGTI may be sexually transmitted or arise after childbirth or abortion. Antibiotic choices are directed by the most common pathogens in each scenario.

If peritonitis or pelvic abscess is suspected, request a surgical opinion while initiating antibiotic therapy.

Clinical features

Sexually transmitted infections

Diagnosis may be difficult, as clinical presentation is variable.

- Suggestive symptoms are: abdominal pain, abnormal vaginal discharge, fever, dyspareunia, menometrorrhagia, dysuria.
- Infection is probable when one or more of the above symptoms are associated with one or more of the following signs: cervical motion tenderness, adnexal tenderness, tender abdominal mass.

Infections after childbirth or abortion

- Most cases present with a typical clinical picture, developing within 2 to 10 days after delivery (caesarean section or vaginal delivery) or abortion (spontaneous or induced):
 - Fever, generally high
 - Abdominal or pelvic pain
 - Malodorous or purulent lochia
 - Enlarged, soft and/or tender uterus
- Check for retained placenta.
- In the early stages, fever may be absent or moderate and abdominal pain may be mild.

Treatment

- Criteria for hospitalisation include:
 - Clinical suspicion of severe or complicated infection (e.g. peritonitis, abscess, septicaemia)
 - Diagnostic uncertainty (e.g. suspicion of extra-uterine pregnancy, appendicitis)
 - Significant obstacles to ambulatory oral treatment
 - No improvement after 48 hours, or deterioration within 48 hours, of outpatient treatment
- All other patients may be treated on an ambulatory basis. They should be reassessed routinely on the third day of treatment to evaluate clinical improvement (decrease in pain, absence of fever). If it is difficult to organise routine follow-up advise patients to return to clinic if there is no improvement after 48 hours of treatment or sooner if their

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Treatment in hospital:
 - ceftriaxone** IM: 250 mg once daily
 - + **doxycycline** PO: 100 mg 2 times daily for 14 days^a
 - + **metronidazole** PO or IV infusion: 500 mg 2 times daily for 14 days
 - Continue triple therapy for 24 to 48 hours after signs and symptoms have improved (resolution of fever, decrease in pain), then continue doxycycline (or erythromycin) + metronidazole to complete 14 days of treatment.
- If an IUD is in place, it should be removed (offer another method of contraception).
- Analgesic treatment according to pain intensity.
- Treatment of the partner: single dose treatment for both gonorrhoea AND chlamydia (as for Urethral discharge), whether or not symptoms are present.

Infections after childbirth or abortion

- Antibiotic therapy: treatment must cover the most frequent causative organisms: anaerobes, Gram negatives and streptococci.
 - Ambulatory treatment (early stages only):
 - amoxicillin/clavulanic acid (co-amoxiclav)** PO for 7 days
 - Use formulations in a ratio of 8:1 or 7:1 exclusively. The dose is expressed in amoxicillin:
 - Ratio 8:1: 3000 mg daily (2 tablets of 500/62.5 mg 3 times daily)
 - Ratio 7:1: 2625 mg daily (1 tablet of 875/125 mg 3 times daily)
 - or
 - amoxicillin** PO: 1 g 3 times daily + **metronidazole** PO: 500 mg 3 times daily doses for 7 days
 - Treatment in hospital:
 - amoxicillin/clavulanic acid (co-amoxiclav)** IV (dose expressed in amoxicillin): 1 g every 8 hours
 - + **gentamicin** IM: 5 mg/kg once daily
 - or
 - ampicillin** IV: 2 g every 8 hours
 - + **metronidazole** IV infusion: 500 mg every 8 hours
 - + **gentamicin** IM: as above

Stop antibiotic therapy 48 hours after resolution of fever and clinical signs and symptoms.
In penicillin-allergic patients, use **clindamycin** IV (900 mg every 8 hours) + **gentamicin** (as above).
- In case of placental retention: perform digital curettage or manual vacuum extraction (refer to the guide Essential obstetric and newborn care, MSF) 24 hours after initiation of antibiotic therapy.
- Analgesic treatment according to pain intensity.
- If the patient's condition deteriorates or if fever persists after 48-72 hours of treatment, consider the possibility of complication requiring additional treatment (e.g. pelvic abscess drainage), otherwise change the antibiotic to ceftriaxone + doxycycline + metronidazole as in hospital-based treatment of sexually transmitted UGTI.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Venereal warts

Venereal warts are benign tumours of the skin or mucous membranes due to certain papilloma viruses (HPV).

Clinical features

- Venereal warts are soft, raised, painless growths, sometimes clustered (cauliflower-like appearance) or macules (flat warts), which are more difficult to discern. Warts can be external (vulva, penis, scrotum, perineum, anus) and/or internal (vagina, cervix, urethra, rectum; oral cavity in HIV infected patients).
- In women, the presence of external warts is an indication for a speculum examination to exclude vaginal or cervical warts. Speculum exam may reveal a friable, fungating tumour on the cervix, suggestive of cancer associated with papilloma virus^a.

Treatment

Choice of treatment depends on the size and location of the warts. Treatment may be less effective, and relapses more frequent, in HIV infected patients.

External warts < 3 cm and vaginal warts

podophyllotoxin 0.5%^b solution may be self-applied by the patient, but in the event of vaginal warts, the treatment must be applied by medical staff.

Explain the procedure to the patient: apply the solution to the warts using an applicator or cotton bud, sparing the surrounding healthy skin, allow to air dry. On vaginal warts, the solution should be allowed to dry before the speculum is withdrawn.

Apply the solution 2 times daily, 3 consecutive days per week, for up to 4 weeks.

Podophyllum preparations are contra-indicated in pregnant^c or breastfeeding women. They should not be applied on cervical, intra-urethral, rectal, oral or extensive warts. Improper use may result in painful ulceration.

External warts > 3 cm; cervical, intra-urethral, rectal and oral warts; warts in pregnant or breastfeeding women

Surgical excision or cryotherapy or electrocoagulation.

Footnotes

(a) Certain types of HPV may cause cancer. Presence of genital warts in women is an indication to screen for precancerous

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Major genital infections (summary)

Last updated: July 2021

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Pathogens/ Infections	Clinical features	Investigations	Treatment
<i>Neisseria gonorrhoeae</i> (gonorrhoea)	<ul style="list-style-type: none"> In women: <ul style="list-style-type: none"> vaginal discharge, cervicitis (mucopurulent cervical discharge), dysuria (50% of infections are asymptomatic); UGTI (salpingitis, endometritis). In men: purulent urethral discharge and sometimes dysuria (5 to 50% of infections are asymptomatic). 	<ul style="list-style-type: none"> Best method is PCR (Xpert), if available. In men (not sensitive enough in women): Gram or methylene blue stain: intracellular diplococci and polymorphonuclear leukocytes (more than 4 per field). 	ceftriaxone IM: 500 mg single dose or, if not available, cefixime PO: 400 mg single dose Treat also for chlamydia. In case of upper genital tract infection, see <u>UGTI</u> .
<i>Chlamydia trachomatis</i> (chlamydia)	<ul style="list-style-type: none"> In women: <ul style="list-style-type: none"> vaginal discharge, cervicitis, and rarely dysuria (> 50% of infections are asymptomatic); UGTI (salpingitis, endometritis). In men: mild urethral discharge and/or dysuria but up to 90% of infections are asymptomatic. 	<ul style="list-style-type: none"> The best method is PCR (Xpert), if available. 	azithromycin PO: 1 g single dose or doxycycline PO ^(a) : 200 mg daily for 7 days Treat also for gonococcal infection (except when a Gram stain in males or PCR shows no <i>N. gonorrhoeae</i>). In case of upper genital tract infection, see <u>UGTI</u> .
<i>Trichomonas vaginalis</i> (trichomoniasis)	<ul style="list-style-type: none"> In women: yellow-green vaginal discharge, sometimes foul smelling, vulvar irritation (10 to 50% of infections are asymptomatic). In men: most infections are asymptomatic. Can produce balanitis, urethritis with mild discharge and sometimes dysuria. 	<ul style="list-style-type: none"> Wet mount of fresh vaginal fluid shows motile trichomonas (low sensitivity). pH of urethral/vaginal fluid > 4.5. 	tinidazole or metronidazole PO: 2 g single dose
Bacterial vaginosis	<p>Diagnosis is made in the presence of 3 of the following 4 signs:</p> <ul style="list-style-type: none"> Homogenous grey-white adherent vaginal discharge 		tinidazole or metronidazole PO: 2 g

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	<ul style="list-style-type: none"> In men: balanitis/balanoposthitis (inflammation of the glans/prepucie, erythema, pruritus, white pustules) and rarely urethritis 	<ul style="list-style-type: none"> pH of vaginal fluid: normal 	<ul style="list-style-type: none"> In men: <p>miconazole 2% cream: 1 application 2 times daily for 7 days</p>
<i>Herpes simplex</i> virus type 2 (genital herpes)	Many asymptomatic carriers. Multiple vesicles on genitals leading to painful ulcerations. In women, affects vulva, vagina and cervix; in males, penis and sometimes urethra. In primary episodes, fever (30%) and lymphadenopathy (50%). Recurrences in 1/3 of infections with shorter and milder symptoms.	Diagnosis by culture, serology and PCR done exclusively at a reference laboratory.	<p>Analgesics, local disinfection. If available, aciclovir PO:</p> <ul style="list-style-type: none"> Primary episode: 1200 mg daily for 7 days, given within 5 days after onset of lesions. Recurrent infections: same dose for 5 days, given within 24 hours after onset of lesions.
<i>Treponema pallidum</i> (syphilis)	Single firm painless genital ulcer, often unnoticed.	RPR/VDRL lack sensitivity and specificity, but may be useful for following treatment effectiveness (decrease in titer) or confirming re-infection (rise in titer). Treponemal tests (TPHA, FTA-ABS, rapid tests such as SD Bioline®) are more sensitive and specific.	<p>benzathine benzylpenicillin IM: 2.4 MIU per injection, single dose (syphilis < 12 months) or once weekly for 3 weeks (syphilis > 12 months or unknown duration) or azithromycin PO: 2 g single dose or erythromycin PO: 2 g daily for 14 days or doxycycline PO^(a): 200 mg daily for 14 days Treat also for chancroid.</p>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

<i>Human papillomavirus (venereal warts)</i>	Soft, raised, painless growths, sometimes clustered (acuminate condyloma) or macules (flat warts). Warts can be external (vulva, penis, scrotum, perineum, anus) and/or internal (vagina, cervix, urethra, rectum; oral cavity in HIV infected patients).	The diagnosis is based on clinical features. It is feasible in the context, the presence of genital warts in women in an indication to screen for pre-cancerous lesions of the cervix (visual inspection with acetic acid, or cervical smear, or other available techniques).	<ul style="list-style-type: none"> External warts < 3 cm and vaginal warts: podophyllotoxin 0.5% External warts > 3 cm; cervical, intra-urethral, rectal and oral warts; warts in pregnant or breastfeeding women: surgical excision or cryotherapy or electrocoagulation.
--	---	---	---

- (a) Doxycycline is contra-indicated in pregnant women. It should not be administered to breast-feeding women if the treatment exceeds 7 days (use erythromycin).
- (b) Ciprofloxacin should be avoided in pregnant women.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Abnormal uterine bleeding (in the absence of pregnancy)

Last updated: October 2021

- Heavy menstrual bleeding or intermenstrual genital bleeding
- In women of childbearing age:
 - assess if the bleeding is pregnancy-related;
 - perform a pregnancy test.

For the management of pregnancy-related bleeding, refer to the guide [Essential obstetric and newborn care](#), MSF.

In all events

- Rapidly assess the severity of bleeding.
- Perform a pelvic examination:
 - speculum examination: determine the origin (vagina, cervix, uterine cavity) and cause of the bleeding; appearance of the cervix; amount and intensity of bleeding;
 - bimanual examination: look for cervical motion tenderness, uterine enlargement or irregularity.
- Assess for recent trauma or surgical history.
- Measure haemoglobin, if possible, to prevent or treat anaemia.
- In the event of signs of shock, see [Shock](#), Chapter 1.
- In the event of heavy bleeding:
 - start an IV infusion of Ringer lactate;
 - monitor vital signs (heart rate, blood pressure);
 - administer^[1] :
tranexamic acid IV: 10 mg/kg (max. 600 mg) every 8 hours. When bleeding has been reduced, switch to
tranexamic acid PO: 1 g 3 times daily, until bleeding stops (max. 5 days).
 - if bleeding persists and/or in case of contraindication to tranexamic acid, administer one of the following two drugs (except if suspicion of cervical or endometrial cancer):
ethinylestradiol/levonorgestrel PO (0.03 mg/0.15 mg tab): one tablet 3 times daily for 7 days
or **medroxyprogesterone acetate** PO: 20 mg 3 times daily for 7 days
- In case of massive haemorrhage and/or lack of response to medical management: surgical management (dilation and curettage, intrauterine balloon, and as a last resort, hysterectomy).
- In the event of referral to a surgical facility, difficult transport conditions may aggravate the bleeding: the patient should have an IV line and/or be accompanied by family members who are potential blood donors.
- If available, POCUS^a : perform FAST to evaluate for free fluid and/or urological abnormalities; perform pelvic views

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Enlarged, irregular uterus: uterine fibroids. In case of failure to respond to medical treatment, surgical management is required. While waiting for surgery or if surgery is not indicated, treat as for functional uterine bleeding.
- Normal uterus and cervix: possible functional uterine bleeding: **tranexamic acid** PO as above. In case of repeated bleeding, it can be combined with an NSAID (**ibuprofen** PO for 3 to 5 days, see [Pain](#), Chapter 1) and/or one of the following long-term treatments:
 - **levonorgestrel intrauterine device**
 - or **ethinylestradiol/levonorgestrel** PO (0.03 mg/0.15 mg tab): one tablet daily
 - or **medroxyprogesterone acetate** IM: 150 mg every 3 months
 - or **medroxyprogesterone acetate** PO^b: 10 mg once daily (up to 30 mg once daily if necessary) for 21 days monthly.

Note: rule out other causes of vaginal bleeding before diagnosing functional uterine bleeding. Consider for example poorly tolerated contraceptive, endometrial cancer in postmenopausal women, genitourinary schistosomiasis in endemic areas (see [Schistosomiasis](#), Chapter 6).

Footnotes

- (a) POCUS should only be performed and interpreted by trained clinicians.
- (b) Unlike the other treatments, this drug has no contraceptive effect.

References

1. American College of Obstetricians and Gynecologists. Management of acute abnormal uterine bleeding in nonpregnant reproductive-aged women. *Obstet Gynecol*. 2013 Apr;121(4):891-6.
<https://www.acog.org/-/media/project/acog/acogorg/clinical/files/committee-opinion/articles/2013/04/management-of-acute-abnormal-uterine-bleeding-in-nonpregnant-reproductive-aged-1.pdf>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 10: Medical and minor surgical procedures

[Dressings](#)

[Treatment of a simple wound](#)

[Burns](#)

[Cutaneous abscess](#)

[Pyomyositis](#)

[Leg ulcers](#)

[Necrotising infections of the skin and soft tissues](#)

[Venomous bites and stings](#)

[Dental infections](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Dressings

The objective of dressing wounds is to promote healing. The procedure includes cleaning, disinfection and protection of the wound while respecting the rules of hygiene.

Not all wounds need to be covered by a dressing (e.g. a clean wound that has been sutured for several days; a small dry wound not requiring sutures).

Equipment

Sterile instruments

- One Kocher or Pean forceps
- One dissecting forceps
- One pair of surgical scissors or one scalpel to excise necrotic tissue and to cut gauze or sutures

Instruments for one dressing for one patient must be wrapped together in paper or fabric (or can be placed in a metallic box) and sterilised together to limit handling and breaks in asepsis. 5 to 10 compresses may be included in this set.

If there are no sterile instruments, a dressing can be done using sterile gloves.

Renewable supplies

- Sterile compresses
- Non-sterile disposable gloves
- Adhesive tape and/or crepe or gauze bandage
- Sterile 0.9% sodium chloride or sterile water
- Depending on the wound: antiseptic (7.5% povidone iodine scrub solution, 10% povidone iodine dermal solution), paraffin compresses, analgesics

Organisation of care

Proper organization of care helps maintain the rules of asepsis and decreases the risk of contamination of the wound or transmission of organisms from one patient to another:

- Assign one room for dressings. It must be cleaned and the waste removed every day. The dressing table must be disinfected after each patient.
- Dressings may be applied at the bedside if the patient's condition requires. Use a clean, disinfected dressing trolley with: on the upper tray, sterile and/or clean material (dressing set, extra compresses, etc.) and on the lower tray, septic material (container for contaminated instruments, sharps disposal container and a container or garbage bag for waste).
- Prepare all the necessary material in a well lit area. If necessary, arrange for an assistant to be present.
- Wear protective glasses if there is a risk of projection from an oozing wound.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- To prevent drug interactions, use the same antiseptic for all care of one patient.

Removal of an old dressing

- Wash hands (ordinary soap) or disinfect them with an alcohol-based hand rub.
- Put on non-sterile gloves and remove the adhesive tape, bandage and superficial compresses.
- Proceed gently with the last compresses. If they stick to the wound, loosen them with 0.9% sodium chloride or sterile water before removal.
- Observe the soiled compresses. If there is significant discharge, a greenish colour or a foul odour, a wound infection is likely.
- Discard the dressing and the non-sterile gloves in the waste container.

Observe the wound

- In the case of an open wound, loss of cutaneous tissue or ulcer, the colour is an indicator of the stage in the healing process:
 - **black** area = necrosis, wet or dry infected eschar
 - **yellow or greenish** area = infected tissue and presence of pus
 - **red** area = granulation, usually a sign of healing (unless there is hypertrophy), however, red edges indicate inflammation or infection
 - **pink** area = process of epithelisation, the final stage of healing that begins at the edges of the wound
- In the case of a sutured wound, the existence of local signs of suppuration and pain requires the removal of one or more sutures to avoid the infection spreading. Local signs include:
 - red, indurated and painful edges
 - drainage of pus between the sutures, either spontaneously or when pressure is applied on either side of the wound
 - lymphangitis
 - sub-cutaneous crepitations around the wound

In any case, if local signs of infection are observed, look for general signs of infection (fever, chills, changes in the overall condition).

Technique for cleaning and dressing of the wound

- Wash hands again or disinfect them with an alcohol-based hand rub.
- Open the dressing set or box after checking the date of sterilisation and that the wrapping is intact.
- Pick up one of the sterile forceps being careful not to touch anything else.
- Pick up the second forceps with the help of the first one.
- Make a swab by folding a compress in 4 using the forceps.
- **Clean sutured wound or clean open wound with red granulation:**
 - clean with 0.9% sodium chloride or sterile water to remove any organic residue; work from the cleanest to the dirtiest areas.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Discard any sharp materials used in an appropriate sharps container and the rest of the waste in a waste container.
- As quickly as possible, soak the instruments in disinfectant.
- Wash hands again or disinfect them with an alcohol-based hand rub.

The principles remain the same if the dressing is done using instruments or sterile gloves.

Subsequent dressings

- Clean, sutured wound: remove the initial dressing after 5 days if the wound remains painless and odourless, and if the dressing remains clean. The decision to re-cover or to leave the wound uncovered (if it is dry) often depends on the context and local practices.
- Infected, sutured wound: remove one or more sutures and evacuate the pus. Change the dressing at least once daily.
- Open, dirty wound: daily cleaning and dressing change.
- Open granulating wound: change the dressing every 2 to 3 days, except if the granulation is hypertrophic (in this case, apply local corticosteroids).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Treatment of a simple wound

A simple wound is a break in the continuity of the skin limited in depth at the sub-cutaneous fatty tissue, that does not affect the underlying structures (muscle, bone, joints, major arteries, nerves, tendons) and without significant loss of tissue.

The goal of treatment is to assure rapid healing of the wound without complications or sequelae. Several basic rules apply:

- rapidly treat wounds, while maintaining the rules of asepsis and the order of the initial procedures: cleaning-exploration-excision;
- identify wounds that need to be sutured and those for which suturing would be harmful or dangerous;
- immediately suture recent, clean, simple wounds (less than 6 hours old) and delay suturing contaminated wounds and/or those more than 6 hours old;
- prevent local (abscess) or general (gas gangrene; tetanus) infections.

Equipment

Instruments

(Figures 1a to 1d)

- One dissecting forceps, one needle-holder, one pair of surgical scissors and one Pean or Kocher forceps are usually enough.
- One or two other artery forceps, a pair of Farabeuf retractors and a scalpel may be useful for a contused or deep wound.

Instruments to suture one wound for one patient must be packaged and sterilised together (suture box or set) to limit handling and breaks in asepsis.

Renewable supplies

- For local anaesthesia: sterile syringe and needle; 1% lidocaine (without epinephrine)
- Sterile gloves, fenestrated sterile towel
- Sterile absorbable and non-absorbable sutures
- Antiseptic and supplies for dressings
- For drainage: corrugated rubber drain or equivalent, nylon suture

Technique

- Settle the patient comfortably in an area with good lighting and ensure all the necessary material is prepared.
- Explain the procedure to the patient and ensure his co-operation.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- If the wound is infected and the patient has general signs of infection (fever, chills, changes in the overall condition) systemic antibiotic therapy may be required. Administer antibiotics at least one hour prior to starting care.

Exploration

- Wash hands and put on sterile gloves.
- Disinfect the wound and surrounding area with 10% povidone iodine.
- Cover the wound with a fenestrated sterile towel.
- Local anaesthetic: infiltrate **1% lidocaine** into the edges of the wound and wait at least 2 minutes for the anaesthetic to take effect.
- Proceed carefully from the superficial to the deepest parts of the wound to explore the extent of the wound, if necessary, aided by an assistant.
- Consider the anatomical location of the wound and look for injury to any underlying structures (the clinical examination of a limb must include evaluation of sensitivity and motor functioning, as well as that of tendons in order to orient surgical exploration):
 - a wound that communicates with a fracture is an open fracture,
 - a wound close to a joint may be a joint wound,
 - a wound on the hands or feet may affect the nerves and/or tendons,
 - a wound close to a major artery may be an arterial wound even if it is no longer bleeding.
- Look for and remove any foreign bodies.
- In the event of significant pain or bleeding, the exploration must be completed in an operating room.

Wound excision

- The goal of the excision is to remove non-viable tissue, which favours the proliferation of bacteria and infection.
- The wound may require little or no excision if it is clean. The excision is more extensive if the wound is bruised, irregular or extensive.
- Limit excision of the skin around the wound, particularly in facial wounds.
- Sub-cutaneous fat and tissue of doubtful viability should be generously excised in order to leave only well vascularised tissue.

Immediate suturing of a simple wound

- Immediate suturing may have serious consequences for the patient if precautions to prevent infection and promote healing are not taken.
- The decision to suture immediately can only be taken after the cleaning, exploration and satisfactory excision, and if the following conditions are met: simple wound, no more than 6 hours old with no devitalised or contused tissue (the wound may be as long as 24 hours old if on the face, scalp, upper limbs or hands).
- Bites (for local treatment see [Rabies](#), Chapter 8) and bullet, shell or mine shrapnel wounds should not be immediately sutured.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

If the wound does not meet the conditions of cleanliness described above, the wound cannot be sutured. It will heal either spontaneously (healing by secondary intention), or will require a skin graft (once the wound is clean) if there is significant loss of tissue.

Figures 1 : Basic instruments



Figure 1a

Kocher forceps,
straight, toothed



Figure 1b

Kelly forceps,
curved, non-toothed



Figure 1c

Small artery forceps,
curved, non-toothed




Figure 1d

Farabeuf retractors

Figures 2 : How to hold instruments

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All




Figure 2a

Always mount a surgical blade using a needle holder.
Change the blade for each new procedure.




Figure 2b

Dissecting forceps should not be held in the palm of the hand,
but rather between the thumb and index finger.
Toothed dissecting forceps should only be used on skin.




Figure 2c

Insert the thumb and the ring finger into the handle of a needle holder (or scissors), and stabilize the instrument using
the index finger.

Figures 3 : Wound debridement

This should be done sparingly, limited to excision of severely contused or lacerated tissue that is clearly becoming necrotic.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All



Figure 3a

Debridement of a contused, ragged wound: straightening of the wound edges with a scalpel. Be conservative in facial wounds.




Figure 3b

Excision of edges of the aponeurosis to prevent necrosis.




Figure 3c

Excision of contused muscle.

Figures 4 : Practising making knots using forceps

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All




Figure 4a

Loop the suture around the needle holder in one direction and remember the direction of the loop. Grasp the loose end with the needle holder and pull it through the loop to make the first knot.
Lower the knot so that it closes the wound.




Figure 4b

The second loop should be in the opposite direction. At least 3 knots are needed to make a suture, alternating from one direction to the other.




Figure 4c

In principle the first knot lies flat.



Figure 4d

Second knot in the opposite direction.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Grasp the loose end with the needle holder.




Figure 4g

Fist flat knot.

Slide the knot towards the wound using the hand holding the loose end while holding the other end with the needle holder.

Tighten the knot without causing tissue ischaemia.




Figure 4h




Figure 4i

Second knot in the opposite direction.

Figures 5 : Particular problems

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All




Figure 5a




Figure 5b

The suture should be as deep as it is wide.




Figure 5c




Figure 5d

The suture is too shallow, the edges are invaginated.




Figure 5e

Poor lining of the edges.





Figure 5f

Do not make the knot directly over the wound.

Figure 6 : Closing a corner



This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Burns

Last updated: August 2022

Burns are cutaneous lesions caused by exposure to heat, electricity, chemicals or radiation. They cause significant pain and may threaten survival and/or compromise function.

Classification of burns

Severe burns: one or more of the following parameters:

- Involving more than 10% of the body surface area (BSA) in children and 15% in adults
- Inhalation injury (smoke, hot air, particles, toxic gas, etc.)
- Major concomitant trauma (fracture, head injury, etc.)
- Location: face, hands, neck, genitalia/perineum, joints (risk of functional deficit)
- Electrical and chemical burns or burns due to explosions
- Age < 3 years or > 60 years or significant co-morbidities (e.g. epilepsy, malnutrition)

Minor burns: involving less than 10% of the BSA in children and 15% in adults, in the absence of other risk factors

Evaluation of burns

Extent of burns

Lund-Browder table – Percentage of body surface area according to age

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Location	< 1 year	1-4 years	5-9 years	10-15 years	Adults
Head	19	17	13	10	7
Neck	2	2	2	2	2
Anterior trunk	13	13	13	13	13
Posterior trunk	13	13	13	13	13
Right buttock	2.5	2.5	2.5	2.5	2.5
Left buttock	2.5	2.5	2.5	2.5	2.5
Perineum/genitalia	1	1	1	1	1
Right upper arm	4	4	4	4	4
Left upper arm	4	4	4	4	4
Right lower arm	3	3	3	3	3
Left lower arm	3	3	3	3	3
Right hand	2.5	2.5	2.5	2.5	2.5
Left hand	2.5	2.5	2.5	2.5	2.5
Right thigh	5.5	6.5	8.5	8.5	9.5
Left thigh	5.5	6.5	8.5	8.5	9.5
Right leg	5	5	5.5	6	7
Left leg	5	5	5.5	6	7
Right foot	3.5	3.5	3.5	3.5	3.5
	--	--	--	--	--

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Apart from first-degree burns (painful erythema of the skin and absence of blisters) and very deep burns (third-degree burns, carbonization), it is not possible, upon initial examination, to determine the depth of burns. Differentiation is possible after D8-D10.

	Superficial burn on D8-D10	Deep burn on D8-D10
Sensation	Normal or pain	Insensitive or diminished sensation
Colour	Pink, blanches with pressure	White, red, brown or black Does not blanch with pressure
Texture	Smooth and supple	Firm and leathery
Appearance	Minimal fibrinous exudate Granulation tissue evident Bleeds when incised	Covered with fibrinous exudate Little or no bleeding when incised
Healing	Heals spontaneously within 5-15 days	<ul style="list-style-type: none"> Very deep burn: always requires surgery (no spontaneous healing) Intermediate burn: may heal spontaneously in 3 to 5 weeks; high risk of infection and permanent sequelae

Evaluation for the presence of inhalation injury

Dyspnoea with chest wall indrawing, bronchospasm, soot in the nares or mouth, productive cough, carbonaceous sputum, hoarseness, etc.

Treatment of severe burns

I. Initial management

On admission

- Ensure airway is patent; high-flow oxygen, even when SpO₂ is normal.
- Establish IV access, through unburned skin if possible (intraosseous access if IV access is not possible).
- Ringer lactate (RL):** 20 ml/kg during the first hour, even if the patient is stable.
- Morphine SC:** 0.2 mg/kg (Step 1 and Step 2 analgesics are not effective).
- In the event of chemical burns: flush with copious amounts of water for 15 to 30 minutes, avoiding contamination of healthy skin; do not attempt to neutralize the chemical agent.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Insert a nasogastric tube if burns involve > 20% of BSA (in the operating room while carrying out dressing procedure).
- Calculate and initiate fluid and electrolyte requirements for the first 24 hours.
- Intensive monitoring: level of consciousness, heart rate, blood pressure, SpO₂, respiratory rate (RR) hourly; temperature and urine output every 4 hours.
- Additional testing: haemoglobin, blood group, urine dipstick test.
- Prepare the patient for the first dressing procedure in the operating room.

Notes:

- Burns do not bleed in the initial stage: check for haemorrhage if haemoglobin level is normal or low.
- Burns alone do not alter the level of consciousness. In the case of altered consciousness, consider head injury, intoxication, postictal state in epileptic patients.
- Clinical manifestations of electrical burns vary significantly according to the type of current. Look for complications (arrhythmia, rhabdomyolysis, neurological disorders).

II. General measures during the first 48 hours

Resuscitative measures

Intravenous replacement fluid to correct hypovolaemia:

Fluid and electrolyte requirements during the first 48 hours according to age

	Children < 12 years	Children ≥ 12 years and adults
0 - 8 h	2 ml/kg x % BSA of RL + maintenance fluid ^(a) per hour x 8 h	2 ml/kg x % BSA of RL
8 - 24 h	2 ml/kg x % BSA of RL + maintenance fluid ^(a) per hour x 16 h	2 ml/kg x % BSA of RL
24 - 48 h	Daily maintenance IV fluid requirements ^(a) minus oral fluids such as milk, broth, gavage feeds (do not include drinking water in the calculation).	40 ml/kg RL minus oral fluids (do not include drinking water in the calculation).

(a) Maintenance fluid: alternate RL and 5% glucose: 4 ml/kg/h for first 10 kg of body weight + 2 ml/kg/h for next 10 kg + 1 ml/kg/h for each additional kg (over 20 kg, up to 30 kg)

Note: increase replacement volumes by 50% (3 ml/kg x % BSA for the first 8 hours) in the event of inhalation injury or electrical burn. For burns > 50% BSA, limit the calculation to 50% BSA.

This formula provides a guide only and should be adjusted according to systolic arterial pressure (SAP) and urine

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

	Non-electrical burns			Electrical burns
	Children < 1 year	Children 1-12 years	Children > 12 years/adults	All ages
AP (mmHg)	SAP \geq 60	SAP 70 to 90 + (2 x age)	SAP \geq 100	Age appropriate SAP
Urine output	1 to 2 ml/kg/h	1 to 1.5 ml/kg/h	0.5 to 1 ml/kg/h	1 to 2 ml/kg/h

In patients with oliguria despite adequate fluid replacement:

dopamine IV: 5 to 15 micrograms/kg/minute by IV pump

or

epinephrine IV: 0.1 to 0.5 micrograms/kg/minute by IV pump

Stop the infusion after 48 hours, if fluid requirements can be met by the oral route or gavage.

Respiratory care

- In all cases: continuous inhalation of humidified oxygen, chest physiotherapy.
- Emergency surgical intervention if necessary: tracheotomy, chest escharotomy.
- Do not administer corticosteroids (no effect on oedema; predisposition to infection). No specific treatment for direct bronchopulmonary lesions.

Analgesia

See [Pain management](#)

Nutrition

Start feeding early, beginning at H8:

- Daily needs in adults
 - calories: 25 kcal/kg + 40 kcal/% BSA
 - proteins: 1.5 to 2 g/kg
- High energy foods (NRG5, Plumpy'nut, F100 milk) are necessary if the BSA is > 20% (normal food is inadequate).
- Nutritional requirements are administered according to the following distribution: carbohydrates 50%, lipids 30%, proteins 20%.
- Provide 5-10 times the recommended daily intake of vitamins and trace elements.
- Enteral feeds are preferred: oral route or nasogastric tube (necessary if BSA > 20%).
- Start with small quantities on D1, then increase progressively to reach recommended energy requirements within 3 days.
- Assess nutritional status regularly (weigh 2 times weekly).
- Reduces energy loss: avoiding droppings, warm environment (20-22 °C), early grafting, management of pain

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Precautions against infection are of paramount importance until healing is complete. Infection is one of the most frequent and serious complications of burns:

- Hygiene precautions (e.g. sterile gloves when handling patients).
- Rigorous wound management (dressing changes, early excision).
- Separate “new” patients (< 7 days from burn) from convalescent patients (\geq 7 days from burn).
- Do not administer antibiotic therapy in the absence of systemic infection.

Infection is defined by the presence of at least 2 of 4 following signs: temperature $>$ 38.5 °C or $<$ 36 °C, tachycardia, tachypnoea, elevation of white blood cell count by more than 100% (or substantial decrease in the number of white blood cells).

- In the event of systemic infection, start empiric antibiotic therapy:

cefazolin IV

Children > 1 month: 25 mg/kg every 8 hours

Adults : 2 g every 8 hours

+ ciprofloxacin PO

Children > 1 month: 15 mg/kg 2 times daily

Adults: 500 mg 3 times daily

- Local infection, in the absence of signs of systemic infection, requires topical treatment with silver sulfadiazine. Not to be applied to children under 2 months.

Other treatments

- **Omeprazole IV** from D1

Children: 1 mg/kg once daily

Adults: 40 mg once daily

- **Tetanus vaccination** (see [Tetanus](#), Chapter 7).

- **Thromboprophylaxis**: low molecular weight heparin SC beginning 48 to 72 hours post-injury.

- **Physiotherapy** from D1 (prevention of contractures), analgesia is necessary.

- **Intentional burns** (suicide attempt, aggression): appropriate psychological follow-up.

III. Local treatment

Regular dressing changes^a prevent infection, decrease heat and fluid losses, reduce energy loss, and promote patient comfort. Dressings should be occlusive, assist in relieving pain, permit mobilisation, and prevent contractures.

Basic principles

- Rigorous adherence to the principles of asepsis.
- Dressing changes require morphine administration in the non-anaesthetised patient.
- The first dressing procedure is performed in the operating room under general anaesthesia, the following in an operating room under general anaesthesia or at the bedside with morphine.

Technique

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Apply a greasy dressing (Jelonet® or petrolatum gauze) using a back and forth motion (do not use a circular movement).
- Cover with a sterile compresses, unfolded into a single layer. Never encircle a limb with a single compress.
- Wrap with a crepe bandage, loosely applied.
- Elevate extremities to prevent oedema; immobilise in extension.

Frequency

- Routinely: every 48 hours.
- Daily in the event of superinfection or in certain areas (e.g. perineum).

Monitoring

- Distal ischaemia of the burned limb is the main complication during the first 48 hours. Assess for signs of ischaemia: cyanosis or pallor of the extremity, dysaesthesia, hyperalgia, impaired capillary refill.
- Monitor daily: pain, bleeding, progression of healing and infection.

IV. Surgical care

Emergency surgical interventions

- Escharotomy: in the case of circumferential burns of arms, legs or fingers, in order to avoid ischaemia, and circumferential burns of chest or neck that compromise respiratory movements.
- Tracheotomy: in the event of airway obstruction due to oedema (e.g. deep cervicofacial burns). Tracheotomy can be performed through a burned area.
- Tarsorrhaphy: in the event of ocular or deep eyelid burns.
- Surgery for associated injuries (fractures, visceral lesions, etc.).

Burn surgery

- Excision-grafting of deep burns, in the operating room, under general anaesthesia, between D5 and D6: excision of necrotic tissue (eschar) with simultaneous grafting with autografts of thin skin. This intervention entails significant bleeding risk, do not involve more than 15% of BSA in the same surgery.
- If early excision-grafting is not feasible, default to the process of sloughing-granulation-reepithelisation. Sloughing occurs spontaneously due to the action of sulfadiazine/ petrolatum gauze dressings and, if necessary, by mechanical surgical debridement of necrotic tissue. This is followed by granulation, which may require surgical reduction in the case of hypertrophy. The risk of infection is high and the process is prolonged (> 1 month).

V. Pain management

All burns require analgesic treatment. Pain intensity is not always predictable and regular assessment is paramount: use a simple verbal scale (SVS) in children > 5 years and adults and NFCS or FLACC scales in children < 5 years (see [Pain](#), Chapter 1).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

In patients with severe burns, oral drugs are poorly absorbed in the digestive tract during the first 48 hours, morphine is administered by SC route.

Acute pain experienced during care

Analgesics are given in addition to those given for continuous pain.

- Significant medical interventions and extensive burns: general anaesthesia in an operating room.
- Limited non-surgical interventions (dressings, painful physiotherapy):
 - Mild to moderate pain, 60 to 90 minutes before giving care:
tramadol PO (see [Pain](#), Chapter 1) rarely allows treatment to be completed comfortably. In the event of treatment failure, use morphine.
 - Moderate or severe pain, 60 to 90 minutes before giving care:
immediate release morphine PO: initial dose of 0.5 to 1 mg/kg; the effective dose is usually around 1 mg/kg, but there is no maximum dose.
or **morphine** SC: initial dose of 0.2 to 0.5 mg/kg; the effective dose is usually around 0.5 mg/kg, but there is no maximum dose.

Note: these doses of morphine are for adults, dosing is the same in children > 1 year, should be halved in children less than 1 year, and quartered in infants less than 3 months.

- Pain management using morphine during dressing changes at the bedside requires:
 - A trained nursing team.
 - Availability of immediate release oral morphine and naloxone.
 - Close monitoring: level of consciousness, RR, heart rate, SpO₂, every 15 min for the first hour following dressing change, then routine monitoring.
 - Assessment of pain intensity and sedation during the intervention and for 1 hour thereafter.
 - Necessary equipment for ventilation by mask and manual suction.
 - Gentle handling of the patient at all times.
- Adjustment of morphine doses for subsequent dressings:
 - If pain intensity (SVS) is 0 or 1: continue with the same dose.
 - If SVS score ≥ 2: increase the dose by 25 to 50%. If pain control remains inadequate, the dressing change should be carried out in the operating room under anaesthesia.
- Take advantage of the residual analgesia following dressing changes to carry out physiotherapy.
- As a last resort (morphine unavailable and no facilities to give general anaesthesia), in a safe setting (trained staff, resuscitation equipment, recovery room), adding **ketamine** IM at analgesic doses (0.5 to 1 mg/kg) reinforces the analgesic effect of the paracetamol + tramadol combination given before a dressing change.

Chronic pain (during the rehabilitation period)

- The treatment is guided by self-evaluation of pain intensity, and utilises paracetamol and/or tramadol. Patients may develop neuropathic pain (see [Pain](#), Chapter 1).
- All other associated pain (physiotherapy, mobilization) should be treated as acute pain.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Footnotes

- (a) Open technique « naked burn patient under a mosquito net » and water immersion therapy are obsolete and should no longer be used.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Cutaneous abscess

A cutaneous abscess is a collection of pus within the dermis or subcutaneous tissue. It is most commonly due to *Staphylococcus aureus*.

Clinical features

- Painful, red, shiny nodule with or without fluctuance; suppuration or surrounding cellulitis (see [Erysipelas and cellulitis](#), Chapter 4).
- Regional adenopathy and fever may be present.
- Complications: osteomyelitis, septic arthritis, septic shock (see [Shock](#), Chapter 1).

Paraclinical investigations

Radiography in case of suspected osteomyelitis or septic arthritis.

Treatment

- Treatment is surgical incision and drainage, under aseptic conditions (i.e. sterile consumables and instruments, antiseptic skin preparation).
- Refer to a surgeon any cutaneous abscess:
 - located in anterior and lateral neck, central triangle of the face, hand, perirectal region, breast, or
 - adjacent to major blood vessels (e.g. femoral artery), or
 - involving joint and bone.
- Antibiotic therapy only if signs of systemic infection, extensive surrounding cellulitis or for individuals with risk factors e.g. immunosuppression or diabetes (for antibiotic therapy, see [Erysipelas and cellulitis](#), Chapter 4).

Equipment

- Sterile scalpel
- Sterile curved, non-toothed artery forceps (Kelly type)
- Sterile disposable gloves and compresses
- Antiseptic solution and 0.9% sodium chloride
- 5 or 10 ml syringe

Anaesthesia

- For small (approximately < 5 cm), well delineated abscess in adults: use local anaesthesia with **1% lidocaine**

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Hold the scalpel between the thumb and middle finger of the dominant hand, the index finger presses on the handle. Hold the abscess between the thumb and index finger of the other hand. The scalpel blade should be perpendicular to the skin.
- The incision is made in a single stroke along the long axis of the abscess. The incision must be long enough for a finger to be inserted.




Figure 8a
Incision with a scalpel

Digital exploration

(Figure 8b)

- Explore the cavity with the index finger, breaking down all loculi (a single cavity should remain), evacuate the pus (and foreign body, if present) and explore to the edges of the cavity.
- The exploration also allows an assessment of the extent of the abscess, the depth, and location with respect to underlying structures (arterial pulsation) or any possible contact with underlying bone. In this last case, seek surgical advice.



Figure 8b
Exploration of the cavity,
breaking down any loculi

Washing

Abundant washing of the cavity using a syringe filled with 0.9% sodium chloride.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All




Figure 8c
Drain fixed to the skin

Dressing

Cover with sterile compresses.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Pyomyositis

Pyomyositis is an infection of the muscle, almost always due to *Staphylococcus aureus*. It most commonly affects the muscles of the limbs and torso. Infections may occur simultaneously in multiple sites.

Risk factors include immunosuppression, concurrent *S. aureus* infection, malnutrition, trauma and injection drug use.

Risk of mortality is significant if treatment is delayed.

Clinical features

- Signs and symptoms:
 - local: exquisite muscle tenderness, oedema giving muscles "woody" texture on palpation.
 - systemic: regional adenopathy and fever.
 - pyomyositis of the psoas muscle: patient keeps hip flexed and experiences pain on hip extension. If the abscess is on the right side, the clinical signs are the same as for appendicitis with pain in the right iliac fossa.
- Complications: septic emboli, endocarditis and septic arthritis, septic shock (see [Shock](#), Chapter 1).

Paraclinical investigations

- POCUS^a : assists in characterisation of abscess; can rule out deep venous thrombosis.
- Radiography: may demonstrate a foreign body, signs of osteomyelitis or osteosarcoma.

Treatment

- Immobilise the limb.
- Systematic antibiotic therapy (see [Erysipelas and cellulitis](#), Chapter 4).
- Adapt analgesics to the pain level (see [Pain](#), Chapter 1).
- Apply compresses soaked in 70% alcohol 2 times daily (max. 3 times daily to prevent burns to the skin) until incision and drainage.
- Treatment is surgical incision and drainage, under aseptic conditions (sterile consumables and instruments, antiseptic skin preparation) following the rules for incision and drainage of abscesses (see [Cutaneous abscess](#), Chapter 10). Muscle abscesses are often deeper than other abscesses. As a result, aspiration with a large bore needle may be necessary to locate the abscess. Needle aspiration is insufficient treatment even if pus is evacuated and should be followed by surgical incision and drainage.
- In case of pyomyositis of the psoas muscle, start antibiotics and refer to a surgeon.

Equipment and anaesthesia

As for [Cutaneous abscess](#), Chapter 10.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Use a forefinger to explore the cavity, break down any loculi and evacuate the pus (Figure 9c).
- Wash abundantly with 0.9% sodium chloride.
- Insert a large drain.
- Fix the drain to the edge of the wound using a single suture. Remove the drain on about the 5th day (Figure 9d).

Figures 9: Surgical incision-drainage of a pyomyositis




Figure 9a
Long incision




Figure 9b
Dissection of the muscle using Kelly forceps, insert closed
then withdraw
with the instrument slightly open




Figure 9c
Exploration and evacuation of pus with the finger



Figure 9d
Drain fixed to the skin

Footnotes

(a) POCUS should only be performed and interpreted by trained clinicians.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Leg ulcers

- Leg ulcers are chronic losses of cutaneous tissue. They are common in tropical regions, resulting from varied aetiologies:
 - vascular: venous and/or arterial insufficiency,
 - bacterial: leprosy, Buruli ulcer (*Mycobacterium ulcerans*), phagedenic ulcer, yaws, syphilis,
 - parasitic: dracunculiasis (Guinea-worm disease), leishmaniasis,
 - metabolic: diabetes,
 - traumatic: trauma is often a precipitating factor combined with another underlying cause.
- The history of the disease and a complete clinical examination (paying particular attention to the neurological examination to determine if there is a peripheral neuropathy caused by leprosy or diabetes) usually leads to an aetiological diagnosis.
- All ulcers may become complicated with either local or regional secondary infections (abscess, lymphadenopathy, adenitis, osteomyelitis, erysipela, pyodermitis), generalised infection (septicaemia), tetanus and after many years of evolution, skin cancer.

Daily local treatment

- Bathe the leg for 10 to 15 minutes in **NaDCC** and rinse in boiled water.
- Remove any necrotic (black) and fibrinous (yellowish) tissue using compresses or excise the tissue with a scalpel.
- Apply:
 - to a clean ulcer, with little discharge: **10% povidone iodine** and vaseline;
 - to a dirty ulcer, with little discharge: **silver sulfadiazine** to a limited area (monitor for systemic adverse effects);
 - to an oozing ulcer: **10% povidone iodine** alone;
 - to an extensive, oozing ulcer or multiple ulcers: diluted **povidone iodine** (1/4 of 10% povidone iodine + 3/4 of 0.9% sodium chloride or clean water) for one minute then rinse with 0.9% sodium chloride or clean water to reduce the risk of transcutaneous iodine absorption.
- Cover with a dry sterile dressing.

Systemic treatment

- Treatment with analgesics in the event of pain: adapt the level and dosage to the individual (see [Pain](#), Chapter 1).
- Give systemic antibiotics in case of:
 - Secondary infection (see [Bacterial skin infections](#), Chapter 4).
 - Phagedenic ulcer (in the early stages, antibiotics may be useful. They are often ineffective in the chronic stages):
doxycycline PO (except in children under 8 years and pregnant or lactating women)
Children 8 years and over: 4 mg/kg once daily

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Tetanus prophylaxis if appropriate (see [Tetanus](#), Chapter 7).
- Skin graft if the ulcer is extensive, clean, red and flat. Skin grafts are often necessary after surgical excision to heal phagedenic and Buruli ulcers.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Necrotising infections of the skin and soft tissues

Invasive infections of the soft tissues: skin, subcutaneous tissue, superficial or deep fascia, muscles. They include necrotising cellulitis, necrotising fasciitis, myonecrosis, gas gangrene, etc.

Clinical presentation depends on the causative organism and the stage of progression. Group A streptococcus is frequently isolated, as are *Staphylococcus aureus*, enterobacteriaceae and anaerobic bacteria including *Clostridium* sp.

Delay in treatment of a minor wound or certain types of wounds (gunshot wounds or stabbings, open fractures or non-sterile intramuscular injections/circumcisions) or certain infections (varicella or omphalitis), favours the development of a necrotising infection. Patient risk factors include immunosuppression, diabetes, malnutrition and advanced age.

A necrotising infection is a surgical emergency and has a high mortality rate.

Clinical features

- Initial signs and symptoms include erythema, oedema and pain disproportionate to appearance of infection. Location depends on the portal of entry. It may be difficult to differentiate necrotising infections from nonnecrotising infections (see [Erysipelas and cellulitis](#), Chapter 4). Systemic signs of infection (fever, tachycardia etc.) may be present.
- Lesions progress rapidly despite antibiotic therapy, with the development of the typical signs of a necrotizing infection: haemorrhagic blisters and necrosis (cold bluish or blackish hypoesthetic macules).
- Signs of late infection: crepitus on palpation and fetid odour (gas gangrene) with signs of severe systemic infection (see [Shock](#), Chapter 1).

Laboratory

- If available, the following tests can help identify an early necrotising infection: white blood cell count $> 15\,000/\text{mm}^3$ or $< 4000/\text{mm}^3$; serum creatinine $> 141 \mu\text{mol/litre}$; serum glucose $> 10 \text{ mmol/litre}$ (180 mg/dl) or $< 3.3 \text{ mmol/litre}$ (60 mg/dl). However, normal results do not exclude a necrotising infection.
- Obtain specimens for bacterial culture in the operating room and blood cultures if possible.

Paraclinical investigations

Radiography: may demonstrate gas in muscles or along the fascia planes. Can rule out foreign body, osteomyelitis or osteosarcoma.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- IV antibiotic therapy for at least 14 days or more depending on clinical response:
cloxacillin + ceftriaxone + clindamycin or amoxicillin/clavulanic acid + clindamycin. For doses, see below.

cloxacillin IV infusion (60 minutes)^a

Children < 40 kg: 50 mg/kg every 6 hours

Children ≥ 40 kg and adults: 3 g every 6 hours

ceftriaxone slow IV (3 minutes) or IV infusion (30 minutes)^b

Children 1 month and over: 100 mg/kg once daily

Adults: 2 g once daily

clindamycin IV infusion (30 minutes)^c

Neonates 0 to 7 days (< 2 kg): 5 mg/kg every 12 hours

Neonates 0 to 7 days (≥ 2 kg): 5 mg/kg every 8 hours

Neonates 8 days to < 1 month (< 2 kg): 5 mg/kg every 8 hours

Neonates 8 days to < 1 month (≥ 2 kg): 10 mg/kg every 8 hours

Children 1 month and over: 10 to 13 mg/kg every 8 hours (max. 2700 mg daily)

Adults: 900 mg every 8 hours

amoxicillin/clavulanic acid (co-amoxiclav) slow IV injection (3 minutes) or IV infusion (30 minutes)^d

Children less than 3 months: 50 mg/kg every 12 hours

Children ≥ 3 months and < 40 kg: 50 mg/kg every 8 hours (max. 6 g daily)

Children ≥ 40 kg and adults: 2 g every 8 hours

- Other treatments:

- Deep vein thrombosis prophylaxis;
- Appropriate management of pain (see [Pain](#), Chapter 1);
- Early nutritional support.

Footnotes

- (a) Cloxacillin powder for injection should be reconstituted in 4 ml of water for injection. Then dilute each dose of cloxacillin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (b) For administration by IV route, ceftriaxone powder should be reconstituted in water for injection only. For administration by IV infusion, dilute each dose of ceftriaxone in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (c) Dilute each dose of clindamycin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (d) Dilute each dose of amoxicillin/clavulanic acid in 5 ml/kg of 0.9% sodium chloride in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride in children 20 kg and over and in adults. Do not dilute in glucose.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Venomous bites and stings

Snake bites and envenomation

- More than 50% of the bites are dry bites, i.e. no envenomation occurred. In the event that venom is injected, the severity of envenomation depends on the species, the amount of venom injected, the location of the bite (bites on the head and neck are the most dangerous) and the weight, general condition and age of the individual (more serious in children).
- It is rare that the snake involved is identified. However, observation of the clinical signs may orient diagnosis and management. Two major syndromes are identified:
 - neurological disorders that evolve towards respiratory muscle paralysis and coma are common manifestations of elapid envenomation (cobra, mamba, etc.);
 - extensive local lesions (intense pain, inflammation with oedema and necrosis) and coagulation abnormalities are common manifestations of viperid or crotalid (rattle snake) envenomation.

Clinical manifestations and management of bites and envenomations are described in the [table](#) below.

- Early diagnosis and monitoring of coagulation abnormalities is based on whole blood clotting tests performed in a dry tube (at the patient's arrival and then every 4 to 6 hours for the first day).

Take 2 to 5 ml of whole blood, wait 30 minutes and examine the tube:

- Complete clotting: no coagulation abnormality
- Incomplete clotting or no clotting: coagulation abnormality, susceptibility to bleeding^a

In the event of coagulation abnormalities, continue to monitor once daily until coagulation returns to normal.

- Aetiological treatment is based on the administration of snake antivenom serum, only if there are clear clinical manifestations of envenomation or coagulation abnormalities are observed.

Antivenom sera are effective, but rarely available (verify local availability) and difficult to store. Antivenom serum should be administered as early as possible: by IV infusion (in 0.9% sodium chloride) if using a poorly purified serum; by slow IV in the event of severe envenomation if the serum is known to be well purified. Repeat antivenom serum administration after 4 or 6 hours if the symptoms of envenomation persist.



For all patients, be prepared for an anaphylactic reaction, which, despite its potential severity (shock), is usually more easily controlled than coagulation disorders or serious neurological disorders.

- In asymptomatic patients (bites without signs of envenomation and with normal coagulation), monitoring must continue for at least 12 hours (24 hours preferred).

Clinical signs and treatment

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Time since bite	Clinical manifestations	Possible aggressor	Treatment
Bite			
0	Fang marks Pain at the site of bite	?	Strict rest, immobilisation of the limb with a splint to slow the diffusion of venom ^(a) . Wound cleansing. Tetanus prophylaxis (Tetanus , Chapter 7). Observe for manifestations of envenomation. At the dispensary level, prepare patient evacuation to a referral centre.
Envenomation			
10-30 minutes	Hypotension, myosis, excessive salivation and sweating, dysphagia, dyspnoea Local paraesthesia, paresis	Elapids	Insert a peripheral IV line. IV antivenom serum as soon as possible.
	Inflammatory syndrome: intense pain, extensive regional oedema	Viperids Crotalids	Insert a peripheral IV line. IV antivenom serum as soon as possible. Analgesics ^(b) . IV or PO ^(b) anti-inflammatories.
30 minutes-5 hours	Cobra syndrome: bilateral eyelid drooping, trismus, respiratory muscle paralysis Shock	Elapids	Intubation and assisted ventilation. See Shock , Chapter 1.
30 minutes-48 hours	Haemorrhagic syndrome: epistaxis, purpura, haemolysis or disseminated intra-vascular coagulation Shock	Viperids Crotalids	Monitor coagulation (blood clotting test in a dry tube). Transfusion of fresh blood in the event of severe anaemia. See Shock , Chapter 1.
6 hours or more	No signs or changes in coagulation (non-venomous snakes or snake bite without envenomation)	?	Reassure the patient. Send him home after 12 hours.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

- In case of clinical evidence of infection only: drainage of any abscess; amoxicillin/clavulanic acid (co-amoxiclav) for 7 to 10 days in case of cellulitis.

Infections are relatively rare, and most often associated with traditional treatment or with nosocomial transmission after unnecessary or premature surgery.

Scorpion stings and envenomation

- In most cases, the sting causes local effects including: pain, oedema, erythema. Management includes strict rest, wound cleansing, analgesics PO, and tetanus prophylaxis (see [Tetanus](#), Chapter 7).

In patients with significant pain, infiltrate the area around the sting with local anaesthetic (**1% lidocaine**). Observe for 12 hours.

- General signs appear in the event of severe envenomation: hypertension, excessive salivation and sweating, hyperthermia, vomiting, diarrhoea, muscle pain, respiratory difficulties, seizures; rarely, shock.

- Aetiological treatment:

The use of scorpion antivenom sera is controversial (most of them are not very effective; they may be poorly tolerated due to insufficient purification).

In practice, in countries where scorpion envenomations are severe (North Africa, the Middle East, Central America and Amazonia), check local availability of antivenom sera and follow national recommendations.

The criteria for administration are the severity of the envenomation, the age of the patient (more severe in children) and the time elapsed since the sting. This should not exceed 2 to 3 hours. If the time elapsed is more than 2 or 3 hours, the benefit of antivenom serum is poor in comparison with the risk of anaphylaxis (in contrast to envenomation by snakes).

- Symptomatic treatment:

- In the event of vomiting, diarrhoea or excessive sweating: prevention of dehydration (oral rehydration salts), especially in children.

- In the event of muscle pain: **10% calcium gluconate** slow IV (children: 5 ml per injection, adults: 10 ml per injection, administered over 10 to 20 minutes).

- In the event of seizures: diazepam may be used with caution; the risk of respiratory depression is increased in envenomated patients (see [Seizures](#), Chapter 1).

Spider bites and envenomation

- Treatment is usually limited to wound cleansing, strict rest, analgesics PO and tetanus prophylaxis (see [Tetanus](#), Chapter 7).

- Severe envenomations are rare. There are two main clinical syndromes:

- Neurotoxic syndrome (black widow spider): severe muscle pain, tachycardia, hypertension, nausea, vomiting, headache, excessive sweating. The signs develop for 24 hours and then resolve spontaneously over a few days.

- ~~Necrotic syndrome (reduces rapidly local tissue lesions, possible necrosis and ulceration; mild general signs)~~

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Local care: remove the embedded sting (bee); clean with soap and water; **calamine** lotion if pruriginous (children and adults: one application 3 to 4 times daily in a thin layer).
- Analgesics if necessary (paracetamol PO).
- In the event of an anaphylactic reaction:

epinephrine (adrenaline) IM

Use *undiluted* epinephrine solution (1:1000 = 1 mg/ml) and a 1 ml syringe graduated in 0.01 ml in children:

Children under 6 years: 0.15 ml

Children from 6 to 12 years: 0.3 ml

Children over 12 years and adults: 0.5 ml

For children, if 1 ml syringe is not available, use a diluted solution, i.e. add 1 mg epinephrine to 9 ml of 0.9% sodium chloride to obtain a 0.1 mg/ml solution (1:10 000):

Children under 6 years: 1.5 ml

Children from 6 to 12 years: 3 ml

Repeat after 5 minutes if no clinical improvement.

In patients with circulatory collapse or those who deteriorate despite receiving IM epinephrine, use IV epinephrine (for doses, see [Anaphylactic shock](#), Chapter 1).

Footnotes

- (a) There can be a considerable delay between the decrease in coagulation factors (less than 30 minutes after the bite) and the first signs of bleeding (other than bleeding at the site of the bite and/or the development of sero-sanguinous blisters), which may appear only 3 days after the bite. Conversely, bleeding may resolve prior to normalization of coagulation parameters.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Dental infections

Infection arising as a secondary complication of an inflammation of the dental pulp. The severity and the treatment of dental infections depend on their evolution: localised to the infected tooth, extended to adjacent anatomical structures or diffuse infections.

Clinical features and treatment

Infection localised to a tooth and its surroundings (acute dental abscess)

- Intense and continuous pain.
- On examination: swelling limited to the gum surrounding the infected tooth. Purulent exudate may be present draining either through the root canal, or through the periodontal ligament (loosening the tooth) or through a gingival fistula. There are no signs of the infection extending to adjacent anatomical structures nor general signs of infection.
- Treatment:
 - Treatment is only surgical (the source of infection is inaccessible to antibiotics): root canal therapy (disinfection of the root canal) if possible or extraction of the tooth.
 - Pain: paracetamol or ibuprofen PO (see [Pain](#), Chapter 1).

Infections extending to adjacent anatomical structures (acute dento-alveolar abscess)

Local spreading of an acute dental abscess into the surrounding bone and tissue.

- Painful gingival and buccal swelling with warm and tender skin, developing into a ripe abscess: intense pain, with trismus, particularly if the infection is in a posterior tooth, presence of general signs (fever, fatigue, cervical lymphadenopathy).
 - In patients with acute gangrenous cellulitis (crepitations on palpation), treat as an infection extending into the cervico-facial tissues (see below).
 - Treatment:
 - First surgical: incision and drainage of the pus or extraction of the tooth.
 - Then antibiotic treatment for 5 days following the procedure:
amoxicillin PO
Children: 25 mg/kg 2 times daily
Adults: 1 g 2 times daily
- Notes:**
- If the dental procedure has to be delayed (local anaesthesia not possible due to inflammation, significant trismus), start an antibiotic therapy, but the dental procedure must be completed in the following days.
- If there is no improvement within 48 to 72 hours after the dental procedure, do not change antibiotic, but start a new procedure on the tooth.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Chapter 11: Mental disorders in adults

[Anxiety](#)

[Insomnia](#)

[Agitation](#)

[Acute confusional state \(delirium\)](#)

[Post-traumatic stress disorder](#)

[Depression](#)

[Psychotic disorders](#)

[Acute psychotic episode](#)

[Chronic psychoses](#)

[Bipolar disorder](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Anxiety

Last updated: November 2021

A patient suffering from anxiety has:

- psychological symptoms: pervasive worries, e.g. fear of having a serious illness, fear with no clearly-defined object or phobias;
- behavioural changes: nervousness, avoidance behaviour, self-isolating tendency, irritability;
- physical symptoms: e.g. dry mouth, “lump in the throat”; sometimes medically unexplained symptoms (e.g. feeling of malaise, hot flashes or chills, diffuse pain);
- concentration difficulties, sleep problems (difficulty getting to sleep, recurrent nightmares).

Anxiety is a common feature in [depression](#), [post-traumatic stress disorder](#) and [psychosis](#). It can also occur in isolation, not associated with any other mental disorders. Anxiety symptoms often occur immediately after a difficult life event. Medically unexplained symptoms are frequent in refugees and people exposed to adversity; in certain cultures they may be the only expression of psychological distress.

Management

Try to determine the source of the anxiety and reassure the patient (without minimising the distress or symptoms). If necessary, use simple relaxation techniques to alleviate the symptoms^a.

If symptoms are exacerbated (e.g., tachycardia, feeling of suffocation, fear of dying or “going crazy,” agitation, or conversely, prostration), it may be necessary to administer **diazepam**: 5 to 10 mg PO or 10 mg IM, to be repeated after one hour if required.

Acute severe anxiety may justify a short course (max. 2 or 3 weeks) of:

diazepam PO: 2.5 to 5 mg 2 times daily; reducing the dose by half in the last few days of treatment

Moderate anxiety lasting more than 2 weeks, administer as first-line treatment:

hydroxyzine PO: 25 to 50 mg 2 times daily (max. 100 mg daily)

or, only if there is no improvement after 1 week, **diazepam** PO: 2.5 to 5 mg 2 times daily for max. 2 weeks.

If symptoms recur after treatment discontinuation, do not resume diazepam or hydroxyzine. Re-evaluate for possible depression or post-traumatic stress disorder.

For generalised anxiety that lasts more than 2 months, and does not improve with psychosocial interventions, an antidepressant should be prescribed (**fluoxetine** or **paroxetine** PO: 20 mg once daily), to be continued for 2 to 3 months after symptoms resolve then, stop gradually over 2 weeks.

Footnotes

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Insomnia

Last updated: November 2021

Complaints may be: difficulty falling or remaining asleep, waking up too early in the morning, nightmares, or fatigue. Symptoms occur at least three times a week for at least one month.

Management

If insomnia is related to an organic cause, treat the cause (e.g. administer analgesics for pain).

If insomnia is related to the use of alcohol, drugs or a medication^a, management depends on the substance involved.

If insomnia is related to a particular life event (e.g. bereavement), a short term treatment with a sedative may be useful:
promethazine PO: 25 mg once daily at bedtime for 7 to 10 days

or, if promethazine is not available, **hydroxyzine** PO: 25 mg once daily at bedtime for 7 to 10 days

or, as a last resort (risk of addiction), **diazepam** PO: 2 to 5 mg once daily at bedtime for 7 days max.

If insomnia persists, re-evaluate the patient. Insomnia is a common feature in depression ([Depression](#)), post-traumatic stress disorder ([Post-traumatic stress disorder](#)) and anxiety disorders ([Anxiety](#)). In such cases, the underlying disorder should be addressed.

Footnotes

- (a) The main drugs known to cause sleep problems are corticosteroids, beta blockers, levodopa/carbidopa, fluoxetine, levothyroxine, etc.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Agitation

Last updated: November 2021

People who have recently experienced violent events, or with anxiety, depression, psychotic disorders or delirium, may have periods of psychomotor agitation.

Agitation is common in acute intoxication (alcohol/psychostimulant drugs) and withdrawal syndrome. Certain drugs may cause agitation (selective serotonin reuptake inhibitors (SSRIs), levodopa, mefloquine, efavirenz, etc.).

Agitation may be accompanied by oppositional, violent or fleeing behaviour.

Management

Clinical evaluation is best performed in pairs, in a calm setting, with or without the person's family/friends, depending on the situation.

It is essential to check for signs of delirium. If present, the priority is to identify the cause and treat it (see [Acute confusional state](#)).

It may be necessary to administer **diazepam** 10 mg PO to reduce the agitation and conduct the clinical exam, without over-sedating the patient.

If the patient is violent or dangerous, urgent sedation is required: **diazepam** IM 10 mg, to be repeated after 30 to 60 minutes if necessary.

Physical restraint should only be used in certain circumstances, strictly following the procedure in place.

Avoid diazepam if agitation is related to acute alcohol intoxication or in case of delirium (risk of respiratory depression). Use haloperidol (see [Acute confusional state](#)).

Alcoholic patients can experience withdrawal symptoms within 6 to 24 hours after they stop drinking. Withdrawal syndrome should be taken into consideration in patients who are hospitalised and therefore forced to stop drinking abruptly. In the early phase (pre-delirium tremens), the symptoms include irritability, a general feeling of malaise, profuse sweating and shaking. Treatment consists in:

diazepam PO (10 mg every 6 hours for 1 to 3 days, then reduce and stop over 7 days)

+ oral hydration (3 litres of water daily)

+ **thiamine** IM or very slow IV (100 mg 3 times daily for at least 3 days)

If the agitation is associated with anxiety, see [Anxiety](#); if associated with psychotic disorders, see [Psychotic disorders](#).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Acute confusional state (delirium)

Last updated: July 2022

Clinical features

The clinical picture includes:

- disorientation in time and space;
- impaired consciousness;
- concentration problems;
- memory impairment.

These symptoms develop rapidly (hours or days), and often fluctuate during the course of the day.

Agitation, delusions, behavioural disorders and hallucinations (often visual) may be associated symptoms.

Management

Delirium almost always has an organic cause:

- Infectious: meningitis, severe malaria, encephalitis, septicaemia, syphilis, AIDS, etc.
- Metabolic: hyper/hypoglycaemia, electrolyte imbalance, niacin (vitamin PP or B₃) or thiamine (vitamin B₁) deficiencies, etc.
- Endocrine: thyroid disorders
- Neurological: epilepsy, raised intracranial pressure, head trauma, meningeal haemorrhage, brain tumour, etc.

Also consider the use of drugs which may cause delirium (opioid analgesics, psychotropic drugs, fluoroquinolones, etc.), use of toxic substances (alcohol/drugs), or withdrawal from these substances.

Delirium requires hospitalisation.

- Treat the underlying cause.
- Provide supportive care (i.e. nutrition, fluid, electrolyte balance); ensure bladder function.
- Ensure that the patient receives only medications appropriate to their needs.
- Treat pain if needed (see [Pain](#), Chapter 1);
- Ensure adequate sensory environment: low lightening, limit noise.

The administration of diazepam may increase delirium. If it is absolutely necessary to sedate an agitated patient, use low dose haloperidol for a short time (7 days or less):

haloperidol PO: 0.5 to 1 mg 2 times daily

or **haloperidol** IM: 0.5 to 1 mg, to be repeated if the patient is still agitated 30 to 60 minutes after the first injection.

If necessary, administer additional doses every 4 hours, do not exceed a total dose of 5 mg daily.

In case of delirium related to alcohol withdrawal (delirium tremens):

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Post-traumatic stress disorder

Last updated: November 2021

An event is “traumatic” when someone has been directly confronted with death, either by seeing another person being killed or seriously injured as the result of violence, or by experiencing serious harm, such as a threat to his/her life or physical integrity (e.g. rape, torture). Exposure to one or several of these events causes feelings of helplessness and horror.

Immediate, transitory symptoms (disorientation, anxiety, sadness, fleeing, etc.) are to be distinguished from secondary, long-lasting problems that appear and/or last several weeks or months after the event: post-traumatic stress, often associated with depression ([Depression](#)), or sometimes acute psychosis ([Psychotic disorders](#)), even in people with no history of psychotic symptoms.

Post-traumatic stress disorder is characterized by three types of psychological response, generally seen in combination^[1].

- **Persistent re-experiencing**

The patient describes:

- images, thoughts or perceptions related to the traumatic experience, which intrude despite efforts to block them out, including at night in the form of distressing dreams;
- flashbacks during which the patient “relives” parts of the traumatic scene.

- **Avoidance**

The patient tries to avoid:

- places, situations and people that might be associated with the trauma;
- having thoughts or feelings related to the trauma; patients may use alcohol, drugs or any psychotropic agents for this purpose.

- **Persistent perceptions of heightened current threat**

Hypervigilance (constant state of alert), exaggerated startle reaction, anxiety, insomnia, poor concentration; sometimes somatic symptoms (sweating, shaking, tachycardia, headache, etc.).

Re-experiencing is highly distressing and causes disorders that may worsen over time; people isolate themselves, behave differently, stop fulfilling their family/social obligations, and experience diffuse pain and mental exhaustion.

Management

Psychological intervention is essential to reduce the suffering, disabling symptoms and social handicaps resulting from PTSD.

It is important to reassure the patient that their symptoms are a normal response to an abnormal event. Sessions should be conducted with tact. The patient should be listened to. Avoid intensely questioning the patient about their

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

References

1. World Health Organization. Post traumatic stress disorder. International Classification of Diseases for Mortality and Morbidity Statistics, Eleventh Revision (ICD-11).
<https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/2070699808> [Accessed 26 January 2021]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Depression

Last updated: July 2022

Depression is characterised by a set of symptoms that have been present at least two weeks and represent a change from previous functioning.

The standard criteria for diagnosis of major depressive disorder are:

- Pervasive sadness and/or a lack of interest or pleasure in activities normally found pleasurable and
- At least four of the following signs:
 - Significant change in appetite or weight
 - Insomnia, especially early waking (or, more rarely, hypersomnia)
 - Psychomotor agitation or retardation
 - Significant fatigue, making it difficult to carry out daily tasks
 - Diminished ability to make decisions or concentrate
 - Feelings of guilt or worthlessness, loss of self-confidence or self-esteem
 - Feelings of despair
 - Thoughts of death, suicidal ideation or attempt

The features of depression can vary according to the patient's culture^a. For example, the depressed patient may express multiple somatic complaints rather than psychological distress. Depression may also manifest itself as an acute psychotic disorder in a given cultural context.

Management

When faced with symptoms of depression, consider an underlying organic cause (e.g. hypothyroidism or Parkinson's disease) or adverse effects from medical treatment (corticosteroids, cycloserine, efavirenz, mefloquine, etc.). Look for a triggering event (e.g. sexual violence, recent childbirth and post-partum depression).

Depressive disorders are the most common mental disorders in patients with severe chronic infectious diseases such as HIV infection or tuberculosis. These disorders should not be neglected, especially as they have a negative impact on adherence to treatment.

Symptoms of depression are common after a major loss (bereavement, forced displacement, etc.). They gradually subside, in most cases, with social support. Psychological support may be useful.

Pharmacological treatment should always be offered, along with counseling, to patients with severe depression (Patient Health Questionnaire-9 (PHQ-9) score > 19; severe functional impairment, psychotic symptoms, and/or suicidal risk).

In patients with moderately severe depression (PHQ-9 score 15-19), pharmacological treatment should be considered if there is no improvement after 3 counselling sessions, or from the outset if patients express a personal preference for

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Preferably use a serotonin reuptake inhibitor (SRI), particularly in older patients. Preferably use fluoxetine, except during pregnancy when sertraline is preferred.

fluoxetine PO: 20 mg on alternate days for one week, then once daily for 3 weeks, then increase the dose if necessary (max. 40 mg daily); use with caution in patients with severe anxiety disorders or who are immobilised (e.g. wounded) or

paroxetine PO: 10 mg once daily for 3 days, then 20 mg once daily for 3 weeks, then increase the dose if necessary (max. 40 mg daily), especially if the depression is accompanied by severe anxiety

or

sertraline PO: 25 mg once daily for 3 days, then 50 mg once daily for 3 weeks, then increase the dose if necessary (max. 100 mg daily)

Assess tolerance and response every week for 4 weeks. If the response is inadequate after 4 weeks at optimal dose or if the SRI is poorly tolerated, replace with another SRI (there is no need for a medication-free interval between the two).

If SRIs are not available, **amitriptyline** PO may be used as an alternative: start with 25 mg once daily at bedtime and gradually increase over 8 to 10 days to 75 mg once daily (max. 150 mg daily). The therapeutic dose is close to the lethal dose; in older patients, reduce the dose by half.

There is a delay of 2 to 3 weeks before the antidepressant effect of SRIs occurs, at least 4 weeks for amitriptyline. During this period, anxiety may be exacerbated and the risk of suicide may increase, especially with fluoxetine. **Hydroxyzine** PO (25 to 50 mg 2 times daily, max 100 mg daily) or **promethazine** PO (25 to 50 mg once daily at bedtime) may be given for the first 2 weeks of treatment. If there is no improvement after 1 week, change to **diazepam** PO (2.5 to 5 mg 2 times daily) for 2 weeks max.

During the first 2 to 4 weeks, do not give the patient more tablets than the quantity required for each week or entrust the treatment to someone in the patient's close entourage that can initially ensure administration of the drug.

Severe depression carries the risk of suicide. Talking to patients about this will not increase the risk of suicide attempt. On the contrary – depressed people are often anxious and ambivalent about suicide and feel relieved when able to talk about it.

If major symptoms have not improved after a month of treatment, increase to the maximum dose and assess after 2 weeks. If there is no improvement, refer the patient to a psychiatrist, if possible; if not, try a different antidepressant.

The treatment should always be stopped gradually over a 4-week period. Inform the patient about problems associated with abrupt treatment discontinuation (very common with paroxetine).

Special situations: pregnant or breast-feeding women

- Pregnancy in a woman under antidepressants:**

Re-evaluate the need to continue treatment. If treatment is still necessary, it is best to continue a treatment that has been effective rather than switching to a different antidepressant. Nevertheless, if the woman plans to breastfeed and is taking fluoxetine, consider switching to another SRI at least 3 weeks before expected delivery to reduce adverse effects in the neonate during breastfeeding. Monitor the neonate the first few days of life for signs

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

- (a) Hence the importance of working with an “informant” (in the anthropological sense of the word) when dealing with unfamiliar cultural contexts.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Psychotic disorders

- [Acute psychotic episode](#)
- [Chronic psychoses](#)
- [Bipolar disorder](#)

Last updated: July 2022

Psychoses are characterised by delusions (the patient is convinced of things that are not real and not accounted for by the person's cultural background), or hallucinations (the patient hears voices that do not exist) and behavioural symptoms (e.g. strange behaviour, agitation, mutism, opposition, fleeing).

Management includes psychosocial support and antipsychotic medication.

Treatment efficacy and prognosis depend largely on the quality of the therapeutic relationship established with the patient and their family.

Keeping the patient at home with outpatient follow-up is preferred if there is no risk of self-harm or harm to others, and if the family is capable of managing the disorder.

Interpretation of psychotic symptoms vary according to the cultural context^a. For example, psychotic disorders may be attributed to charms or to ancestor intervention. Therapeutic approach should take those beliefs into account.

Patients are usually already under "traditional" treatments, this should not be seen as an obstacle to conventional medical treatment.

Footnotes

- (a) Hence the importance of working with an "informant" (in the anthropological sense of the word) when dealing with unfamiliar cultural contexts.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute psychotic episode

Last updated: July 2022

An acute psychotic episode can be a one-time occurrence, usually of sudden onset, or can occur repeatedly, or it may be the early phase of chronic psychosis. It can occur following an adverse life event (e.g. loss, acute stress or trauma). In postpartum psychosis, delusions are frequently related to the mother-child relationship.

Before prescribing antipsychotic medication, consider the possibility of an underlying organic cause (see [Acute confusional state \(delirium\)](#)) or substance use; check and record blood pressure, heart rate, weight.

Antipsychotic treatment is the same as for [chronic psychoses](#) (haloperidol or risperidone) and should last at least 3 months. After 3 months, if the patient is stable, stop the treatment gradually over 4 weeks, monitoring for potential relapse. If the acute episode lasted more than 3 months, continue antipsychotic treatment for at least 2 years.

For severe anxiety or agitation, a [short-course anxiolytic](#) or sedative treatment may be added to the antipsychotic treatment, at the beginning of treatment.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Chronic psychoses

Last updated: July 2022

Chronic psychoses (schizophrenia, paranoid psychosis, etc.) are defined by specific clinical characteristics and their long-term nature.

Schizophrenia is characterized by delusions, disorganized thinking, hallucinations, depersonalisation, loss of motivation, diminished emotional expression, impaired cognition, abnormal behaviour and neglected hygiene. Such patients are often very anxious.

The goal of treatment is to reduce symptoms and improve social and occupational functioning. It offers real benefits, even if chronic symptoms persist (tendency toward social isolation, possible relapses and periods of increased behavioural problems, etc.).

Before prescribing antipsychotic medication, consider the possibility of an underlying organic cause (see [Acute confusional state \(delirium\)](#)) and use of substances. Check and record blood pressure, heart rate and weight.

Treatment should last at least one year, possibly for life, particularly in patients with schizophrenia. Uncertainty about the possibility of follow-up at one year or beyond is no reason not to treat. However, it is better not to start pharmacological treatment for patients who have no family/social support (e.g. homeless), provided they do not have severe behavioural disorders.

Only prescribe one antipsychotic at a time. To limit the risk of adverse effects, start treatment at a low dose and gradually increase until the minimum effective dose is reached. In older patients, reduce the dose by half, whichever medication is used.

Haloperidol is the first-line antipsychotic. Preferably use oral haloperidol with a view to switching to long-acting haloperidol (haloperidol decanoate) if the patient is likely to need long-term treatment (e.g. patients with schizophrenia).

haloperidol PO: start with 0.5 mg 2 times daily for 3 days then 1 mg 2 times daily until the end of the first week; increase to 2.5 mg 2 times daily the second week. After 2 weeks, assess if the treatment is well tolerated and effective. If it is not effective, check adherence; if necessary increase to 5 mg 2 times daily (max. 15 mg daily).

If haloperidol is not available, contraindicated or poorly tolerated, possible alternative are:

risperidone PO: 1 mg 2 times daily for one week, then 2 mg 2 times daily for one week; if necessary, increase to 3 mg 2 times daily as of the third week (max. 10 mg daily).

or

chlorpromazine PO (especially if a sedative effect is required):

25 to 50 mg once daily in the evening for one week; if necessary, increase to 50 mg in the morning and 100 mg in the evening for one week; if necessary, 100 mg 3 times daily as of the third week.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

- If the patient is not under antipsychotic treatment:

haloperidol PO 5 mg + **promethazine** PO 25 mg, to be repeated after 60 minutes if necessary. After a further 60 minutes, if necessary administer **promethazine** IM 50 mg.

In case of hostile or aggressive behaviour, use IM route (same dose), to be repeated after 30 minutes if necessary; after a further 30 minutes, if necessary, administer **promethazine** IM 50 mg.

High doses of haloperidol can induce extrapyramidal symptoms, add biperiden if necessary.

- If the patient is already under antipsychotic treatment:

diazepam PO or IM: 10 mg to be repeated after 60 minutes if necessary

Do not combine two antipsychotics.

For long-term treatment (e.g. patients with schizophrenia) a long-acting antipsychotic drug can be used once the patient has been stabilised on oral treatment. The dosage depends on the oral dose the patient is taking. The switch from oral to a long-acting antipsychotic should be gradual, according to a specific protocol. For information, at the end of the transition period from oral to long-acting antipsychotic, the dose of haloperidol decanoate IM administered every 3 to 4 weeks is approximately:

Daily dose of haloperidol PO	Monthly dose of haloperidol decanoate IM ^(a)
2.5 mg	25 mg
5 mg	50 mg
10 mg	100 mg
15 mg	150 mg

(a) If haloperidol decanoate is not available, **fluphenazine** IM: 12.5 to 50 mg/injection every 3 to 4 weeks.

For a patient on **risperidone** PO: gradually decrease the dose of risperidone by slowly introducing haloperidol PO then, once the patient is stabilised, change to haloperidol decanoate every 3 to 4 weeks as above.

Special situations: pregnant or breast-feeding women

- In the event of pregnancy in a woman taking antipsychotics: re-evaluate the need to continue the treatment. If treatment is still necessary, administer the minimal effective dose and avoid combination with an anticholinergic (biperiden or trihexphenidyl). Monitor the neonate for extrapyramidal symptoms during the first few days of life.
- First symptoms of psychosis during pregnancy: start with the lowest dose of haloperidol and only increase slowly if necessary.
- Post-partum psychosis: if the woman is breast-feeding, haloperidol should be preferred.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Bipolar disorder

Last updated: July 2022

Bipolar disorder is characterised by alternating manic and depressive episodes^a, generally separated by “normal” periods lasting several months or years.

Episodes of mania are characterised by elation, euphoria and hyperactivity accompanied by insomnia, grandiose ideas, and loss of social inhibitions (sexual, in particular).

Depressive episodes are often severe, with significant risk of suicide.

Look for family history of similar symptoms (particularly suicide), very frequent in bipolar patients.

Pharmacological treatment:

- Episodes of mania are treated with **haloperidol** PO: 5 mg once daily for 3 days, then 7.5 mg for one week; if necessary, increase by increments of 2.5 mg per week (max. 15 mg daily).

Possible alternatives:

risperidone PO: 2 mg once daily; if necessary, increase in increments of 1 mg per week (max. 6 mg daily).

or

olanzapine PO: 10 mg once daily for 3 days; if necessary, increase in increments of 5 mg per week (max. 20 mg daily).

If there is improvement after one week of treatment, continue with the same dose for at least 8 weeks after remission of symptoms.

- **Diazepam** PO (5 to 10 mg daily) can be added during the first 2 to 3 weeks.

- If symptoms do not resolve after 2 weeks of antipsychotic treatment at maximum tolerated dose (and 2 different antipsychotics have been tried), add a mood stabiliser:

valproic acid PO: 200 mg 2 times daily (Week 1) then 400 mg 2 times daily (Week 2) then 500 mg 2 times daily

(Week 3). This is usually sufficient to stabilise the patient; if necessary the dose may be increased by 500 mg weekly (max. 1000 mg 2 times daily).

or

carbamazepine PO: 100 mg 2 times daily (Week 1) then 200 mg 2 times daily (Week 2) then 200 mg 3 times daily (Week 3). This is usually sufficient to stabilise the patient; if necessary the dose may be increased by 200 mg weekly (max. 1200 mg daily).

- Treatment should be continued until at least 8 weeks after complete remission of symptoms. Assess together with the patient the benefits and risks of pursuing long-term treatment.
- If it is decided to discontinue antipsychotic treatment, medication should be stopped gradually, monitoring for possible relapse.
- Depressive episodes are treated as for depression (see [Depression](#)).

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Valproic acid is not recommended in women of childbearing age. If it is necessary to start treatment, use carbamazepine.

If a woman of childbearing age is already taking valproic acid, switch to carbamazepine by gradually decreasing the dose of valproic acid over a period of 2 weeks (do not stop treatment abruptly) while gradually starting carbamazepine. If a woman becomes pregnant or is planning pregnancy it is essential to contact a specialist to re-evaluate whether the treatment is still necessary and adjust the dose if needed.

Footnotes

- (a) "Unipolar forms" are characterized by recurring episodes of depression.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Chapter 12: Other conditions

[Sickle cell disease](#)

[Diabetes type 2 in adults](#)

[Essential hypertension in adults](#)

[Heart failure in adults](#)

[Chronic heart failure](#)

[Acute heart failure \(acute pulmonary oedema\)](#)

[Endemic goitre and iodine deficiency](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Sickle cell disease

Homozygous sickle cell disease (SCD) is a life-threatening genetic disorder of haemoglobin (Hb). The abnormal Hb (HbS) results in the distortion of red blood cells into a sickle shape leading to increased destruction (haemolysis), an increase in blood viscosity and obstruction of capillaries (vaso-occlusion).

SCD is common in sub-Saharan Africa (1 to 3% of births), on the American continent, in India and in the Mediterranean basin.

Clinical features

- Symptoms generally begin after 6 months of age.
- Major signs: recurrent painful crises, chronic anaemia, splenomegaly and frequently, growth retardation and malnutrition in children.
- Serious acute life threatening complications such as stroke, overwhelming infections and acute chest syndrome.
- In populations in whom the disease is frequent, diagnosis is suggested by a family history of similar clinical signs.

Major acute manifestations

Painful vaso-occlusive crises (VOC)

- Children under 2 years present with the hand-foot syndrome or dactylitis (acute pain and swelling in the hands or feet).
- Children older than 2 years and adults present with acute pain affecting the back, chest, abdomen (can resemble an acute abdomen) and extremities.
- Young children may have non-specific signs of a VOC: refusal to walk, irritability, lack of appetite, crying, whimpering or moaning when touched, etc.
- Look for an associated infection that might have precipitated the VOC.
- In case of bony pain in a single location, unresponsive to analgesics (or a persistent limp in a child) associated with fever and erythema or swelling, consider an osteomyelitis.

Fever

Look for infection: in particular pneumonia, cellulitis, meningitis, osteomyelitis and sepsis (patients are particularly susceptible to infections especially due to pneumococcus, meningococcus and *Haemophilus influenzae*); malaria.

Acute severe anaemia

- The chronic anaemia is often complicated by acute severe anaemia with gradually appearing fatigue, pallor of the conjunctivae and palms, shortness of breath, tachycardia, syncope or heart failure.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Most often ischaemic (due to vaso-occlusion in cerebral vessels) but a stroke can also be haemorrhagic.
- Sudden loss of motor function or aphasia, in children and in adults.
- Signs can resemble meningitis and cerebral malaria: headache, photophobia, vomiting, stiff neck, alteration of consciousness and neurologic signs or rarely seizures.

Acute chest syndrome (ACS)

- Chest pain, tachypnoea, respiratory distress, hypoxia; fever (more frequent in children); pulmonary infiltrate on chest x-ray. Often preceded by a VOC.
- Complications: multiorgan failure (lung, liver, kidney).

Priapism

Painful prolonged erection in the absence of sexual stimulation, also occurring in young boys. Risk of necrosis and irreversible erectile dysfunction.

Laboratory and other investigations

Diagnosis

- Hb electrophoresis confirms the diagnosis but is often unavailable.
- If not available, a positive Emmel test (or sickling test) in the presence of clinical signs of sickle cell disease supports the diagnosis.

Other examinations

Tests	Indications
Haemoglobin	<ul style="list-style-type: none"> • At the time of diagnosis and annually (frequently 7 to 9 g/dl). • In case of VOC, fever, acute anaemia (≤ 5 g/dl or drop in Hb ≥ 2 g/dl below the patient's baseline), stroke, ACS. • For monitoring of transfused patients.
Platelets	<ul style="list-style-type: none"> • At the time of diagnosis and annually. • In case of acute anaemia (thrombocytopenia - platelet count $\leq 100\,000/\text{mm}^3$ if splenic sequestration).
Urine dipstick	<ul style="list-style-type: none"> • In case of fever: look for a urinary tract infection. • In case of acute severe anaemia: look for haemoglobinuria.
Malaria test	In case of VOC, fever, acute anaemia or stroke.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Painful vaso-occlusive crisis (VOC)

- Moderate pain (at home):
 - Generous oral hydration (water, soup, juice, coconut water): minimum 100 ml/kg daily in children and 50 ml/kg daily in adults (2.5 to 3 litres daily);
 - Warm compresses (application of cold is contra-indicated);
 - Level 1 (paracetamol and ibuprofen) and level 2 (tramadol) analgesics;
 - If pain is not controlled at home within 24 hours, seek medical attention.
- Severe pain or pain not controlled at home (in hospital):
 - PO hydration (as above); if the patient is unable to drink sufficiently, IV hydration ([Appendix 1](#)); in the event of dehydration, treat according to the degree of dehydration (see [Dehydration](#), Chapter 1);
 - Level 3 analgesics (morphine);
 - Do not give routine antibiotics in the absence of fever; do not transfuse for VOC.

For the treatment of pain according to intensity, see [Pain](#) (Chapter 1).

Fever and infection

- Admit to hospital:
 - All children less than 2 years;
 - Children with fever $\geq 38.5^{\circ}\text{C}$ and adults with fever $\geq 39.5^{\circ}\text{C}$; patients who are critically ill appearing^a or have acute anaemia.
- PO or IV hydration ([Appendix 1](#)).
- Treat malaria if present.
- Treat bacterial infections according to cause.
- Treat all patients with respiratory symptoms for pneumonia and ACS.

In case of osteomyelitis:

ceftriaxone slow IV^b injection (3 minutes) or IV infusion (30 minutes)

Children < 40 kg: 50 mg/kg every 12 hours

Children ≥ 40 kg and adults: 2 g every 12 hours

+ **cloxacillin** IV infusion (60 minutes)^c

Children < 40 kg: 50 mg/kg every 6 hours

Children ≥ 40 kg and adults: 3 g every 6 hours

Administer IV therapy for at least 14 days. Then if the patient has improved, change to the oral route for an additional 14 days of treatment with a combination of:

ciprofloxacin PO

Children < 35 kg: 15 mg/kg 2 times daily

Children ≥ 35 kg and adults: 500 mg 2 times daily

+ **amoxicillin/clavulanic acid** PO (see below)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

Strictly necessary

Functional

Performance Cookies

Submit

Accept All

Children ≥ 40 kg and adults:

Ratio 8:1: 3000 mg daily (2 tab of 500/62.5 mg 3 times daily)

Ratio 7:1: 2625 mg daily (1 tab of 875/125 mg 3 times daily)

Patients over 2 years without acute anaemia can continue treatment as outpatients.

Patients under 2 years or with acute anaemia or who cannot be monitored and treated at home by their family should complete PO antibiotherapy in hospital.

- If the patient is not improving, continue ceftriaxone until the patient is afebrile, then, change to PO treatment.
Monitor for acute anaemia.

Acute severe haemolysis

- Admit to hospital.
- Treat malaria if present.
- Transfuse packed red blood cells ^{d,e} if Hb < 5 g/dl or drop of 2 g/dl below the patient's baseline. Target a Hb level of 9 g/dl.
 - Start with 10 to 15 ml/kg in 3 to 4 hours. For information, 10 ml/kg of packed red blood cells usually raise the Hb by 2.5 g/dl.
 - Repeat the Hb. If a second transfusion is needed, check for signs of fluid overload before starting the transfusion.
 - Measure Hb and perform urine dipstick in the following days. Further transfusions may be necessary if haemolysis is ongoing.

Aplastic crisis

- Admit to hospital.
- Treat an associated bacterial infection if present.
- Transfuse as for haemolysis. Repeat the Hb every other day. An increasing reticulocyte count and a gradual increase of the Hb indicate improvement. Follow patient until they have reached their baseline Hb.

Splenic sequestration

- Admit to hospital.
- Treat hypovolaemic shock if present.
- Monitor the size of the spleen.
- Transfuse if Hb < 5 g/dl, target a Hb level of 7 to 8 g/dl maximum.
- Administer ceftriaxone as above.
- After clinical improvement, monitor for relapse (follow the size of the spleen).

Note: splenectomy is contra-indicated (high operative mortality).

Stroke

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Acute chest syndrome

- Admit to hospital.
- Measure SpO₂ and administer oxygen as in stroke.
- PO hydration as for a VOC; if the patient is unable to drink sufficiently, IV hydration ([Appendix 1](#)) while monitoring for fluid overload; in the event of fluid overload, administer one dose of furosemide IV (see [Dehydration](#), Chapter 1).
- Antibiotics:
ceftriaxone slow IV^b injection (3 minutes) or IV infusion (30 minutes) for 7 to 10 days
Children < 20 kg: 50 mg/kg once daily (max. 2 g daily)
Children ≥ 20 kg and adults: 1 to 2 g once daily
+ **azithromycin** PO for 5 days
Children: 10 mg/kg once daily (max. 500 mg daily)
Adults: 500 mg on D1 then 250 mg once daily from D2 to D5
- Transfuse if symptoms are unresponsive to antibiotics and Hb < 9 g/dl.
- If wheezing is present treat with:
salbutamol aerosol (100 micrograms/puff)
Children and adults: 2 to 4 puffs with a spacer every 10 to 30 minutes as needed
- Encourage deep breathing (incentive spirometry hourly).
- Treat pain (see [Pain](#), Chapter 1).

Priapism

- PO hydration as for a VOC; IV hydration if necessary ([Appendix 1](#)) and treat dehydration if present (see [Dehydration](#), Chapter 1).
- Encourage urination, apply warm compresses, treat pain.
- Erection > 4 hours: consider transfusion and refer to surgery.

Prevention of complications

Certain complications can be avoided with appropriate health education of patients/families, routine preventive care and regular follow-up.

Education of patients (including children) and families

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our [Privacy Policy](#)

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Basic knowledge

- | | |
|--|---|
| <ul style="list-style-type: none">• Disease• Treatment• Monitoring | Chronic, necessarily transmitted by both parents, non-contagious.
Routine (see below) and symptomatic (pain).
Size of the spleen, temperature, baseline Hb. |
|--|---|

Major precipitating factors of a painful crisis and how to prevent them

- | | |
|--|--|
| <ul style="list-style-type: none">• Cold• Excessive heat• Tight clothing• Dehydration• Excessive effort• Infections | <p>Wear warm clothing, avoid bathing in cold water.
For example, avoid going out at mid-day.
Wear wide comfortable clothing without elastics.
Drink plenty of fluids.
Moderate physical activity is beneficial.
Follow routine treatments (including vaccination).</p> |
|--|--|

Principal complications requiring the patient to seek urgent medical advice

- Pain unresponsive to analgesia after 24 hours or severe from the start.
- Any fever (do not treat at home).
- Respiratory problems (cough, difficulty breathing, chest pain).
- Diarrhoea/vomiting and inability to drink.
- Dehydration (dark, infrequent urine).
- Anaemia (pale or yellow conjunctivae, pale palms, enlarged spleen).

Routine preventive care

- Prevention of pneumococcal infections

phenoxyethylpenicillin (penicillin V) PO until age 15 years (at least until 5 years):

Children < 1 year: 62.5 mg 2 times daily

Children 1 to < 5 years: 125 mg 2 times daily

Children 5 to 15 years: 250 mg 2 times daily

- Immunization

Ensure that the child's immunisations are up to date; if not, administer catch up vaccines:

**Children
< 5 years**

- DTP, hepatitis B, polio, measles, *H. influenzae* type B vaccines
- Pneumococcal conjugate vaccine (PCV13 or, if not available, PCV10)
- Meningococcal conjugate vaccine in endemic areas
- At 2 years: pneumococcal 23-valent polysaccharide vaccine, at least 8 weeks after the last

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Children ≥ 1 year and adults: 5 mg once daily

- Malaria chemoprophylaxis (if malaria prevalence ≥ 5%)

mefloquine PO

Children 6 months to 5 years and > 5 kg: 5 mg base/kg once weekly

Do not use to treat malaria.

- Provide nutritional support at hospital discharge.

Routine follow-up of patients

- Between crises, for information:

Children < 5 years: every 1 to 3 months;

Children ≥ 5 years and adults: every 3 to 6 months.

- After a crisis: as often as necessary, according to the clinical course.

Footnotes

- (a) Critically ill appearing child: weak grunting or crying, drowsy and difficult to arouse, does not smile, disconjugate or anxious gaze, pallor or cyanosis, general hypotonia.
- (b) For administration by IV route, ceftriaxone powder should be reconstituted in water for injection only. For administration by IV infusion, dilute each dose of ceftriaxone in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (c) Cloxacillin powder for injection should be reconstituted in 4 ml of water for injection. Then dilute each dose of cloxacillin in 5 ml/kg of 0.9% sodium chloride or 5% glucose in children less than 20 kg and in a bag of 100 ml of 0.9% sodium chloride or 5% glucose in children 20 kg and over and in adults.
- (d) Always inquire how many transfusions a patient has previously received (risk of iron overload).
- (e) Do not transfuse whole blood if possible (risk of fluid overload).
- (f) Iron is contraindicated in patients who have received multiple transfusions. Avoid combined preparations of iron and folic acid.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Diabetes type 2 in adults

Diabetes is a metabolic disorder that leads to hyperglycaemia.

Type 2 diabetes usually occurs in adults and accounts for 90% of diabetes cases worldwide.

Type 2 diabetes can lead to acute complications, as well as chronic complications that result in serious organ damage (cardiovascular events; diabetic retinopathy, neuropathy, nephropathy).

Clinical features

- Few or no symptoms; symptoms of hyperglycaemia may be present: polyuria (frequent urination) and polydypsia (excessive thirst and drinking).
- In rare cases, patients may present with severe hyperglycaemia (impaired consciousness, coma or acute dehydration).

Diagnosis

- Look for diabetes in the event of:
 - symptoms of hyperglycaemia;
 - cardiovascular disorders: stroke, myocardial infarction, hypertension;
 - peripheral neuropathies, foot ulcers, absence of tendon reflexes or peripheral pulse.
- Diagnosis is made on one of the following results^[1]:

	Fasting blood glucose ^(a)	Random blood glucose ^(b)	Glycated Hb ^(c)
Symptomatic patient	1 fasting blood glucose ≥ 7 mmol/litre (or ≥ 126 mg/dl)	1 random blood glucose ≥ 11 mmol/litre (or ≥ 200 mg/dl)	1 HbA1c ≥ 6.5%
Asymptomatic patient	2 fasting blood glucose ≥ 7 mmol/litre on 2 samples collected at 2 different times ^(d)	1 random glucose ≥ 11 mmol/litre followed by 1 fasting blood glucose ≥ 7 mmol/litre	2 HbA1c ≥ 6.5% on 2 samples collected at 2 different times ^(e)

(a) Fasting blood glucose test: performed on patient that has fasted at least 8 hours

(b) Random blood glucose test: performed at any moment of the day.

(c) Glycated Hb (HbA1c) reflects average glycaemia over around 3 months.

(d) For example. interval of at least one or more days.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Fasting blood glucose < 8.3 mmol/litre (or < 150 mg/dl) or HbA1c between 7 and 7.5.

The closer blood glucose levels remain to these values, the more cardiovascular complications are prevented or delayed.

Depending on the context (healthcare provision) or patient profile (elderly patient, history of severe hypoglycaemia or long-standing poorly controlled diabetes), fasting blood glucose < 10 mmol/litre (or < 180 mg/dl) or HbA1c of around 8 are acceptable.

Blood glucose should not fall < 4.5 mmol/litre (or < 80 mg/dl) or HbA1c < 6.5.

Lifestyle and dietary advice^a

- Avoid sugared foods and drinks (but no excessive restriction of carbohydrates).
- High fibre intake; limit animal fats and alcohol.
- Physical activity.
- Weight control. If BMI ≥ 25, try to reduce weight by 5 to 10%.
- Stop smoking.

Pharmacological treatment

First-line treatment **metformin PO^b**.

The usual dose is 1 to 2 g daily. For information:

Week 1: 500 mg once daily in the morning at breakfast

Week 2: 500 mg 2 times daily (morning and evening) during meals

Increase in increments of 500 mg per week as long as the drug is well tolerated (max. 2 g daily, i.e. 1 g morning and evening)^[3].

If glycaemic control is not achieved, administer metformin in combination with a sulfonylurea.

Sulfonylurea doses are adjusted in increments to avoid the risk of hypoglycaemia, based on blood glucose results.

- In patients under 60, **glibenclamide PO**:

The usual dose is 5 mg 2 times daily. For information:

Week 1: 2.5 mg once daily in the morning at breakfast

Week 2: 5 mg once daily in the morning at breakfast

Increase in increments of 2.5 mg weekly until fasting blood glucose reaches target levels (max. 15 mg daily).

- In patients over 60, **gliclazide PO** (immediate release tablet):

The usual dose is 40 to 80 mg 2 times daily. For information:

Weeks 1 and 2: 40 mg once daily in the morning at breakfast

Increase in increments of 40 mg every 2 weeks (weeks 3 and 4: 80 mg once daily in the morning at breakfast) until fasting blood glucose reaches target levels (max. 240 mg daily, i.e. 120 mg morning and evening).

If glycaemic control is not achieved with the combination of metformin + a sulfonylurea, continue metformin but replace the sulfonylurea with **intermediate-acting insulin SC**: start with 0.2 IU/kg at bedtime. The dose is adjusted after measuring fasting blood glucose in the morning. Once blood glucose levels have stabilized, test levels once weekly then after each consultation. Doses of 1 IU/kg/day or more may be necessary to reach glycaemic targets. If

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Morning blood glucose	Action
< 4 mmol/litre < 70 mg/dl	Treat hypoglycaemia (see <u>Hypoglycaemia</u> , Chapter 1). Reduce daily dose of insulin by 2 to 4 units. Maintain new dose for 4 days. Check blood glucose after 4 days, readjust dose if glycaemic target has not been reached. Check blood glucose again after 4 days and repeat the process until glycaemic target is reached.
≥ 4 and < 8.3 mmol/litre ≥ 70 and < 150 mg/dl	Do not change dose.
≥ 8.3 and < 11 mmol/litre ≥ 150 and < 200 mg/dl	Increase daily dose of insulin by 2 units. Check blood glucose after 4 days, readjust dose if glycaemic target has not been reached. Check blood glucose again after 4 days and repeat the process until glycaemic target is reached.
≥ 11 and < 16,5 mmol/litre ≥ 200 and < 300 mg/dl	Increase daily dose of insulin by 4 units. Check blood glucose after 4 days, readjust dose if glycaemic target has not been reached. Check blood glucose again after 4 days and repeat the process until glycaemic target is reached.
≥ 16.5 mmol/litre ≥ 300 mg/dl	Perform dipstick analysis for ketones. Treat hyperosmolar hyperglycaemia or ketoacidosis if present.

Example for a man weighing 79 kg:

Start with 16 IU per day ($79 \text{ kg} \times 0.2 \text{ IU}$).

On D4, blood glucose is 14.6 mmol/litre. Add 4 IU (daily dose of insulin is 20 IU).

On D8, blood glucose is 10.4 mmol/litre. Add 2 IU (daily dose of insulin is 22 IU).

On D12, blood glucose is 6.1 mmol/litre. Glycaemic target is reached.

Surveillance and monitoring

Laboratory surveillance

- Patients on oral hypoglycemic agents: blood glucose test once a month to begin with, then during monitoring visits.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Annual check-up: check for cardiovascular and neurological complications, evaluate renal function (serum creatinine and proteinuria dipstick test), examination of teeth and gums.
- Management of diabetes complications.

Patient education

- Lifestyle and dietary measures (diet, physical activity, etc.).
- Patients on sulfonylurea or insulin therapy: signs of hypoglycaemia/hyperglycaemia and management.
- Patients on insulin therapy: auto-administration (schedule, injection sites and techniques); storage of insulin; self-monitoring of blood glucose and adjustment of doses in patients using glucometers.
- Patients with sensory neuropathy or peripheral arterial disease: autoexamination of feet; prevention of foot lesions.

Footnotes

- (a) These measures concern all patients regardless of medication prescribed. They can be sufficient alone to normalize blood glucose levels in certain patients.
- (b) If metformin is contraindicated or not tolerated, replace with a sulfonylurea.

References

1. Partners in Health. Chronic care integration for endemic non-communicable diseases, Chapter 7, Table 7.1. PIH, Boston, 2013.
https://www.pihi.org/sites/default/files/2017-07/PIH_NCD_Handbook.pdf.pdf [Accessed 13 June 2018]
2. American Diabetes Association. Glycemic targets. Diabetes Care 2017 Jan; 40 (Supplement 1): S48-S56.
<https://doi.org/10.2337/dc17-S009> [Accessed 13 June 2018]
3. Joint Formulary Committee. British National Formulary (online) London: BMJ Group and Pharmaceutical Press.
<http://www.medicinescomplete.com> [Accessed 18 June 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Essential hypertension in adults

Hypertension (or high blood pressure - HBP) is defined as elevated blood pressure (BP) at rest that persists over time i.e. measured 3 times during 3 separate consultations over a period of three months.

Essential hypertension is defined as HBP of undetermined cause (the large majority of cases).

The global overall prevalence of HBP in adults aged 25 and over is around 40%. [1]

Serious complications of HBP can be acute (hypertensive encephalopathy, left-sided heart failure, acute renal failure) or delayed i.e. occur after a long period during which HBP has not been controlled (stroke, ischaemic heart disease, peripheral arterial disease, chronic renal impairment).

For pregnancy-induced hypertension, see [Essential obstetric and newborn care](#), MSF.

Clinical features

- HBP thresholds:

HBP classification	Blood pressure (BP) in mmHg	
	Systolic (SBP)	Diastolic (DBP)
Mild	140 or over	90 or over
Moderate	160 or over	100 or over
Severe	180 or over	110 or over

- Severe HBP is defined more by the presence of serious end-organ damage than the blood pressure reading:
 - Uncomplicated hypertensive crisis:
SBP \geq 180 and/or DBP \geq 110 and some symptoms (moderate headaches, epistaxis, dizziness, tinnitus, eye floaters) but no signs of end-organ damage;
 - Hypertensive emergency:
SBP \geq 180 and/or DBP \geq 110 and signs of end-organ damage:
 - ▷ intense headaches, nausea/vomiting, confusion, seizures, coma in the event of hypertensive encephalopathy;
 - ▷ dyspnoea, chest pain in the event of heart failure or cardiac ischaemia;
 - ▷ rapid and/or irregular heart rate in the event of heart failure;
 - ▷ anuria, oliguria in the event of renal impairment.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- ECG and echocardiogram to look for signs of heart failure, coronary disease, or arrhythmia.

Long-term treatment

- The goal of treatment is to lower BP. Target BP are:
 - SBP < 140 and/or DBP < 90
 - SBP < 140 and/or DBP < 80 in diabetic patients
 - SBP < 150 and/or DBP < 90 in patients aged > 80 years
- In patients with mild HBP (SBP \geq 140 and/or DBP \geq 90) without associated cardiovascular disorders or stroke or diabetes, start with lifestyle and dietary advice.
- Pharmacological treatment is indicated in the following cases:
 - SBP \geq 160 and/or DBP \geq 100;
 - HBP associated with cardiovascular disorder, stroke or diabetes;
 - HBP not controlled by lifestyle and dietary changes alone.

Lifestyle and dietary advice

Recommended for all hypertensive patients:

- Reduce calorie and salt intake.
- Regular physical activity.
- Weight loss if BMI \geq 25.
- Stop smoking and alcohol consumption.

Pharmacological treatment

Start with a monotherapy. One of four classes of antihypertensive drugs can be chosen as first line treatment, according to the patient's characteristics (e.g. age, contra-indications, etc.). For information:

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Patient with no comorbidities	Patient with comorbidities
Thiazide diuretic	After a stroke: thiazide diuretic
Patient > 65 years: thiazide diuretic or calcium channel blocker	Diabetic patient: angiotensin converting enzyme (ACE) inhibitor or beta blocker if concomitant cardiovascular disorder
Patient with black skin: thiazide diuretic or calcium channel blocker (avoid ACE inhibitors)	Renal impairment: ACE inhibitor

Thiazide diuretic:
hydrochlorothiazide PO: 12.5 to 25 mg once daily in the morning (max. 25 mg daily)

Angiotensin converting enzyme inhibitor:
enalapril PO: start with 5 mg once daily. Gradually increase, every 1 to 2 weeks, according to BP, up to 10 to 20 mg once daily (max. 40 mg daily).
In elderly patients or patients taking a diuretic or patients with renal impairment: start with 2.5 mg once daily.

Calcium channel blocker:
amlodipine PO: 5 mg once daily. Increase to 10 mg once daily if necessary (max. 10 mg daily).
In elderly patients or patients with hepatic impairment: start with 2.5 mg once daily.

Beta-blocker: (contra-indicated in patients with asthma)
bisoprolol PO: 5 to 10 mg once daily in the morning
Do not stop treatment abruptly (risk of malaise, angina).

In patients with no comorbidity start with a thiazide diuretic and check BP after 4 weeks of treatment.
If the treatment has been correctly taken but there is no improvement after 4 weeks, add a second antihypertensive drug.
After 4 weeks of bitherapy, reevaluate. If the patient's BP remains too high, consider triple-therapy.
In diabetic patients, if there is no improvement after 4 weeks of AEC inhibitor treatment taken correctly, add a calcium channel blocker.
In patients with a cardiac disorder (heart failure or coronary heart disease), bitherapy is usually necessary from the start (AEC inhibitor + beta-blocker).

Surveillance and monitoring

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

- Lifestyle and dietary advice.
- Treatment observance: do not stop treatment abruptly, particularly if taking beta blockers (risk of malaise, angina).
- Consultation in the event of epistaxis, tinnitus, eye floaters; adverse effects of treatment (e.g. cough with AEC inhibitors, erectile dysfunction with beta blockers, oedema with calcium channel blockers).

Treatment of hypertensive crisis

Uncomplicated hypertensive crisis

Most frequent. Reassure the patient and prescribe rest. Check BP a few days later to start or adapt treatment.

Hypertensive emergency

Treat in an intensive care unit.

- Hypertensive encephalopathy:

The aim is to reduce BP by 10 to 15% within the first hour and to not reduce it more than 25% during the first 24 hours.

labetalol IV (contra-indicated in patients with asthma^b:

20 mg over at least 1 minute. Administer another dose after 10 minutes if BP has not decreased. If necessary, 40 mg doses are administered every 10 minutes until hypertension is controlled (max. 300 mg total dose).

- Stroke: do not try to decrease BP during the first 3 days unless SBP is ≥ 220 and/or DBP ≥ 120 (in this event administer labetalol).
- Acute pulmonary oedema: see [Acute heart failure](#).

Footnotes

- (a) Consider secondary hypertension caused by medications being taken, mainly NSAID, corticosteroids, opioids, oral estroprogestogens, etc. Treatment, in this event, consists in stopping or replacing the causative drug.
- (b) In patient with asthma, hydralazine IV: 5 to 10 mg diluted in 10 ml of 0,9% sodium chloride administered by slow IV, to be repeated after 20 to 30 minutes if necessary.

References

1. World Health Organization. Media center. High blood pressure: a public health problem, 2018. <http://www.emro.who.int/media/world-health-day/public-health-problem-factsheet-2013.html> [Accessed 12 September 2018]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Heart failure in adults

Heart failure (HF) is defined as the inability of the heart to maintain adequate cardiac output.

It is a serious condition, particularly frequent in people over 70 years.

There are two types:

- chronic HF: gradual onset of signs of HF;
- acute HF: sudden onset of life-threatening HF (cardiogenic acute pulmonary oedema or shock), in most cases in patients with known cardiopathy.

• [Chronic heart failure](#)

• [Acute heart failure \(acute pulmonary oedema\)](#)

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

[Submit](#)

[Accept All](#)

Chronic heart failure

Clinical features

- Left-sided HF (left ventricle failure; most frequent form)
Fatigue and/or progressive onset of dyspnoea, occurs on exertion and then at rest, accentuated by the decubitus position, preventing the patient from lying down; peripheral oedema.
- Right-sided HF (right ventricle failure)
Oedema of the lower limbs, hepatomegaly, jugular vein distention, hepatojugular reflux; ascites in advanced stages.
- Global HF (failure of both ventricles)
Left and right-sided signs; right-sided signs are often the most prominent.

Evaluate severity of HF^[1]:

Class I	No limitation of physical activity. No symptoms during ordinary physical activity.
Class II	Slight limitation of physical activity. Comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnoea.
Class III	Marked limitation of physical activity. Comfortable at rest. Less than ordinary activity causes fatigue, palpitation, or dyspnoea.
Class IV	Unable to carry on any physical activity without discomfort. Symptoms of heart failure at rest.

Identify causative or aggravating factors:

- Coronary or valvular heart disease, hypertension, viral or toxic cardiopathy, pericarditis.
- Anaemia, American trypanosomiasis, rheumatic fever, diabetes, thyroid disorder, drug/alcohol addiction.

Paraclinical investigations

- Cardiac ultrasound: if available, method of choice to confirm cardiopathy.
- Electrocardiogram (ECG): can diagnose left ventricular cardiomyopathy (left ventricular hypertrophy and/or left bundle branch block) or arrhythmia and particularly atrial fibrillation (AF or Afib) or signs of myocardial ischemia or infarction.
- Chest x-ray: can exclude lung disease in patients with dyspnoea or can show cardiomegaly or pleural effusion (often bilateral) and alveolar-interstitial syndrome.
- Blood test: full blood count, ionogram, serum creatinine.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Weight loss if BMI ≥ 25 .

Treatment of fluid retention

furosemide PO: start with 20 mg once daily; increase if necessary, according to clinical response (certain patients need doses of 80 mg 1 to 2 times daily) then reduce once oedema decrease (20 to 40 mg once daily).

The reabsorption of oedema can sometimes be slow, taking up to 2 to 3 weeks.

The gradual worsening of HF may require an increase in dosage. Lifelong treatment with diuretics is not always necessary.

In the event of resistant oedema, add **hydrochlorothiazide** PO (25 mg 1 to 2 times daily for a few days) but only in hospital settings and monitoring renal function.

Long-term (lifelong) treatment

- ACE inhibitors are the first line treatment. Start with low doses, especially in patients with hypotension, renal impairment, hyponatraemia.

While increasing the dose monitor: drug tolerance (dry cough), blood pressure (the systolic BP should remain above > 90 mmHg), serum potassium and creatinine levels.

In patients taking diuretics, reduce the dose of the diuretic if possible while introducing ACE inhibitors (risk of hypotension if the patient is on high doses of diuretics).

enalapril PO:

Week 1: 2.5 mg once daily for 3 days then 5 mg once daily

Week 2: 10 mg once daily for 3 days then 20 mg once daily

The effective dose is usually 20 mg once daily (or 10 mg 2 times daily). Doses of 10 mg daily are sometimes enough; conversely, doses of 40 mg daily (maximum) are sometimes necessary.

- Once the patient has been stable for at least 2 weeks taking ACE inhibitors and in the absence of any contraindications (asthma, hypotension, bradycardia, conduction disorders, particularly atrio-ventricular heart blocks), add a beta blocker.

bisoprolol PO: start with a low dose and gradually increase as long as the drug is well tolerated (monitor for signs of worsening HF, blood pressure, heart rate).

Week 1: 1.25 mg once daily

Week 2: 2.5 mg once daily

Week 3: 3.75 mg once daily

Weeks 4 to 8: 5 mg once daily

If insufficient:

Weeks 9 to 12: 7.5 mg once daily

As from week 13: 10 mg once daily (max. 10 mg daily)

In the event of temporarily worsening HF, hypotension or bradycardia, readjust doses of associated treatments and reduce the dose of bisoprolol or gradually stop treatment (stopping abruptly can lead to acute deterioration of the patient's condition). Once the patient is stabilized, re-increase/recommence bisoprolol.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- Digitalis glycosides: administer with caution, in intensive care unit (the therapeutic dose is close to the toxic dose), only in patients with AF with rapid ventricular response confirmed by ECG: no visible P waves, irregularly irregular QRS complex (120-160).

Treatment of causative or aggravating factors

According to the cause.

Surveillance and monitoring

Laboratory surveillance

According to treatment (ACE, diuretic, etc.).

Clinical monitoring

- Once stabilised, consultations once a month for the first 6 months, then individualised frequency of consultations depending on the patient's characteristics.
- Routine consultations: weight curve, BP, progress of signs (dyspnoea, oedema, etc.).
- Monitoring of comorbidities and causative or aggravating.

Patient education

- Lifestyle and dietary measures (diet, weight control, physical activity adapted to the patient's capacity, etc.).
- Warning signs (shortness of breath or oedema of the lower limbs, serious adverse effects of treatment) and management (timely/urgent medical consultation).

References

1. Chop WM, Jr. Extending the New York Heart Association classification system. JAMA. 1985;254:505.

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Acute heart failure (acute pulmonary oedema)

Last updated: April 2021

Clinical features

- Sudden onset or exacerbation of dyspnoea
- Fatigue, increased time to recover after exercise
- Bilateral peripheral oedema
- Cold extremities
- Elevated jugular venous pressure
- On auscultation: bilateral pulmonary crepitations and/or extra heart sound (gallop rhythm)

Signs of severity:

- Severe respiratory distress (intercostal retractions, nasal flaring, see-saw breathing, $\text{SpO}_2 < 90\%$, etc.), cyanosis, profuse sweating, confusion
- Systolic blood pressure $< 90 \text{ mmHg}$ (cardiogenic shock)
- Rapid and excessive increase in arterial blood pressure (hypertensive emergency)
- Heart rate (HR) $> 130/\text{minute}$ or $< 40/\text{minute}$
- Respiratory rate (RR) $> 30/\text{minute}$ or $< 12/\text{minute}$
- Chest pain if underlying cardiac ischemia

Paraclinical investigations

Diagnosis is mainly clinical.

- ECG: look for signs of myocardial ischemia or arrhythmia.

If available:

- Chest x-ray: signs vary depending on the severity of pulmonary oedema. In early stage, dilation of vessels in upper lobes then perihilar haze and thickening of septa. In advanced stage, prominent opacities in hilar and perihilar regions and pleural effusion. Can exclude other lung disease, such as pulmonary infection.
- POCUS^a:
 - Perform 12-zone lung exam to evaluate for signs of bilateral pulmonary oedema and/or pleural effusions.
 - Perform 5-view cardiac exam to evaluate for signs of acute volume overload and/or decreased cardiac function.
- Monitoring: full blood count, electrolytes, serum creatinine; cardiac troponins if available.

Treatment

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

- hypoxia.
- Insert an IV line.
 - If there are signs of volume overload (and/or in case of hypertensive emergency): **furosemide** IV, 40 to 80 mg, may be repeated if necessary according to urine output, signs of respiratory distress and SpO₂. If the patient was already taking furosemide at doses of > 40 mg, administer pre-existing dose by IV route.
 - Add a short-acting nitrate (vasodilator) if systolic blood pressure is > 100 mmHg. The aim is to gradually lower systolic blood pressure to near-baseline value. If the patient's baseline value is unknown, for information, lower systolic blood pressure to 120-150 mmHg and the diastolic pressure to under 110 mmHg.

isosorbide dinitrate sublingual (5 mg tablet)

5 mg per dose; if necessary up to 2 doses taken 10 minutes apart

or

isosorbide dinitrate IV (10 ml ampoule, 1 mg/ml)

2 mg (= 2 ml) by slow IV injection (over 2 minutes) then if necessary 2 to 10 mg/hour by continuous infusion with an electric syringe pump

or

glyceryl trinitrate sublingual (0.5 mg tablet)

0.5 mg per dose; if necessary up to 3 doses taken 5 minutes apart

- Non-invasive ventilation using continuous positive airway pressure (CPAP) is recommended in patients with persistent hypoxaemia, unless contraindicated (e.g. impaired consciousness) and on condition that appropriate monitoring is available.
- Monitoring: HR, RR, BP, SpO₂, mental status, urine output.

Subsequent treatment depends on the underlying pathology ([chronic heart failure](#), [hypertension](#), acute coronary syndrome, etc.).

Footnotes

- (a) POCUS should only be performed and interpreted by trained clinicians.

References

1. Ponikowski P et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. Eur J Heart Fail. 2016;18(8):891-975.
<https://academic.oup.com/eurheart/article/37/27/2129/1748921> [Accessed 23 March 2021]
2. Ezekowitz, Justin A. et al. 2017 Comprehensive Update of the Canadian Cardiovascular Society Guidelines for the Management of Heart Failure. Can J Cardiol 2017;33:1342-1433.
<https://www.onlinejc.ca/action/showPdf?pii=S0828-282X%2817%2930973-X> [Accessed 23 March 2021]

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Endemic goitre and iodine deficiency

Goitre is an enlargement of the thyroid gland. Endemic goitre occurs in iodine-deficient areas. Goitre can also be caused or aggravated by the regular consumption of goitrogens such as manioc, cabbage, turnips, millet etc.

Goitre is an adaptive process: iodine is essential for the production of thyroid hormones; iodine deficiency impairs thyroid hormone synthesis; to compensate, the thyroid gland increases in volume. Thyroid function usually remains normal.

As well as the development of goitre, iodine deficiency in pregnant women has serious consequences for the child (foetal and perinatal mortality, physical and mental retardation, cretinism). These risks must be prevented by providing iodine supplementation in iodine-deficient areas.

Clinical features

- The WHO proposes a simplified classification based on the significance of goitre:
 - Group 0: normal thyroid, no palpable or visible goitre
 - Group 1: enlarged thyroid, palpable but not visible when the neck is in the normal position
 - Group 2: thyroid clearly visible when the neck is in the normal position
- Possible mechanical complications (rare): compression, deviation of the trachea or of the oesophagus.

Prevention and treatment

The objective of prevention is to reduce the consequences of iodine deficiency in neonates and children. Supplying iodised salt through national programmes is the recommended method of prevention.

For prevention in populations living in iodine deficient areas where iodised salt is not available and for curative treatment of patients with goitre: use **iodised oil**, according to national protocols. For information (according to the WHO):

Population	Iodised oil PO once yearly (190 mg capsule)
Children under 1 year	1 capsule
Children from 1 to < 6 years	2 capsules
Children from 6 to 15 years	3 capsules

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Appendices

Appendix 1. Normal daily maintenance IV fluids in children > 1 month

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

[Submit](#)

[Accept All](#)

Appendix 1. Normal daily maintenance IV fluids in children > 1 month

Last updated: January 2021

Indications

Basic hydration needs^a for patients unable to drink sufficiently. After 48 hours, it is essential to provide nutrition to the patient orally or by nasogastric tube and to gradually reduce IV fluids.

 This protocol should not be used for surgical or burns patients, those with renal, cardiac disease or diabetic ketoacidosis.

Fluid to be administered

The fluid of choice in children is **Ringer lactate-Glucose 5% (RL-G5%)**. Use a premixed solution if available. If not, add 50 ml of G50% to 500 ml of RL or 100 ml of G50% to 1000 ml of RL. If RL is not available, use 0.9% sodium chloride instead.

For ease of prescription and administration, the daily volumes and rates in drops per minute have been rounded off.

Weight	Volume /24 hours	Rate ^(*) (paediatric infusion set 1 ml = 60 drops)
3 to < 4 kg	350 ml/24 h	15 drops/min
4 to < 5 kg	450 ml/24 h	19 drops/min
5 to < 6 kg	550 ml/24 h	23 drops/min
6 to < 7 kg	650 ml/24 h	27 drops/min
7 to < 8 kg	750 ml/24 h	31 drops/min
8 to < 9 kg	850 ml/24 h	35 drops/min

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
 Functional
 Performance Cookies

Submit

Accept All

Weight	Volume /24 hours	Rate ^(*) (paediatric infusion set 1 ml = 60 drops)	Rate (standard infusion set 1 ml = 20 drops)
20 to < 22 kg	1500 ml/24 h	63 drops/min	21 drops/min
22 to < 26 kg	1600 ml/24 h	67 drops/min	22 drops/min
26 to < 30 kg	1700 ml/24 h	71 drops/min	24 drops/min
30 to < 35 kg	1800 ml/24 h	75 drops/min	25 drops/min
≥ 35 kg	2000 ml/24 h	83 drops/min	28 drops/min

(*) In a paediatric infusion set, the number of drops per minute is equal to the number of ml per hour.
For example: 15 drops/min = 15 ml/hour

Footnotes

- (a) Daily needs are calculated according the following formula:
 Children 0-10 kg: 100 ml/kg per day
 Children 11-20 kg: 1000 ml + (50 ml/kg for every kg over 10 kg) per day
 Children > 20 kg: 1500 ml + (20-25 ml/kg for every kg over 20 kg) per day
 Adults: 2 litres per day

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

Main references

Websites consulted between June 2019 and December 2022

British National Formulary (BNF) and British National Formulary for Children (BNFc)

MedicinesComplete

Martindale. The Complete Drug Reference

MedicinesComplete

UpToDate. Evidence-based clinical decision support resource

BMJ Group. BMJ Best Practice.

La revue *Prescrire*

Centre belge d'information pharmacothérapeutique (CBIP)

<http://www.cbjp.be/fr/start>

Centers for Disease Control and Prevention

<http://www.cdc.gov/DiseasesConditions/>

Cochrane Library

World Health Organization

<http://www.who.int/publications/en/>

This site uses cookies to enhance your browsing experience. By clicking "Accept All", you agree to the storing of cookies on your device. Read our Privacy Policy

- Strictly necessary
- Functional
- Performance Cookies

Submit

Accept All

