



ESTABLISHED 1760

NEW YORK OFFICE
200 N. 42ND ST.

P. Lorillard Company
CIGARETTES - TOBACCO
RESEARCH DIVISION

2525 East Market Street, Greensboro, N.C.

June 7, 1960

Dr. C. O. Jensen
Research Supervisor
Research Division
P. Lorillard Company, Inc.
2525 E. Market Street
Greensboro, North Carolina

Dear Sir:

This report is to inform you of the results obtained by application of the smoke from cigarettes treated with one percent citric acid to frog cilia.

The treated cigarettes were machine made from Old Gold King Size tobacco which had been sprayed with a solution of citric acid in methanol. The citric acid used amounted to one percent by weight of the tobacco sprayed. The methanol was removed from the tobacco by evaporation prior to the making of the cigarettes.

Control cigarettes were made using the same tobacco sprayed with the same volume of methanol with no citric acid added.

The smoke from these two groups of cigarettes was applied to the frog cilia as previously described [May 20, 1960].

The average of ten tests with the control cigarettes showed an increase in rate of 6.8 percent at three minutes after exposure, and a decrease of 77 percent at sixteen minutes after exposure.

The average of seven tests with the 1% citric acid treated cigarettes showed an increase in rate of 47 percent at three minutes after exposure, and an increase of 20 percent at sixteen minutes after exposure.

It will be noted that in a previous report concerning the effect of cigarettes treated with ten percent by weight of citric acid, the average effect at the end of sixteen minutes was an increase in rate of nine percent. This does not indicate that decreasing amounts of citric



01190307