

Complex Number Law Of Sines

<https://github.com/Nazgand/nazgandMathBook>

Mark Andrew Gerads: Nazgand@Gmail.Com

June 12, 2023

Abstract

The goal of this paper is to appreciate The Law of Sines rewritten in the form of complex numbers.

0.1 The obvious single variable case where $z \in \mathbb{C}$, $\neg[z \in \{0, 1\}]$:

$$\frac{\sin(\Im(\ln(z)))}{|1-z|} = \frac{\sin(\Im(\ln(1-\bar{z})))}{|z|} = \sin(\Im(\ln(z) + \ln(1-\bar{z}))) \quad (0.1)$$

For $x \in \mathbb{R}$,

$$\sin(\Im(\ln(z) + \ln(1-\bar{z}))) = \sin\left(\Im\left(\ln\left(z - |z|^2\right)\right)\right) = \sin\left(\Im\left(\ln\left(z|z|^x - |z|^{2+x}\right)\right)\right) \quad (0.2)$$

A nice substitution is $x \rightarrow -2$:

$$\sin\left(\Im\left(\ln\left(z|z|^x - |z|^{2+x}\right)\right)\right) = \sin\left(\Im\left(\ln\left(z|z|^{-2} - 1\right)\right)\right) = \sin(\Im(\ln(\bar{z}^{-1} - 1))) \quad (0.3)$$

0.2 The distinct 3 variable case where $\{z_0, z_1, z_2\} \subset \mathbb{C}$, $z_0 \neq z_1$, $z_1 \neq z_2$, $z_2 \neq z_0$:

$$\frac{\sin\left(\Im\left(\ln\left(\frac{z_2-z_0}{z_1-z_0}\right)\right)\right)}{|z_2-z_1|} = \frac{\sin\left(\Im\left(\ln\left(\frac{z_0-z_1}{z_2-z_1}\right)\right)\right)}{|z_0-z_2|} = \frac{\sin\left(\Im\left(\ln\left(\frac{z_1-z_2}{z_0-z_2}\right)\right)\right)}{|z_1-z_0|} \quad (0.4)$$

is a formula with a symmetry rotating of the 3 variables, that is, substituting $z_0 \rightarrow z_1, z_1 \rightarrow z_2, z_2 \rightarrow z_0$. Furthermore, swapping 2 variables negates the value, so

$$\left| \frac{\sin\left(\Im\left(\ln\left(\frac{z_2-z_0}{z_1-z_0}\right)\right)\right)}{z_2-z_1} \right| \quad (0.5)$$

is invariant under any permutation of the 3 variables.